



The diagnostic utility of stabilized blood for detection of foot-and-mouth disease virus RNA by RT-qPCR

S. Fontél, Kristina; Bøtner, Anette; Belsham, Graham; Lohse, Louise

Publication date:
2014

[Link back to DTU Orbit](#)

Citation (APA):

S. Fontél, K., Bøtner, A., Belsham, G., & Lohse, L. (2014). *The diagnostic utility of stabilized blood for detection of foot-and-mouth disease virus RNA by RT-qPCR*. Poster session presented at EuFMD - Open Session of the European Commission, Dubrovnik, Croatia.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The diagnostic utility of stabilized blood for detection of foot-and-mouth disease virus RNA by RT-qPCR

Kristina S. Fontél, Anette Bøtner, Graham J. Belsham, Louise Lohse*

National Veterinary Institute, Technical University of Denmark, Lindholm, DK-4771 Kalvehave, Denmark

In Europe, clinical signs indicative of foot-and-mouth disease (FMD), would immediately lead to collection of blood and relevant organ material for further laboratory examination for this vesicular disease virus. Today, the first line system for detection of virus in the sample material is real time RT-PCR (RT-qPCR). The aim of this study was to investigate the diagnostic utility of stabilized blood for detection of FMDV RNA in this system.

Materials and Methods

EDTA-stabilized and unstabilized blood (serum) samples were collected from pigs and cattle during experimental studies. The cattle experiment included 13 animals (5 inoculated and 8 contacts) infected with serotype O FMDV. The pig experiment included 14 animals (4 inoculated and 10 contacts) infected with serotype A FMDV. All samples collected at days 0-15 post inoculation (dpi) were analysed after robotic extraction of RNA using RT-qPCR and the sensitivity of detection of FMDV RNA in the two different types of blood sample was

compared.

Results

In the cattle experiment, 13/13 animals developed clinical signs (indicative of FMD) and viral RNA was detected in serum as well as in EDTA-stabilized blood samples from all animals (see fig. 1a and fig 1b for

inoculated and contact animals, respectively).

In the pig experiment, 10/14 animals developed clinical signs and viral RNA was also detected in both sample types (see fig. 2a and 2b for inoculated and contact animals, respectively).

blood, however, for EDTA-stabilized blood with slightly lower sensitivity. In the present set-up, it can therefore not be advised to use EDTA-stabilized blood for individual monitoring of FMDV infection in infected animals, however, on a herd basis (large sample size) this material will provide a clear picture of the overall FMDV infection status.

Results from these experiments showed a very similar profile of RNA detection but with, in general, slightly reduced sensitivity for EDTA-stabilized blood compared to serum.

Discussion

In this study, viral RNA from FMDV could be detected in both unstabilized and EDTA-stabilized




Figure 1a: Comparison of FMDV RNA by Ct-value in EDTA-blood and serum, inoculated cattle




Figure 1b: Comparison of FMDV RNA by Ct-value in EDTA-blood and serum, contact cattle




Figure 2a: Comparison of FMDV RNA by Ct-value in EDTA-blood and serum, inoculated pigs




Figure 2b: Comparison of FMDV RNA by Ct-value in EDTA-blood and serum, contact pigs

