Time Resolved Broadband Terahertz Relaxation Dynamics of Electron in Water

Wang, Tianwu; Iwaszczuk, Krzysztof; Cooke, David G.; Jepsen, Peter Uhd

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Wang, T., Iwaszczuk, K., Cooke, D. G., & Jepsen, P. U. (2014). *Time Resolved Broadband Terahertz Relaxation Dynamics of Electron in Water*. Abstract from 39th International Conference on Infrared, Millimeter, and Terahertz Waves, Tucson, AZ, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Time Resolved Broadband Terahertz Relaxation Dynamics of Electron in Water

Tianwu Wang¹, Krzysztof Iwaszczuk¹, David. G. Cooke², and Peter Uhd Jepsen¹

¹the Department of Photonics Engineering, Technical University of Denmark, DK-2800 Kongens Lyngby, Denmark

² the Department of Physics, McGill University, Montreal, Quebec, Canada

Abstract—we investigated the transient response of the solvated electron ejected by photodetachment from potassium ferrocyanide in water using time resolved terahertz spectroscopy (TSTS). We find that the measured frequency dependent conductivity can be well described by a Drude-Smith model, supplemented by a Lorentz model oscillating near 5 THz.

I. INTRODUCTION

THE electron solvation response in water has been a topic of intense experimental and theoretical investigations for many years[1–3]. It is well known that after an electron is injected into water by directly ionization of water molecule or photodetachment of anions through charge transfer to solvent (CTTS), the quasi-free electron will become trapped and relax to a fully solvated state. However, the detailed picture of electron relaxation process still needs further investigation.

In this paper we will investigate the electron relaxation process by directly measuring the optical conductivity using time resolved terahertz spectroscopy (TRTS)[4, 5]. TRTS is not only useful for characterizing charge carriers under steady state conditions, but is also qualified for nonequilibrium measurements. In the experiment we use air photonics terahertz time domain spectroscopy (THz-TDS) setup [6], which gives usable bandwidth up to 12 THz. The electron in water is injected by photodetachment from $K_4Fe(CN)_6$ using UV pump at 267 nm. The anion $[Fe(CN)_6]^{4-}$ has a high quantum yield with one photon absorption at 267 nm. The 267 nm UV pulse with pulse energy of 30 μJ is generated by frequency tripling part of the fundamental 800 nm pulse. The water sample is prepared by using a gravity driven free flowing liquid film with thickness of 30 μm . To avoid the anion oxidation and water vapor absorption line in the terahertz region, the whole set up is placed in a continuously nitrogen purged box. Ultrabroadband THz transients are generated and detected by a two-color femtosecond-induced air plasma and air biased coherent detection, respectively. To be able to measure the weak signal induced by the pump, an avalanche photodiode (APD) together with a boxcar averager is used in the detection. Time traces of the differential and reference signals were measured simultaneously.

II. RESULTS

Figure 1 shows the pump induced differential THz waveform transmitting through the water film. The whole relaxation process happens in a few hundred fs, followed by a nonzero component lasting more than 10 ps. This is likely due to the slow recombination rate of electron with their donor anion.

The frequency-dependent optical conductivity of electrons

in water 260 fs after excitation is plotted in the inset of Fig. 1. Our measured conductivity data can be well described with a Drude-Smith model plus a Lorentz model oscillating near 5 THz. The conductivity is given by

$$\tilde{\sigma}(\omega) = \frac{\omega_p^2 \epsilon_0 \tau_1}{1 - i\omega\tau_1} \left(1 + \frac{c}{1 - i\omega\tau_1}\right) + \frac{\omega_p^2 \epsilon_0 \omega}{\omega/\tau_2 + i(\omega_0^2 - \omega^2)} \quad (1)$$

Where ω_p is the plasma frequency, ω_0 is the resonance frequency, τ is the carrier scattering time and c is a constant describing the degree of electron confinement.

In summary, we have shown the frequency resolved optical conductivity of electrons in water 260 fs after excitation, which is well fitted with Drude-Smith model plus a Lorentz model resonating near 5 THz.

Fig. 1. Negative differential THz transmission of water film as a function of pump-probe delay time. The inset shows the complex conductivity of water at $\tau_p = +260$ fs (the black solid dot data is the real part of the conductivity and the circle dot data is the imaginary part of the conductivity; the red line and black line are corresponding fitting to Drude-Smith model plus Lorentz model with fitting parameters: $\omega_p/2\pi = 7.8$ THz, $\omega_p/2\pi = 0.93$ THz, $\omega_0/2\pi = 4.7$ THz, $\tau_1 = 4.77$ fs, $\tau_2 = 64$ fs, $c = -0.8717$).

REFERENCES

- [1] V. Lenchenkov, J. Kloepper, V. Vilchiz, and S. E. Bradforth, "Electron photodetachment from $[Fe(CN)_6]^{4-}$: photoelectron relaxation and geminate recombination," *Chem. Phys. Lett.*, vol. 342, pp. 277–286, 2001.
- [2] R. Jimenez, G. R. Fleming, P. V. Kumar, M. Maroncelli, "Femtosecond solvation dynamics of water," *Nature*, vol. 369, pp. 471–473, 1994.
- [3] K. R. Siefermann, Y. Liu, E. Lugovoy, O. Link, M. Faubel, U. Buck, B. Winter, and B. Abel, "Binding energies, lifetimes and implications of bulk and interface solvated electrons in water," *Nat. Chem.*, vol. 2, no. 4, pp. 274–279, Apr. 2010.
- [4] P. U. Jepsen, D. G. Cooke, and M. Koch, "Terahertz spectroscopy and imaging - Modern techniques and applications," *Laser Photonics Rev.*, vol. 5, no. 1, pp. 124–166, Jan. 2011.
- [5] D. G. Cooke, F. C. Krebs, and P. U. Jepsen, "Direct Observation of Sub-100 fs Mobile Charge Generation in a Polymer-Fullerene Film," *Phys. Rev. Lett.*, vol. 108, no. 5, p. 056603, Jan. 2012.
- [6] T. Wang, P. Klarskov, and P. U. Jepsen, "Ultrabroadband THz Time-Domain Spectroscopy of a Free-flowing Water Film," *IEEE Trans. Terahertz Sci. Technol.*, 2014.