



New constraints on Earth's radial conductivity structure

Püthe, C.; Kuvshinov, A.; Olsen, Nils

Published in:
Geophysical Research Abstracts

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Püthe, C., Kuvshinov, A., & Olsen, N. (2014). New constraints on Earth's radial conductivity structure. *Geophysical Research Abstracts*, 16, Article EGU2014-7474-1.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



New constraints on Earth's radial conductivity structure

Christoph Püthe (1), Alexey Kuvshinov (1), Nils Olsen (2), and Terence Sabaka (3)

(1) Institute of Geophysics, ETH Zürich, Switzerland (christoph.pueth@erdw.ethz.ch), (2) DTU Space, Technical University of Denmark, Denmark, (3) Goddard Space Flight Center, NASA, USA

We present a new model of Earth's radial (1-D) conductivity structure at depths between 10 km and the core-mantle boundary. It is based on CM5, the latest version in the Comprehensive Model series that has been derived using 13 years (September 2000 to September 2013) of magnetic data collected by the three satellites Oersted, CHAMP and SAC-C and at the global network of geomagnetic observatories. CM5 describes contributions due to sources in core, lithosphere, ionosphere and magnetosphere (and corresponding induced parts) in form of spherical harmonic expansion (SHE) coefficients.

Removing predictions of the core, lithospheric and ionospheric field contributions as given by CM5 from the observations, we determine time series of the dominating external and induced SHE coefficients of the magnetic potential due to the magnetospheric ring current. Scalar Q-responses are estimated from these coefficients. An iterative approach is used to correct the estimated responses for 3-D effects arising from lateral heterogeneities in the top 10 km. The corrected Q-responses are converted to C-responses; the latter are subsequently inverted for the layered 1-D mantle conductivity profile with the Newton method. The Hessian matrix of the misfit function, which is derived analytically, is used to estimate confidence limits for the conductivity of each layer. The resulting conductivity-depth profile is compared to 1-D conductivity models of Earth's mantle recovered in previous studies.