Phytotoxicity of Sodium Fluoride and Uptake of Fluoride to Willow Trees

Clausen, Lauge Peter Westergaard; Gosewinkel Karlson, Ulrich; Trapp, Stefan

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Clausen, L. P. W., Gosewinkel Karlson, U., & Trapp, S. (2014). *Phytotoxicity of Sodium Fluoride and Uptake of Fluoride to Willow Trees*. Poster session presented at 11th International conference of Phytotechnologies, Heraklion, Greece.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Phytotoxicity of Sodium Fluoride and Uptake of Fluoride to Willow Trees

Lauge Peter Westergaard Clausen^{*1}, Ulrich Gosewinkel Karlson² and Stefan Trapp¹

¹Department of Environmental Engineering, Technical University of Denmark

²Department of Environmental Science, Aarhus University

*Email: lpwc@env.dtu.dk

Intro and scope

Fluoride (F) is present in pristine air, soil and water. As exposure to F through food and beverages is unavoidable, the effects of F on human and mammalian health have been intensely investigated. Due to continued use of super phosphate fertilizers, F accumulates in agricultural soils. However, little is known about the phytotoxicity of sodium fluoride (NaF) and the uptake of F when exposed through roots.

The aim of this study is to:

- 1) Assess the phytotoxicity of NaF to willows when taken up through the roots
- 2) Determine the uptake of F to willows when taken up through roots
- 3) Describe the uptake of F to willows with a non-linear mass balance model

Experimental setup – the willow tree tox test

- 40 cm willow sticks (*Salix viminalis*) were pre-grown and transferred to Erlenmeyer flasks
- Trees were exposed to solutions of various NaF concentrations
- Test duration: 96 h
- Normalized Relative Transpiration (NRT) used as test parameter:

$$NRT(C, t)(\%) = \frac{\frac{1}{n} \sum_{i=1}^n T_i(C, t) / T_i(C, 0)}{\frac{1}{m} \sum_{j=1}^m T_j(0, t) / T_j(0, 0)} \times 100$$

Where,
C) Conc.; t) Time period; T) Abs. transpiration; i and j) Replicate no.;
n) No. of replicates; m) No. of controls see Trapp et al (2008)

Phytotoxicity results

- Toxic effects observed within 48 h for conc. >100 mg F L⁻¹
- Clear dose-response relationship
- No statistically significant difference between toxicity of 50 mg F L⁻¹ and 100 mg F L⁻¹
- Wilting of the trees exposed to 400 mg F L⁻¹

Model description and output

$$0 = \underbrace{\frac{K_M Q C_W}{M_R} + \frac{Q C_W C_R}{M_R}}_{\text{Uptake by transp.}} + \underbrace{\frac{K_M Q C_R}{M_R K_{RW}} - \frac{C_R^2 Q}{M_R K_{RW}}}_{\text{Translocation}} - \underbrace{K_M k_R C_R - k_R C_R^2 - v_{\max} C_R}_{\text{Enzymatic removal}}$$

where,
K_M) Half saturation constant;
Q) Transpiration; C_W) Conc. in external solution; K_R) Mass of roots; C_R) Conc. in roots; K_{RW}) Conc. ratio root/water;
k_R) Root growth rate and v_{max}) Max. enzymatic removal rate see Trapp et al (2008)*

The model assumes:

- Steady state
- Passive uptake of F with the transpiration stream
- Enzymatic removal described by Michaelis-Menten kinetics

Model parameterization

- Maximal enzymatic removal rate, v_{max}: 9 g F kg⁻¹ d⁻¹
- Breakthrough point (BTP): 210 mg F L⁻¹ – On a molar basis the same as obtained by Trapp et al. (2008)* for chloride (Cl)
- Uptake slope approx. 1 for C > BTP – unhindered transport with water

Parameter	Value	Unit	Origin		
Conc. In solution, C_W	0-400	mg L ⁻¹	Measured		
Root mass, m_R	0.001	Kg	Estimated		
Transpiration stream, Q	0.04	L d ⁻¹	Measured ave.		
Half-saturation constant, K_M	2	g L ⁻¹	Fitted		
Max. Enzymatic removal rate, v_{max}	8.992	g kg ⁻¹ d ⁻¹	Fitted		
Root growth rate, k_R	0	d ⁻¹	Estimated		
	Equation	Valid conc.	R ²	n	P
Regression 1	$y = 0.128x - 5.186$	$C < 209.5$	0.764	16	$<< 0.001$
Regression 2	$y = 1.005x - 210.5$	$C > 209.5$	0.790	8	< 0.005

* Trapp S, Felicova D, Rasmussen N F, Bauer-Gottwein P, 2008, Plant uptake of NaCl in relation to enzyme kinetics and toxic effects, Environ. Exp. Bot., 64, 1-7

Conclusions

- NaF is phytotoxic at levels of EC₁₀: 38.0±34.2; EC₂₀: 59.6±40.7 and EC₅₀: 128±51 (values in mg F L⁻¹ ±95% confidence interval)
- Uptake of F can be described by a non-linear mass balance model assuming enzymatic removal
- At external concentrations above 210 mg F L⁻¹ the willows can no longer pump out F from the plant cells and start to accumulate F
- Enzymatic removal of F and Cl from plant cells is likely to be managed by the same "pump"-system

For more details, see: Lauge PW Clausen, Ulrich G Karlson and Stefan Trapp, 2014, Phytotoxicity of Sodium Fluoride and Uptake of Fluoride to Willow Trees, International Journal of Phytoremediation (In press)