



Dynamic flow method to study the CO₂ loading capacity of amino acid salt solutions

Lerche, Benedicte Mai; Stenby, Erling Halfdan; Thomsen, Kaj

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Lerche, B. M., Stenby, E. H., & Thomsen, K. (2011). *Dynamic flow method to study the CO₂ loading capacity of amino acid salt solutions*. Poster session presented at 25th European Symposium on Applied Thermodynamics, Saint Petersburg, Russian Federation.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

PI-19. Dynamic flow method to study the CO₂ loading capacity of amino acid salt solutions

Lerche B.M.¹, Stenby E.H.², Thomsen K.¹

1 - CERE (Center for Energy Resources Engineering) Department of Chemical and Biochemical Engineering, DTU (Technical University of Denmark) Søtofts plads building 229, 2800 Kgs. Lyngby, DK

2 - CERE (Center for Energy Resources Engineering) Department of Chemistry, DTU (Technical University of Denmark) Søtofts plads building 229, 2800 Kgs. Lyngby, DK

bml@kt.dtu.dk

Introduction

Due to a number of advantages amino acid salt solutions have emerged as alternatives to the alkanolamine solvents for the chemical absorption of CO₂ from flue gas. The use of amino acids in CO₂ capture is a bio-mimetic process, as it is similar to CO₂ binding by proteins in the blood, such as hemoglobin. Amino acid salt solutions have the same amine functionality as alkanolamines, and are thus expected to behave similar towards CO₂ in flue gas. Despite rising interest, few studies have been performed so far on amino acids as CO₂ absorbents. [1]

Studying the CO₂ loading capacity of amino acid salt solutions

For the purpose of studying the CO₂ loading capacity of amino acid salt solutions, we have developed an experimental set-up (Figure 1) based on a dynamic analytical mode, with analysis of the effluent gas.




Figure 1: Experimental set-up to study the CO₂ loading capacity of amino acid salt solutions.

The aim has been to mimic, the actual conditions for CO₂ absorption from a coal fired power plant, as closely as possible. We describe the construction and validation of the set-up. Validation data were produced using aqueous solutions of mono-ethanolamine

(MEA), and compared to modelling data, as well as experimental data from literature. Following this validation, the CO₂ loading capacity of aqueous solutions of the potassium salts of selected amino acids was examined. These experiments were performed at a partial pressure of CO₂ close to 10 kPa, and a total pressure around 100 kPa, and a temperature close to 298 K.

References

[1] Benedicte Mai Lerche, Erling H. Stenby and Kaj Thomsen. "CO₂ Capture from Flue Gas using Amino Acid Salt Solutions". *Proceedings from Risoe International Energy Conference 2009*.