



Coupling ecosystems exposure to nitrogen and species sensitivity to hypoxia: modelling marine eutrophication in LCIA

Cosme, Nuno Miguel Dias; Koski, Marja; Hauschild, Michael Zwicky

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Cosme, N. M. D., Koski, M., & Hauschild, M. Z. (2015). *Coupling ecosystems exposure to nitrogen and species sensitivity to hypoxia: modelling marine eutrophication in LCIA*. Abstract from ICES Annual Science Conference 2015, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Coupling ecosystems exposure to nitrogen and species sensitivity to hypoxia: modelling marine eutrophication in LCIA

Nuno Cosme (1), Marja Koski (2), Michael Zwicky Hauschild (1)

(1) Quantitative Sustainability Assessment, Department of Management Engineering, Technical University of Denmark, Kongens Lyngby, Denmark; (2) Section of Marine Ecology and Oceanography, Technical University of Denmark, Charlottenlund, Denmark. Presenter contact details: nmcdc@dtu.dk, Phone +45 45 25 47 29

Summary

Characterisation modelling in Life Cycle Impact Assessment (LCIA) quantifies impacts of anthropogenic emissions by applying substance-specific impact potentials, or Characterisation Factors (CF), to the amount of substances emitted. Nitrogen (N) emissions from human activities enrich coastal marine ecosystems and promote planktonic growth that may lead to marine eutrophication impacts. Excessive algal biomass and dissolved oxygen (DO) depletion typify the ecosystem response to the nutrient input. The present novel method couples a mechanistic model of coastal biological processes that determines the ecosystem response (exposure) to anthropogenic N enrichment (eXposure Factor, XF [$\text{kgO}_2\cdot\text{kgN}^{-1}$]) with the sensitivity of species exposed to oxygen-depleted waters (Effect Factor, EF [(PAF) $\cdot\text{m}^3\cdot\text{kgO}_2^{-1}$], expressed as a Potentially Affected Fraction (PAF) of species). Thus, the coupled indicator (XF*EF, [(PAF) $\cdot\text{m}^3\cdot\text{kgN}^{-1}$]) represents the potential impact on benthic and demersal marine species caused by N inputs. Preliminary results range from 2 (PAF) $\cdot\text{m}^3\cdot\text{kgN}^{-1}$ (Central Arctic Ocean) to 94 (PAF) $\cdot\text{m}^3\cdot\text{kgN}^{-1}$ (Baltic Sea). Comparative contributions per country or watersheds can also be obtained. Further adding environmental fate modelling of N emissions completes the CF for eutrophying emissions making it a useful contribution for sustainability assessment of human activities, as applied in Life Cycle Assessment (LCA).

Introduction

In Life Cycle Impact Assessment (LCIA) substance-specific factors (Characterisation Factors, CF) are multiplied by the amount of substances emitted from human activities to the environment in order to estimate their potential impact. LCIA modelling frameworks typically encompass environmental fate of the emissions, exposure of the receiving ecosystems, and effects on the exposed biota. Nitrogen (N) emissions from anthropogenic sources, mainly agriculture and combustion processes, enrich coastal marine ecosystems where they promote the growth of planktonic biomass. Ecosystem responses involve excessive algal growth and dissolved oxygen (DO) depletion in bottom waters after degradation of sinking organic matter. The DO depletion pathway was addressed in combination with the potential effects on exposed species. The loss of species diversity is the damage dimension ultimately assessed as a function of N emitted.

The present study aims at integrating the processes that describe the ecosystem exposure to nutrient enrichment and the resulting effect on the exposed species. Together these characterise the marine eutrophication phenomenon, which is seen as a syndrome of ecosystem responses to the increased availability of the limiting nutrient in the photic zone (Cloern 2013; Cosme et al. submitted).

Material and Methods

This novel method estimates the ecosystem response to anthropogenic N inputs and their effects on exposed species. It couples (i) a mechanistic model of relevant marine biological processes that determines the ecosystems exposure to N inputs (delivering an eXposure Factor, XF), with (ii) an effect estimation model based on species sensitivity to hypoxia (delivering an Effect Factor, EF).

The XF [$\text{kgO}_2\cdot\text{kgN}^{-1}$] quantifies the nitrogen-to-oxygen 'conversion' potential. It is based on the carbon pump concept (Ducklow et al. 2001) of production, sinking, and degradation of organic carbon that leads to benthic

DO consumption after aerobic respiration by heterotrophic bacteria, with site-dependent parameterisation. The Large Marine Ecosystems (LME) biogeographical classification system (Sherman and Alexander 1986) was adopted to address the spatial variation of some of the modelled parameters and the XF results. The method is fully described by Cosme et al. (submitted) (see also poster R:24 by Cosme et al., this volume).

The EF $[(\text{PAF}) \cdot \text{m}^3 \cdot \text{kgO}_2^{-1}]$ builds on a probabilistic Species Sensitivity Distribution (SSD) methodology. It quantifies potential losses of species richness caused by an environmental stressor (DO depletion) expressed as a Potentially Affected Fraction (PAF) of species (Posthuma et al. 2002). The EFs were estimated at a spatial resolution of five climate zones (polar, subpolar, temperate, subtropical, and tropical) and then assigned to the corresponding 66 LMEs (see also poster R:26 by Cosme and Hauschild, this volume).

The coupled indicator $(\text{XF} \cdot \text{EF}, [(\text{PAF}) \cdot \text{m}^3 \cdot \text{kgN}^{-1}])$ represents the potential impact on benthic and demersal marine species caused by N-enrichment, modulated by the specific biological response. The oxygen depletion and sensitivity to it in each affected spatial unit is quantified at a LME resolution.

Results and Discussion

Preliminary results of the coupled indicator range from 2 $(\text{PAF}) \cdot \text{m}^3 \cdot \text{kgN}^{-1}$ in the Central Arctic Ocean to 94 $(\text{PAF}) \cdot \text{m}^3 \cdot \text{kgN}^{-1}$ in the Baltic Sea, i.e. spatial variation of a factor ≈ 42 among LMEs. The damage dimension of the impacts can be converted into $[\text{species} \cdot \text{kgN}^{-1}]$ by applying global or site-dependent species densities $[\text{species} \cdot \text{m}^{-3}]$ if available.

The XFs obtained show spatial differentiation in agreement with the current understanding of the biological and oceanographic processes, and is consistent with primary productivity patterns and hotspots (Behrenfeld and Falkowski 1997; Chassot et al. 2010). The coupled indicator is an essential component of CFs modelling in LCIA for marine eutrophication impacts and may also be useful for ecosystems health management. Further adding a fate model of the anthropogenic N emissions (Azevedo et al. 2013) will deliver comparative contributions from distinct countries or watersheds, thus being useful to the assessment of the sustainability of human activities with eutroifying emissions, as applied in Life Cycle Assessment (LCA) of product systems.

References

- Azevedo, L. B., Cosme, N., Hauschild, M. Z., Henderson, A. D., Huijbregts, M. A. J., Jolliet, O., Larsen, H. F., and van Zelm, R. 2013. Recommended assessment framework, method and characterisation and normalisation factors for ecosystem impacts of eutrophying emissions: phase 3 (report, model and factors). Project report, 243827 FP7- ENV-2009-1 LC-IMPACT. 154 pp.
- Blanchard, J. L., Jennings, S., Holmes, R., Harle, J., Merino, G., Allen, J. I., Holt, J., Dulvy, N. K., and Barange, M. 2012. Potential consequences of climate change for primary production and fish production in large marine ecosystems. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, 367: 2979–89.
- Chassot, E., Bonhommeau, S., Dulvy, N. K., Mélin, F., Watson, R., Gascuel, D., and Le Pape, O. 2010. Global marine primary production constrains fisheries catches. *Ecol. Lett.*, 13: 495–505.
- Cloern, J. E. 2013. Eutrophication. Retrieved from <http://www.eoearth.org/view/article/152690/> (accessed 6.18.14).
- Cosme, N., Koski, M., and Hauschild, M. Z. Exposure factors for marine eutrophication impacts assessment. *Submitted to Ecological Modelling*.
- Ducklow, H., Steinberg, D., and Buesseler, K. 2001. Upper Ocean Carbon Export and the Biological Pump. *Oceanography*, 14: 50–58.
- Posthuma, L., Suter II, G. W., and Traas, T. P. (Eds.) 2002. Species Sensitivity Distributions in Ecotoxicology, Environmental and Ecological Risk Assessment. Lewis Publishers, CRC Press. 616 pp.
- Sherman, K., and Alexander, L. M. (Eds.) 1986. Variability and Management of Large Marine Ecosystems. Westview Press Inc., Boulder, CO. 319 pp.