



Unified treatment of coupled optical and acoustic phonons in piezoelectric cubic materials

Willatzen, Morten; Wang, Zhong Lin

Published in:
Physical Review B

Link to article, DOI:
[10.1103/PhysRevB.92.224101](https://doi.org/10.1103/PhysRevB.92.224101)

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Willatzen, M., & Wang, Z. L. (2015). Unified treatment of coupled optical and acoustic phonons in piezoelectric cubic materials. *Physical Review B*, 92, [224101]. <https://doi.org/10.1103/PhysRevB.92.224101>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Unified treatment of coupled optical and acoustic phonons in piezoelectric cubic materials

Morten Willatzen

*Technical University of Denmark, Department of Photonics Engineering, Kgs-Lyngby 2800, Denmark
and Beijing Institute of Nanoenergy and Nanosystems, Chinese Academy of Sciences, Beijing, China*

Zhong Lin Wang

*School of Materials Science & Engineering Georgia Institute of Technology, Atlanta, Georgia 30332, USA
and Beijing Institute of Nanoenergy and Nanosystems, Chinese Academy of Sciences, Beijing, China*

(Received 21 September 2015; published 1 December 2015)

A unified treatment of coupled optical and acoustic phonons in piezoelectric cubic materials is presented whereby the lattice displacement vector and the internal ionic displacement vector are found simultaneously. It is shown that phonon couplings exist in pairs only; either between the electric potential and the lattice displacement coordinate perpendicular to the phonon wave vector or between the two other lattice displacement components. The former leads to coupled acousto-optical phonons by virtue of the piezoelectric effect. We then establish three new conjectures that entirely stem from piezoelectricity in a cubic structured material slab. First, it is shown that isolated optical phonon modes generally cannot exist in piezoelectric cubic slabs. Second, we prove that confined acousto-optical phonon modes only exist for a discrete set of in-plane wave numbers in piezoelectric cubic slabs. Third, it is shown that coupled acousto-optical phonons do not exist at the longitudinal-optical (LO) phonon frequency where the dielectric constant vanishes.

DOI: [10.1103/PhysRevB.92.224101](https://doi.org/10.1103/PhysRevB.92.224101)

PACS number(s): 63.20.-e, 77.65.-j

I. INTRODUCTION

Optical and acoustic phonons play a key role in the dissipation mechanisms of quantum electronic systems and determine the thermal and electrical properties of solids. While electron-phonon interactions usually lead to performance degradation of devices, they may also open up possibilities to tune device characteristics by virtue of material and geometry design and appropriate control of ambient properties as well as external parameters. Recent examples of phonon tuning applications include the realization of population inversion in InGaAs quantum dot systems [1–3] and phonon-assisted gain in semiconductor quantum dot masers [4–6]. It was demonstrated in Refs. [7,8] that phonon reservoirs significantly influence dephasing mechanisms [9–11] in quantum dot cavity systems with strong implications for potential applications in quantum information technology (QIT) [12]. Dephasing is a limiting factor for QIT since it destroys the entanglement between light and matter [7]. Some other recent phenomena where optical and acoustic phonon interactions are central include dissipation mechanisms in graphene and consequences for new graphene-based electronic and optical devices [13–15].

Piezoelectricity is important for applications and the understanding of phonon interactions in 3D and 2D semiconductor materials lacking inversion symmetry. A recent discussion of piezoelectric applications in 2D (monolayer) MoS₂ can be found in Ref. [16]. Acoustic phonon limited mobility in 2D MoS₂ was addressed theoretically in Ref. [17]. Graphene-based devices using GaAs as a substrate reveal significant influence at low temperatures on relaxation mechanisms due to piezoelectric surface acoustic phonon scattering originating from the substrate [18–21]. Another example is piezoelectric surface acoustic generation and manipulation that show

promising potential in both quantum phononics [22,23] and plasmonics [24,25].

In this paper we present a detailed treatment of phonons in semiconductors and a full accounting of crystal anisotropy and inversion-asymmetry effects including piezoelectricity for structures based on, e.g., zincblende or wurtzite symmetries such as GaAs, GaN, and ZnO materials or 2D materials including MoS₂ and BN [26]. It is demonstrated in a first unified approach that optical and acoustic phonons are inherently coupled in piezoelectric materials, and we compare with the existing literature in the field where optical and acoustic phonons are treated separately [27–31]. This coupling of optical and acoustic phonons is important for the understanding of electron-phonon interaction selection rules in piezoelectric materials and, e.g., Raman scattering processes. By solving the fully coupled system we obtain, in one step, the acoustic and optical phonon modes and determine to what extent they are coupled. It will be shown that in piezoelectric cubic materials acoustic phonon couplings exist entirely between two of the lattice displacement coordinates (u_x and u_z) similar to the case of nonpiezoelectric cubic materials. An additional pair coupling is shown to exist between the third lattice displacement coordinate (u_y) and the electric potential (ϕ) by virtue of the piezoelectric effect. Hence optical phonons are generally always coupled to acoustic phonons. Choosing a cubic piezoelectric slab as an example, we demonstrate that only coupled acousto-optical phonon modes are allowed unless the phonon in-plane wave-number component vanishes. We show that coupled acousto-optical phonons cannot exist at the LO phonon frequency where the dielectric constant vanishes. Further, it is proven that confined acousto-optical phonon modes cannot exist at any frequency in a piezoelectric slab. These findings are both qualitatively and quantitatively different from the standard nonpiezoelectric result.

II. THEORY

From the elastic equations and the Maxwell-Poisson equation of a piezoelectric cubic material, we obtain the following set of differential equations in the lattice displacements u_x, u_y, u_z and the electric potential ϕ [32]:

$$\left(-q_x^2 c_{11} + c_{44} \frac{\partial^2}{\partial z^2} + \rho \omega^2\right) u_x + i q_x (c_{12} + c_{44}) \frac{\partial u_z}{\partial z} = 0, \quad (1)$$

$$\left(-q_x^2 c_{44} + c_{44} \frac{\partial^2}{\partial z^2} + \rho \omega^2\right) u_y + 2i q_x e_{14} \frac{\partial \phi}{\partial z} = 0, \quad (2)$$

$$i q_x (c_{12} + c_{44}) \frac{\partial u_x}{\partial z} + \left(-q_x^2 c_{44} + c_{11} \frac{\partial^2}{\partial z^2} + \rho \omega^2\right) u_z = 0, \quad (3)$$

$$2i q_x e_{14} \frac{\partial u_y}{\partial z} + \epsilon_{11} \left(q_x^2 - \frac{\partial^2}{\partial z^2}\right) \phi = 0, \quad (4)$$

where c_{11}, c_{12}, c_{44} are the three independent stiffness components and $e_{14}, \rho, \epsilon_{11}$, and q_x are the piezoelectric constant, the mass density, the permittivity, and the phonon wave-number component in the slab plane, respectively. We have followed the common choice where the x axis is directed along the phonon in-plane wave-number component, i.e., $q_y = 0$ [29].

It is clear from the structure of the differential equations above that two solution types (I and II) for u_x and u_z are possible. For the type-I modes, we have [apart from a factor $\exp(i\omega t)$]

$$u_x = \sum_{j=1}^2 A_{x,j} \cos \gamma_j z, \quad u_z = \sum_{j=1}^2 A_{z,j} \sin \gamma_j z, \quad (5)$$

and for type-II modes

$$u_x = \sum_{j=1}^2 B_{x,j} \sin \gamma_j z, \quad u_z = \sum_{j=1}^2 B_{z,j} \cos \gamma_j z. \quad (6)$$

The above two types correspond to compressional (dilatational) and flexural modes, respectively, and they are known as Rayleigh-Lamb modes in the case of an isotropic slab.

Similarly, two solution forms exist for acousto-optical phonon modes. For type-I modes

$$u_y = A_{y,1} \cos(\delta_1 z) + A_{y,2} \cosh(\delta_2 z), \quad (7)$$

$$\phi = A_{\phi,1} \sin(\delta_1 z) + A_{\phi,2} \sinh(\delta_2 z), \quad (8)$$

and for type-II modes

$$u_y = B_{y,1} \sin(\delta_1 z) + B_{y,2} \sinh(\delta_2 z), \quad (9)$$

$$\phi = B_{\phi,1} \cos(\delta_1 z) + B_{\phi,2} \cosh(\delta_2 z). \quad (10)$$

We note that the phonon wave vector z components γ_j and δ_j ($j = 1, 2$) are obtained from the secular equation of the differential equation system in Eqs. (1)–(4) [32]. Combining

Eqs. (1) and (5) we obtain for the type-I acoustic modes

$$A_{z,j} = \frac{c_{44} \gamma_j^2 + q_x^2 c_{11} - \rho \omega^2}{i \gamma_j q_x (c_{12} + c_{44})} A_{x,j} \quad (j = 1, 2). \quad (11)$$

For the slab problem with boundaries at $\pm a/2$ the stress tensor fulfills

$$T_{xz}(z = \pm a/2) = 0, \quad T_{zz}(z = \pm a/2) = 0. \quad (12)$$

Use of the constitutive equations and Eq. (5) gives for type-I modes

$$\sum_{j=1}^2 -\gamma_j A_{x,j} \sin\left(\gamma_j \frac{a}{2}\right) + i q_x A_{z,j} \sin\left(\gamma_j \frac{a}{2}\right) = 0, \quad (13)$$

$$\sum_{j=1}^2 i q_x c_{12} A_{x,j} \cos\left(\gamma_j \frac{a}{2}\right) + \gamma_j c_{11} A_{z,j} \cos\left(\gamma_j \frac{a}{2}\right) = 0. \quad (14)$$

Solving the 4×4 determinantal equation in the four unknowns $A_{x,1}, A_{x,2}, A_{z,1}, A_{z,2}$ leads to the dispersion relations (q_x, ω_n) for type-I acoustic phonons as well as the coupled (u_x, u_z) phonon modes. A similar procedure can be used to determine the type-II phonon modes.

Next, consider the coupled $u_y - \phi$ phonon modes. Combining instead Eqs. (4), (7), and (8), we obtain for the type-I modes acousto-optical phonon modes

$$2i q_x e_{14} [-\delta_1 A_{y,1} \sin(\delta_1 z) + \delta_2 A_{y,2} \sinh(\delta_2 z)] + \epsilon_{11} (q_x^2 + \delta_1^2) A_{\phi,1} \sin(\delta_1 z) + \epsilon_{11} (q_x^2 - \delta_2^2) A_{\phi,2} \sinh(\delta_2 z) = 0 \quad (j = 1, 2). \quad (15)$$

The appropriate acoustic boundary condition is

$$T_{yz}(z = \pm a/2) = c_{44} \frac{\partial u_y}{\partial z} + e_{14} \frac{\partial \phi}{\partial x} \Big|_{z=\pm a/2} = 0, \quad (16)$$

giving

$$-c_{44} \delta_1 A_{y,1} \sin\left(\delta_1 \frac{a}{2}\right) + c_{44} \delta_2 A_{y,2} \sinh\left(\delta_2 \frac{a}{2}\right) + e_{14} i q_x A_{\phi,1} \sin\left(\delta_1 \frac{a}{2}\right) + e_{14} i q_x A_{\phi,2} \sinh\left(\delta_2 \frac{a}{2}\right) = 0. \quad (17)$$

Two electric boundary conditions are needed since the electric field outside the slab must be taken into account as well. For vacuum environments, the Maxwell-Poisson equation simply reads, if $|z| > a/2$,

$$-\epsilon_0 \nabla^2 \phi = \epsilon_0 \left(q_x^2 - \frac{\partial^2}{\partial z^2}\right) \phi = 0, \quad (18)$$

with the solution

$$\phi(z) = \phi_+ e^{-|\mathbf{q}_\parallel|z} e^{i\mathbf{q}_\parallel \cdot \mathbf{r}} \quad \text{if } z > a/2, \quad (19)$$

$$\phi(z) = \phi_- e^{|\mathbf{q}_\parallel|z} e^{i\mathbf{q}_\parallel \cdot \mathbf{r}} \quad \text{if } z < -a/2. \quad (20)$$

Continuity of the transverse electric field at $z = \pm a/2$ yields for type-I acousto-optical phonon modes

$$\phi_+ = \frac{A_{\phi,1} \sin\left(\delta_1 \frac{a}{2}\right) + A_{\phi,2} \sinh\left(\delta_2 \frac{a}{2}\right)}{e^{-|\mathbf{q}_\parallel|a/2}}, \quad \phi_- = -\phi_+. \quad (21)$$

Imposing continuity of the normal electric displacement at the slab boundaries gives for type-I modes

$$\begin{aligned} & -\epsilon_{11} \left[\delta_1 A_{\phi,1} \cos\left(\delta_1 \frac{a}{2}\right) + \delta_2 A_{\phi,2} \cosh\left(\delta_2 \frac{a}{2}\right) \right] \\ & + e_{14} i q_x \left[A_{y,1} \cos\left(\delta_1 \frac{a}{2}\right) + A_{y,2} \cosh\left(\delta_2 \frac{a}{2}\right) \right] \\ & = \epsilon_0 \phi_+ |\mathbf{q}_{\parallel}| e^{-|\mathbf{q}_{\parallel}| a/2}. \end{aligned} \quad (22)$$

The above set of equations completely specifies the type-I acousto-optical phonon modes. We note that the procedure for determining type-II modes goes along the same steps. Since the electric field $\mathbf{E} = -\nabla\phi$, the ionic displacement $\mathbf{u}_{\text{ion}} = \mathbf{u}_+ - \mathbf{u}_-$ can be found from the Born-Huang equations [33]

$$\mathbf{u}_{\text{ion}} = -\sqrt{\frac{\Omega}{m_r}} \frac{\sqrt{\epsilon(0) - \epsilon(\infty)} \omega_{TO}}{\omega^2 - \omega_{TO}^2} \mathbf{E}, \quad (23)$$

where Ω and m_r are the unit cell volume and reduced ion mass, respectively. Hence, the present coupled phonon formalism allows determination of the lattice displacement \mathbf{u} and the internal ion displacement \mathbf{u}_{ion} simultaneously.

Finally, to completely specify the phonon mode solutions, coupled normalization conditions are needed:

$$\begin{aligned} & \int d\mathbf{r} \left[\sqrt{\frac{m_r}{\Omega}} \mathbf{u}_{\text{ion}}^*(\mathbf{r}, t) \right] \left[\sqrt{\frac{m_r}{\Omega}} \mathbf{u}_{\text{ion}}(\mathbf{r}, t) \right] \\ & + \int d\mathbf{r} [\sqrt{\rho} \mathbf{u}^*(\mathbf{r}, t)] [\sqrt{\rho} \mathbf{u}(\mathbf{r}, t)] = \frac{\hbar}{2\omega_n}, \end{aligned} \quad (24)$$

where ω_n is the phonon mode frequency. An important result that distinguishes piezoelectric materials from nonpiezoelectric materials can now be proved. We will show that at the LO phonon frequency where the permittivity is zero coupled acousto-optical phonon modes cannot be excited. In contrast, in nonpiezoelectric media confined optical phonon modes with an arbitrary in-plane wave number q_x exist at the LO phonon frequency.

First, if $\epsilon_{11} = 0$, Eq. (4) shows that

$$u_y = A_y e^{iq_x x}. \quad (25)$$

Then Eqs. (2) and (16) yield

$$\phi = (\phi_0 + \phi_1 z) e^{iq_x x}, \quad \phi_0 = \phi_1 = 0. \quad (26)$$

Continuity of the transverse electric field component guarantees further that

$$\phi_+ = \phi_- = 0, \quad (27)$$

and finally, from continuity in the normal electric displacement, we obtain

$$A_y = 0. \quad (28)$$

Hence, both the electric potential ϕ and the phonon u_y component must be zero simultaneously. In other words, it is not possible to excite coupled $u_y - \phi$ modes at the LO phonon frequency. It is well known that confined LO phonon modes exist at the LO phonon frequency. We will now prove that confined coupled acousto-optical $u_y - \phi$ modes cannot exist in piezoelectric media except at certain discrete q_x wave-number values.

A confined phonon mode is characterized by an electric field that vanishes at the slab interfaces. Thus, coupled $u_y - \phi$ solutions can be sought quite generally in the form (type I modes)

$$u_y = A_y \cos\left(\frac{m\pi}{a} z\right) e^{iq_x x}, \quad \phi = A_{\phi} \sin\left(\frac{m\pi}{a} z\right) e^{iq_x x}, \quad (29)$$

where $m = 2, 4, 6, \dots$. From Eq. (2) we have

$$\left(-q_x^2 c_{44} - \left(\frac{m\pi}{a}\right)^2 c_{44} + \rho \omega^2\right) A_y + 2iq_x e_{14} \left(\frac{m\pi}{a}\right) A_{\phi} = 0, \quad (30)$$

and from Eq. (4)

$$-2iq_x e_{14} \left(\frac{m\pi}{a}\right) A_y + \epsilon_{11} \left(q_x^2 + \left(\frac{m\pi}{a}\right)^2\right) A_{\phi} = 0. \quad (31)$$

Hence the possible mode frequencies are

$$\omega_m = \sqrt{\frac{c_{44}}{\rho} \left[q_x^2 + \left(\frac{m\pi}{a}\right)^2 + \frac{4q_x^2 e_{14}^2 \left(\frac{m\pi}{a}\right)^2}{\epsilon_{11} c_{44} \left(q_x^2 + \left(\frac{m\pi}{a}\right)^2\right)} \right]^{1/2}}. \quad (32)$$

It follows immediately from the preceding equations that Eq. (16) is fulfilled. Continuity in the transverse electric field and normal displacement yields $\phi_+ = 0$ and

$$A_{\phi} = \frac{iq_x e_{14}}{\epsilon_{11} (-1)^{m/2} \left(\frac{m\pi}{a}\right)} A_y. \quad (33)$$

Combining Eqs. (31) and (33) requires

$$2(-1)^{m/2} \left(\frac{m\pi}{a}\right)^2 = q_x^2 + \left(\frac{m\pi}{a}\right)^2, \quad (34)$$

which is possible only at certain discrete q_x values $\{q_{x,m}\}$. Hence we have proven the conjecture that confined coupled $u_y - \phi$ phonon modes cannot exist in piezoelectric media except at certain discrete wave-number values, in contrast to the case of nonpiezoelectric media where confined optical phonon modes exist for any in-plane wave number q_x .

III. DISCUSSIONS

Consider first the influence of anisotropy on the phonon dispersion for the coupled $u_x - u_z$ lattice displacement components in a cubic piezoelectric GaAs slab. The physical properties of GaAs are $c_{11} = 11.88 \times 10^{10}$ Pa, $c_{12} = 5.38 \times 10^{10}$ Pa, $c_{44} = 5.94 \times 10^{10}$ Pa, $\rho = 5.318 \times 10^3$ kg/m³, $e_{14} = 0.154$ C/m², $\epsilon(0) = 12.9\epsilon_0$, $\hbar\omega_{LO} = 33.2$ meV, $\hbar\omega_{TO} = 36.1$ meV, $\epsilon_{\infty} = \epsilon(0) \frac{\omega_{TO}^2}{\omega_{LO}^2}$.

Since all cubic materials are anisotropic, the isotropic relation

$$c_{44} = \frac{c_{11} - c_{12}}{2} \quad (35)$$

used for GaAs in, e.g., Ref. [29] is an approximation. In fact, the degree of anisotropy, defined as

$$\alpha = \left| \frac{\frac{c_{11} - c_{12}}{2} - c_{44}}{c_{44}} \right|, \quad (36)$$




FIG. 1. (Color online) Dispersion relations for the coupled $u_x - u_z$ acoustic phonon modes of a 2 nm GaAs slab. (Upper plot) GaAs and (lower plot) GaAs but using the isotropic assumption: $c_{44} = \frac{c_{11} - c_{12}}{2} = 3.25 \times 10^{10}$ Pa. The first axis is $q_x a$ and the second axis the phonon band energy in meV.

is large in GaAs and equal to 45%. Hence, substantial deviations in the phonon modes and dispersion relations due to anisotropy must be expected for GaAs. In passing, we note that it follows from the results above that unit-cell inversion asymmetry and piezoelectricity do not influence the $u_x - u_z$ phonon modes and dispersion.

In Fig. 1 the coupled $u_x - u_z$ dispersion curves are shown for a 2 nm GaAs slab. Evidently, the phonon mode energies change markedly when including the full anisotropy (by more than 40% for the first band energy at $q_x a = 0$) and so does the shape of the dispersion curves.

Next, let us consider the dispersion curves for coupled $u_y - \phi$ acousto-optical phonon modes.

In Fig. 2 the $u_y - \phi$ dispersion curves are shown for a 2 nm GaAs slab. It is evident that the coupled acousto-optical phonon modes are strongly influenced by anisotropy, as in the case with the coupled acoustic $u_x - u_z$ phonons, leading




FIG. 2. (Color online) Dispersion relations for the coupled $u_y - \phi$ acousto-optical phonon modes of a 2 nm GaAs slab. (Upper plot) GaAs including piezoelectricity and anisotropy and (lower plot) GaAs but using the isotropic non-piezoelectric assumption: $c_{44} = \frac{c_{11} - c_{12}}{2} = 3.25 \times 10^{10}$ Pa and $e_{14} = 0$.

to a difference of about 40% in the first band energy near $q_x a = 0$. Moreover, a close inspection of the dispersion curves around the LO phonon energy of 36.1 meV reveals that, when piezoelectricity is included and for any strictly positive $q_x a$ value, no band crosses the forbidden LO phonon energy, in agreement with the proof given above.

It must be noted that interface phonons are not captured in the plots as they correspond to imaginary q_x values, but the procedure for finding them is the same as for nonpiezoelectric and isotropic media [27].

A combined treatment of coupled acoustic and optical phonons in piezoelectric cubic media was presented that allows determination of the lattice displacement vector and the internal ionic displacement vector simultaneously. It was demonstrated that phonons generally exist in pairs due to couplings between either the lattice displacement component perpendicular to the phonon wave vector (u_y coordinate) and

the electric potential ϕ or the two other lattice displacement components (u_x and u_z). Three new conjectures were derived for piezoelectric cubic slabs: (1) isolated optical phonon modes cannot exist unless the in-plane phonon wave number vanishes ($q_x = 0$); (2) we proved that confined acousto-optical phonon modes only exist in piezoelectric cubic slabs for a discrete set of in-plane wave numbers; (3) at the LO phonon frequency coupled acousto-optical phonons cannot exist. We point out

that the present method can be generalized along the same lines to other piezoelectric materials such as piezoelectric hexagonal (wurtzite) materials.

ACKNOWLEDGMENT

The authors are grateful to Professor Antti-Pekka Jauho for comments and a critical reading of the manuscript.

-
- [1] J. H. Quilter, A. J. Brash, F. Liu, M. Glässl, A. M. Barth, V. M. Axt, A. J. Ramsay, M. S. Skolnick, and A. M. Fox, *Phys. Rev. Lett.* **114**, 137401 (2015).
 - [2] A. E. Siegman, *Lasers* (University Science Books, Oxford, 1986).
 - [3] M. Glässl, A. M. Barth, and V. M. Axt, *Phys. Rev. Lett.* **110**, 147401 (2013).
 - [4] M. J. Gullans, Y.-Y. Liu, J. Stehlik, J. R. Petta, and J. M. Taylor, *Phys. Rev. Lett.* **114**, 196802 (2015).
 - [5] M. Sargent, M. Scully, and W. Lamb, *Laser Physics* (Perseus Books Group, New York, 1978).
 - [6] Y. Y. Liu, J. Stehlik, C. Eichler, M. J. Gullans, J. M. Taylor, and J. R. Petta, *Science* **347**, 285 (2015).
 - [7] P. Kaer, T. R. Nielsen, P. Lodahl, A.-P. Jauho, and J. Mørk, *Phys. Rev. Lett.* **104**, 157401 (2010).
 - [8] A. Nysteen, P. Kaer, and J. Mørk, *Phys. Rev. Lett.* **110**, 087401 (2013).
 - [9] A. Auffèves, J. M. Gerard, and J. P. Poizat, *Phys. Rev. A* **79**, 053838 (2009).
 - [10] T. Yoshie, A. Scherer, J. Heindrickson, G. Khitrova, H. M. Gibbs, G. Rupper, C. Ell, O. B. Shchekin, and D. G. Deppe, *Nature (London)* **432**, 200 (2004).
 - [11] H. J. Carmichael, R. J. Brecha, M. G. Raizen, H. J. Kimble, and P. R. Rice, *Phys. Rev. A* **40**, 5516 (1989).
 - [12] E. Knill, R. Laflamme, and G. J. Milburn, *Nature (London)* **409**, 46 (2001).
 - [13] M. W. Graham, S.-F. Shi, D. C. Ralph, J. Park, and P. L. McEuen, *Nat. Phys.* **9**, 103 (2009).
 - [14] F. D. Natterer, Y. Zhao, J. Wyrick, Y.-H. Chan, W.-Y. Ruan, M.-Y. Chou, K. Watanabe, T. Taniguchi, N. B. Zhitenev, and J. A. Stroschio, *Phys. Rev. Lett.* **114**, 245502 (2015).
 - [15] D. N. Basov, M.M. Fogler, A. Lanzara, F. Wang, and Y. Zhang, *Rev. Mod. Phys.* **86**, 959 (2014).
 - [16] W. Wu, L. Wang, Y. Li, F. Zhang, L. Lin, S. Niu, D. Chenet, X. Zhang, Y. Hao, T. F. Heinz, J. Hone, and Z. L. Wang, *Nature (London)* **514**, 470 (2014).
 - [17] K. Kaasbjerg, K. S. Thygesen, and A.-P. Jauho, *Phys. Rev. B* **87**, 235312 (2013).
 - [18] S. H. Zhang, W. Xu, F. M. Peeters, and S. M. Badalyan, *Phys. Rev. B* **89**, 195409 (2014).
 - [19] M. Woszczyzna, M. Friedemann, M. Gtz, E. Pesel, K. Pierz, T. Weimann, and F. J. Ahlers, *Appl. Phys. Lett.* **100**, 164106 (2012).
 - [20] S. Fratini and F. Guinea, *Phys. Rev. B* **77**, 195415 (2008).
 - [21] V. Perebeinos, S. V. Rotkin, A. G. Petrov, and P. Avouris, *Nano Lett.* **9**, 312 (2009).
 - [22] A. Clerk, *Nat. Phys.* **8**, 256 (2012).
 - [23] M. V. Gustafsson, P. V. Santos, G. Johansson, and P. Delsing, *Nat. Phys.* **8**, 338 (2012).
 - [24] M. Farhat, S. Guenneau, and H. Bagci, *Phys. Rev. Lett.* **111**, 237404 (2013).
 - [25] J. Schiefele, J. Pedros, F. Sols, F. Calle, and F. Guinea, *Phys. Rev. Lett.* **111**, 237405 (2013).
 - [26] K. H. Michel and B. Verberck, *Phys. Rev. B* **80**, 224301 (2009).
 - [27] J. J. Licari and R. Evrard, *Phys. Rev. B* **15**, 2254 (1977).
 - [28] K. L. Kliewer and R. Fuchs, *Phys. Rev.* **144**, 495 (1966); **150**, 573 (1966).
 - [29] N. Bannov, V. Aristov, V. Mitin, and M. A. Stroschio, *Phys. Rev. B* **51**, 9930 (1995).
 - [30] B. K. Ridley, *Electrons and Phonons in Semiconductor Multilayers*, 2nd ed. (Cambridge University Press, New York, 2009).
 - [31] M. A. Stroschio and M. Dutta, *Phonons in Nanostructures* (Cambridge University Press, Cambridge, 2001).
 - [32] Details will be presented elsewhere.
 - [33] M. Born and K. Huang, *Dynamical Theory of Crystal Lattices* (Oxford University Press, Oxford, 1954).