



## Equilibrium sampling for a thermodynamic assessment of contaminated sediments

**Mayer, Philipp; Nørgaard Schmidt, Stine; Mäenpää, Kimmo; Witt, Gesine; Schäfer, Sabine; Gidley, Phil; Jahnke, Annika**

*Publication date:*  
2015

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Mayer, P. (Author), Nørgaard Schmidt, S. (Author), Mäenpää, K. (Author), Witt, G. (Author), Schäfer, S. (Author), Gidley, P. (Author), & Jahnke, A. (Author). (2015). Equilibrium sampling for a thermodynamic assessment of contaminated sediments. 2D/3D (physical products), Technical University of Denmark, DTU Environment.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/283720827>

# Equilibrium sampling for a thermodynamic assessment of contaminated sediments

CONFERENCE PAPER · NOVEMBER 2015

---

READS

64

7 AUTHORS, INCLUDING:



[Philipp Mayer](#)

Technical University of Denmark

153 PUBLICATIONS 3,283 CITATIONS

[SEE PROFILE](#)



[Kimmo Mäenpää](#)

University of Eastern Finland

25 PUBLICATIONS 239 CITATIONS


[SEE PROFILE](#)

# Equilibrium sampling for a thermodynamic assessment of contaminated sediments


Philipp Mayer<sup>1</sup>, Stine N. Schmidt<sup>1</sup>, Kimmo Mäenpää<sup>2</sup>,  
Gesine Witt<sup>3</sup>, Sabine Schäfer<sup>4</sup>, Phil Gidley<sup>5</sup> & Annika Jahnke<sup>6</sup>

<sup>1</sup>Technical University of Denmark, <sup>2</sup>University of Eastern Finland, <sup>3</sup>Hamburg University of Applied Sciences, Germany, <sup>4</sup>German Federal Institute of Hydrology, <sup>5</sup> USACE-ERDC, USA & <sup>6</sup>Stockholm University, Sweden (now UFZ, Germany)

## Classical extraction





## Equilibrium sampling device (ESD)



# 1) Matrix-SPME with disposable fibers (2000)



1. Equilibrate thin polymer with sample



2. Measure  $C_{PDMS}$

$$3. C_{free} = C_{PDMS} / K_{PDMS,water}$$

Mayer et al, ES&T, 2000, V 34: 5177-5183

Mayer et al., ES&T, 2003, V37: 184A-191A

## 2) Silicone coated vials (2008)


# Chemistry Central Journal

Methodology

### Determining the chemical activity of hydrophobic organic compounds in soil using polymer coated vials



Fredrik Reichenberg<sup>1,2</sup>, Foppe Smedes<sup>3</sup>, Jan-Åke Jönsson<sup>2</sup> and Philipp Mayer\*<sup>1</sup>

Address: <sup>1</sup>Department of Environmental Chemistry and Microbiology, National Environmental Research Institute, University of Aarhus, PO Box 358, 4000 Roskilde, Denmark, <sup>2</sup>Division of Analytical Chemistry, Lund University, PO Box 124, S-221 00 Lund, Sweden and <sup>3</sup>Ministry of Transport, Public Works and Water management, National Institute for Coastal and Marine Management/RIKZ, PO Box 207, 9750 AE Haren, Netherlands



Drawing of equilibrium sampling device (ESD) for determination of thermodynamic activity by partitioning to micrometer thin polydimethylsiloxane (PDMS).

# Coated vials with multiple coating thickness




Confirms:

1. equilibrium
2. negligible depletion
3. no surface artefacts/abrasion



3)

## Passive Equilibrium Sampler for in Situ Measurements of Freely Dissolved Concentrations of Hydrophobic Organic Chemicals in Sediments

Gesine Witt,<sup>\*,†</sup> Susann-Cathrin Lang,<sup>†</sup> Dagny Ullmann,<sup>†</sup> Gotja Schaffrath,<sup>†</sup> Detlef Schulz-Bull,<sup>‡</sup> and Philipp Mayer<sup>§</sup>



- **in-situ deployment of sampler**
- **rinse with water and wipe dry**
- **store fiber in GC injection liner**
  
- **insert fiber with automated injection liner exchange (ALEX)**
- **thermal desorption**
- **MS identification/ quantification**





# Measurement endpoints of Equilibrium Sampling

PCB concentrations in polymer  $C_{\text{polymer} \rightleftharpoons \text{sed}}$  can be converted into:

1) The effective concentration driving diffusion

$$C_{\text{free}} \text{ (g L}^{-1}\text{)} = \frac{C_{\text{polymer} \rightleftharpoons \text{sed}} \text{ (g kg}^{-1}\text{)}}{K_{\text{polymer}:\text{water}} \text{ (L kg}^{-1}\text{)}} \quad (\text{since 2000})$$

2) The chemical activity (energetic state) driving equilibrium partitioning

$$a \text{ (unit less)} \cong \frac{C_{\text{free}} \text{ (g L}^{-1}\text{)}}{S_L \text{ (g L}^{-1}\text{)}} \quad (\text{since 2006})$$

**3) The thermodynamic potential for bioaccumulation**


$$C_{\text{lip} \rightleftharpoons \text{sed}} \text{ (g kg}^{-1}\text{)} = C_{\text{polymer} \rightleftharpoons \text{sed}} \text{ (g kg}^{-1}\text{)} \times K_{\text{lip:pol}} \text{ (kg kg}^{-1}\text{)} \quad (\text{since 2011})$$

Case Studies focusing on  
PCBs &  $C_{lip \rightleftharpoons sed}$

1)

## Sensitive Equilibrium Sampling To Study Polychlorinated Biphenyl Disposition in Baltic Sea Sediment



Annika Jahnke,<sup>\*,†</sup> Philipp Mayer,<sup>‡</sup> and Michael S. McLachlan<sup>†</sup>



## Silicone passive equilibrium samplers as 'chemometers' in eels and sediments of a Swedish lake†

Cite this: *Environ. Sci.: Processes Impacts*, 2014, 16, 464


Annika Jahnke,<sup>\*a</sup> Philipp Mayer,<sup>bc</sup> Michael S. McLachlan,<sup>a</sup> Håkan Wickström,<sup>d</sup> Dorothea Gilbert<sup>c</sup> and Matthew MacLeod<sup>a</sup>



3)


# Equilibrium Sampling to Determine the Thermodynamic Potential for Bioaccumulation of Persistent Organic Pollutants from Sediment

Annika Jahnke,<sup>\*,†</sup> Matthew MacLeod,<sup>†</sup> Håkan Wickström,<sup>‡</sup> and Philipp Mayer<sup>§</sup>




# Equilibrium Sampling to Determine the Thermodynamic Potential for Bioaccumulation of Persistent Organic Pollutants from Sediment

Annika Jahnke,<sup>\*,†</sup> Matthew MacLeod,<sup>†</sup> Håkan Wickström,<sup>‡</sup> and Philipp Mayer<sup>§</sup>



# Equilibrium Sampling to Determine the Thermodynamic Potential for Bioaccumulation of Persistent Organic Pollutants from Sediment

Annika Jahnke,<sup>\*,†</sup> Matthew MacLeod,<sup>†</sup> Håkan Wickström,<sup>‡</sup> and Philipp Mayer<sup>§</sup>






# Equilibrium sampling of polychlorinated biphenyls in River Elbe sediments – Linking bioaccumulation in fish to sediment contamination



Sabine Schäfer<sup>a,\*</sup>, Catherine Antoni<sup>a,1</sup>, Christel Möhlenkamp<sup>a</sup>, Evelyn Claus<sup>a</sup>, Georg Reifferscheid<sup>a</sup>, Peter Heininger<sup>a</sup>, Philipp Mayer<sup>b</sup>

1. ex situ equilibrium sampling of sediment from 10 locations in River Elbe (Germany)
2.  $C_{lipid-normalized}$  in common bream from independent monitoring program



1. Cross validation of two monitoring programs
2. Bioaccumulation in common bream tightly linked to  $C_{lip=sed}$ 
  - $R^2 > 0.9$  (linear scale!) & very similar slopes between sites




5)

ET&C, 2015, V 34: 2463–2474 (DOI: 10.1002/etc.3099)

## Fate of polychlorinated biphenyls in a contaminated lake ecosystem: Combining equilibrium passive sampling of sediment and water with total concentration measurements of biota

Kimmo Mäenpää, Matti Leppänen, Kaisa Figueiredo, Philipp Mayer, Dorothea Gilbert, Sirpa Herve, Annika Jahnke, Carmen Gil-Allué, Jarkko Akkanen, Inna Nybom, Auli Kostamo



# Quantitative Thermodynamic Exposure Assessment (Q-TEA) Supporting Resilient Contaminated Sediment Site Restoration

**USACE-ERDC (Lead):** Todd S. Bridges, Philip T. Gidley, Guilherme Lotufo, Carlos Ruiz, Alan J. Kennedy,

**DTU:** Philipp Mayer, Stine Nørgaard Schmidt

**Sediment Solutions:** Upal Ghosh


**USA-EPA:** Robert M. Burgess

**Northeastern University:** Loretta Fernandez,



# Cross validation of equilibrium sampling

1. Coated jars produced at DTU & BfG
2. Equilibrium sampling of New Bedford sediment at BfG & ERDC
3. Instrumental analysis at BfG & ERDC



# Conclusions and take home messages

## Equilibrium sampling of sediments

- equilibrium sampling feasible even for PCBs
- can yield  $C_{\text{free}}$ , chemical activity &  $C_{\text{lip} \rightleftharpoons \text{sed}}$

## $C_{\text{lip} \rightleftharpoons \text{sed}}$

- in all studies so far  $\geq C_{\text{lipid,normalized}}$
- thermodynamic potential for bioaccumulation

Sediments have high PCB levels, not only in concentration but also thermodynamically!

## **Acknowledgements**

This research was financially supported by

- EU Commission (OSIRIS, NOMIRACLE & MODELPROBE)
- Danish Council for Independent Research in Technology and Production Sciences (FTP)
- Swedish Research Council FORMAS
- German Federal Ministry for the Environment, Nature Conservation, Building and Nuclear Safety
- German Research Council (DFG)
- Academy of Finland
- US Strategic Environmental Research and Development Program (SERDP) (14 ER03-035/ER-2431).