



## Chemical herder effectiveness as oil spill response tool in ice-infested water

van Gelderen, Laurens; Adamopoulou, E.; Fritt-Rasmussen, Janne; Jomaas, Grunde

*Published in:*  
Proceedings of Arctic Frontiers 2016

*Publication date:*  
2016

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
van Gelderen, L., Adamopoulou, E., Fritt-Rasmussen, J., & Jomaas, G. (2016). Chemical herder effectiveness as oil spill response tool in ice-infested water. In *Proceedings of Arctic Frontiers 2016*

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Chemical herder effectiveness as oil spill response tool in ice-infested water

---

Laurens van Gelderen<sup>a</sup>, Eirini Adamopoulou<sup>a</sup>, Janne Fritt-Rasmussen<sup>b</sup>, Grunde Jomaas<sup>a</sup>

<sup>a</sup>*Department of Civil Engineering, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark*

<sup>b</sup>*Department of Bioscience, Aarhus University, 4000 Roskilde, Denmark*

## Abstract

The effectiveness of the chemical herding of crude oil, an enhancement for oil spill response methods, was studied in ice-infested waters to evaluate the applicability to oil spills in Arctic waters. The average slick thickness, surface coverage distribution of the slicks, as well as the burning efficiency of a North Sea crude oil were studied after herding to determine the herder effectiveness as a function of the ice coverage, amount of herder, wind speed and use of free floating or fixed ice. Experiments were performed in a small scale laboratory setup, featuring a 1.0x1.0x0.5 m<sup>3</sup> water basin (1 m<sup>2</sup> surface area), and in an outdoor test basin of 4.0x4.0x0.05 m<sup>3</sup> (16 m<sup>2</sup> surface area) in Sisimiut, Greenland. The chemical herder, which is a surfactant that spreads out to form a monolayer, was distributed equally (150 µL/m<sup>2</sup> oil) among the edges of the water surface after the crude oil had spread out, to compact and thicken the oil slick. Objects on the water surface were shown to have a negative effect on this process, distributing up to 25% of the total oil among additional small oil slicks as compared to the formation of a single oil slick on open water. The average slick thickness of the herded oil reached up to approximately 6 mm and the oil slicks could easily be ignited. However, the formation of additional slicks reduced the burning efficiency by at least 8%, as some of the smaller slicks did not participate in the burning. Burning times were also increased significantly as separate slicks needed to be ignited one at a time. The results showed that while the herding process is effective because it thickens the oil slick to an ignitable thickness, its effectiveness was reduced with increasing ice concentrations on the water surface. Current herding and *in-situ* burning logistics would need to adapt to the formation and burning of multiple oil slicks in ice-infested waters common in the Arctic environment in order to achieve maximum efficiency.