Squeezed light in optomechanical systems

Harris, G. I.; Taylor, M. A.; Hoff, Ulrich Busk; Janousek, J.; Daria, V.; Knittel, J.; Kerdoncuff, Hugo; Hage, B.; Andersen, Ulrik Lund; Bachor, H.-A.

Total number of authors:

11

Published in:

Proceedings of Frontiers in Optics (FiO)

Publication date:

2012

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Harris, G. I., Taylor, M. A., Hoff, U. B., Janousek, J., Daria, V., Knittel, J., Kerdoncuff, H., Hage, B., Andersen, U. L., Bachor, H.-A., & Bowen, W. P. (2012). Squeezed light in optomechanical systems. In *Proceedings of Frontiers in Optics (FiO)* (pp. FW5C.4). Optical Society of America.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Squeezed light in optomechanical systems

G. I. Harris¹, M. A. Taylor¹, U. Busk-Hoff³, J. Janousek², V. Daria², J. Knittel¹, H. Kerdoncuff³, B. Hage², U. L. Andersen³, H.-A. Bachor², and W. P. Bowen¹

¹Centre of Excellence in Engineered Quantum Systems, University of Queensland, St. Lucia, Queensland 4072, Australia

²Department of Quantum Science, Australian National University, Canberra, ACT 0200, Australia

³Department of Physics, Technical University of Denmark, Building 309, 2800 Lyngby, Denmark

Presenting author e-mail address: glen.harris@uqconnect.edu.au

Abstract: Squeezed light enhanced optomechanical measurements are demonstrated in both intra-cavity and biological contexts, with respective enhancements of 1.0 and 2.7 dB. Quantum enhanced microrheology of the cytoplasm of a yeast cell is thereby realized.

OCIS codes: (270.0270) Quantum optics, (230.4685) Optical microelectromechanical devices, (120.4880) Optomechanics.

1. Introduction

The quantum nature of light places a fundamental limit on the sensitivity of optical measurements. In circumstances with constrained optical power, this limit may only be surpassed using non-classical resources. Despite the promise of non-classical resources, to date the sole example of a real application is in interferometric gravity wave detection where the optical power is constrained due to absorptive heating[1]. Much broader and widely discussed applications are possible in the areas of quantum optomechanics, where ultimately a non-classical light field interacts with a non-classical mechanical oscillator, and biological sensing[2], where low light levels are often required to avoid damaging the specimen. Here we report two distinct experiments which use squeezed light to, for the first time, achieve quantum enhanced sensitivity in biological and cavity optomechanical contexts, with 2.7 and 1.0 dB of enhancement achieved, respectively. The biological experiments, particularly, allow quantum enhanced microrheology to be performed on the cytoplasm of a *Saccharomyces cerevisiae* yeast cell, revealing subdiffusive motion. This provides a pathway towards microrheology of cell mechanics and the cytoskeleton at high frequencies, where motion amplitudes are beneath the sensitivity of current technology.

2. Microcavity experiment

As shown in Fig. 1, phase squeezed light was injected, on resonance, into a silicon chip based microtoroidal resonator. Microtoroidal resonators exhibit high quality mechanical modes, with an example shown in Fig. 1 (right), which have been cooled close to their quantum ground state[3]. The output field from the microtoroid was interfered with a bright local oscillator field on a 50/50 beam splitter. The non-classical photon correlations inherent to phase squeezing allowed the mechanical motion of the microtoroid to be transduced with 1 dB higher sensitivity than the quantum noise limit introduced by quantization of light.

Fig. 1. Left: schematic of cavity optomechanical squeezing experiment. SQZ: squeezer. Center right: SEM image of microtoroid. Far right: microtoroid mechanical mode.

3. Optical tweezers experiment

The apparatus used to demonstrate sub-quantum noise limited sensitivity in a biological context is shown in Fig.~1. A specimen was suspended in water within a sample chamber formed by two microscope coverslips, and trapped with a dual beam optical trap at 1064 nm. An orthogonally polarized amplitude squeezed local oscillator field was spatially engineered using a phase plate for maximum sensitivity to motion of the specimen (see vertical panels on the left of Fig. 2) and injected into an optical tweezers. A further probe laser field, coherent with the squeezed local oscillator field, was injected transversely to the optical axis of the tweezers. Scattering of this field interfered with

the squeezed field, encoding position information about the specimen which could be retrieved via direct detection. Apart from the deleterious effects of optical losses and spatial distortion in the objectives and specimen, a major technical challenge was to minimize exposure to noise sources at the sub-kHz frequencies relevant to biological motion. This was achieved by stroboscopically pulsing the probe field at 3.522-MHz, which had the effect of mixing up the motion into the region of strongest squeezing. This technique should be broadly applicable to squeezed light enhanced sensors.

Motion sensitivity surpassing the quantum noise limit by up to 2.7-dB and 2.4-dB was achieved, respectively, for trapped silica beads and lipid granules within a yeast cell; in both cases allowing the mean squared displacement of the specimen to be more accurately measured than is possible with classical light. The mean squared displacement measurements for yeast with (orange) and without (blue) squeezing are shown in Fig. 2 (*right*), clearly demonstrating the superiority of measurements with squeezed light. The lipid granule exhibits sub-diffusive motion as it interacts with the cells cytoplasm, consistent with recent classical measurements with a different yeast strain[4].

Fig.2. *Left*: experimental schematic, vertical panels show the probe (red) and scattered (blue) fields, illustrating dependence of detected intensity on particle position. *Right*: mean squared displacement measurements on lipid particle within a yeast cell. Orange and blue respectively indicate measurement results and uncertainty with and without squeezed light. Dashed line: expected mean squared displacement for diffusive motion.

4. Conclusion

We have demonstrated optomechanical transduction sensitivity surpassing the quantum noise limit in both cavity optomechanical systems, and optical tweezers. These experiments represent enabling steps towards new practical sensors, including force measurement beyond the standard quantum limit, measurements of the non-Einsteinian Brownian motion[5], and low light level high bandwidth sensing of biological dynamics within living cells[6].

5. Acknowledgements

This research was funded by the ARC Centre of Excellence CE110001013 and Discovery Project DP0987146. Device fabrication was undertaken within the Queensland Node of the Australian Nanofabrication Facility.

6. References

- [1] H. Vahlbruch *et al.* *Class. Quantum Grav.* **27** 084027 (2010)
- [2] N. Treps *et al.* *Phys. Rev. Lett.* **88**, 203601 (2002)
- [3] E. Verhagen, S. Deléglise, S. Weis, A. Schliesser, T. J. Kippenberg, *Nature* **482** 63 (2012).
- [4] I. M. Tolic-Norrellykke *et al.* *Phys. Rev. Lett.* **93**, 078102 (2004).
- [5] R. Huang, I. Chavez, K. M. Taute, B. Lukic, S. Jeney, M. G. Raizen, and E.-L. Florin, *Nature Phys.* **7** 576580 (2011).
- [6] S. Yamada, D. Wirtz, and S. C. Kuo, *Biophys. J.* **78**, 17361747 (2000).