

Toxicovenomics and antivenom profiling of the Eastern green mamba snake (Dendroaspis angusticeps)

Lauridsen, Line P.; Laustsen, Andreas Hougaard; Lomonte, Bruno; Gutiérrez, José María

Published in: Journal of Proteomics

Link to article, DOI: 10.1016/j.jprot.2016.02.003

Publication date: 2016

Document Version Peer reviewed version

Link back to DTU Orbit

Citation (APA): Lauridsen, L. P., Laustsen, A. H., Lomonte, B., & Gutiérrez, J. M. (2016). Toxicovenomics and antivenom profiling of the Eastern green mamba snake (*Dendroaspis angusticeps*). *Journal of Proteomics*, *136*, 248-261. https://doi.org/10.1016/j.jprot.2016.02.003

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

2	
3	Toxicovenomics and antivenom profiling of the Eastern green mamba
4	snake (Dendroaspis angusticeps)
5	
6	
7	Line P. Lauridsen ¹ , Andreas H. Laustsen ² , Bruno Lomonte ³ , José María Gutiérrez ³
8	
9	¹ Department of Systems Biology, Technical University of Denmark, Denmark
10	² Department of Drug Design and Pharmacology, Faculty of Health and Medical
11	Sciences, University of Copenhagen, Denmark
12	³ Instituto Clodomiro Picado, Facultad de Microbiología, Universidad de Costa Rica,
13	San José, Costa Rica
14	
14 15	
14 15 16	Running title: Proteomics of green mamba venom
14 15 16 17	Running title: Proteomics of green mamba venom
14 15 16 17 18	Running title: Proteomics of green mamba venom
14 15 16 17 18 19	Running title: Proteomics of green mamba venom
14 15 16 17 18 19 20	Running title: Proteomics of green mamba venom
14 15 16 17 18 19 20 21 22	Running title: Proteomics of green mamba venom
14 15 16 17 18 19 20 21 22 23	Running title: Proteomics of green mamba venom Address for correspondence: Dr José María Gutiérrez
14 15 16 17 18 19 20 21 22 23 24	Running title: Proteomics of green mamba venom Address for correspondence: Dr José María Gutiérrez Professor
14 15 16 17 18 19 20 21 22 23 24 25	Running title: Proteomics of green mamba venom Address for correspondence: Dr José María Gutiérrez Professor Instituto Clodomiro Picado
14 15 16 17 18 19 20 21 22 23 24 25 26	Running title: Proteomics of green mamba venom Address for correspondence: Dr José María Gutiérrez Professor Instituto Clodomiro Picado Facultad de Microbiología
14 15 16 17 18 19 20 21 22 23 24 25 26 27 28	Address for correspondence: Dr José María Gutiérrez Professor Instituto Clodomiro Picado Facultad de Microbiología Universidad de Costa Rica Sen Jacá COSTA PICA
 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 	Address for correspondence: Dr José María Gutiérrez Professor Instituto Clodomiro Picado Facultad de Microbiología Universidad de Costa Rica San José, COSTA RICA iose quiterrez/murca o cr
14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30	Address for correspondence: Dr José María Gutiérrez Professor Instituto Clodomiro Picado Facultad de Microbiología Universidad de Costa Rica San José, COSTA RICA jose.gutierrez@ucr.ac.cr

32 Abstract

A toxicovenomic study was performed on the venom of the green mamba, 33 Dendroaspis angusticeps. Forty-two different proteins were identified in the venom of 34 D. angusticeps, in addition to the nucleoside adenosine. The most abundant proteins 35 belong to the three-finger toxin (3FTx) (69.2%) and the Kunitz-type proteinase inhibitor 36 37 (16.3%) families. Several sub-subfamilies of the 3FTxs were identified, such as Orphan Group XI (Toxin F-VIII), acetylcholinesterase inhibitors (fasciculins), and aminergic 38 toxins (muscarinic toxins, synergistic-like toxins, and adrenergic toxins). Remarkably, 39 40 no α -neurotoxins were identified. Proteins of the Kunitz-type proteinase inhibitor family include dendrotoxins. Toxicological screening revealed a lack of lethal activity 41 in all RP-HPLC fractions, except one, at the doses tested. Thus, the overall toxicity 42 depends on the synergistic action of various types of proteins, such as dendrotoxins, 43 44 fasciculins, and probably other synergistically-acting toxins. Polyspecific antivenoms manufactured in South Africa and India were effective in the neutralization of venom-45 induced lethality. These antivenoms also showed a pattern of broad immunorecognition 46 of the different HPLC fractions by ELISA and immunoprecipitated the crude venom by 47 gel immunodiffusion. The synergistic mechanism of toxicity constitutes a challenge for 48 the development of effective recombinant antibodies, as it requires the identification of 49 the most relevant synergistic toxins. 50

51 (197 words)

Keywords: *Dendroaspis angusticeps*; Green mamba; Snake venom; Proteomics;
Toxicovenomics: Antivenoms.

54

55 Biological significance

Envenomings by elapid snakes of the genus *Dendroaspis*, collectively known as 56 mambas, represent a serious medical problem in sub-Saharan Africa. The development 57 of novel antivenoms and of recombinant neutralizing antibodies demands the 58 identification of the most relevant toxins in these venoms. In this study, a bottom-up 59 approach was followed for the study of the proteome of the venom of the Eastern green 60 mamba, D. angusticeps. Forty-two different proteins were identified, among which the 61 three-finger toxin (3FTx) family, characteristic of elapid venoms, was the most 62 abundant, followed by the Kunitz-type proteinase inhibitor family. In addition, several 63 other protein families were present in the venom, together with the nucleoside 64 adenosine. No α -neurotoxins were identified within the family of 3FTxs in the venom 65 of D. angusticeps, in contrast to the venom of D. polylepis, in which α -neurotoxins are 66 largely responsible for the toxicity. With one exception, HPLC fractions from D. 67 68 angusticeps venom did not kill mice at the doses tested. This underscores that the 69 toxicity of the whole venom is due to the synergistic action of various components, such as fasciculins and dendrotoxins, and probably other synergistically-acting toxins. Thus, 70 the venoms of these closely related species (D. angusticeps and D. polylepis) seem to 71 have different mechanisms to subdue their prey, which may be related to different prey 72 preferences, as D. angusticeps is predominantly arboreal, whereas D. polylepis lives 73 mostly in open bush country and feeds mainly on mammals. It is therefore likely that 74 the predominant clinical manifestations of human envenomings by these species also 75 differ, although in both cases neurotoxic manifestations predominate. Polyspecific 76 antivenoms manufactured in South Africa and India were effective in the neutralization 77 of venom-induced lethality in mice and showed a pattern of broad immunorecognition 78 of the various venom fractions. It is necessary to identify the toxins responsible for the 79

synergistic mode of toxicity in this venom, since they are the targets for the
development of recombinant antibodies for the treatment of envenomings.

82

83 **1. Introduction**

The Eastern green mamba (Dendroaspis angusticeps) is a highly venomous 84 elapid found primarily in southeastern Africa (Figure 1). First described by Smith in 85 1848 [1], D. angusticeps is a relatively small mamba species, averaging 1.4 m in length. 86 Due to its arboreal, shy, and elusive nature, human envenomings are less frequent than 87 those inflicted by the more territorial Dendroaspis polylepis (black mamba) [2,3]. Adult 88 specimens of D. angusticeps have a brilliant emerald to lime green coloration, providing 89 them with an excellent camouflage in their natural habitat of the tropical rainforests in 90 the coastal lowlands of Southeast Africa [4]. D. angusticeps is, however, also found in 91 areas with coastal bush, dune, and montane forest [5], as well as in closer proximity to 92 humans, when residing in farm trees, such as citrus, mango, coconut, and cashew [6]. 93 94 Due to its color and habitat, D. angusticeps is often mistaken for a harmless tree snake, 95 why people often do not take proper precaution [2,3]. D. angusticeps preferably preys 96 on warm-blooded animals, such as rodents, bats, birds, and nestlings, but also on eggs [4]. 97

Despite a low number of human envenomings reported, but due to its potent 98 neurotoxic venom, D. angusticeps is classified as a category 1 snake, which is the 99 highest level of medically important snakes, according to the WHO [7]. Furthermore, its 100 high abundance, particularly in Kenya, Tanzania, Mozambique, Malawi, eastern 101 Zimbabwe, and the Republic of South Africa, makes this a snake of high 102 epidemiological relevance [7]. Severe envenomings by D. angusticeps can lead to rapid 103 104 mortality within only 30 minutes of a bite [6]. The typical clinical manifestations include swelling of the bitten area, dizziness, nausea, difficult breathing, irregular 105 heartbeat, and respiratory paralysis [6]. These life-threatening symptoms may escalate 106 107 rapidly, but deaths are rare when effective antivenom is administered timely [6].

Given the medical importance of D. angusticeps, it is necessary to have a 108 thorough understanding of the composition of its venom, as well as of the underlying 109 110 mechanisms for venom pathophysiology in human victims. Furthermore, preclinical assessment of antivenoms is critical for predicting efficacy of snakebite envenoming 111 112 therapy, which may be used to guide clinicians in the treatment of snakebites by D. 113 angusticeps. Currently, only the SAIMR Polyvalent Snake Antivenom from the South African Vaccine Producers is claimed to be effective against *D. angusticeps*, although it 114 115 is possible that other polyvalent antivenoms raised against the venoms of other mamba 116 species may be effective in neutralization of *D. angusticeps* venom.

The venom of *D. angusticeps* has not undergone a full proteomics evaluation, 117 and its quantitative protein composition is not known. Nevertheless, several biochemical 118 119 and pharmacological studies have been performed on different toxins from D. 120 angusticeps venom [8-11]. These studies report that this venom contains several 121 neurotoxins, such as the fasciculins [10] and dendrotoxins [8,9], which are unique to the 122 Dendroaspis genus [12,13]. This venom also contains a number of other toxins of the 123 three-finger toxin family (3FTx), such as muscarinic toxins, adrenergic toxins, and synergistic-type toxins [14–16]. 124

The dendrotoxins, of structural similarity to the Kunitz-type serine protease 125 126 inhibitors, target the presynaptic voltage-gated potassium channels with high specificity, facilitating the release of acetylcholine from the presynaptic nerve terminals, causing 127 excitatory activity [17,18]. Other important neurotoxins of the 3FTx family present in 128 129 D. angusticeps venom are the fasciculins, which prolong the presence of acetylcholine in the neuromuscular junction by inhibiting acetylcholinesterase, leading to muscle 130 fasciculations [10]. Although the venom composition of D. angusticeps has not been 131 132 elucidated, a study of the venom of the closely related and more feared relative, D. 133 *polylepis* (black mamba), has recently been reported [19]. According to this study, *D*. 134 *polylepis* venom is dominated by α -neurotoxins from the 3FTx family and dendrotoxins 135 (BPTI-type/Kunitz type protease inhibitors). It is therefore of relevance to study the 136 venom proteome of *D. angusticeps* in order to identify similarities and differences with 137 that of *D. polylepis*.

138 Toxicovenomics defines the recent convergence between toxicological evaluation of toxins and venomics [20,21]. Together with antivenomics, this tool may 139 help provide a better understanding of D. angusticeps venom, the relative importance of 140 different proteins for toxicity, and how venom toxicity may best be abrogated. While 141 previous investigations of D. angusticeps have focused on the biochemical and 142 pharmacological features of the toxins, recent advances in the field of venomics and 143 144 antivenomics facilitate development of novel antivenoms through rational and 145 knowledge-based interpretation of pharmacological relevant toxins [22].

Here, we report the first full toxicovenomics analysis of *D. angusticeps*, a quantitative estimation of its venome, and a preclinical and immunochemical assessment of three antivenoms against *D. angusticeps* venom.

149

150 2. Materials and Methods

151 2.1 Snake venom

Venom from *D. angusticeps* was obtained from Latoxan SAS, Valence, France,
from a pool of 50 specimens collected in Tanzania.

154

155 2.2 Venom separation by reverse-phase HPLC and SDS-PAGE

The 'snake venomics' analytical strategy [23] involving fractionation of crude venom by a combination of RP-HPLC and SDS-PAGE separation steps, was followed.

Venom (2 mg) was dissolved in 200 µL of water containing 0.1% trifluoroacetic acid 158 (TFA; solution A) and separated by RP-HPLC (Agilent 1200) on a C₁₈ column (250 x 159 160 4.6 mm, 5 µm particle; Teknokroma). Elution was carried out at 1 mL/min by applying a gradient towards solution B (acetonitrile, containing 0.1% TFA): 0% B for 5 min, 0-161 15% B over 10 min, 15–45% B over 60 min, 45–70% B over 10 min, and 70% B over 9 162 163 min, as previously described [24]. Fractions, collected manually, were dried in a vacuum centrifuge, redissolved in water, reduced with 5% β-mercaptoethanol at 100 °C 164 165 for 5 min, and further separated by SDS-PAGE in 15% gels. Proteins were stained with colloidal Coomassie blue G-250, and gel images were acquired on a ChemiDoc® 166 recorder using ImageLab[®] software (Bio-Rad). 167

168

169 2.3 Protein identification by tandem mass spectrometry of tryptic peptides

170 Protein bands were excised from the polyacrylamide gels and subjected to 171 reduction (10 mM dithiothreitol), alkylation (50 mM iodoacetamide), and overnight ingel digestion with sequencing grade trypsin (Sigma), in 50 mM ammonium bicarbonate 172 at 37 °C. Tryptic peptides were extracted with 50% acetonitrile containing 1% TFA, and 173 analyzed by MALDI-TOF-TOF on an AB4800-Plus Proteomics Analyzer (Applied 174 Biosystems). Peptides were mixed with an equal volume of saturated α -cyano-175 hydroxycinnamic acid (in 50% acetonitrile, 0.1% TFA), and spotted (1 μ L) onto an 176 Opti-TOF 384-well plate, dried, and analyzed in positive reflector mode. TOF spectra 177 178 were acquired using 500 shots at a laser intensity of 3000. Selection of the ten most 179 intense precursor ions was done automatically and their TOF/TOF fragmentation 180 spectra were acquired using 500 shots at a laser intensity of 3900. External calibration in each run was performed with CalMix[®] standards (ABSciex) spotted onto the same 181 182 plate. Resulting spectra were searched against the UniProt/SwissProt database for Serpentes (20150217) using ProteinPilot[®] v.4 and the Paragon[®] algorithm (ABSciex) at $\geq 95\%$ confidence, or, in few cases, manually interpreted and the deduced sequences searched using BLAST (<u>http://blast.ncbi.nlm.nih.gov</u>) for protein family assignment by similarity.

187

188 2.4 Relative protein abundance estimations

Relative abundance of the venom proteins was estimated by integrating the areas 189 of their chromatographic peaks at 215 nm, roughly corresponding to peptide bond 190 abundance, using the ChemStation[®] software (Agilent) [23]. In the case where HPLC 191 192 peaks contained several electrophoretic bands, their percentage distributions were assigned by densitometry, using ImageLab[®] (Bio-Rad). Finally, for electrophoretic 193 bands in which more than one protein was identified by MALDI-TOF-TOF, their 194 195 percentage distributions were estimated on the basis of the corresponding intensities of 196 the intact protein ions, as observed in the nESI-MS analysis. Intensities lower than 5% 197 (relative to the major protein ions in such mixtures) were considered as traces.

198

199 2.5 Adenosine analysis

The presence of the nucleoside adenosine was determined by spiking a sample of 200 2 mg of venom with 10 µg of adenosine and separating it by RP-HPLC as described in 201 section 2.2. If the adenosine coincided with a peak already present in a crude venom 202 sample (as judged by the increment in the height of the peak), and if this venom peak 203 204 showed an ESI-MS spectrum essentially identical to adenosine, the identity of venom component was judged to be adenosine. Further confirmation of the molecular identity 205 of adenosine was obtained by acquiring its collision-induced dissociation MS/MS 206 207 spectrum in positive mode, using the Enhanced Product Ion tool of Analyst v1.5 in the QTrap3200 mass spectrometer, to show the expected reporter ion transition $268 \rightarrow 136$. Nucleoside abundance was estimated by deriving un-spiked nucleoside concentration from integrating the areas of both spiked and un-spiked chromatographic peaks.

211

212 2.6 In vitro enzymatic activities

213 2.6.1. Phospholipase A₂ activity

Assay of PLA₂ activity was carried out using the monodisperse synthetic 214 chromogenic substrate 4-nitro-3-octanoyloxybenzoic acid (NOBA) [25]. Twenty-five 215 216 μ L of solution containing various amounts of venom were mixed with 200 μ L of 10 mM 217 Tris, 10 mM CaCl₂, 0.1 M NaCl, pH 8.0, and 25 µL of NOBA to achieve a final 218 substrate concentration of 0.32 mM. Plates were incubated at 37 °C for 60 min, and 219 absorbances were recorded at 405 nm in a microplate reader. For comparative purposes, 220 the activities of the venoms of D. polylepis and the viperid snake Bothrops asper were 221 also assessed.

222

223 2.6.2 Proteinase activity

Proteinase activity was assayed by adding 20 µg of venom to 100 µL of 224 azocasein (10 mg/mL in 50 mM Tris-HCl, 0.15 M NaCl, 5 mM CaCl₂ buffer, pH 8.0), 225 226 and incubated for 90 min at 37 °C. The reaction was terminated by addition of 200 μ L of 5% trichloroacetic acid, and after centrifugation (5 min, 6000 g), 150 µL of 227 supernatants were mixed with 100 µL of 0.5 M NaOH, and absorbances were recorded 228 229 at 450 nm. The absorbance of azocasein incubated with distilled water alone was used as a blank, being subtracted from all readings [26]. For comparative purposes, the 230 activities of the venoms of D. polylepis and the viperid snake Bothrops asper were also 231 assessed. 232

234 2.7 Toxicological profiling

235 2.7.1 Animals

In vivo assays were performed in CD-1 mice of both sexes, provided by Instituto Clodomiro Picado, following protocols approved by the Institutional Committee for the Use and Care of Animals (CICUA), University of Costa Rica. Mice were provided food and water *ad libitum*.

240

241 2.7.2 Toxicity of crude venom and isolated venom fractions

The lethality of the whole venom and venom fractions was tested by intravenous (i.v.) injection in groups of four mice (18–20 g body weight). Various amounts of venom or venom fractions were dissolved in phosphate-buffered saline (PBS; 0.12 M NaCl, 0.04 M sodium phosphate buffer, pH 7.2) and injected in the caudal vein, in a volume of 100 μ L. Deaths occurring within 24 h were recorded, and LD₅₀s were calculated by probits [27], using the BioStat[®] software (AnalySoft).

The acute toxicity of venom fractions was initially screened by selecting a dose 248 based on fraction abundance in the venom and assuming a venom yield of 75 mg for D. 249 (http://snakedatabase.org/pages/LD50.php#legendAndDefinitions), 250 angusticeps 251 Laustsen et al.'s Toxicity Score [20], and 50 kg as the weight of a human being. On this basis, a cutoff dose (mg/kg) was selected and tested for each fraction. Fractions that 252 253 were not lethal at this dose (corresponding to a Toxicity Score below 7) were considered 254 as having insignificant acute toxicity, whereas fractions which did kill mice at this level were further evaluated, and precise $LD_{50}s$ were determined for them. 255

256

257 2.8 Antivenoms

233

Polyspecific antivenoms from the following manufacturers were used: (a) 258 SAIMR (South African Institute for Medical Research) Polyvalent Snake Antivenom 259 from South African Vaccine Producers (Pty) Ltd (batch number BC02645, expiry date 260 07/2016); (b) Snake Venom Antivenom (Central Africa) from VINS Bioproducts Ltd 261 262 (batch 12AS13002, expiry date 04/2017); (c) Snake Venom Antivenom (African) from 263 VINS Bioproducts Ltd (batch 13022, expiry date 01/2018). In addition, the monospecific Micrurus nigrocinctus Anticoral Antivenom from Instituto Clodomiro 264 Picado (batch 5310713ACLQ, expiry date 07/2016) was used for comparison in certain 265 experiments. 266

267

268 2.9 Immunoreactivity of antivenoms against crude venom and venom fractions by ELISA

Wells in MaxiSorp plates (NUNC, Roskilde, Denmark) were coated overnight 269 270 with 1.0 µg of each HPLC venom fraction, or crude venom, dissolved in 100 µL PBS. 271 After a washing step, wells were blocked by adding 100 μ L PBS containing 2% (w:v) bovine serum albumin (BSA, Sigma), and incubated at room temperature for 1 h. Plates 272 were then washed five times with PBS. A dilution of each antivenom in PBS + 2% BSA273 was prepared. 100 µL of antivenom solution was added to each well in triplicates and 274 incubated for 2 h. Plates were then washed five times with PBS. 100 µL of a 1:2000 275 276 dilution of conjugated antibody (Sigma A6063, rabbit anti-horse IgG (whole molecule)alkaline phosphatase in PBS + 1% BSA) was then added to each well. The plates were 277 278 incubated for 2 h, and then washed five times with FALC buffer (0.05 M Tris, 0.15 M NaCl, 20 µM ZnCl₂, 1 mM MgCl₂, pH 7.4). Development of color was attained by 279 addition of 100 µL p-nitrophenyl phosphate (1 mg/mL in 9.7% v/v diethanolamine 280 buffer, pH 9.8) and absorbances at 405 nm were recorded (Multiskan FC, Thermo 281 282 Scientific).

Agarose was dissolved in 30 mL of PBS to attain 1% concentration, and poured into a Petri dish. Six holes were punched in the gel, and 50 μ l of antivenom was placed in the center well, while 30 μ l of solutions of *D. angusticeps* and *D. polylepis* venoms were added to the surrounding wells at variable concentrations (0.5, 1 and 2 μ g/ μ L). After overnight incubation at room temperature, agarose gels were photographed using a ChemiDoc[®] recorder and ImageLab[®] software.

292

293 2.10 Neutralization studies with antivenoms

294 Mixtures containing a fixed amount of venom and variable dilutions of 295 antivenoms were prepared using PBS as diluent and incubated at 37 °C for 30 min. 296 Controls contained PBS instead of antivenom. Aliquots of 100 µL of the solutions, 297 containing 4 LD₅₀s of venom (64 μ g/mouse), were then injected i.v. into groups of four 298 mice (18-20 g). Deaths occurring within 24 h were recorded for determining the neutralizing capacity of antivenoms. Neutralization was expressed as the Median 299 Effective Dose (ED₅₀) of antivenom, defined as the ratio mg venom/mL antivenom at 300 301 which 50% of the injected mice were protected. The ED_{50} s as estimated by probits, as described in Section 2.6.2. 302

303

304 3.0 Results and Discussion

305 3.1 Venomics

306 SDS-PAGE separation of venom proteins revealed similarities and differences 307 between the venoms of *D. angusticeps* and *D. polylepis* (Figure 2). Both venoms

showed predominantly low molecular mass bands, in addition to a number of bands of a 308 wide range of molecular masses, including some large proteins with molecular masses 309 310 above 100 kDa. When SDS-PAGE was run under non-reducing conditions, the venom 311 of D. angusticeps showed more bands than that of D. polylepis. In particular, D. 312 angusticeps venom presented three bands of molecular masses between 18 and 22 kDa, 313 which were absent in the venom of D. polylepis. In turn, D. polylepis venom had a band of 37 kDa, absent in *D. angusticeps* venom. In contrast, with the exception of a 25 kDa 314 315 band in the venom of *D. angusticeps*, the majority of these intermediate molecular mass 316 bands were not observed in reduced gels, indicating that these bands were comprised of higher order protein complexes. 317

A bottom-up proteomic characterization of D. angusticeps venom was carried 318 319 out. Using RP-HPLC, the venom was resolved into 29 fractions, where the first three 320 eluting from the column did not contain proteins as evidenced by electrophoresis. After 321 SDS-PAGE separation, the remaining 26 fractions were resolved into 63 bands (Figure 3), of which 59 resulted in positive identifications upon in-gel digestion and MALDI-322 323 TOF-TOF analysis, whereas 4 remained unknown. In total, 42 different proteins were identified within these bands (Table 1). In certain cases, exemplified by fraction number 324 5, the bands separated by SDS-PAGE contained the same protein in both monomer and 325 dimer forms. 326

Fractions 1-3 did not contain proteins according to SDS-PAGE. Due to its high abundance, fraction 1 was analyzed by direct infusion using nESI-MS/MS, which revealed a component with a molecular mass of 268 Da. Upon collision-induced dissociation, this ion produced a fragment of 136 Da, corresponding to the characteristic transition of adenosine. Furthermore, spiking with adenosine as described in section 2.5, provided an estimation that 0.75% of the chromatographic signal of the venom corresponded to this nucleoside. Presence of a substantial amount of adenosine was also
observed in the venom of *D. polylepis* [19]. Adenosine might play an auxiliary role in
the toxicity of mamba venoms owing to its vasodilatory effect, as previously suggested
[19].

The overall protein composition of *D. angusticeps* venom was determined by 337 338 assigning the identified proteins to families and expressing these as percentages of total protein content (Figure 4). The most abundant components belong to the three-finger 339 340 toxin family (3FTx; 69.2%) and the family of Kunitz-type proteinase inhibitors, which includes the dendrotoxins (KUN; 16.3%). The 3FTxs in elapid venoms all share a 341 common structural architecture with a distinct protein fold, comprising between 60 and 342 343 80 amino acids in length, containing a small, globular, hydrophobic core with four or 344 five conserved disulfide bridges, from which three β -stranded loops extend [28–30]. 345 This makes this group of toxins resemble three outstretched fingers [31]. Despite the 346 common structural motif, a diverse array of functions has been associated with 3FTxs 347 [32].

All 3FTxs found in *D. angusticeps* venom belong to the short chain subfamily, but attained to different sub-subfamilies (Figure 4). The majority of 3FTxs in this venom belong to the Orphan Group XI (from Toxin FV-III), whose function has not yet been established [29], followed by aminergic toxins (Muscarinic toxin 2, Muscarinic toxin 4, Synergistic-like protein, and Adrenergic toxins) [16]. A further 8.4% of 3FTxs were attained to fasciculins (all from the acetylcholinesterase inhibitory sub-subfamily), which are unique to *D. angusticeps* [33].

Interestingly, the proteomic analysis of the 3FTxs of *D. angusticeps* venom did not reveal the presence of α -neurotoxins, perhaps the most studied 3FTxs from elapid venoms. α -neurotoxins bind with high affinity to the nicotinic cholinergic receptor at the motor end-plate of the neuromuscular junction, causing a blockage in neuromuscular transmission and flaccid paralysis, generally inducing death by respiratory failure [34]. α -neurotoxins show the highest Toxicity Score values among the fractions of *D*. *polylepis* venom [19]. Their absence in the venom of *D*. *angusticeps* marks a significant difference between these two mamba venoms and suggests that the predominant mechanisms for prey immobilization in these venoms might be different.

Another type of neurotoxins unique to the *Dendroaspis* genus, and found in our 364 365 proteomic analysis of D. angusticeps venom, is comprised by the dendrotoxins, which are homologous to Kunitz-type serine proteinase inhibitors [35]. Dendrotoxins interact 366 367 and inhibit the presynaptic voltage-gated potassium channels, thus exerting a facilitatory effect associated with excitability [18,35]. The venom of D. angusticeps has a lower 368 relative content of Kunitz-type proteinase inhibitors, but a higher content of 3FTxs, 369 370 when compared to the venom of D. polylepis [19]. The combined action of the fasciculins and dendrotoxins results in enhanced skeletal muscle excitability and 371 372 contraction, probably leading to respiratory arrest.

Other protein families found in lower proportions in the venom of D. 373 angusticeps include metalloproteinases (SVMP; 6.7%), cysteine-rich secretory proteins 374 (CRISP; 2.1%), and traces of Galactose-binding lectins (GAL; < 0.5%), peptidases 375 (PEP; < 0.1%), hyaluronidases (HYA; < 0.3%), and nerve growth factors (NGF < 0.1%) 376 (Figure 4). An extremely low PLA₂ activity was observed in vitro for D. 377 378 angusticeps venom (Figure 5A), in agreement with previous findings [36]. The 379 proteomic analysis, however, did not identify any PLA₂ in this venom, implying that such enzyme would be present only in trace amounts. Alternatively, the very low PLA₂ 380 activity recorded for this venom may correspond to low levels of non-specific 381 382 hydrolysis of the NOBA synthetic substrate caused by other enzymes. The negligible

content of PLA₂s in *Dendroaspis* venoms contrasts with the characteristic high amounts 383 384 and activity of this enzyme in many other elapid venoms [37,38]. Also, despite the presence of 6.7% of SVMPs in the venom proteome, very low proteinase activity was 385 386 observed for *D. angusticeps* venom when using azocasein as substrate (Figure 5B). This 387 observation mirrors the negligible activity described for *D. polylepis* venom [19]. It is 388 likely that Dendroaspis SVMPs have a restricted substrate specificity, as occurs in SVMPs from other elapid venoms [39,40]. 389

- 390
- 391

3.2. Toxicity of venom fractions

Toxicity testing was performed for most venom fractions (Table 2). Using the 392 393 Toxicity Score defined by Laustsen et al. [20], a cut-off Toxicity Score value of 7, 394 below which a fraction would be deemed to not be of medical relevance for lethality, 395 was chosen for screening the fractions. From Table 2 it is evident that the vast majority 396 of the fractions did not induce lethality in mice when tested individually. Only fraction 397 8 (containing Rho-elapitoxin-Da1b and Fasciculin-2) was shown to be lethal at the 398 doses tested, with an LD₅₀ of 0.58 mg/kg (95% confidence limits: 0.17-1.23 mg/kg) and a toxicity score of 10.9. A previous study showed that an 'angusticeps-type' toxin, 399 400 which corresponds to a fasciculin, induced respiratory arrest in mice within minutes 401 after an i.v. injection of a dose of 1 mg/kg, and also caused cardiovascular alterations 402 [41]. Nevertheless the Toxicity Score of fraction 8 contrasts with the overall Toxicity 403 Score of 117.6 for the whole venom, suggesting that different toxins in *D. angusticeps* 404 venom may act in a synergistic manner, thereby potentiating each other's toxic effects, leading to higher toxicity for whole venom. To further investigate the possible 405 synergism between toxins in the venom, fractions 4-12 were combined in equivalent 406 407 amounts (according to mass), and the LD₅₀ was determined to be 1.36 mg/kg (95%

confidence limits: 0.96-1.66 mg/kg), corresponding to a Toxicity Score of 51.7, 408 providing further evidence for the presence of synergism. The identity of the toxins 409 410 acting synergistically is presently unknown; however, it is suggested that fasciculins and 411 dendrotoxins, and probably other synergistically acting proteins, might be involved in 412 this phenomenon. It should be kept in mind that the solvents used in RP-HPLC 413 separation, particularly acetonitrile, denature some venom components, especially SVMPs; thus, the toxicity of SVMP fractions cannot be assessed with our approach. 414 Nevertheless, elapid SVMPs are unlikely to play a key role in lethality. In support of 415 416 this, it was previously shown that the LD_{50} of D. polylepis venom was not significantly altered after incubating venom with RP-HPLC solvents [19]. 417

418 Despite its lack of α -neurotoxins, the venom of *D. angusticeps* is quite effective in killing mice rapidly after injection, as observed in our toxicity experiments with 419 420 crude venom, where the controls receiving 4 LD₅₀s of venom on average died within 10 minutes. Previous studies highlighted two main toxic activities when D. angusticeps 421 whole venom is tested in experimental systems. On various nerve-muscle preparations, 422 423 this venom augmented the responses to indirect stimulation [35], possibly due to the combined action of dendrotoxins and fasciculins. Then, prolonged exposure to higher 424 425 venom concentrations resulted in failure of muscle contraction. Additionally, the venom 426 induced hypotension in various animal models, an effect that was blocked by the muscarinic cholinergic antagonist atropine [42]. This effect could be caused by the 427 428 3FTxs, previously characterized from this venom, that act on muscarinic cholinergic 429 and adrenergic receptors [14–16]. Thus, the combined action of the various neurotoxin 430 types present in D. angusticeps venom may result in a complex series of neuromuscular 431 and cardiovascular effects, which result in effective prey immobilization in the absence of the action of α -neurotoxins. This toxicological scenario, and the existence of 432

433 synergistic effects, complicates the selection of the most relevant toxins towards which
434 antibodies should be raised in order to abrogate venom toxicity. This challenging task
435 demands the identification of the most relevant synergistic toxins.

436

437 *3.3 Immunoprofiling and neutralizing ability of antivenoms*

438 Three polyspecific antivenoms, which are distributed in sub-Saharan Africa, were investigated for their ability to neutralize *D. angusticeps* venom and their ability to 439 recognize both whole venoms and venom fractions. The SAVP antivenom showed the 440 highest neutralizing ability against D. angusticeps venom, with an ED_{50} (mg venom 441 neutralized per mL antivenom) of 4.0 mg/mL (95% confidence limits: 1.7-10.0 mg/mL). 442 443 VINS African antivenom also neutralized the lethal activity of the venom, with an ED_{50} of 2.4 mg/mL (95% confidence limits: 1.4-4.0 mg/mL). On the other hand, VINS 444 445 Central African antivenom failed to neutralize D. angusticeps venom at the lowest 446 venom/antivenom ratio tested (1.0 mg venom/mL antivenom). These results bear a 447 relationship with the fact that the venom of D. angusticeps is included in the immunization mixture for the manufacture of SAVP antivenom, whereas the two VINS 448 antivenoms do not include this venom during immunization. The two VINS products 449 do, however, include the venoms of other Dendroaspis species, according to their leaflet 450 451 information. Gel immunodiffusion tests of the three antivenoms indeed revealed that cross-reactive antigens between D. angusticeps and D. polylepis venoms exist, 452 evidenced by the SAVP antivenom, which produced the strongest precipitin bands with 453 454 identity or partial identity patterns (Figure 6). Cross-reactivity between at least some components of these two venoms would explain the neutralization obtained with the 455 VINS antivenom, despite these being produced without using D. angusticeps venom. It 456 would be relevant to perform detailed studies on the antigenic relationships of the main 457

458 toxicologically-relevant components of *Dendroaspis* venoms, such as the various types of 3FTxs and dendrotoxins, in order to have a knowledge base for selecting the venoms 459 or toxins to be used for preparing antivenoms. Interestingly, gel immunodiffusion 460 results, regarding the intensity of precipitates, showed a better correlation with the 461 462 neutralization potencies observed for the three antivenoms compared to their ELISA 463 titration curves against immobilized crude venoms, which showed only minor differences in binding among them (Figure 7). Although the SAVP antivenom displays 464 a slightly stronger binding when comparing the three antivenoms on the basis of 465 volume, differences are less evident when the antivenoms are evaluated based on their 466 protein concentrations (Figure 7). In general, solid-phase immunoassays of antivenoms 467 against crude venoms do not always predict their neutralizing efficacy, as antibodies 468 469 may bind to highly immunogenic venom components that may not have a key role in 470 toxicity.

471 To further investigate the immunorecognition patterns of the antivenoms, binding of their antibodies to the different venom fractions was measured by ELISA. 472 473 From Figure 8, it is evident that a somewhat similar recognition pattern exists for the different antivenoms. However, not only does the SAVP antivenom in general display 474 stronger binding to the venom fractions compared to the VINS antivenoms, but SAVP 475 476 antivenom also shows a much stronger binding to the fractions in the first part of the 477 chromatogram (4-10), containing the 3FTxs and the dendrotoxins. These findings, based 478 on the use of immobilized venom fractions rather than crude venoms, better agree with 479 the *in vivo* neutralization studies described above.

480 Observations performed on mice injected with mixtures of venom and 481 antivenom in the neutralization experiments revealed that, at some venom/antivenom 482 ratios, mice were protected from death, but nevertheless showed evident manifestations

of toxicity, such as reduced mobility (without paralysis) and congestion of the eyes. 483 This suggests that toxins responsible for these effects are not fully neutralized, at some 484 of the tested venom/antivenom ratios. Since these toxins may play an important role in 485 envenomings, it would be relevant to assess whether these non-lethal manifestations of 486 487 toxicity are neutralized or not in the evaluation of an antivenom. For instance, in the 488 case of SAVP antivenom, complete neutralization of lethality and of these additional manifestations was observed at a venom/antivenom ratio of 1.0 mg/mL. In contrast, at 489 490 ratios of 2 and 3 mg venom/mL antivenom, lethality was abrogated, but reduced mobility and eye congestion were present to some extent. At ratios of 4 mg venom/mL 491 antivenom and higher, lethality was not completely neutralized. Similar observations 492 493 were performed with VINS African antivenom, whereby complete neutralization of 494 lethality and the other effects was achieved at 0.5 mg venom/mL antivenom, whereas at 495 1 mg/mL the additional effects were observed, and lethality was abrogated. These 496 findings underscore the relevance of identifying the most relevant toxins in the venom 497 of D. angusticeps in order to ensure that neutralizing antibodies against them are 498 included in heterologous or recombinant antivenoms in the future.

499

500 **4.0 Concluding remarks and outlook**

The venom proteome of *D. angusticeps* was characterized by a bottom-up approach. It shows a predominance of 3FTxs and Kunitz-type proteinase inhibitors, with additional less abundant components of various protein families. A remarkable feature of this venom is the absence of α -neurotoxins, in sharp contrast with the venom of the closely related species *D. polylepis*. The toxicity analysis of RP-HPLC fractions revealed that only one fraction was lethal to mice at the doses tested, and that the lethality of whole venom was much higher than what would be expected based on the

lethality of individual fractions. This highlights the presence of synergism between 508 various venom components, such as dendrotoxins, fasciculins, and probably aminergic 509 510 3FTxs of various types. South African polyvalent antivenom and one Indian antivenom 511 were effective in the neutralization of venom lethality, in agreement with a pattern of immunorecognition of the various RP-HPLC fractions. On the basis of the synergism 512 513 observed in the overall toxicity of this venom, the development of an effective combination of recombinant neutralizing antibodies demands the identification of the 514 most relevant synergistic toxins that need to be neutralized - a task that awaits future 515 research efforts. 516

517

518 Acknowledgments

The authors thank Julián Fernández, Instituto Clodomiro Picado, for his 519 520 collaboration. We also thank the Department of Drug Design and Pharmacology, 521 University of Copenhagen, the Department of Systems Biology, Technical University of 522 Denmark, and Instituto Clodomiro Picado, Universidad de Costa Rica, for supporting 523 the research. Finally, the financial support of the following foundations is greatly acknowledged: Erik Birger Christensens Legat, Dansk Tennis Fond, Augustinus 524 525 Fonden, Knud Højgaards Fond, Oticon Fonden, Vera & Carl Johan Michaelsens legat, Frants Allings Legat, and Rudolph Als Fondet. 526

527

528 **References**

- A. Smith, Illustrations of the Zoology of South Africa- Invertebratae, in: Reptilia,
 London: Smith, Elder and Co, 1848.
- 531 [2] V.F.M. FitzSimons, Snakes of Southern Africa, London: Macdonald, 1962.
- 532 [3] P.S. Hodgson, T.M. Davidson, Biology and treatment of the mamba snakebite,
 533 Wilderness Environ. Med. 7 (1996) 133–145.
- G. V. Haagner, D.R. Morgan, The captive propagation of the Eastern green
 mamba *Dendroaspis angusticeps*, Int. Zoo Yearb. 28 (1989) 195–199.
- 536 [5] B. Branch, Bill Branch's Field Guide Snakes Reptiles Southern Africa, Ralph
 537 Curtis Publisher, 1988.
- 538 [6] S. Spawls, B. Branch, The Dangerous Snakes of Africa: Natural History, Species
 539 Directory, Venoms, and Snakebite, Blandford, 1995.
- 540 [7] World Health Organization, WHO guidelines for the production, control and
- 541 regulation of snake antivenom immunoglobulins, Geneva, World Health
- 542 Organization. (2010). Available in http://
- 543 www.who.int/bloodproducts/snake_antivenoms/ snakeantivenomguide/en/.
- 544 [8] A.L. Harvey, Presynaptic effect of toxins, Int Rev Neurobiol 32. (1990) 201–239.
- A.L. Harvey, A. L., Anderson, Dendrotoxins Snake Toxins that Block Potassium
 Channels and Facilitate Neurotransmitter release, Int. Encycl. Pharmacol. Ther.
- 547 Sect. 134. (1991) 131–164.
- 548 [10] E. Cerveñansky, C., Dajas, F., Harvey, A. L., E. Karlsson, Fasciculins,
- anticholinesterase toxins from mamba venoms: biochemistry and pharmacology,
 Int. Encycl. Pharmacol. Ther. (1991) 303–321.
- [11] A. Adem, A. Asblom, G. Johansson, P.M. Mbugua, E. Karlsson, Toxins from the
 venom of the green mamba *Dendroaspis angusticeps* that inhibit the binding of

- quinuclidinyl benzilate to muscarinic acetylcholine receptors., Biochim. Biophys.
 Acta. 968 (1988) 340–345.
- [12] R.J.R. McCleary, R.M. Kini, Non-enzymatic proteins from snake venoms: A
 gold mine of pharmacological tools and drug leads, Toxicon. 62 (2013) 56–74.
- E.G. Rowan, A.L. Harvey, Snake toxins from Mamba Venoms: Unique Tools dor
 the Physiologist, Acta Chim. Slov. 58a (2011) 689–692.
- 559 [14] D. Jerusalinsky, E. Kornisiuk, P. Alfaro, J. Quillfeldt, A. Ferreira, V.E. Rial, et
 al., Muscarinic toxins: novel pharmacological tools for the muscarinic
- 561 cholinergic system., Toxicon. 38 (2000) 747–61.
- A. Maïga, G. Mourier, L. Quinton, C. Rouget, C. Gales, C. Denis, et al., G
 protein-coupled receptors, an unexploited animal toxin targets: Exploration of
 green mamba venom for novel drug candidates active against adrenoceptors.,
 Toxicon. 59 (2012) 487–96.
- 566 [16] G. Blanchet, G. Collet, G. Mourier, N. Gilles, C. Fruchart-Gaillard, E. Marcon, et
- 567al., Polypharmacology profiles and phylogenetic analysis of three-finger toxins
- from mamba venom: case of aminergic toxins., Biochimie. 103 (2014) 109–17.
- 569 [17] A.L. Harvey, A.J. Anderson, Dendrotoxins: snake toxins that block potassium
 570 channels and facilitate neurotransmitter release., Pharmacol. Ther. 31 (1985) 33–
 571 55.
- A.L. Harvey, B. Robertson, Dendrotoxins: structure-activity relationships and
 effects on potassium ion channels., Curr. Med. Chem. 11 (2004) 3065–72.
- A.H. Laustsen, B. Lomonte, B. Lohse, J. Fernández, J.M. Gutiérrez, Unveiling
 the nature of black mamba (*Dendroaspis polylepis*) venom through venomics and
 antivenom immunoprofiling: Identification of key toxin targets for antivenom
 development., J. Proteomics. 119 (2015) 126–142.

- 578 [20] A.H. Laustsen, B. Lohse, B. Lomonte, M. Engmark, J.M. Gutiérrez, Selecting
- key toxins for focused development of elapid snake antivenoms and inhibitors
- 580 guided by a Toxicity Score., Toxicon. 104 (2015) 43–5.
- 581 doi:10.1016/j.toxicon.2015.07.334.
- J.J. Calvete, B. Lomonte, A bright future for integrative venomics., Toxicon. 107
 (2015) 159–162.
- J.M. Gutiérrez, B. Lomonte, L. Sanz, J.J. Calvete, D. Pla, Immunological profile
 of antivenoms: preclinical analysis of the efficacy of a polyspecific antivenom
 through antivenomics and neutralization assays., J. Proteomics. 105 (2014) 340–
 50.
- J.J. Calvete, Proteomic tools against the neglected pathology of snake bite
 envenoming, Expert Rev Proteomics. (2011) 739–758.
- 590 [24] B. Lomonte, W.C. Tsai, J.M. Ureña-Diaz, L. Sanz, D. Mora-Obando, E.E.
- 591 Sánchez, et al., Venomics of new world pit vipers: Genus-wide comparisons of
- venom proteomes across *Agkistrodon*, J. Proteomics. 96 (2014) 103–116.
- 593 [25] M. Holzer, S. Mackessy, An aqueous endpoint assay of snake venom
 594 phospholipase A2, Toxicon. (1996) 1149–55.
- W. Wang, C. Shih, T. Huang, A novel P-I class metalloproteinase with broad
 substrate-cleaving activity, agkislysin, from *Agkistrodon acutus* venom, Biochem
 Biophys Res Commun. (2004) 224–230.
- 598 [27] D. Finney, Statistical Methods in Biological Assay, London: Charles Griffin and
 599 Company Limited, 1971.
- [28] V. Tsetlin, Snake venom alpha-neurotoxins and other "three-finger" proteins.,
 Eur. J. Biochem. 264 (1999) 281–6.
- 602 [29] R.M. Kini, R. Doley, Structure, function and evolution of three-finger toxins:

603	Mini proteins	with multiple	e targets, Toxicon.	56 (2010)	855-867.
-----	---------------	---------------	---------------------	-----------	----------

- 604 [30] T. Endo, N. Tamiya, Current view on the structure-function relationship of postsynaptic neurotoxins from snake venoms, Pharmacol. Ther. 34 (1987) 403-
- 51. http://www.ncbi.nlm.nih.gov/pubmed/3324114 (accessed January 5, 2016). 606
- [31] R.M. Kini, Evolution of three-finger toxins a versatile mini protein scaffold., 607 Acta Chim. Slov. 58 (2011) 693-701. 608
- [32] K. Sunagar, T.N.W. Jackson, E.A.B. Undheim, S.A. Ali, A. Antunes, B.G. Fry, 609
- Three-fingered RAVERs: Rapid Accumulation of Variations in Exposed 610
- Residues of snake venom toxins., Toxins (Basel). 5 (2013) 2172-208. 611
- E. Karlsson, P.M. Mbugua, D. Rodriguez-Ithurralde, Fasciculins, 612 [33]
- anticholinesterase toxins from the venom of the green mamba Dendroaspis 613 614 angusticeps., J. Physiol. (Paris). 79 (1984) 232-40.
- 615 [34] B.F. Utkin, Y. Sunagar, K., Jackson, T.N.W. Reeks, B. Fry, Three-finger toxins 616 (3FTxs), in: Venomous Reptiles. Their Toxins, Oxford University Press, Oxford,
- 2015: pp. 215-227. 617

605

- [35] J.C. Barrett, A.L. Harvey, Effects of the venom of the green mamba, 618
- Dendroaspis angusticeps on skeletal muscle and neuromuscular transmission, Br. 619 J. Pharmacol. 67 (1979) 199-205. 620
- [36] N.H. Tan, A. Arunmozhiarasi, G. Ponnudurai, A comparative study of the 621
- biological properties of *Dendroaspis* (mamba) snake venoms., Comp. Biochem. 622
- Physiol. C. 99 (1991) 463-466. 623
- [37] D. Petras, L. Sanz, Á. Segura, M. Herrera, M. Villalta, D. Solano, et al., Snake 624
- venomics of African spitting cobras: Toxin composition and assessment of 625
- congeneric cross-reactivity of the Pan-African EchiTAb-Plus-ICP antivenom by 626
- antivenomics and neutralization approaches, J. Proteome Res. 10 (2011) 1266-627

1280. 628

[38] J. Fernández, N. Vargas-Vargas, D. Pla, M. Sasa, P. Rey-Suárez, L. Sanz, et al., 629 Snake venomics of *Micrurus alleni* and *Micrurus mosquitensis* from the 630 Caribbean region of Costa Rica reveals two divergent compositional patterns in 631 New World elapids., Toxicon. 107 (2015) 217-33. 632 633 [39] Q.-Y. Sun, J. Bao, Purification, cloning and characterization of a metalloproteinase from Naja atra venom., Toxicon. 56 (2010) 1459-69. 634 [40] M. Ito, J. Hamako, Y. Sakurai, M. Matsumoto, Y. Fujimura, M. Suzuki, et al., 635 Complete amino acid sequence of kaouthiagin, a novel cobra venom 636 metalloproteinase with two disintegrin-like sequences., Biochemistry. 40 (2001) 637 4503-11. 638 639 [41] C.Y. Lee, S.Y. Lee, Y.M. Chen, A study on the cause of death produced by 640 angusticeps-type toxin F7 isolated from eastern green mamba venom., Toxicon. 641 24 (1986) 33-40. O.H. Osman, M. Ismail, M.F. El-Asmar, Pharmacological studies of snake 642 [42] (Dendroaspis angusticeps) venom., Toxicon. 11 (1973) 185–92. 643 [43] UniProtKB - P00982 (VKTHD DENAN), (n.d.). 644 http://www.uniprot.org/uniprot/P00982 (accessed October 1, 2015). 645 [44] UniProtKB - P00980 (VKTHA DENAN), (n.d.). 646 http://www.uniprot.org/uniprot/P00980 (accessed October 1, 2015). 647 648

649 Figure legends

650

Figure 1: (A) Dendroaspis angusticeps **(B)** Distribution of *D*. angusticeps in Africa.

652

Figure 2: SDS-PAGE comparing crude venom of *Dendroaspis angusticeps* and *D. polylepis* under non-reduced (A) and reduced (B) conditions. Various amounts of each venom were separated in 15% gels and stained with Coomassie Blue G-250. Molecular mass markers (M) are labeled to the right, in kDa.

Figure 3: Separation of *Dendroaspis angusticeps* venom proteins using RP-HPLC (**A**), followed by SDS-PAGE (**B**). Two mg of venom were fractionated on a C_{18} column and eluted with an acetonitrile gradient (dashed line), as described in Methods. Further separation of protein fractions was performed by SDS-PAGE under reducing conditions. Molecular weight markers (M) are indicated in kDa. Coomassie-stained bands were excised, in-gel digested with trypsin, and subjected to MALDI-TOF/TOF analysis for assignment to protein families, as shown in Table 1.

664

Figure 4: Composition of the Dendroaspis angusticeps venom proteome according to 665 protein families (A) and three-finger toxin sub-subfamilies (B), expressed as 666 percentages of total protein content. KUN: Bovine pancreatic trypsin inhibitors/Kunitz 667 inhibitors (dendrotoxins); 3FTx: Three-finger toxins; SVMP: Metalloproteinases; 668 GAL: Galactose-binding lectins; PEP: Peptidases; HYA: Hyaluronidases; KTC: 669 Prokineticins; NGF: Nerve growth factors. CRISP: Cysteine-rich secretory proteins. 670 *Proteins in this fraction (Mambalgins) are not classified to a sub-subfamily; however 671 they are known to inhibit acid sensing ion channels. MIX: Fractions of different 672

673 members of the 3FTx family for which percentages were not determined; sub-674 subfamilies in this group include: Aminergic toxin, Antiplatelet toxin, Orphan group XI, 675 and Acid sensing ion channel inhibitor.

676

Figure 5: (A) Comparison of the phospholipase A_2 activity of 20 µg of the venoms of *Dendroaspis angusticeps, Dendroaspis polylepis,* and *Bothrops asper,* on 4-nitro-3octanoyloxybenzoic acid synthetic substrate. (B) Comparison of the proteolytic activity of 40 µg of venoms of *D. angusticeps, D. polylepis,* and *B. asper,* on azocasein substrate. Venoms from both species of *Dendroaspis* show extremely low phospholipase A_2 and proteinase activities. Each bar represents mean \pm SD of triplicates.

684

Figure 6: Gel immunodiffusion assay of antivenoms against the venoms of *Dendroaspis angusticeps* (Da) and *Dendroaspis polylepis* (Dp). Antivenoms (50 μ L) were added to the central wells, and solutions of various concentrations of venoms (30 μ L) were added to peripheral wells. (A): VINS African antivenom. (B): VINS Central Africa Antivenom. (C): SAVP antivenom. (D): *Micrurus nigrocinctus* antivenom.

690

Figure 7: ELISA titrations of antivenoms against immobilized crude venoms of *Dendroaspis angusticeps* (A and C) and *Dendroaspis polylepis* (B and D) SAVP:
SAIMR Polyvalent Snake Antivenom from South African Vaccine Producers, VINS
African: Snake Venom Antiserum (African) from VINS Bioproducts Ltd., VINS
Central Africa Snake Venom Antiserum (Central Africa) from VINS Bioproducts Ltd.
Normal horse serum was used as a negative control. Each point represents mean ± SD

697 of triplicate wells. Antivenom titrations are represented as volumetric dilutions in A and698 B, or as protein concentrations in C and D.

699

Figure 8: ELISA-based immunoprofiling of antivenoms against HPLC fractions of 700 Dendroaspis angusticeps venom. Binding of the equine antibodies to the immobilized 701 venom fractions was detected as described in Methods. Normal horse serum was used as 702 a negative control. For identification of venom fractions see Table 2. (A) SAVP: 703 SAIMR Polyvalent Snake Antivenom from South African Vaccine Producers. (B): 704 VINS African: Snake Venom Antiserum (African) from VINS Bioproducts Ltd., VINS 705 706 Central Africa Snake Venom Antiserum (Central Africa) from VINS Bioproducts Ltd. 707 Each bar represents mean \pm SD of triplicate wells.

708

709

710

.

Peak*	%	Mass (kDa) [▼]	Peptide ion		MS/MS-derived sequence **	Conf (%)	Sc	Protein family ***	Related
			m/z z	Z					
4.i	2.7	11.8	2014.9 1815.8 1045.6		AKQCLPFDYSGCGGNANR QCLPFDYSGCGGNANR KIPSFYYK	99 99 98.5	13 15 6	BPT1/k	KUN
4. ii			1501.7		ASIPAFYYNWAAK	66	14	BPTI/	KUN
5a	0.5	16.1	1030.5 1143.6 1709.8 1356.5		PAFYYNQK IPAFYYNQK CYDKIPAFYYNQK FDWSGCGGNSNR	66 66 66	7 14 12	BPTI	/KUN
5b	4.3	11.7	1030.5 1236.6 1143.6 1709.8 1356.5		PAFYYNQK LCILHR(N ^{da})PGR IPAFYYNQK CYDKIPAFYYNQK FDWSGCGGNSNR	66 66 66 66	11 10 12 12 19	BPT	I/KUN
6a.i	1.3	12.9	2039.9 2196.0		SIGGVTTEDCPAGQNVCFK SIGGVTTEDCPAGQNVCFKR	66 66	17 20	3FTx	
6a.ii			1253.6	-	MIWTYDGVIR	66	9	3FTx	
6b.i	0.9	12.1	1143.6 1709.8 1356.5		IPAFYYNQK CYDKIPAFYYNQK FDWSGCGGNSNR	66 66	11 10	BPT	T/KUN
6b.ii	trace		1253.6		MIWTYDGVIR	66	12	3FT	×
6b.iii	0.9		2196.0		SIGGVTTEDCPAGQNVCFKR	66	Ξ	3FT	×
7a	0.5	16.4	1253.6		MIWTYDGVIR	99	7	3F	Гх

sele	Tal
ecte	ble
d pe	1: A
ptid	ssig
e ioi	nme
ns fr	ent o
om	of the
in-g	e RF
el tr	-HF
ypsi	PLC
n-di	isol
gest	ated
ed p	frac
rotei	tion
aq u	s of
unds	Den
•	ıdro
	aspi
	s an
	tsn8
	icep
	s ve
	nom
	to]
	prote
	ein f
	ami
	lies l
	by ∿
	1AL
	DI-
	TOF
	-TC
)F of

10 b	10a	9c.ii	9c.i	č	46	9a	8b.ii	8b.i			8a	7c.ii	/c.1		76	7b.i	
1.6	0.6	6.6	4.4	i	32	1.8		3.4			3.0	trace	4.0			1.1	
12.9	15.7		11.0	į	12.9	15.8		11.5			13.7		12.0	2		14.0	
1304.5 1344.6 1566.7	1557.8	1216.4	1507.7	1685.8 1344.6	1557.7	1507.7	1253.6	2426.2	1685.8 1557.7	1304.3 1344.6	1517.6	1356.6	2821.2 1253.6 1409.7		1253 6	1557.7	
	1	1	-	<u> </u>		-	-	-				-		<u> </u>	_	1	
CCTSPDKCNY TMCYSHTTTSR GCGCPPGDDYLEVK	AILTNCGENSCYR	CL(E ^{dh})FTYGGCK	SIGGISTEECAAGQK	AILTNCGENSCYRK	AILTNCGENSCYR	SIGGISTEECAAGQK	MIWTYDGVIR	DTIFGITTQNCPAGQNLCFIR	AILTNCGENSCYRK AILTNCGENSCYR	TMCYSHTTTSR	GCGCPPGDDNLEVK	FDWSGCGGNSNR	MIWTYDGVIRR MIWTYDGVIRR		MIWTYDGVIR	AILTNCGENSCYR	
66 66	68.8	99	99	93.6 50.2	66	66	97.7	66	66 66	99 66	99	99	86 66 66		00	66	
9 12 18	5	11	6	4 6 i	12	8	6	13	15 17	10	7	Ξ	16 10	5	10	10	
3FTx	3FTx	BPTI/KUN	3FTx		3FTx	3FTx	3FTx	3FTx			3FTx	BPTI/KUN	3F1X		3FTv	3FTx	
Fasciculin-1 D. angusticeps; P0C1Y9	Dendrotoxin A (fragm) D. angusticeps; Q9PS08	D. angusticeps, F17090 Protease inhibitor 1 W. Aegyntia: C11C50	Synergistic-like protein	D. angusticeps; P0C1Y9	Fasciculin-1	Synergistic-like protein D. angusticens: P17696	Thrombostatin D. angusticeps; P81946	Rho-elapitoxin-Da1b D. angusticeps; P86419		D. angusuceps, FUC1ZU	Fasciculin-2	Alpha-dendrotoxin D. angusticeps; P00980	D. angusticeps; P81946	D. angusticeps; P18328	D. angusticeps; Q9PS08 Thromhostatin	Dendrotoxin A (fragm)	D. angusticeps; P81946

12c.i	12b.ii	12b.i	12a	11c.ii	11c.i	11b.iii	11b.ii	11b.i	11a.ii	11a.i	10с.іі	10c.i	
trace	3.0	trace	1.2	trace	9.0	trace	trace	6.5	trace	1.3		4.6	
10.1		11.1	15.3		10.4			11.5		15.1		10.8	
1645.9 1669.8	1441.7	1294.7 2724.4 2209.1 2296.1	1645.9	2434.1	1288.7 1281.6 2035.0	1557.8	2724.4	1288.7	1356.6	1288.7 2035.0	1253.7	1165.7 1507.8	1685.8 1557.7
	1		1	1		1	1	-	1		1		
WQPPWYCKEPVR CLPFLFSGCGGNANR	FCYHNIGMPFR	GTCCAVSLWIK VCTPVGTSGEDCHPASHKIPFSGQR MHHTCPCAPNLACVQTSPK VCTPVGTSGEDCHPASHKIPF	WQPPWYCKEPVR	GCGCPTAMWPYQTECCKGDR	EMLVAIHCCR EMLVAIHCCR GCGCPSKEMLVAIHCCR	AILTNCGENSCYR	VCTPVGTSGEDCHPASHKIPFSGQR	EMLVAIHCCR	FDWSGCGGNSNR	EMLVAIHCCR GCGCPSKEMLVAIHCCR	MIWTYDGVIR	MGPKLYDVSR SIGGISTEECAAGQK	AILTNCGENSCYRK AILTNCGENSCYR
66 66	66	99 99 99 91.8	80.2	99	66 66	66	99	99	96.7	99 99	99	66 66	66 66
11 15	11	12 13 26	S	17	14 12	6	10	11	6	10 11	7	7 13	14 16
BPTI/KUN	3FTx	КТС	BPTI/KUN	3FTx	3FTx	3FTx	KTC	3FTx	BPTI/KUN	3FTx	3FTx	3FTx	
Kunitz-type calcicludine D. angusticeps; P81658	Mambalgin-3 D. angusticeps; C0HJB0	Toxin MIT1 <i>D.polylepis</i> , P25687	Kunitz-type calcicludine D. angusticeps; P81658	Toxin S4C8 <i>D. jamesoni</i> ; P25683	Toxin F-VIII D. angusticeps; P01404	Dendrotoxin A (fragm) D. angusticeps; Q9PS08	D: ungusinceps, 101404 Toxin MIT1 D nobilaris: D25687	Toxin F-VIII	Alpha-dendrotoxin D. angusticeps; P00980	Toxin F-VIII D. angusticeps; P01404	Thrombostatin D. angusticeps; P81946	Synergistic-like protein D. angusticeps; P17696	

14b.ii 1.2		14b.i 0.1 15.4	14a 0.4 19.1	13ciii	13c.ii 2.9****	13c.i trace 15.4	13b.iii	13b.ii [.]****	13b.i trace 19.8	13a.iii	13a.ii 0.4****	13a.i trace 24.4	12c.ii 3.5	
1288.6	2034.9	1797.9 1669.8 1645.8 1164.5		1441.7	2034.9 1288.6	1669.8 1645.8	1441.7	1288.6 2034.9	1669.8 1645.8	1441.7	1288.6	1669.8	2034.9 1288.7	1797.9
•	1			-			-				1	1		1
EMLVAIHCCR	GCGCPSKEMLVAIHCCR	KCLPFLFSGCGGNANR CLPFLFSGCGGNANR WQPPWYCKEPVR WQPPWYCK	Negative	FCYHNIGMPFR	GCGCPSKEMLVAIHCCR EMLVAIHCCR	CLPFLFSGCGGNANR WQPPWYCKEPVR	FCYHNIGMPFR	EMLVAIHCCR GCGCPSKEMLVAIHCCR	CLPFLFSGCGGNANR WQPPWYCKEPVR	FCYHNIGMPFR	EMLVAIHCCR	CLPFLFSGCGGNANR	GCGCPSKEMLVAIHCCR EMLVAIHCCR	KCLPFLFSGCGGNANR
99	66	99 99 95.1		99	66 66	66 66	99	66 66	66 66	99	99	99	66 66	98.7
11	13	16 17 13 5		16	16 9	13 12	15	9 19	10 9	13	11	7	13 14	7
2 E T v	3FTx	BPTI/KUN		3FTx	3FTx	BPTI/KUN	3FTx	3FTx	BPTI/KUN	3FTx	3FTx	BPTI/KUN	3FTx	
D. angusuceps; P01404 Mambalgin-3	Toxin F-VIII	Kunitz-type calcicludine D. angusticeps; P81658	D. angusticeps; C0HJB0	Mambalgin-3	Toxin F-VIII D. angusticeps; P01404	Kunitz-type calcicludine D. angusticeps; P81658	Mambalgin-3 D. angusticeps; C0HJB0	Toxin F-VIII D. angusticeps; P01404	Kunitz-type calcicludine D. angusticeps; P81658	D. angusticeps; C0HJB0	D angusticans: B01404	Kunitz-type calcicludine	Toxin F-VIII D. angusticeps; P01404	

16c.iii	16c.ii	16c.i	16b.iii	16b.ii	16b.i		16a	15c.iii	15c.ii	15c.i	15b	15a
		1.0	trace		0.5****	0.1	trace		1.1	0.1	1.1	0.1
		16.2			18.5		21.2			15.0	18.3	23.5
1291.7	2329.2 1124.7	2358.2 2197.1 2253.0	1669.8	1288.7	996.6 2197.1	996.6 2197.1	1097.6 1781.9 1413.7	1441.7	2034.9 1288.6	1797.9 1669.8 1645.8 1010.5 1164.5		1669.8
-			-					-				1
NIWTFDNIIR	SIFGITTENCPDGQNLCFKK KWYYIVPR	SIFGITTEDCPDGQNLCFKR SIFGITTEDCPDGQNLCFK GCAATCPIPENYDSIHCCK	CLPFLFSGCGGNANR	EMLVAIHCCR	WYYIVPR YSDITWGCAATCPKPTNVR	WYYIVPR YSDITWGCAATCPKPTNVR	NPNPVPSGCR HWNSYCTTTHTFVK CRNPNPVPSGCR	FCYHNIGMPFR	GCGCPSKEMLVAIHCCR EMLVAIHCCR	KCLPFLFSGCGGNANR CLPFLFSGCGGNANR WQPPWYCKEPVR FQTIGECR WQPPWYCK	Negative	CLPFLFSGCGGNANR
97	66 66	66 66	99	99	99 97.7	98.6 65.7	99 99 80.7	66	66 66	99 99 95.4 86.5		66
8	14 9	13 17 21	6	9	10 8	6 9	6 20	13	7 15	15 18 14 5		7
3FTx	3FTx	3FTx	BPTI/KUN	3FTx	3FTx	FTx	NGF 3	3FTx	3FTx	BPTI/KUN		3FTx
Dendroaspin	Muscarinic toxin 4 D. angusticeps; Q9PSN1	Toxin AdTx1 D. angusticeps; P85092	<i>D. ungusticeps</i> , 101404 Kunitz-type calcicludine <i>D. angusticeps</i> ; P81658	Toxin F-VIII D annuticent: D01404	Muscarinic toxin 4 D. angusticeps; Q9PSN1	Muscarinic toxin 4 D. angusticeps; Q9PSN1	Uncharact.prot (frag) <i>O.hannah</i> ; V8NP13	Mambalgin-3 D. angusticeps; C0HJB0	Toxin F-VIII D. angusticeps; P01404	Kunitz-type calcicludine D. angusticeps; P81658		Kunitz-type calcicludine <i>D. angusticeps</i> ; P81658

22a.111	20a.11	25a.i	24b	24a.iii	24a.ii	24a.i	23b	20a.11	23a.i	22.iii	22.ii	22.i	21	20d	
		0.6	0.9			0.3	0.3		0.6			1.1	2.1	0.7	
		71.9	50.0			74.4	51.3		69.5			70.7	29.9	15.2	
1087.6	1852.9	1881.8	1256.7 2053.0	2053.1	1508.9	1967.0 1838.9	1256.7 2053.0	1300.0	1234.7	1508.9	2053.9	1234.7	2551.3 1193.7 1349.8 1537.7	1346.6 1319.9 1390.7	1319.9
1	1	-		-	1			-		1	-			<u> </u>	-
EHQEYLLR	TDIVSPPVCGNYFVEVG	NGHPCQNNQGYCYNR	VTL(D ^{III})LFGKWR TKPAYQFSSCSVQEHQR	TKPAYQFSSCSVQEHQR	RNPQCILNKPLR	KYIEFYVVVDNKMYR YIEFYVVVDNKMYR	VTL(D ^{III})LFGKWR TKPAYQFSSCSVQEHQR	INFLUCTION	VTLNLFGEWR	RNPQCILNKPLR	TKPAYQFSSCSVQEHQR	VTLNLFGEWR	YLYVCQYCPAGNIIGSIATPYK QIVDKHNALR QIVDKHNALRR NMLQMEWNSDAAQ	TCEENSCYKR SLPKIPLIIIGR GCGCPLTLPFLR	SLPKIPLIIIGR
99	66	99	66 66	97.7	99	96 66	66 66	99	00 66	99	99	99	99 99 62.1 99	66 66 66	99
10	15	14		7	13	6 9	13 12	13	10	11	11	12	14 7 21	10 12	16
MP	MP	MP	4 MP	MP	MP	MP	MP	IVIF	MP	MP	MP	MP	CRISP	3FTx	
SVMP atrase-A.	SVMP	SVMP-Hop-50 (Fragm) H.bungaroides R4G719	SVMP 1 <i>M. fulvius</i> ; U3EPC7	SVMP 1 SVMP 1 M. fulvius; U3EPC7	SVMP ussurin Glovdius ussuriensis 0787D9	Scutatease-1 Notechis scutatus; B5KFV7	SVMP 1 M. fulvius U3EPC7	S VINF Bothrops erythromelas; Q8UVG0	SVMP Echis coloratus; E9JG68	SVMP	Macrowpera teetina, Q20222 SVMP 1 M. fulvius U3EPC7	SVMP	CRiSP Micropechis ikaheka; A0A024AX20	Toxin C13S1C1 D. angusticeps; P18329	

26f	26e	26d.iii	26d.ii	26d.i	26c.ii	26c.i					26b	26a.iii	208.11	16 0 :::	26a.i	25c	25b.ii		25h.i	
0.1	0.1			0.3		0.6					0.2				0.2	0.5		C. H	0.2	
21.1	22.8			35.9		39.3					54.4				70.1	22.5			52.2	
1118.6	1118.7 1026.5	1234.7	1811.9	1759.9	1234.7	1811.9	2780.4	1243.7	2032.0	1904.0	2412.2	1607.9	1852.9	1881.8	2009.9		2053.0 1256.7	1110.0	11186	
-			<u> </u>	-	1	-	-	-		-	1	1	1	-	-		<u>,</u> ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	· -	_	
LVLNTFQAGR	LVLNTFQAGR WGDEQVH(K ¹⁶)	VTLNLFGEWR	YIEFYVVVDNEMYK	VYEMVNALNTMYRR	VTLNLFGEWR	YIEFYVVVDNEMYK	APMYPNEPFLVFWNAPTTQCQLR	NDQLIWLWR	KHSDSNAFLHLFPESFR	HSDSNAFLHLFPESFR	TFHGLGVIDWENWRPQWDR	GATVGLAYVGSLC(N ^{da})PK	TDIVSPPVCGNYFVEVG	NGHPCQNNQGYCYNR	NGHPCQNNQGYCYNRK	Negative	TKPAYQFSSCSVQEHQR VTL(D ⁱⁿ)LFGKWR		LVINTEOAGR	
99	99 98.3	<u>66</u>	99	66	66	66	99	99	66	66	99	66	99	99	99		98.5		66	
12	12 12	9	12	9	11	12	17	11	Ξ	12	12	15	16	14	9		8 9	, 5	10	
GAL	GAL	MP	MP	MP	MP	MP		НҮА			HYA	MP	MP		MP		MP		GAL	
Galectin (Frag) <i>O.hannah</i> ; V8NHB1	Galectin (Frag) <i>O.hannah</i> ; V8NHB1	SVMP Echis coloratus; E9JG68	SVMP-Sut-51 (Frag) Suta fasciata; R4FIX4	SVMP mocarhagin	SVMP Echis coloratus; E9JG68	SVMP-Sut-51 (Frag)	Hyaluronidase <i>O.hannah</i> ; V8P1Z9			M. fulvius; U3FYQ4	Hyaluronidase	Hemiaspis signata; R4G2W9	S V Mr Ovophis okinavensis;U3TBS9 SVMP-Hem-2 (Frao)	H.bungaroides; R4G719	SVMP-Hop-50 (Fragm)		SVMP 1 <i>M. fulvius</i> ; U3EPC7	O.hannah; V8NHB1	Galectin (Frag)	Naja atra; D5LMJ3

*** Protein family abbreviations: 3FTx: three-finger toxin; A₂; CRISP: cysteine-rich secretory protein; HYA: hyaluronidase BPTI/KUN: bovine

pancreatic trypsin inhibitor/Kunitz inhibitor; MP: Metalloproteinase; KTC: prokineticin; GAL: galactose binding galectin

they were of similar mass, however the percentage was attained to the 3FTx in calculating total venom composition. ****It was not possible to determine the specific percentages of the two proteins of the three finger toxin family using ESI in these bands, as

9	∞	7	6	S	4		Whole ve	Pe
15.9 (2:1:2 1	6.3	6.2	3.0 (1:1 m	4.8	2.7	70.3	nom 100	ık %
 BPTI/Kunitz inhibitor nix) Protease inhibitor 1 W. aegyptia; C1IC50 	3FTx Fasciculin-2 <i>D.angusticeps</i> ; P0C1Z0 Thrombostatin <i>D.angusticeps</i> ; P81946 Rho-elapitoxin-Da1b <i>D.angusticeps</i> ; P86419	3FTx Thrombostatin <i>D.angusticeps</i> ; P81946	 BPTI/Kunitz inhibitor Alpha-dendrotoxin D. angusticeps; P00980 3FTx Muscarinic toxin 2 D. angusticeps; P18328 	BPTI/Kunitz inhibitor Alpha-dendrotoxin <i>D. angusticeps</i> ; P00980	BPTI/Kunitz inhibitor Delta-dendrotoxin <i>D. angusticeps</i> ; P00982 Long epsilon-dendrotox.R55 <i>D. angusticeps</i> ; Q7LZS8	Fractions 4-12 (equal amounts according to mass)		Protein family
>2.38	0.58 (0.17-1.23)	>0.92	>0.45	>0.71	>0.40	1.36 (0.96-1.66)	0.85 (0.61-1.23)	LD ₅₀ (mg/kg) (95 % conf.)
	Fasciculin 2 >20 [33]			23 [44]	Delta- dendrotoxin: 15 [43]		1.13*	Reported LD ₅₀ (mg/kg)
4	10.9	4	4	4	4	51.7	117.6	Toxicity score ¹ % / LD ₅₀ (kg/mg)

Table 2: Lethality and Toxicity Score of RP-HPLC fractions of the venom of D. angusticeps

15	14	13	12	11	10	
2.4	1.7	2.1	7.7 (2:2:1 mix)	16.8	6.8	
3FTx Mambalgin-3 <i>D.angusticeps</i> ; C0HJB0 Toxin F-VIII <i>D.angusticeps</i> ; P01404	3FTx Mambalgin-3 <i>D.angusticeps</i> ; C0HJB0 Toxin F-VIII <i>D.angusticeps</i> ; P01404 BPTUKunitz inhibitor Kunitz-type calcicludine <i>D.angusticeps</i> ; P81658	3FTx Mambalgin-3 <i>D.angusticeps</i> ; C0HJB0 Toxin F-VIII <i>D.angusticeps</i> ; P01404	3FTx Mambalgin-3 <i>D.angusticeps</i> ; C0HJB0 Toxin F-VIII <i>D.angusticeps</i> ; P01404 BPTUKunitz inhibitor Kunitz-type calcicludine <i>D.angusticeps</i> ; P81658	3FTx Toxin F-VIII D.angusticeps; P01404	3FTx Fasciculin-1 <i>D.angusticeps</i> ; P0C1Y9 Synergistic-like protein <i>D.angusticeps</i> ; P17696 Thrombostatin <i>D.angusticeps</i> ; P81946 Dendrotoxin A (fragm) <i>D.angusticeps</i> ; Q9PS08	3FTx Fasciculin-1 <i>D.angusticeps</i> ; P0C1Y9 Synergistic-like protein <i>D.angusticeps</i> ; P17696
>0.35	>0.26	>0.64	× 	>2.52	>2.11	
4	4	4	4	<i>_</i>	<3.3	

the tab	² Mix i	CD-1 1	¹ Toxic	*: http	21	20	19	18	17	16
ole.	mice by i.v. inject ndicates that the f	mice by i.v. inject	the second s	://snakedatabase.	2.1	1.4	1.0	3.4	1.6	3.0
	raction did not contain a pure, isolated toxin, but instead a mix	on. In the case of crude venom, the % abundance was 100% action did not contain a pure, isolated toxin, but instead a mi		org/pages/LD50.php#legendAndDefinitions	CRiSP CRiSP <i>Micropechis ikaheka</i> ; A0A024AX20	3FTx Toxin C13S1C1 <i>D.angusticeps</i> ; P18329	3FTx Toxin C13S1C1 <i>D.angusticeps</i> ; P18329	3FTx Toxin AdTx1 <i>D.angusticeps</i> ; P85092	3FTx Toxin F-VIII <i>D.angusticeps</i> ; P01404 Muscarinic toxin 4 <i>D.angusticeps</i> ; Q9PSN1	3FTx Toxin F-VIII <i>D.angusticeps</i> ; P01404 Toxin AdTx1 <i>D.angusticeps</i> ; P85092 Muscarinic toxin 4 <i>D.angusticeps</i> ; Q9PSN1
	ture of 2–4 different toxins in varia		n divided by its estimated median l		>0.31	>0.21	>0.15	>0.50	>0.24	>0,44
	ble ratios indicated in		ethal dose (LD_{50}) for		4	<	<7	^	4	4





Figure 2 Click here to download high resolution image



Figure 3 Click here to download high resolution image





Figure 5 Click here to download high resolution image





Figure 7 Click here to download high resolution image



Figure 8 Click here to download high resolution image





*Conflict of Interest Click here to download Conflict of Interest: Conflict of interest statement.docx