



Danger in the reef

proteome, toxicity, and neutralization of the venom of the olive sea snake, *Aipysurus laevis*

Laustsen, Andreas Hougaard; Gutiérrez, José Maria; Redsted Rasmussen, Arne; Engmark, Mikael; Gravlund, Peter; Sanders, Kate L.; Lohse, Brian; Lomonte, Bruno

Published in:
Toxicon

Link to article, DOI:
[10.1016/j.toxicon.2015.07.008](https://doi.org/10.1016/j.toxicon.2015.07.008)

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Laustsen, A. H., Gutiérrez, J. M., Redsted Rasmussen, A., Engmark, M., Gravlund, P., Sanders, K. L., Lohse, B., & Lomonte, B. (2015). Danger in the reef: proteome, toxicity, and neutralization of the venom of the olive sea snake, *Aipysurus laevis*. *Toxicon*, 107(Part B), 187-196. <https://doi.org/10.1016/j.toxicon.2015.07.008>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 Danger in the reef: Proteome, toxicity, and neutralization of the venom of
2 the olive sea snake, *Aipysurus laevis*

3

4 Andreas H. Laustsen¹, José María Gutiérrez², Arne R. Rasmussen³, Mikael Engmark⁴,
5 Peter Gravlund⁵, Kate L. Sanders⁶, Brian Lohse¹, Bruno Lomonte²

6

7 ¹ Department of Drug Design and Pharmacology, Faculty of Health and Medical
8 Sciences, University of Copenhagen, Denmark

9 ² Instituto Clodomiro Picado, Facultad de Microbiología, Universidad de Costa Rica,
10 San José, Costa Rica

11 ³Royal Danish Academy of Fine Arts, School of Conservation, Denmark

12 ⁴Department of Systems Biology, Technical University of Denmark, Denmark

13 ⁵National Aquarium of Denmark (Den Blå Planet), Denmark

14 ⁶School of Earth & Environmental Sciences, University of Adelaide, Australia

15

16 **Running title:** *Aipysurus laevis* venom proteomics

17

18 **Keywords:** *Aipysurus laevis*; olive sea snake; snake venom; proteomics; toxicity;
19 venomics

20

21 **Address for correspondence:**

22 Dr Bruno Lomonte
23 Professor
24 Instituto Clodomiro Picado
25 Facultad de Microbiología
26 Universidad de Costa Rica
27 San José, COSTA RICA
28 bruno.lomonte@ucr.ac.cr

29

30

31 **Highlights**

32

- 33 • The venom proteome of the olive sea snake, *Aipysurus laevis*, is presented
- 34 • Most abundant venom components are phospholipases A₂ and short neurotoxins
- 35 • Lethality screening coupled to abundance estimation identified the medically
- 36 relevant toxins
- 37 • BioCSL Sea Snake Antivenom neutralizes venom lethality
- 38 • ICP Anti-Coral Antivenom cross-recognizes phospholipases A₂ from *A. laevis*
- 39 venom

40

41 **Abstract**

42 Four specimens of the olive sea snake, *Aipysurus laevis*, were collected off the
43 coast of Western Australia, and the venom proteome was characterized and
44 quantitatively estimated by RP-HPLC, SDS-PAGE, and MALDI-TOF-TOF analyses. *A.*
45 *laevis* venom is remarkably simple and consists of phospholipases A₂ (71.2%), three-
46 finger toxins (3FTx; 25.3%), cysteine-rich secretory proteins (CRISP; 2.5%), and traces
47 of a complement control module protein (CCM; 0.2%). Using a Toxicity Score, the
48 most lethal components were determined to be short neurotoxins. Whole venom had an
49 intravenous LD₅₀ of 0.07 mg/kg in mice and showed a high phospholipase A₂ activity,
50 but no proteinase activity *in vitro*. Preclinical assessment of neutralization and ELISA
51 immunoprofiling showed that BioCSL Sea Snake Antivenom was effective in cross-
52 neutralizing *A. laevis* venom with an ED₅₀ of 821 µg venom per mL antivenom, with a
53 binding preference towards short neurotoxins, due to the high degree of conservation
54 between short neurotoxins from *A. laevis* and *Enhydrina schistosa* venom. Our results
55 point towards the possibility of developing recombinant antibodies or synthetic
56 inhibitors against *A. laevis* venom due to its simplicity.

57 (176 words)

58

59 1. Introduction

60 The viviparous sea snakes are a diverse clade of more than 60 species that are
61 phylogenetically nested within the front-fanged Australo-Melanesian terrestrial elapids
62 (Hydrophiinae) (Rasmussen et al., 2011). They are highly aquatic and occupy most
63 shallow-marine habitats throughout the tropical and subtropical Indo-West Pacific, yet
64 are estimated to share a common ancestor dated at only 6–8 million years ago (Sanders
65 et al., 2008; Lukoschek et al., 2012). The amphibious sea kraits (Hydrophiinae:
66 Laticauda) represent an independently aquatic and earlier diverging lineage that is the
67 sister to terrestrial and viviparous marine hydrophiines (Keogh, 1998; Scanlon and Lee,
68 2004; Sanders et al., 2008). Two major clades are recognised within the viviparous
69 marine group: An ‘Aipysurus’ lineage comprising ten species found primarily in the
70 Australo-Papuan region, and a ‘Hydrophis’ lineage containing at least 50 species
71 distributed throughout the Indo-West Pacific (Rasmussen et al., 2011).

72 In the *Aipysurus* group, the olive sea snake, *Aipysurus laevis*, has a large
73 muscular head and is the most robustly built and longest species recorded, reaching
74 more than 170 cm in total length (Smith, 1926; Cogger, 1975). *A. laevis* has been
75 recorded from Aru Archipelago and Kai Islands (Indonesia) in the west and from the
76 northern coast of Australia and southern coast of New Guinea (Timor Sea and Arafura
77 Sea) to New Caledonia in the east (Coral Sea) (Cogger, 1975; Ineich and Rasmussen,
78 1997; Sanders et al., 2014). *A. laevis* is found in shallow marine habitats – coral reefs as
79 well as sandy, rocky, and mud-bottom habitats, and is often one of the most abundant
80 species throughout its range (Cogger, 1975; Lukoschek et al., 2007; Sanders et al.,
81 2014). It hunts primarily in crevices on the sea floor, and the following fish families
82 have been found as prey items in *A. laevis*: Acanthuridae, Apogonidae, Carangidae,
83 Clupeidae, Engraulidae, Labridae, Lutjanidae, Pempheridae, Pomacentridae, Scaridae,

84 Scorpaenidae and Serranidae (McCosker, 1975; Voris and Voris, 1983). Fish eggs,
85 crabs, shrimp and pelecypod (Limidae) have also been found in stomach content
86 (McCosker, 1975; Voris and Voris, 1983).

87 During mating season *A. laevis* is more prone to defensive attacks than at other
88 times of the year (Heatwole, 1975). However, normally *A. laevis* will ignore a diver
89 even if the diver approaches quite close (Heatwole, 1975). *A. laevis* has up to at least 5
90 mm long fangs and the venom is known for being extremely toxic (Limpus, 1978;
91 Minton, 1983; Mackessy and Tu, 1993; Greer, 1997). *A. laevis* is commonly caught as
92 by-catch, and commercial trawler fishers and recreational fishers handling nets are
93 therefore the typical bite victims of *A. laevis*.

94 The venoms of sea snakes, typically containing α -neurotoxins and
95 phospholipases A₂ (PLA₂s), are known to be generally more potent than the venoms
96 from terrestrial snakes in terms of lethality (Minton, 1983; Takasaki, 1998). In contrast
97 to the latter, however, only few studies have been focused on determining the
98 comprehensive composition of sea snake venoms by means of proteomic analyses, i.e.
99 venomics. The venom of *A. laevis* has been shown to be neurotoxic, nephrotoxic, and
100 myotoxic in mice, causing acute renal tubular degeneration, proliferative
101 glomerulonephritis, local muscle degeneration, necrosis, enlarged spleen, inflammation,
102 and lymphadenopathy (Zimmerman et al., 1992a, 1992c; Ryan and Yong, 1997, 2002).
103 Regarding the venom components of *A. laevis*, a total of four short-chain neurotoxin
104 isoforms with minor amino acid sequence variations (P19958, P19959, P19960, and
105 P32879) and one PLA₂ (P08872) have been fully sequenced (Maeda and Nobuo, 1976;
106 Ducancel et al., 1988, 1990). The short α -neurotoxins display a high affinity towards
107 the acetylcholine receptor (Ishikawa et al., 1977), which is in agreement with the very
108 low LD₅₀ observed for the whole venom (Tamiya, 1973; Maeda and Nobuo, 1976).

109 Toxicity of the venom has additionally been tested in different fish species, showing
110 variations in responses (Berman, 1983; Zimmerman et al., 1990, 1992a, 1992c). It has
111 been suggested that several components of the venom may act in a synergistic manner
112 to potentiate toxic effects (Ryan and Yong, 1997). Finally, antivenoms raised against
113 tiger snake (*Notechis scutatus*) or common sea snake (*Enhydrina schistosa*) venoms
114 have been shown to have some cross-reactivity towards the venom of *A. laevis*,
115 although the efficacies of these antivenoms are lower than against the venoms of
116 homologous species (Baxter and Gallichio, 1974).

117 Aiming to further develop understanding of sea snake venoms and to expand
118 knowledge of venom intra-species variability, this study presents the proteomic analysis
119 of the venom of *A. laevis*, together with an assessment of variability in three different
120 specimens, and of toxicity of all its main protein components in mice. In addition, the
121 ability to cross-recognize and neutralize *A. laevis* venom was evaluated for two
122 antivenoms against coral snakes and sea snakes.

123

124 **2. Materials and Methods**

125 *2.1 Snake venom*

126 *Aipysurus laevis* venom was obtained from four specimens (“Mifisto”,
127 “Medusa”, “His”, and “Nessi”) kept at the National Aquarium, Den Blå Planet,
128 Denmark. All specimens were collected at night by Kate L. Sanders from a boat using
129 spotlights and dip nets. The boat was operating at shallow water close to Broome,
130 Australia. The venom, collected in the National Aquarium of Denmark, was
131 immediately frozen, lyophilized, and kept at -20 °C. In order to assess individual
132 variability, a small sample of venom from each snake was kept separated, while the
133 remaining material was pooled.

134

135 *2.2 Venom separation by reverse-phase HPLC and SDS-PAGE*

136 The pooled venom of *A. laevis* was fractionated by sequential RP-HPLC and
137 SDS-PAGE separation steps, following the ‘snake venomics’ analytical strategy
138 (Calvete, 2011) under conditions described previously (Lomonte et al., 2014). Venom
139 load for the RP-HPLC step on C₁₈ (4.6 x 250 mm column, 5 µm particle diameter;
140 Teknokroma) was 2 mg. Protein fractions were monitored at 215 nm, manually
141 collected, dried by vacuum centrifugation, and electrophoretically separated under
142 reducing conditions. Resulting bands were stained with colloidal Coomassie blue G-
143 250, and digitally recorded on a ChemiDoc[®] imager using ImageLab[®] software (Bio-
144 Rad).

145

146 *2.3 Protein identification by tandem mass spectrometry of tryptic peptides*

147 Protein bands were excised from gels, destained with 50% acetonitrile in 25 mM
148 ammonium bicarbonate, and then subjected to reduction (10 mM dithiothreitol),
149 alkylation (50 mM iodoacetamide), and overnight in-gel digestion with sequencing
150 grade trypsin (Sigma), in 50 mM ammonium bicarbonate at 37°C. The resulting tryptic
151 peptides were extracted with 50% acetonitrile containing 1% trifluoroacetic acid (TFA),
152 and analyzed by MALDI-TOF-TOF on an AB4800-Plus Proteomics Analyzer (Applied
153 Biosystems), under conditions previously described (Lomonte et al., 2014). In each run,
154 CalMix[®] standards (ABSciex) spotted onto the same plate were used as external
155 calibrants. Resulting spectra were searched against the UniProt/SwissProt database
156 using ProteinPilot[®] v.4 and the Paragon[®] algorithm (ABSciex) for protein identification
157 at ≥ 95% score confidence, or manually interpreted. Few peptide sequences with lower
158 confidence scores were manually searched using BLAST (<http://blast.ncbi.nlm.nih.gov>)

159 for protein similarity and assignment to protein families.

160

161 2.4 Relative protein abundance estimations

162 Areas of the RP-HPLC chromatographic peaks at 215 nm were integrated using
163 ChemStation[®] (Agilent) in order to estimate relative protein abundances (Calvete,
164 2011). For peaks containing several electrophoretic bands, percentage distributions were
165 assigned by densitometry, using ImageLab[®] (Bio-Rad).

166

167 2.5 Phospholipase A₂ and proteolytic enzyme activities

168 Enzymatic activities of *A. laevis* venom were tested comparatively with samples
169 obtained from other elapid snakes (*Dendroaspis polylepis*, *Naja kaouthia*; obtained
170 from Latoxan, France; and *Micrurus nigrocinctus*, obtained from Instituto Clodomiro
171 Picado) or the viperid *Bothrops asper* (Instituto Clodomiro Picado). PLA₂ activity was
172 assayed on the chromogenic 4-nitro-3-octanoyloxybenzoic acid (NOBA) synthetic
173 substrate, as described (Lomonte et al., 2015). Venoms (20 µg, dissolved in 25 µL of
174 10 mM Tris, 10 mM CaCl₂, 0.1 M NaCl, pH 8.0, buffer) were mixed with 200 µL of the
175 same buffer and 25 µL of NOBA to achieve a final substrate concentration of 0.32 mM.
176 Plates were incubated for 60 min at 37 °C, and absorbance was recorded at 405 nm in a
177 microplate reader. Proteolytic activity was determined on azocasein, according to Wang
178 et al. (2004). Venoms (40 µg, dissolved in 50 mM Tris-HCl, 0.15 M NaCl, 5 mM CaCl₂
179 buffer, pH 8.0) were added to 100 µL of azocasein (10 mg/mL in the same buffer), and
180 incubated for 90 min at 37 °C. The reaction was stopped by addition of 200 µL of 5%
181 trichloroacetic acid, and after centrifugation (5 min, 13,000 rpm), 150 µL of
182 supernatants were mixed with 100 µL of 0.5 M NaOH, and absorbance was recorded at
183 450 nm. All samples in these assays were run in triplicate wells, and controls of solvents

184 without venoms were included.

185

186 *2.7 Lethality screening*

187 Lethality assays were conducted in CD-1 mice, supplied by Instituto Clodomiro
188 Picado, following protocols approved by the Institutional Committee for the Use and
189 Care of Animals (CICUA), University of Costa Rica. The lethality of the whole venom
190 and fractions or isolated toxins was tested by intravenous (i.v.) injection in groups of
191 four mice (18–20 g body weight). Various amounts of venom or fractions/toxins were
192 dissolved in phosphate-buffered saline (PBS; 0.12 M NaCl, 0.04 M sodium phosphate
193 buffer, pH 7.2), and injected in the caudal vein, using a volume of 100 μ L. Deaths
194 occurring within 24 h were recorded, and the LD₅₀ values were calculated by probits
195 (Finney, 1971), using the BioStat[®] software (AnalySoft).

196 The toxicity of venom fractions was initially screened by selecting a dose based on
197 fraction abundance. The dose was selected to assess whether the fraction would score
198 above or below 1 according to the Toxicity Score defined by Laustsen et al. (2015a) as
199 the toxin abundance (%) divided by its LD₅₀. Fractions that were not lethal at this dose
200 (yielding a Toxicity Score <1) were considered as having insignificant toxicity, whereas
201 fractions, which did kill mice at this level, were further evaluated, and LD₅₀ values were
202 determined for them.

203

204 *2.8 Myotoxicity of phospholipases A₂*

205 A pool of all the PLA₂ fractions was prepared, and doses of 30 μ g, dissolved in
206 50 μ L PBS, were injected intramuscularly, either in the right gastrocnemius, the thigh or
207 the soleus, to groups of five mice (18-20 g). In another experiment, mice received 30 μ g
208 of the PLA₂ fractions in the soleus muscle. Injection of PBS was used for the control

209 group. Blood was collected after 3 h from the tip of the tail into heparinized capillaries.
210 Plasma creatine kinase (CK) activity was determined using an UV kinetic assay (CK-
211 Nac[®], Analyticon). After blood collection, mice were sacrificed by CO₂ inhalation and a
212 sample of muscles were obtained and immediately fixed in 10% formalin solution. After
213 routine processing, tissues were embedded in paraffin, sectioned, and stained with
214 hematoxylin-eosin for histological observation. In addition, in order to assess the acidic
215 or basic nature of the various PLA₂s of the venom, chromatographic peaks 9-18 were
216 analyzed by zone electrophoresis under native conditions, using a 1% agarose gel
217 dissolved in 0.1 M Tris, 0.3 M glycine, pH 8.6 buffer. The gel was run at 75 V for 90
218 min, and protein migration was detected by Coomassie R-250 staining.

219

220 2.9 Antivenom neutralization studies

221 Two antivenoms were used: (a) BioCSL Sea Snake Antivenom, manufactured
222 by BioCSL Limited (Melbourne, Victoria, Australia) (batch 054908201; expiry date:
223 04/2015); (b) Monospecific *Micrurus nigrocinctus* Anticoral Antivenom from Instituto
224 Clodomiro Picado (batch 5310713ACLQ, expiry date 07/2016), for comparison.
225 Mixtures containing a fixed amount of venom and several dilutions of antivenoms were
226 prepared using PBS as diluent, and incubated at 37 °C for 30 min. Controls included
227 venom incubated with PBS instead of antivenom. Aliquots of 100 µL of the solutions,
228 containing 4×LD₅₀ of venom (11.2 µg/mouse) were then injected i.v. into groups of four
229 mice (18-20 g). Deaths occurring within 24 h were recorded for assessing the
230 neutralizing capacity of antivenoms. Neutralization was expressed as the Median
231 Effective Dose (ED₅₀) of antivenom, defined as the ratio µg venom/mL antivenom at
232 which 50% of the injected mice were protected. ED₅₀s were estimated by probits, as
233 described in Section 2.7.

234

235 *2.10 Antivenom immunoprofiling by ELISA*

236 Wells in MaxiSorp plates (NUNC, Roskilde, Denmark) were coated overnight
237 with 0.6 µg of each HPLC venom fraction, dissolved in 100 µL PBS. Then, wells were
238 blocked by adding 100 µL PBS containing 2% (w:v) bovine serum albumin (BSA,
239 Sigma) at room temperature for 1 h, and washed five times with PBS. A dilution of each
240 antivenom in PBS containing 2% BSA was prepared such that the protein concentration
241 was 86 µg/mL (as measured by their absorbance at 280 nm on a NanoDrop[®] 2000c
242 instrument, Thermo Scientific), and 100 µL were added to the wells for 2 h. After five
243 washings with PBS, 100 µL of a 1:2000 dilution of rabbit anti-horse IgG (whole
244 molecule)-alkaline phosphatase conjugated antibodies (Sigma A6063, in PBS, 2% BSA)
245 was added to each well for 2 h, and then washed five times with FALC buffer (0.05 M
246 Tris, 0.15 M NaCl, 20 µM ZnCl₂, 1 mM MgCl₂, pH 7.4). Color was developed by
247 adding 100 µL of *p*-nitrophenyl phosphate (1 mg/mL in 9.7% v/v diethanolamine
248 buffer, pH 9.8), and the absorbances at 405 nm were recorded at several time intervals
249 in a microplate reader (Multiskan FC, Thermo Scientific).

250

251

252 **3.0 Results and Discussion**253 *3.1 Venomics*

254 A detailed proteomics characterization was performed on the pooled venom
255 from *A. laevis*. From 20 fractions resolved by RP-HPLC, 35 peptidic bands were
256 obtained after SDS-PAGE separation (Fig.1). By in-gel digestion and MALDI-TOF-
257 TOF analysis, 99.2% of the protein bands could be assigned to toxin families. As shown
258 in Fig.2, the predominant family of proteins in this venom corresponds to PLA₂s

259 (71.2%), followed by a significant proportion of three-finger toxins (3FTx; 25.3%). A
260 small amount of cysteine-rich secretory proteins (CRISP; 2.5%) and traces of a
261 complement control module protein (CCM; 0.2%) were also detected. These results
262 highlight the simple protein family composition of *A. laevis* venom, which essentially
263 relies on a relatively small group of PLA₂ and 3FTx isoforms to exert its trophic role.
264 Also, these findings are in agreement with the trend emerging from recent proteomic
265 studies on sea snake venoms, which have revealed that their venoms are much simpler
266 than their terrestrial elapid counterparts in terms of the number of dominant protein
267 families and diversification of isoforms, typically within the PLA₂ and 3FTx families
268 (Fry et al., 2003; Li et al., 2005). Thus far, sea snake venom proteomes have been
269 deciphered for *Hydrophis cyanocinctus* (Calvete et al., 2012) and *Pelamis platura*
270 (Lomonte et al., 2014). Similar to these, *A. laevis* venom contains few toxin families.
271 However, in contrast to the venoms of *P. platura* and *H. cyanocinctus*, where the main
272 toxin families are three-finger toxins (50% and 81% of all venom proteins, respectively)
273 followed by PLA₂s (33% and 19% of all venom proteins, respectively), *A. laevis* venom
274 displays the opposite relative venom composition, being dominated by PLA₂s (71.2%)
275 followed by 3FTxs (25.3%). Furthermore, whereas the three-finger toxins of *A. laevis*
276 venom are all short neurotoxins, *P. platura* and *H. cyanocinctus* venoms contain both
277 short and long neurotoxins. The current findings on *A. laevis* venom composition differ
278 from a previous study, where three of the short neurotoxins were reported to represent
279 22%, 33%, and 21% of the venom (76% altogether), respectively (Maeda and Tamiya,
280 1976). The reasons behind these discrepant results are difficult to determine, although
281 they may reflect possible intraspecies differences in venom composition in specimens
282 collected in different geographical locations: Maeda and Tamiya (1976) used *A. laevis*
283 collected from Ashmore Reef, which is separated from our collection localities near

284 Broome by more than 600 km of mostly unsuitable (deep water) habitat. Potentially, the
285 observed differences could also be explained by interspecific hybridization, which has
286 previously been observed for *A. laevis* and closely related *A. fuscus* on Ashmore Reef,
287 where hybrid individuals closely resemble *A. laevis* in phenotype (Sanders et al., 2014).
288 In similarity with the predominance of PLA₂s over 3FTxs herein reported for *A. laevis*
289 venom, a transcriptomic study on the venom glands of *Aipysurus eydouxii* revealed the
290 existence of as many as sixteen unique PLA₂ transcripts, in contrast to a single
291 transcript corresponding to a 3FTx (Li et al., 2005). This could suggest that both
292 *Aipysurus* species share the same venom compositional predominance. However,
293 assessment of this possibility would require a direct examination of the *A. eydouxii*
294 venom proteins, in addition to its venom gland transcripts.

295 Individual variations of toxin expression in snake venoms are not uncommon
296 (Chippaux et al., 1991). To investigate the possible individual variability in *A. laevis*,
297 samples from three specimens (“Mifisto”, “Medusa”, and “Nessi”) were compared by
298 RP-HPLC (Fig.3). This analysis revealed that some qualitative variation in toxin
299 expression was indeed present, although most fractions did not show significant
300 deviation in abundance between specimens or pooled venom.

301 Unlike several terrestrial elapids (Aird, 2002; Laustsen et al., 2015a),
302 nucleosides were not detected in *A. laevis* venom. On the other hand, its high content of
303 PLA₂s suggests that this venom might induce myotoxicity, as previously shown in
304 experimental studies (Zimmerman et al., 1992c; Ryan and Yong, 1997, 2002). Systemic
305 myotoxicity, i.e. rhabdomyolysis, with myoglobinuria characterizes envenomings by
306 some species of sea snakes in humans (Reid, 1961), and is responsible for acute kidney
307 injury. However, when a pool of PLA₂ fractions of *A. laevis* venom was tested for
308 myotoxicity in mice, only a mild effect was observed, as judged by increments in

309 plasma CK activity. Mice receiving PBS had CK activity of 215 ± 10 U/L, and mice
310 injected in the gastrocnemius, thigh, or soleus muscles with 30 μ g of the PLA₂ fraction
311 pool had plasma CK activities of 926 ± 160 U/L, 1196 ± 119 U/L, and 764 ± 182 U/L
312 (mean \pm SEM), respectively. Increments in CK were significant only when the PLA₂
313 fraction pool was injected in the gastrocnemius and thigh muscles ($p < 0.05$). Thus, *A.*
314 *laevis* PLA₂s only induced a mild myotoxic effect. In agreement, histological analysis of
315 the soleus muscle 3 h after injection of PLA₂ fraction pool showed only few scattered
316 necrotic fibers (Fig.4). These observations contrast with the prominent increment in
317 plasma CK activity described for other elapid venoms, such as that of *Micrurus*
318 *nigrocinctus* (Fernández et al., 2011). By using native zone electrophoresis, it was
319 observed that all PLA₂ fractions (peaks 9-18) migrated towards the anode, indicating
320 that they were of acidic nature (not shown). This observation could explain the low
321 myotoxic effect of the PLA₂ pool tested, since commonly PLA₂s having potent
322 myotoxic effects are of a basic nature (Montecucco et al., 2008). Our results suggest
323 that myotoxicity is unlikely to be a significant effect in envenomings by *A. laevis*.

324 In agreement with its proteomic composition showing an abundance of PLA₂s,
325 high PLA₂ activity of the venom was confirmed *in vitro* (Fig.5A), whereas no
326 proteinase activity was detected (Fig.5B), in line with the absence of these enzymes in
327 the venom proteome. Three-finger toxins were shown to represent the second major
328 group of venom proteins in terms of abundance (25.3%), and all of them were identified
329 as short neurotoxin isoforms (Table 1), previously characterized by Maeda and Tamiya
330 (1976) and Ducancel et al. (1990). These short neurotoxins have been shown to bind
331 with high affinity to nicotinic receptors at the motor end-plate of muscle fibers, leading
332 to flaccid paralysis, which may result in respiratory failure and death (Maeda and
333 Nobuo, 1976; Ducancel et al., 1990).

334 All venom fractions were examined for acute toxicity in CD1 mice, and LD₅₀
335 values were determined for most of those having a Toxicity Score below 1 (Table 2).
336 All fractions containing short neurotoxins (fractions 1-4) and some fractions containing
337 PLA₂s (fractions 5-18) induced lethality in mice, although the LD₅₀ values of the short
338 neurotoxins were 10-40 fold lower than those of the PLA₂s. Evaluated on the basis of
339 their Toxicity Score, the short neurotoxins of *A. laevis* venom are the most relevant
340 toxins to target in order to counteract the main clinical manifestations of the venom. The
341 venom of *A. laevis* is remarkably simple compared to terrestrial elapids, such as
342 *Dendroaspis polylepis* (Laustsen et al., 2015a) and *Naja kaouthia* (Laustsen et al.,
343 2015b), which display a more diverse arsenal of toxins, although also being dominated
344 by only two main toxin families.

345 The concept of a Toxicity Score for acute toxicity was presented for the first
346 time in Laustsen et al. (2015a), and this score can be used to rank the importance of the
347 individual toxins for acute toxicity in the given *in vivo* model (typically rodents). By
348 examining the difference between the Toxicity Score of whole venom and the
349 Accumulated Toxicity Score for all venom components (the sum of the Toxicity Scores
350 for all the for the individual venom components), an indication of how the toxins in
351 whole venom interact can be deduced. For a venom displaying synergism, the Toxicity
352 Score for whole venom will be higher than the sum of the Toxicity Score for the
353 individual components, since the synergistic effects between toxins will lead to an
354 increased potency of the venom. For *A. laevis* there seems to be an indication that the
355 Toxicity Score of whole venom (TS = 676) is almost the double of the Accumulated
356 Toxicity Score of the venom components (ATS = 357) (Table 2), indicating that
357 synergistic effects may exist. This observation is supported by previous studies
358 indicating the presence of synergism (Ryan and Yong, 1997), which is quite fascinating

359 given the simplicity of this venom, being dominated by only a few very similar isoforms
360 of short neurotoxins responsible for the main neurotoxic effects.

361

362 3.2 Venom neutralization and antivenom profiling

363 The ability of BioCSL Sea Snake Antivenom and ICP Anti-Coral Antivenom to
364 neutralize *A. laevis* venom was investigated in CD-1 mice. The BioCSL Sea Snake
365 Antivenom was effective in neutralizing lethality with an ED₅₀ of 821 µg venom per mL
366 antivenom (95% confidence limits: 478–1439 µg/mL), whereas no neutralization was
367 observed for ICP Anti-Coral Antivenom at a level of 100 µg venom per mL. Our
368 observations are in agreement with previous findings on the ability of BioCSLSea
369 Snake Antivenom to neutralize the neuromuscular blocking activity of *A. laevis* and
370 other sea snake venoms (Chetty et al., 2004). To further investigate the antivenoms,
371 both BioCSL Sea Snake Antivenom and ICP Anti-Coral Antivenom were profiled by
372 ELISA to determine the extent of binding between antivenom antibodies and toxins in
373 *A. laevis* venom (Fig.6). Two general trends present themselves: BioCSL Sea Snake
374 Antivenom displays significantly higher binding to fractions containing short
375 neurotoxins (fractions 1-4), whereas the ICP Anti-Coral Antivenom displays either
376 similar or even increased binding against PLA₂ containing fractions (fractions 5-18).
377 This finding further supports that the short neurotoxins are responsible for the main
378 toxic effects of *A. laevis* venom.

379 The underlying reason for the differences in binding preference between the two
380 antivenoms may be explained by the venom compositions of *Micrurus nigrocinctus* and
381 *Enhydrina schistosa*, which are used in the immunization mixtures of BioCSL Sea
382 Snake Antivenom and ICP Anti-Coral Antivenom, respectively (Fig.7A). It must be
383 noted, however, that it is not unlikely that horses hyper-immunized with several

384 different snake venoms were used for production of BioCSL Sea Snake Antivenom, and
385 that the “monovalence” of this antivenom is primarily due to the horses being boosted
386 with *E. schistosa* venom immediately before bleeding (Chetty et al., 2004; O’Leary and
387 Isbister, 2009; Herrera et al., 2014). Therefore, unexpected cross-reactivity is not an
388 unlikely event. *E. schistosa* venom has a high abundance of 3FTxs with a high degree of
389 conservation relative to the short neurotoxins found in *A. laevis* (Fig.7B), and it is
390 therefore not surprising that the BioCSL Sea Snake Antivenom has a strong preference
391 for fractions 1-4, containing short neurotoxins from *A. laevis*. In comparison, the PLA₂s
392 found in *M. nigrocinctus* venom are not more similar to the PLA₂s found in *A. laevis*
393 venom than the PLA₂ reported for *E. schistosa* venom (Fig.7C) (Fohlman and Eaker,
394 1977). However, it is speculated that the much higher abundance of PLA₂s in the
395 immunization mixture used for producing ICP Anti-Coral Antivenom in itself drives the
396 immunological response towards a stronger recognition against PLA₂s in general. Given
397 that BioCSL Sea Snake Antivenom readily cross-recognizes the neurotoxic components
398 having the highest Toxicity Scores, and since this antivenom was shown to neutralize
399 whole venom in rodents, BioCSL Sea Snake Antivenom should be useful for treating
400 human snakebite accidents inflicted by *A. laevis*.

401 The venom of *A. laevis* is remarkably simple. It could therefore be feasible to
402 develop modern antivenoms based on human(ized) monoclonal antibodies or peptide-
403 based inhibitors against this venom, since it is likely that only few antibodies are needed
404 to obtain its full neutralization. The degree of conservation, especially in the clinically
405 relevant short neurotoxins is high, and it is therefore likely that a potent, cross-reactive
406 antibody or peptide-based inhibitor capable of neutralizing all of these components can
407 be developed.

408

409 **4.0 Concluding remarks and outlook**

410 A proteomic analysis and functional study of *A. laevis* venom was carried out,
411 revealing that this venom is remarkably simple and dominated by PLA₂s (71.2% of
412 venom protein content) followed by short neurotoxins of the three-finger toxin family
413 (25.3% of venom protein content). Also, cysteine-rich secretory proteins (CRISP) and a
414 complement control module (CCM) were detected. Based on thorough toxicity testing
415 of the individual fractions obtained from whole venom, the most relevant toxins to
416 target for an effective antivenom against acute toxicity are the short neurotoxins. Based
417 on their Toxicity Scores, the toxins present in *A. laevis* venom seem to interact in a
418 slightly synergistic manner, possibly due to the short neurotoxins all targeting the
419 nicotinic receptors at the motor end-plate of muscle fibers. BioCSL Sea Snake
420 Antivenom was capable of neutralizing *A. laevis* venom in CD-1 mice when venom and
421 antivenom were preincubated and administered by i.v. injection. ELISA-based
422 immunoprofiling indicated that the BioCSL Sea Snake Antivenom has a binding
423 preference for short neurotoxins. Therefore, this antivenom should be of clinical use for
424 treating bites inflicted by *A. laevis*. Finally, given the simplicity of *A. laevis* venom, a
425 potential for developing a modern antivenom based on human(ized) monoclonal
426 antibodies or peptide-based inhibitors may be a possibility in the future.

427

428 **Acknowledgments**

429 The authors thank Dr. Ken Winkel and Dr. David Williams (University of
430 Melbourne, Australia) for kindly providing the sample of BioCSL Sea Snake
431 Antivenom. The authors further thank the Department of Drug Design and
432 Pharmacology (University of Copenhagen), the Instituto Clodomiro Picado
433 (Universidad de Costa Rica), and Den Blå Planet for supporting the research. Finally,

434 we thank the following foundations for financial support: Drug Research Academy
435 (University of Copenhagen), Novo Nordisk Fonden, Dansk Tennis Fond Oticon
436 Fonden, Knud Højgaards Fond, Rudolph Als Fondet, Henry Shaws Legat, Læge
437 Johannes Nicolai Krigsgaard of Hustru Else Krogsgaards Mindelegat for Medicinsk
438 Forskning og Medicinske Studenter ved Københavns Universitet, Lundbeckfonden,
439 Torben of Alice Frimodts Fond, Frants Allings Legat, Christian og Ottilia Brorsons
440 Rejselegat for Yngre Videnskabsmænd- og kvinder, and Fonden for Lægevidenskabens
441 Fremme.

442

443 **Ethical statement**

444 The authors declare that there are no conflicts of interest related to this study.
445 J.M. Gutiérrez and B. Lomonte work at the Instituto Clodomiro Picado (Universidad de
446 Costa Rica), where the anti-coral snake antivenom used in this study is produced.
447 Sources that provided financial support were not involved in the collection, analysis, or
448 interpretation of data, nor in writing the report and submitting it for publication.

449

450

451 **References**

- 452 Aird, S.D., 2002. Ophidian envenomation strategies and the role of purines. *Toxicon* 40,
453 335–393.
- 454 Baxter, E.H., Gallichio, H.A., 1974. Cross-neutralization by tiger snake (*Notechis*
455 *scutatus*) antivenene and sea snake (*Enhydrina schistosa*) antivenene against
456 several sea snake venoms. *Toxicon* 12, 273–278.
- 457 Berman, D.M., 1983. The toxicities of snake venoms to goldfish (*Carassius auratus*) and
458 the susceptibilities of reef fish and crabs to olive sea snake (*Aipysurus laevis*)
459 venom. *Toxicon* 21, 37–40.
- 460 Burns, G., Heatwole, H., 1998. Home range and habitat use of the olive sea snake,
461 *Aipysurus laevis*, on the Great Barrier Reef, Australia. *J. Herpetol.*, 350–358.
- 462 Calvete, J.J., Ghezellou, P., Paiva, O., Matainaho, T., Ghassempour, A., Goudarzi, H.,
463 Fred, K., Sanz, L., Williams, D.J., 2012. Snake venomomics of two poorly known
464 Hydrophiinae: comparative proteomics of the venoms of terrestrial
465 *Toxicocalamus longissimus* and marine *Hydrophis cyanocinctus*. *J. Proteomics*
466 75, 4091–4101.
- 467 Calvete, J.J., 2011. Proteomic tools against the neglected pathology of snake bite
468 envenoming. *Expert Rev. Proteomics* 8, 739–758.
- 469 Chetty, N., Du, A., Hodgson, W.C., Winkel, K.D., Fry, B.G., 2004. The *in vitro*
470 neuromuscular activity of Indo-Pacific sea snake venoms: efficacy of two
471 commercially available antivenoms. *Toxicon* 44, 193–200.
- 472 Chippaux, J.P., Williams, V., White, J., 1991. Snake venom variability: methods of
473 study, results and interpretation. *Toxicon* 29, 1279–1303.

- 474 Cogger, H.G., 1975. Sea snakes of Australia and New Guinea. In: The Biology of Sea
475 Snakes (Dunson, W.A., Ed.), pp. 59–139. University Park Press, Baltimore
476 London & Tokyo.
- 477 Ducancel, F., Guignery-Frelat, G., Boulain, J.C., Ménez, A., 1990. Nucleotide sequence
478 and structure analysis of cDNAs encoding short-chain neurotoxins from venom
479 glands of a sea snake (*Aipysurus laevis*). *Toxicon* 28, 119–123.
- 480 Ducancel, F., Guignery-Frelat, G., Bouchier, C., Ménez, A., Boulain, J.C., 1988.
481 Sequence analysis of a cDNA encoding a PLA₂ from the sea-snake *Aipysurus*
482 *laevis*. *Nucleic Acids Res.* 16, 9048–9048.
- 483 Fernández, J., Alape-Girón, A., Angulo, Y., Sanz, L., Gutiérrez, J.M., Calvete, J.J.,
484 Lomonte, B., 2011. Venomic and antivenomic analyses of the Central American
485 coral snake, *Micrurus nigrocinctus* (Elapidae). *J. Proteome Res.* 10, 1816–1827.
- 486 Fohlman, J., Eaker, D., 1977. Isolation and characterization of a lethal myotoxic
487 phospholipase A from the venom of the common sea snake *Enhydrina schistosa*
488 causing myoglobinuria in mice. *Toxicon* 15, 385–393.
- 489 Fry, B.G., Wüster, W., Ramjan, S.F.R., Jackson, T., Martelli, P., Kini, R.M., 2003.
490 Analysis of Colubroidea snake venoms by liquid chromatography with mass
491 spectrometry: evolutionary and toxinological implications. *Rapid Comm. Mass*
492 *Spec.* 17, 2047–2062.
- 493 Greer, A.E., 1997. The biology and evolution of Australian snakes. Sydney New South
494 Wales, Australia: Surrey Beatty and Sons.
- 495 Heatwole, H., 1975. Attacks by sea snakes on divers. In: W. A. Dunson (Ed.), The
496 Biology of Sea Snakes. University Park Press, Baltimore, pp. 503–516.
- 497 Heatwole, H., 1975. Sea snakes of the Gulf of Carpentaria. In: W. A. Dunson (Ed.), The
498 Biology of Sea Snakes. London & Tokyo, Baltimore, pp. 145–149.

- 499 Herrera, M., Paiva, O.K., Pagotto, A.H., Segura, Á., Serrano, S.M., Vargas, M., Villalta,
500 M., Jensen, S.D., León, G., Williams, D.J., Gutiérrez, J.M., 2014.
501 Antivenomic characterization of two antivenoms against the venom of the
502 Taipan, *Oxyuranus scutellatus*, from Papua New Guinea and Australia. *Am. J.*
503 *Trop. Med. Hyg* 91, 887–894.
- 504 Ineich, I., Rasmussen, A.R., 1997. Sea snakes from New Caledonia and the Loyalty
505 Islands (Elapidae, Laticaudinae and Hydrophiinae). *Zoosystema* 19, 185–192.
- 506 Ishikawa, Y., Menez, A., Hori, H., Yoshida, H., Tamiya, N., 1977. Structure of snake
507 toxins and their affinity to the acetylcholine receptor of fish electric organ.
508 *Toxicon* 15, 477–488.
- 509 Keogh, J.S., 1998. Molecular phylogeny of elapid snakes and a consideration of their
510 biogeographic history. *Biological Journal of the Linnean Society* 63, 177–203.
- 511 Laustsen, A.H., Lomonte, B., Lohse, B., Fernández, J., Gutiérrez, J.M., 2015a.
512 Unveiling the the nature of black mamba (*Dendroaspis polylepis*) venom
513 through venomics and antivenom immunoprofiling: identification of key toxin
514 targets for antivenom development. *J. Proteomics* 119, 126–142.
- 515 Laustsen, A.H., Gutiérrez, J.M., Lohse, B., Rasmussen, A.R., Fernández, J., Milbo, C.,
516 Lomonte, B., 2015b. Snake venomics of monocled cobra (*Naja kaouthia*) and
517 investigation of human IgG response against venom toxins. *Toxicon* 99, 23–35.
- 518 Li, M., Fry, B.G., Kini, R.M., 2005. Putting the brakes on snake venom evolution: the
519 unique molecular evolutionary patterns of *Aipysurus eydouxii* (Marbled sea snake)
520 phospholipase A₂ toxins. *Mol. Biol. Evol.* 22, 934–941.
- 521 Limpus, C.J., 1975. Coastal sea snakes of subtropical Queensland waters (230 to 280
522 south latitude). In: W. A. Dunson (Ed), *The Biology of Sea Snakes*. University Park
523 Press, London, pp. 174–182.

- 524 Limpus, C.J., 1978. The venom apparatus and venom yields of sub-tropical Queensland
525 Hydrophiidae. In: *Toxins. Animal, Plant and Microbial* Pergamon Press, Oxford,,
526 pp. 39–70.
- 527 Lomonte, B., Tsai, W.C., Ureña-Díaz, J.M., Sanz, L., Mora-Obando, D., Sánchez, E.E.,
528 Fry, B.G., Gutiérrez, J.M., Gibbs, H.L., Calvete, J.J., 2014a. Venomics of New
529 World pit vipers: genus-wide comparisons of venom proteomes across
530 Agkistrodon. *J. Proteomics* 96, 103–116.
- 531 Lomonte, B., Pla, D., Sasa, M., Tsai, W.C., Solórzano, A., Ureña-Díaz, J.M.,
532 Fernández-Montes, M.L., Mora-Obando, D., Sanz, L., Gutiérrez, J.M., Calvete, J.J.,
533 2014b. Two color morphs of the pelagic yellow-bellied sea snake, *Pelamis platura*,
534 from different locations of Costa Rica: snake venomics, toxicity, and neutralization
535 by antivenom. *J. Proteomics* 103, 137–152.
- 536 Lomonte, B., Mora-Obando, D., Fernández, J., Sanz, L., Pla, D., Gutiérrez, J.M.,
537 Calvete, J.J., 2015. First crotoxin-like phospholipase A₂ complex from a non-
538 rattlesnake venom: nigroviriditoxin, from the arboreal Central American snake
539 *Bothriechis nigroviridis*. *Toxicon* 93, 144–154.
- 540 Lukoschek, V., Keogh, J.S., Avise, J.C., 2012. Evaluating fossil calibrations for dating
541 phylogenies in light of rates of molecular evolution: a comparison of three
542 approaches. *Systematic Biol.* 61, 22-43.
- 543 Lukoschek, V., Waycott, M., Marsh, H. 2007. Phylogeography of the olive sea snake,
544 *Aipysurus laevis* (Hydrophiinae) indicates Pleistocene range expansion around
545 northern Australia but low contemporary gene flow. *Mol. Ecol.* 16, 3406–3422.
- 546 Maeda, N., Tamiya, N., 1976. Isolation, properties and amino acid sequences of three
547 neurotoxins from the venom of a sea snake, *Aipysurus laevis*. *Biochem. J.* 153,
548 79–87.

- 549 Mackessy, S.P., Tu, A.T., 1993. Biology of the sea snakes and biochemistry of their
550 venoms. In: A. T. Tu (Ed.), Toxin-related diseases: poisons originating from
551 plants, animals and spoilage. Oxford & IBH Publishing Co. PVT. LTD.,
552 England. U. K., pp. 305–351.
- 553 McCosker, J.E., 1975. Feeding behavior of Indo-Australian Hydrophiidae. In: W. A.
554 Dunson (Ed.), The Biology of Sea Snakes. London & Tokyo, Baltimore, pp.
555 217–232.
- 556 Minton, S.A., 1983. Lethal toxicity of venoms of snakes from the Coral Sea. *Toxicon*
557 21, 901–902.
- 558 Montecucco, C., Gutiérrez, J.M., Lomonte, B., 2008. Cellular pathology induced by
559 snake venom phospholipase A₂ myotoxins and neurotoxins: common aspects
560 of their mechanisms of action. *Cell. Mol. Life Sci.* 65, 2897–2912.
- 561 O'Leary, M.A., Isbister, G.K., 2009. Commercial monovalent antivenoms in Australia
562 are polyvalent. *Toxicon* 54, 192–195.
- 563 Rasmussen, A.R., 1997. Systematics of sea snakes: a critical review. *Symp. Zool. Soc.*
564 London 70, 15–30.
- 565 Rasmussen, A.R., Murphy, J.C., Ompi, M., Gibbons, J.W., Uetz, P., 2011. Marine
566 Reptiles. *PLoS One* 6, 1–12.
- 567 Reid, H.A., 1961. Myoglobinuria and sea-snake-bite poisoning. *Br. Med. J.* 1, 1284–
568 1289.
- 569 Ryan, S.E., Yong, J.L., 2002. Acute myotoxic and nephrotoxic effects of *Aipysurus*
570 *laevis* venom following intramuscular injection in mice. *Exp. Toxicol. Pathol.*
571 54, 61–67.
- 572 Ryan, S., Yong, J., 1997. The nephrotoxicity of fractionated components of *Aipysurus*
573 *laevis* venom. *Exp. Toxicol. Pathol.* 49, 47–55.

- 574 Sanders, K.L., Lee, M.S.Y., Leys, R., Foster, R., Keogh, J.S., 2008. Molecular
575 phylogeny and divergence dates for Australasian elapids and sea snakes
576 (Hydrophiinae): evidence from seven genes for rapid evolutionary radiations. *J.*
577 *Evol. Biol.* 21, 682–695.
- 578 Sanders, K.L., Rasmussen, A.R., Guinea, M., 2014. High rates of hybridisation reveal
579 fragile reproductive barriers between endangered Australian sea snakes. *Biol.*
580 *Conserv.* 171, 200–208.
- 581 Scanlon, J.D., Lee, M.S.Y., 2004. Phylogeny of Australasian venomous snakes
582 (Colubroidea, Elapidae, Hydrophiinae) based on phenotypic and molecular
583 evidence. *Zool. Scripta* 33, 335–366.
- 584 Smith, M.A., 1926. Monograph of the sea-snakes (Hydrophiidae): Printed by order of
585 the Trustees of the British museum (Natural History) London.
- 586 Takasaki, C., 1998. The toxinology of sea snake venoms. *Toxin Reviews* 17, 361–372.
- 587 Tamiya, N., 1973. Lethality of sea snake venoms. *Toxicon* 12, 85–87.
- 588 Voris, H.K., Voris, H.H., 1983. Feeding strategies in marine snakes: an analysis of
589 evolutionary, morphological, behavioral and ecological relationships. *Amer.*
590 *Zoologist* 23, 411–425.
- 591 Wang, W.J., Shih, C.H., Huang, T.F., 2004. A novel P-I class metalloproteinase with
592 broad substrate-cleaving activity, agkislysin, from *Agkistrodon acutus* venom.
593 *Biochem. Biophys. Res. Comm.* 324, 224–230.
- 594 Zimmerman, K.D., Heatwole, H., Davies, H.I., 1992. Survival times and resistance to
595 sea snake (*Aipysurus laevis*) venom by five species of prey fish. *Toxicon* 30,
596 259–264.

- 597 Zimmerman, K.D., Heatwole, H., 1992. Ventilation rates in three prey fish species
598 treated with venom of the olive sea snake, *Aipysurus laevis*. *Comp. Biochem.*
599 *Physiol.* 102C, 421–425.
- 600 Zimmerman, S.E., Heatwole, H.H., Andreopoulos, P.C., Yong, L.C.J., 1992.
601 Proliferative glomerulonephritis in mice induced by sea snake (*Aipysurus laevis*)
602 venom. *Exp. Toxicol. Pathol.* 44, 294–300.
- 603 Zimmerman, K.D., Gates, G.R., Heatwole, H., 1990. Effects of venom of the olive sea
604 snake, *Aipysurus laevis*, on the behaviour and ventilation of three species of prey
605 fish. *Toxicon* 28, 1469–1478.
- 606

607 **Figure legends**

608

609 **Figure 1:** Separation of *A. laevis* (A) venom proteins by RP-HPLC (C), followed by
610 SDS-PAGE (B). Two mg of venom were fractionated on a C₁₈ column and eluted with
611 an acetonitrile gradient (dashed line), as described in Methods. Fractions were further
612 separated by SDS-PAGE under reducing conditions. Molecular weight markers (Mw)
613 are indicated in kDa. Coomassie-stained bands were excised, in-gel digested with
614 trypsin, and subjected to MALDI-TOF/TOF analysis for assignment to protein families,
615 as shown in [Table 1](#).

616

617 **Figure 2:** Composition of *A. laevis* venom according to protein families, expressed as
618 percentages of the total protein content. 3FTx: three-finger toxin; PLA₂: phospholipase
619 A₂; CRISP: cysteine-rich secretory protein; CCM: complement control module. (see
620 [Table 1](#)).

621

622 **Figure 3:** Comparison of the chromatographic profiles of the venoms from three
623 individual *A. laevis* specimens (“Mifisto”, “Medusa”, and “Nessi”). The patterns of the
624 individual venoms are similar, although some differences do occur in abundance for
625 certain fractions. Fraction peaks representing more than 1% of total venom protein,
626 which have an abundance deviating more than 50% from the pool, are marked with *.

627

628 **Figure 4:** Light micrographs of sections of mouse soleus muscles collected 3 h after
629 injection of either phosphate-buffered saline (PBS) solution (A) or 30 µg of a pool of all
630 the PLA₂ fractions of the venom of *A. laevis* (B) (see Methods for experimental details).
631 A normal histological pattern is observed in (A), whereas few scattered necrotic muscle

632 fibers (arrows) are observed in **(B)**, thus evidencing the mild myotoxic activity of this
633 venom. Hematoxylin-eosin staining. Bar represents 100 μm .

634

635 **Figure 5:** **(A)** Comparison of the phospholipase A₂ activity between the venoms of
636 *Aipysurus laevis*, *Dendroaspis polylepis*, *Naja kaouthia*, and *Micrurus nigrocinctus*. *A.*
637 *laevis* displays high enzymatic activity, although lower than *M. nigrocinctus*. **(B)**
638 Comparison of the proteolytic activity between the venoms of *A. laevis*, *D. polylepis*, *N.*
639 *kaouthia*, *M. nigrocinctus*, and *Bpithrops asper*, evaluated on azocasein. *A. laevis* shows
640 negligible proteinase activity.

641

642 **Figure 6:** ELISA-based immunoprofiling of antivenoms (**CSL Sea Snake:** BioCSL Sea
643 Snake Antivenom from BioCSL Limited, **ICP Micrurus:** Monospecific *Micrurus*
644 *nigrocinctus* Anticoral Antivenom from Instituto Clodomiro Picado, for comparison,
645 and a negative control (**Horse negative:** normal serum from non-immunized horses
646 from Instituto Clodomiro Picado) to all fractions of *A. laevis* venom separated by RP-
647 HPLC (see Methods for details). For identification of venom fractions refer to [Table 2](#).
648 Each bar represents mean \pm SD of triplicate wells.

649

650 **Figure 7:** **(A)** Comparison between the venom profiles and similarities of toxins from
651 *Aipysurus laevis*, *Enhydrina schistosa* (*Hydrophis schistosa*), and *Micrurus*
652 *nigrocinctus*. *E. schistosa* venom is used for production of BioCSL Sea Snake
653 Antivenom, while *M. nigrocinctus* venom is used for production of ICP's Monospecific
654 *M. nigrocinctus* Anticoral Antivenom. *Venom composition is based on venomics
655 studies of *M. nigrocinctus* ([Fernández et al., 2011](#)) and the reported study of *Hydrophis*
656 *cyanocinctus* ([Calvete et al., 2012](#)), as such studies have not been performed on *E.*

657 *schistosa*. Notice that the PLA₂ content of the venoms differ. The green frames
658 highlight observations used in discussion of [Fig.6](#) (see text). **(B)** Alignment of all
659 known three-finger toxins (3FTx) from *A. laevis* with the most similar toxins from *M.*
660 *nigrocinctus* and *E. schistosa*. A high degree of conservation exists between short
661 neurotoxins from *A. laevis* and the most similar *E. schistosa* toxin, possibly explaining
662 the observed cross-reactivity of BioCSL Sea Snake Antivenom. **(C)** Alignment of the
663 single known PLA₂ from *A. laevis* with the most similar toxins from *M. nigrocinctus*
664 and *E. schistosa*, showing only a limited degree of conservation.

Table 1: Assignment of the RP-HPLC isolated fractions of *Aipysurus laevis* venom to protein families by MALDI-TOF-TOF of selected peptide ions from in-gel trypsin-digested protein bands.

Peak	%	Mass (kDa) [▼]	Peptide ion	M/S/Ms-derived sequence*	Conf (%)	Sc	Protein family**	Related protein, code	
			m/z	Z					
1	21.8	10	1564.8	1	TTTDCADDSCYBK	99	9	3FTx	short neurotoxin D
			1450.8	1	XTCCNBSSBPK	99	19		<i>Aipysurus laevis</i> , P19960
			1300.8	1	GCGCPBVBPGXK	99	18		
2a	0.3	15	1564.5	1	TTTDCADDSCYBK	99	8	3FTx	short neurotoxin D
			1450.6	1	XTCCNBSSBPK	99	14		<i>Aipysurus laevis</i> , P19960
			1300.6	1	GCGCPBVBPGXK	99	16		
2b	0.3	10	1300.6	1	GCGCPBVBPGXK	99	9	3FTx	short neurotoxin D <i>Aipysurus laevis</i> , P19960
3a	1.0	15	1436.6	1	TTTDCADDSCYK	99	12	3FTx	short neurotoxin
			1564.7	1	TTTDCADDSCYBK	99	11		<i>Aipysurus laevis</i> , P19960
			1450.7	1	XTCCNBSSBPK	99	18		
3b	1.0	10	1564.7	1	TTTDCADDSCYBK	99	9	3FTx	short neurotoxin
			1436.6	1	TTTDCADDSCYK	99	8		<i>Aipysurus laevis</i> , P19960
			1450.7	1	XTCCNBSSBPK	99	13		
4a	0.5	15	1300.7	1	GCGCPBVBPGXK	99	14		
			1300.7	1	GCGCPBVBPGXK	99	14		
			1436.6	1	TTTDCADDSCYK	65.7	6		short neurotoxin <i>Aipysurus laevis</i> , P19960
4b	0.4	10	1436.6	1	TTTDCADDSCYK	99	10	3FTx	short neurotoxin
			1450.7	1	XTCCNBSSBPK	99	14		<i>Aipysurus laevis</i> , P19960
			1300.7	1	GCGCPBVBPGXK	99	13		
5a	0.1	29	1564.7	1	TTTDCADDSCYBK	96.8	8		
			1758.8	1	NXYBFDNMXXBCANK	99	10	PLA ₂	phospholipase A ₂
			1895.8	1	AHDDCYGVAED(N ⁴⁸)GCSPK	99	26		<i>Aipysurus eydouxi</i> , ~Q5DNC9
5b	0.3	18	1774.8	1	NXYBFDN(N ⁶⁰)XBCANK	99	11		
			1758.9	1	NXYBFDNMXXBCANK	99	14	PLA ₂	phospholipase A ₂
			1870.9	1	CYCGWGGSGTPVDAXDR	99	14		<i>Aipysurus eydouxi</i> , ~Q5DNE1
			2995.5	1	ATWHYMDYGCYCGWGGSGTPVDAXDR	99	19		
			2337.1	1	MDYGCYCGWGGSGTPVDAXDR	99	19		

6	0.5	18	2091.0	1	YGCCYCGWGGSGTTPVDAXDR	99	17		
			1758.9	1	NXXYBFDNMXBCANK	99	13	PLA ₂	phospholipase A ₂
			1870.9	1	CYCGWGGSGTTPVDAXDR	99	17		<i>Aipysurus eydouxii</i> , ~Q5DNE1
			2994.5	1	ATWHYMDYGCYCGWGGSGTTPVDAXDR	99	21		
7	0.2	-	2337.1	1	MDYGCYCGWGGSGTTPVDAXDR	99	19		
			2091.0	1	YGCCYCGWGGSGTTPVDAXDR	99	21		
8	0.4	-	-	-	-	-	-	unknown	-
9	9.6	10	2091.0	1	DYGCYCGAGGSGTTPVDAXDR	99	11	PLA ₂	β -bungarotoxin chain A ₂ <i>Bungarus caeruleus</i> , ~Q8QFW3
10	6.2	10	2337.1	1	MDYGCYCGWGGSGTTPVDAXDR	99	11	PLA ₂	phospholipase A ₂
			2995.5	1	ATWHYMDYGCYCGWGGSGTTPVDAXDR	99	18		<i>Aipysurus eydouxii</i> , ~Q5DNE1
			1710.9	1	NXXYBFDN(M th)XBCANK	99	14		
			2091.0	1	YGCCYCGWGGSGTTPVDAXDR	99	17		
11a	3.1	23	1870.9	1	CYCGWGGSGTTPVDAXDR	91	9		
			1758.9	1	NXXYBFDNMXBCANK	99	14	PLA ₂	phospholipase A ₂ <i>Aipysurus eydouxii</i> , ~Q5DNE1
			2912.3	1	ATWHY(M th)DYGCYCGSGGSGTTPVDAXDR	99	12	PLA ₂	phospholipase A ₂
11b	9.1	15	1998.8	1	R	98	8		
			2896.3	1	VHDDCYGV AEDNGCYPK	95.4	8		<i>Aipysurus eydouxii</i> , ~Q5DND8
			1774.8	1	ATWHYMDYGCYCGSGGSGTTPVDAXDR	67.2	7		
11c	7.2	10	2091.0	1	YGCCYCGWGGSGTTPVDAXDR	99	10	PLA ₂	phospholipase A ₂
			2895.4	1	ATWHYMDYGCYCGSGGSGTTPVDAXDR	99	14		<i>Aipysurus eydouxii</i> , ~Q5DNE1
			1710.9	1	NXXYBFDN(M th)XBCANK	99	10		
12a	2.7	15	2896.3	1	ATWHYMDYGCYCGSGGSGTTPVDAXDR	97.3	6	PLA ₂	phospholipase A ₂
			1774.8	1	NXXYBFDN(M th)XBCANK	95.3	6		<i>Aipysurus eydouxii</i> , ~Q5DNE1
12b	3.3	10	2835.4	1	WTXYSWBCTENVPTCNSESGCBK	99	15	PLA ₂	phospholipase A ₂
			2896.4	1	ATWHYMDYGCYCGSGGSGTTPVDAXDR	99	23		<i>Aipysurus laevis</i> , P08872
			1213.6	1	CFEAAPYNNK	99	16		
			1758.9	1	NXXYBFDNMXBCANK	99	18		
13a	1.7	20	1894.8	1	AHDDCYGV AEDNGCSPK	99	23		<i>Aipysurus eydouxii</i> , ~Q5DNE1
			1758.8	1	NXXYBFDNMXBCANK	99	9	PLA ₂	phospholipase A ₂
			1213.6	1	CFEAAPYNNK	99	12		<i>Aipysurus laevis</i> , P08872
			2013.9	1	AHDDCYGV AEDNGCYPK	99	20		

13b	9.3	10	1758.9	1	NXXYBFDNMXXBCANK	99	16	PLA ₂	phospholipase A ₂
			2994.5	1	ATWHYYMDYGCYCGWGGSGTTPVDAXDR	99	22		<i>Aipysurus eydouxii</i> , ~Q5DNE1
			1870.9	1	CYCGWGGSGTTPVDAXDR	99	19		
			2337.1	1	MDYGCYCGWGGSGTTPVDAXDR	99	24		
			2091.0	1	YGCYCGWGGSGTTPVDAXDR	99	24		
14a	0.6	20	2970.3	1	ATWHYITDYGCYCGBGSSGTPVDEXDR	99	12	PLA ₂	phospholipase PLA-2
			1352.5	1	THDDCYGEAEK	99	17		<i>Notechis scutatus</i> , ~Q45Z32
14b	1.9	10	2994.4	1	ATWHYMDYGCYCGWGGSGTTPVDAXDR	99	14	PLA ₂	phospholipase A ₂
									<i>Aipysurus eydouxii</i> , ~Q5DNE1
15a	0.2	15	1970.8	1	AHDDCYGV AEDNGCYPK	99	17	PLA ₂	phospholipase A ₂
									<i>Aipysurus eydouxii</i> , ~Q5DNE1
15b	0.1	10	1970.8	1	AHDDCYGV AEDNGCYPK	99	19	PLA ₂	phospholipase A ₂
			2995.4	1	ATWHYMDYGCYCGWGGSGTTPVDAXDR	80	7		<i>Aipysurus eydouxii</i> , ~Q5DNE1
16a	0.1	15	-	-	-	-	-	unknown	-
16b	0.1	10	2050.0	1	YGCYCGSGSGTTPVDEXDR	99	12	PLA ₂	PLA ₂ -Den-12
			1774.9	1	NXXYBFDN(M ⁹³)XBCANK	92.8	8		<i>Denisonia devisi</i> , ~R4G7G2
17	3.6	10	2905.5	1	SVWDFITNYGCYCGSGSGTTPVDEXDR	99	16	PLA ₂	PLA ₂ -9
			2050.0	1	YGCYCGSGSGTTPVDEXDR	99	17		<i>Micrurus fulvius</i> , ~U3FFYN8
18	11.6	10	2970.4	1	(S ⁹⁸)VWDFITNYGCYCGSGSGTTPVDEXDR	99	16	PLA ₂	PLA ₂ -9
			1352.6	1	(T ⁶)HDDCYGEAEK	99	21		<i>Micrurus fulvius</i> , ~U3FFYN8
19a.i	0.4	37	1614.9	1	XGEEVITXGCNYGFR	99	11	CCM	complement decay-accelerating factor transmembrane isoform
									<i>Ophiophagus hannah</i> , ~V8NMM67
19a.ii		37	1777.0	1	YXXYVCBYPAGNXR	99	15	CRISP	CRISP
									<i>Hydrophis hardwickii</i> , ~AAL54918
19b	1.6	20	1161.5	1	YNNDFSNCK	99	10	CRISP	CRISP-Aca-1
			1719.8	1	YXXYVCBYPAGNXR	99	13		<i>Acanthocephalus wellsi</i> , ~R4FJD0
19c	0.4	15	1776.8	1	YXXYVCBYPAGNXR	99	20	CRISP	CRISP
									<i>Hydrophis hardwickii</i> , ~AAL54918
19d	0.3	10	1776.8	1	YXXYVCBYPAGNXR	99	15	CRISP	CRISP
									<i>Hydrophis hardwickii</i> , ~AAL54918
20	0.1	-	-	-	-	-	-	unknown	-

* Cysteine residues are carbamidomethylated. Confidence (Conf) and Score (Sc) values are calculated by the Paragon[®] algorithm of ProteinPilot[®]. ▼ : reduced SDS-PAGE mass estimations, in kDa. X: Leu/Ile; B: Lys/Gln; Z: pyroglutamate (2-oxo-pyrrolidone carboxylic acid). Possible, although unconfirmed/ambiguous amino acid modifications suggested by the automated identification software are shown in parentheses, with the following abbreviations: ^{ox} : oxidized; ^{da} : deamidated; ^{dt} : dethiomethyl; ^{pa} : propionamide; ^{fo} : formylated.

** Protein family abbreviations: 3FTx: three-finger toxin; PLA₂: phospholipase A₂; CRISP: cysteine-rich secretory protein; CCM: complement control module.

Table 2: LD₅₀ values of *Aipysurus laevis* venom and the RP-HPLC isolated fractions

Peak	%	Protein family	LD ₅₀ (95% C.L.)	Toxicity Score ¹ % / LD ₅₀ (kg/mg)
Whole venom	100		0.15 (0.08-0.25)	676
1	21.8	3FTX: short neurotoxin	0.07 (0.04-0.15)	334
2	0.6	3FTX: short neurotoxin	0.18 (0.01-0.69)	3.3
3	2.0	3FTX: short neurotoxin	0.13 (0.09-0.44)	15.3
4	0.9	3FTX: short neurotoxin	0.28 (0.14-0.99)	3.2
5	0.3	PLA ₂ : Phospholipase A ₂	>0.3	<1
6	0.5	PLA ₂ : Phospholipase A ₂	>0.5	<1
7	0.2	Unknown	>0.25	<1
8	0.4	Unknown	>0.5	<1
9	9.6	PLA ₂ : Phospholipase A ₂	>10	<1
10	6.3	PLA ₂ : Phospholipase A ₂	>7.5	<1
11	19.4	PLA ₂ : Phospholipase A ₂	>20	<1
12	6.0	PLA ₂ : Phospholipase A ₂	>6	<1

13	11.0	PLA ₂ : Phospholipase A ₂	>10	<1.1
14	2.5	PLA ₂ : Phospholipase A ₂	>2.5	<1
15	0.3	PLA ₂ : Phospholipase A ₂	>0.3	<1
16	0.2	PLA ₂ : Phospholipase A ₂	>0.25	<1
17	3.6	PLA ₂ : Phospholipase A ₂	3.05 (1.92-4.67)	1.2
18	11.6	CCM	>>2.5	<5
19	2.7	CRISP	>0.5	<1
20	0.1	CRISP	N.t.	N.t.

*<http://snakedatabase.org/pages/LD50.php#legendAndDefinitions>

¹Toxicity Score was defined as the ratio of protein fraction abundance (%) in the venom divided by its estimated median lethal dose (LD₅₀) for CD-1 mice by i.v. injection.

²Mix indicates that the fraction did not contain a pure, isolated toxin, but instead a mixture of 2-4 different toxins in variable ratios indicated in the table.

N.t. : not tested

Figure 1




Figure 2




Figure 3




Figure 4





Figure 5




Figure 6





Figure 7



Graphical Abstract

