



Prevention of foot-and-mouth disease in cattle using a prime-boot-vaccination strategy

Gullberg, Maria; Lohse, Louise; Bøtner, Anette; McInerney, Gerald; Burman, Alison; Jackson, Terry; Polacek, Charlotta; Belsham, Graham

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Gullberg, M., Lohse, L., Bøtner, A., McInerney, G., Burman, A., Jackson, T., Polacek, C., & Belsham, G. (2016). *Prevention of foot-and-mouth disease in cattle using a prime-boot-vaccination strategy*. Abstract from 19th European Study group on the molecular Biology of Picornaviruses (Europic 2016), Switzerland.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

PREVENTION OF FOOT-AND-MOUTH DISEASE IN CATTLE USING A PRIME-BOOST-VACCINATION STRATEGY

Maria Gullberg¹, Louise Lohse¹, Anette Bøtner¹, Gerald M. McInerney², Alison Burman³, Terry Jackson³, Charlotta Polacek¹ and Graham J. Belsham^{1*}

¹DTU National Veterinary Institute, Technical University of Denmark, Lindholm, Kalvehave, Denmark.

²Department of Microbiology, Tumor and Cell Biology, Karolinska Institutet, Stockholm, Sweden.

³The Pirbright Institute, Pirbright, Woking, Surrey, U.K.

Foot-and-mouth disease (FMD) is one of the most economically important infectious diseases of production animals globally. Vaccination can help to control this disease, however, current vaccines are imperfect. They are made using chemically inactivated FMD virus (FMDV) that is produced in mammalian cell culture under high containment. Here, we have expressed the FMDV capsid protein precursor (P1-2A) of strain O1 Manisa alone or with the FMDV 3C protease (3C^{pro}) using a “single cycle” packaged alphavirus self-replicating RNA based on Semliki Forest virus (SFV). When the FMDV P1-2A was expressed with 3C^{pro} then processing of the FMDV capsid precursor protein is observed within cells and the proteins assemble into empty capsid particles. In cattle vaccinated once with these rSFV-FMDV vectors alone, anti-FMDV antibodies were elicited but the immune response was insufficient to give protection against FMDV challenge. However, the prior vaccination with these vectors resulted in a much stronger immune response against FMDV post-challenge and the viremia observed was decreased in level and duration. In subsequent experiments, cattle were sequentially vaccinated with a rSFV-FMDV followed by recombinant FMDV empty

capsid particles, or *vice versa*, prior to challenge. Animals given a primary vaccination with the rSFV-FMDV vector and then boosted with FMDV empty capsids showed a strong anti-FMDV antibody response prior to challenge. Following challenge with FMDV, the cattle were protected against disease and no FMDV RNA was detected in their sera. Initial inoculation with empty capsids followed by the rSFV-FMDV was much less effective at combating the FMDV challenge and a large post-challenge boost to the level of anti-FMDV antibodies was observed and clinical disease occurred. This prime-boost system, using reagents that can be generated outside of high-containment facilities, offers significant advantages to achieve control of FMD by vaccination.