Immune gene expression in the spleen of chickens experimentally infected with *Ascaridia galli*

Dalgaard, Tina S.; Skovgaard, Kerstin; Norup, Liselotte R.; Pleidrup, Janne; Permin, Anders; Schou, Torben W.; Vadekær, Dorte Fink; Jungersen, Gregers; Juul-Madsen, Helle R.

Published in:
Veterinary Immunology and Immunopathology

Link to article, DOI:
[10.1016/j.vetimm.2015.01.003](https://doi.org/10.1016/j.vetimm.2015.01.003)

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Dalgaard, T. S., Skovgaard, K., Norup, L. R., Pleidrup, J., Permin, A., Schou, T. W., Vadekær, D. F., Jungersen, G., & Juul-Madsen, H. R. (2015). Immune gene expression in the spleen of chickens experimentally infected with *Ascaridia galli*. *Veterinary Immunology and Immunopathology*, 164(1-2), 79-86.
<https://doi.org/10.1016/j.vetimm.2015.01.003>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Immune gene expression in the spleen of chickens experimentally infected with**
2 ***Ascaridia galli***

3

4

5 Tina S. Dalgaard^{*1}, Kerstin Skovgaard², Liselotte R. Norup¹, Janne Pleidrup¹, Anders Permin³,
6 Torben W. Schou⁴, Dorte F. Vadekær², Gregers Jungersen² and Helle R. Juul-Madsen¹

7

8 ¹Department of Animal Science, Aarhus University, Blichers Alle 20, DK-8830 Tjele

9 ²National Veterinary Institute, Division of Veterinary Diagnostics and Research, Technical
10 University of Denmark, Bülow'svej 27, DK-1870 Frederiksberg C

11 ³National Food Institute, Technical University of Denmark, Mørkhøj Bygade 19, DK-2860 Søborg

12 ⁴Department of Environment and Toxicology, DHI, Agern Allé 5, DK-2970 Hørsholm

13 *Corresponding author: Tina.Dalgaard@agrsci.dk

14

15

16 **1. Introduction**

17 *Ascaridia galli* is a gastrointestinal nematode infecting chickens (Permin et al., 1999; Permin et al.,
18 1997). Substitution of traditional cages with alternative rearing systems in modern poultry
19 production has led to an increase in the prevalence of *A. galli* and recent reports from Denmark
20 and neighbouring countries show that the majority of chickens kept in free-range systems are
21 indeed infected with *A. galli* (Jansson et al., 2010; Kaufmann et al., 2011; Permin et al., 1999).
22 Infection with *A. galli* may directly contribute to economic losses due to higher feed conversion
23 rates/reduced weight gain and decreased egg production (Permin and Ranvig, 2001; Skallerup et
24 al., 2005). In severe cases, *A. galli* infections are furthermore associated with increased mortality
25 (Das et al., 2010; Gauly et al., 2005; Kilpinen et al., 2005; Permin et al., 2006), increased
26 susceptibility to secondary infections (Dahl et al., 2002; Eigaard et al., 2006; Permin et al., 2006;
27 Saif et al., 2003), impaired vaccine responses (Pleidrup et al., 2014) and even migration of worms
28 into eggs of laying hens (Fioretti et al., 2005; Reid et al., 1973). Previously, *A. galli* control has been
29 based on synthetic anthelmintics, but concerns about parasite drug resistance and left-over
30 residues in food products call for alternative disease control strategies (Sangster, 1999). An
31 attractive alternative is vaccination, but no successful *A. galli* vaccines have yet been developed.

32 Natural acquired immunity is described for avian coccidiosis, another important parasitic disease.
33 Thus, trickle immunization may induce immunity against homologous *Eimeria* challenge (Brake et
34 al., 1997; Joyner and Norton, 1973). Extensive *Eimeria* studies have been performed in order to
35 understand host protective immune responses and aid vaccine development (Lillehoj et al., 2007).
36 Natural acquired immunity against *A. galli* is less well described, but reports exist on variability in
37 disease susceptibility. The outcome of infection may e.g. be influenced by age (Idi et al., 2004;

38 Tongson and McCraw, 1967) and host genetics (Herd and McNaught, 1975; Kaufmann et al., 2011;
 39 Permin and Ranvig, 2001). Estimated heritabilities for resistance/susceptibility to *A. galli* infections
 40 suggest that selective breeding for disease resistance may be possible (Gauly et al., 2002;
 41 Kaufmann et al., 2011; Schou et al., 2003). In addition, several reports describe the presence of
 42 very small larvae (with so called arrested development) in the late stages of an *A. galli* infection
 43 and acquired immunity was suggested to be related to this phenomenon (Chamanza et al., 1999a;
 44 Ferdushy et al., 2014; Herd and McNaught, 1975). Interestingly, Herd et al. (1975) reported that
 45 the proportion of larvae with arrested development was very low in chickens treated with an
 46 immunosuppressive agent. In general, it appears that development of anti-helminthic vaccines is
 47 far more challenging than the development of vaccines directed against viral and bacterial
 48 pathogens. This is in part due to their complex life cycles and the changing host-pathogen
 49 interactions occurring during different stages of helminth infections. Thus, a detailed
 50 understanding of anti-helminth immunity is essential for future disease control.

51 The life cycle of *A. galli* is direct, starting with embryonation of shedded eggs in litter or soil. After
 52 10-20 days infective L3 stage larvae are found within the parasite eggs (Permin et al., 1997). When
 53 ingested by chickens, the *A. galli* eggs hatch within the first 24 hours either in the proventriculus or
 54 the duodenum of the host (Idi et al., 2004; Saif et al., 2003). After three to nine days the larvae
 55 enter their histotrophic phase where they move deeper into the mucosal layers of the intestine
 56 (Luna-Olivares et al., 2012; Saif et al., 2003; Tugwell and Ackert, 1952). Larvae recovery from the
 57 intestinal wall during the first week of infection was highest in the anterior part of the jejunum,
 58 but after day 7 post infection (p.i.) larvae was also found in the posterior part of the jejunum
 59 (Ferdushy et al., 2013). A high infection dose of parasite eggs may lead to a prolonged histotrophic
 60 phase, but usually young adult worms return to the intestinal lumen by day 17-30 of age during

61 which period co-existence of larvae in the intestinal wall and young worms in the intestinal
62 content is seen (Ferdushy et al., 2013; Herd and McNaught, 1975; Katakam et al., 2010). Recently,
63 Luna-Olivares et al. (2012) suggested that “mucosal phase” may be a more appropriate term than
64 “histotrophic phase” (lamina propria invasive) as the larvae may not penetrate as deep into the
65 intestinal tissue as originally thought. They reported that most larvae were observed in the lumen
66 (but in close contact with the epithelium) (63%) followed by “within epithelium” (32%) and only
67 few in the lamina propria (5%). However, only the very early time-point 3 days p.i. was
68 investigated and it is uncertain what happened later in the histotrophic/mucosal phase. However,
69 Katakam et al. (2010) was able to recover all larvae by an EDTA method, i.e. no additional larvae
70 were recovered when applying additional pepsin digestion after EDTA incubation of intestinal
71 samples taken 2 weeks p.i. indicating that lamina propria associated larvae are few also at this
72 time point.

73 The chicken spleen works as a secondary lymphoid organ where innate and adaptive immune
74 responses are efficiently mounted. It is hypothesized that the avian spleen plays an even more
75 important immunological role than in mammals as avian lymphatic vessels and lymph nodes are
76 poorly developed. The aim of this study was to investigate systemic immunological responses at
77 different stages of an *A. galli* infection by comparing gene expression profiles in spleen tissue
78 between infected and control chickens at week 2, 6 and 9 post infection (p.i.).

79

80 **2. Materials and Methods**

81 *2.1. Animals*

82 In the experiment, chickens of mixed gender from the Aarhus University L133 were used. Line 133
83 is of White Leghorn origin and contains only birds with the major histocompatibility complex
84 (MHC) haplotype B13. Water and commercial chicken feed were supplied *ad libitum*. The lighting
85 period was 12 h daily, and the chickens were kept at a temperature of 21°C. All experimental
86 chickens were produced from MHC-characterized parents, and the MHC haplotypes of the
87 offspring were confirmed by genotyping the LEI0258 microsatellite locus (McConnell et al., 1999)
88 by PCR-based fragment analysis as earlier described (Dalgaard et al., 2005). Some birds in the
89 current experiment were shared with an already published experiment (Pleidrup et al., 2014).

90 2.2. Experimental outline

91 Experimental chickens were divided into two treatment groups; 1) negative control chickens and
92 2) chickens subjected to *A. galli* infection that were kept in separate rooms of the chicken facility.
93 At 4 weeks of age, chickens in group 2 were orally infected with 1750 embryonated *A. galli* eggs
94 recovered from female worm uteri obtained from naturally infected commercial hens and
95 embryonated in H2SO4 as described in Permin et al. (1997). Sixteen animals from each group were
96 used for weekly blood sampling and seven other animals from each group were sacrificed at week
97 2, 6 and 9 p.i. for spleen collection. At week 6 and 9 p.i. faecal samples were collected before
98 sacrificing the chickens. Licence to conduct the animal experiment was obtained from the Danish
99 Ministry of Justice, Animal Experimentation Inspectorate by Helle R. Juul-Madsen. The experiment
100 was conducted according to the ethical guidelines

101 2.3. *A. galli*-specific IgG ELISA

102 Blood samples from infected animals were taken at weeks 0, 6, 7, 8, 9 p.i. and from negative
103 controls at week 0, 6, 9 p.i. and serum was used for detection of *A. galli*-specific IgG antibodies as
104 earlier described (Norup et al., 2013).

105 2.4. Faecal *A. galli* egg excretion

106 Faecal samples were obtained from *A. galli*-infected chickens before sacrificing them for spleen
107 sampling at weeks 6 and 9 p.i. Faeces was not sampled from chickens sacrificed 2 weeks p.i. as
108 adult egg secreting worms are not developed until week 5-8 p.i. (Permin and Hansen, 1998). The
109 faecal samples were examined for the presence of *A. galli* eggs using a modified McMaster
110 counting technique (Henriksen and Aagaard, 1976; Permin et al., 1997) with a detection limit of 20
111 eggs per gram faeces (EPG).

112

113 2.5. RNA extraction

114 After collection, spleens were sectioned (triangular cross-sectional slice from upper part) and
115 identical samples from each chicken were immediately placed in RNAlater (Ambion/Life
116 Technologies), kept overnight at 4°C and then at -20°C until further processing. Amounts of 7 to 15
117 mg tissue were homogenised on a TissueLyzer LT (Qiagen), and RNA isolation and DNA digestion
118 was done using the NucleoSpin 96 RNA kit (Macherey-Nagel) according to the manufacturer's
119 instructions. RNA quality was controlled on a 1 % agarose gel and the RNA concentration and
120 purity were determined using a NanoDrop spectrophotometer (Saveen and Werner AB).

121

122 2.6. cDNA synthesis and pre amplification of mRNA

cDNA synthesis and preamplification was performed as described previously (Skovgaard et al., 2013). Extracted total RNA was converted into cDNA by reverse transcription of 480 ng RNA using the QuantiTECT Reverse Transcription kit (Qiagen), cDNA was diluted 1:5 in low EDTA TE-buffer (VWR – Bie & Berntsen) prior to preamplification. Preamplification was performed using TaqMan PreAmp Master Mix (Applied Biosystems) and a 200 nM pooled primer mix was prepared combining each primer used in the present study. TaqMan PreAmp Master Mix (5 µl) was mixed with 2.5 µl 200 nM pooled primer mix and 2.5 µl diluted cDNA, and incubated at 95°C for 10 min and 16 cycles of 95°C for 15 sec and 60°C for 4 min. 16 U of Exonuclease I (New England BioLabs) was added to the preamplified cDNA, thermal cycling conditions were set to 37°C for 30 min followed by 80°C for 15 min. Preamplified cDNA was diluted 1:10 in low EDTA TE-buffer (VWR – Bie & Berntsen) before qPCR. Primers were designed using Primer3 (<http://bioinfo.ut.ee/primer3-0.4.0/>) as described in (Skovgaard et al., 2010), and purchased from Sigma-Aldrich. Primer sequences, efficiencies and amplicon length are shown in Table 1.

136

2.7. qPCR

Gene expression mRNA was analysed by quantitative real-time PCR (qPCR) performed in Dynamic Array Integrated Fluidic Circuits (Fluidigm) following the protocol described previously (Skovgaard et al., 2013). The following cycle parameter was used: 2 min at 50°C, 10 min at 95°C, followed by 35 cycles with denaturing for 15 sec at 95°C and annealing/ elongation for 1 min at 60°C. Melting curves were generated after each run to confirm a single PCR product (from 60°C to 95°C, increasing 1°C/ 3 sec). Reactions were performed in duplicates (cDNA replicates). Non template controls (NTC) were included to indicate potential problems with non-specific amplification or

145 sample contaminations. Non-reverse transcriptase controls were included to assess potential DNA
146 contamination.

147

148 Expression data (Cq values) were acquired using the Fluidigm Real-Time PCR Analysis software
149 3.0.2 (Fluidigm) and exported to GenEx (MultiD) for data pre-processing including interplate
150 correction, correction for PCR efficiency for each primer assay individually, normalising to six
151 highly stable reference genes, and averaging of cDNA technical repeats. Using GeNorm (17) and
152 NormFinder (18), glyceraldehyde-3-phosphate dehydrogenase (GAPDH), β 2 microglobulin (B2M),
153 peptidylprolyl isomerase A (PPIA), hypoxanthine phosphoribisyl transferase I (HRPT1), TATA-box
154 binding protein (TBP), and tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation
155 protein, zeta polypeptide (YWHAE) were identified as the most stably expressed reference genes
156 out of eight candidates. For each primer assay, the mean relative expression level of the control
157 group was scaled to one during data transformation $\log_2$ (Cq) to linear scale. Gene expression data
158 were $\log_2$ -transformed before testing for normal distribution, Student t test was used to analyse
159 normally distributed data, while the non-parametric test (Wilcoxon–Mann–Whitney test) was
160 used when data was non-normal distributed. Gene expression was considered significantly
161 different if the *P* value was less than 0.05 and the relative expression was greater than 2.0.
162 Experimental practice and reporting have been performed according to the Minimum Information
163 for Publication of Quantitative Real-Time PCR Experiments (MIQE) guidelines (Bustin et al., 2009).

164

165 **3. Results and Discussion**

166 According to earlier studies, week 2 p.i. represents the mucosal phase of the *A. galli* larvae
167 whereas at weeks 6 and 9 adult worms are present in the intestinal lumen. In the present

168 experiment only 43 % of the animals (data not shown) shedded *A. galli* eggs in faeces at week 6
169 p.i., and we hypothesise that the *A. galli* worms are young and have just recently started
170 producing eggs. Presumably some larvae are also still present in the mucosa at this time point as
171 earlier reported by Ferdushi et al. (2013). In contrast, 73 % of the chickens (data not shown)
172 shedded *A. galli* eggs in faeces at week 9 p.i. and with a higher mean EPG per animal than at week
173 6 (Figure 1a). Thus, this time point may represent more mature adult worms. None of the chickens
174 in the *A. galli*-free group tested EPG positive at any time-point during the experiment (data not
175 shown). Additional chickens allocated to blood sampling were sero-negative at the day of infection
176 (data not shown). Chickens in the blood sampled *A. galli*-inoculated group had seroconverted by
177 week 6 p.i. and showed positive titres of *A. galli*-specific serum IgG throughout the rest of the
178 experiment. Chickens from the blood sampled negative control group were tested at weeks 6 and
179 9 p.i. and were found to be sero-negative at both time-points (Fig. 1.b). A systemic humoral
180 immune response is reported by others as early as 2 weeks p.i., but serum titres do not appear to
181 correlate with egg excretion or worm burden (Marcos-Atxutegi et al., 2009; Norup et al., 2013;
182 Schwarz et al., 2011).

183

184 In order to understand systemic molecular response mechanisms in different stages of an *A. galli*
185 infection we studied gene expression profiles in spleen sampled 2, 6 and 9 weeks after the
186 experimental infection. Twelve genes (representing inflammatory cytokines, antimicrobial
187 peptides, acute phase proteins, soluble pattern recognition receptors and T cell signature
188 cytokines) were differentially expressed ($P < 0.05$) at at least one of the three analysed time points
189 after the *A. galli* infection compared to the control group.

190

191 3.1. (Pro-)Inflammatory cytokines

192 Only few studies have been published concerning innate immune responses towards *A. galli* in
193 chickens and focus has been on local responses in the small intestine. Thus, a single study reports
194 increased numbers of mast cells in the chicken jejunum 2 weeks post *A. galli* infection (Darmawi et
195 al., 2013). Another study reports increased numbers of presumably heterophils in the jejunum 3
196 days after an *A. galli* infection (Luna-Olivares et al., 2012). Interestingly, a genetic association
197 study indicated that chicken IFN- γ gene variants may influence *A. galli* susceptibility (Luhken et al.,
198 2011). In the present study, we analysed the expression of inflammatory cytokines in the spleen
199 (Table 2). Surprisingly, the expression of IFN- α , IL-1 β , IL-12 β and IL-18 was up regulated at week 6
200 p.i., but not at week 2 p.i. or week 9 p.i. The IL-8 expression was up regulated at week 2 as well as
201 week 6 p.i. in *A. galli*-infected chickens. Despite structural differences most avian cytokines display
202 conserved functions compared to their mammalian counterparts (Staeheli et al., 2001), and roles
203 in the chicken inflammatory response have been described for IL-8, IL-1 β , IL-18, IL-12 β (Balu and
204 Kaiser, 2003; Barker et al., 1993; Laurent et al., 2001; Schneider et al., 2000; Weining et al., 1998;
205 Withanage et al., 2004). Also chicken IFN- α (ChIFN-I) was identified to have a similar function to
206 the mammalian counterpart as a potent antiviral agent (Schultz et al., 1995; Sick et al., 1996). It is
207 now accepted that IFN- α in mammals beside its antiviral properties shows additional
208 immunomodulating effects. Although little is known of IFN- α 's role in parasite infections,
209 treatment of helminth disease in mice has been attempted with recombinant IFN- α (Godot et al.,
210 2003).

211 3.2. Antimicrobial peptides

212 The expression of DEF β 1 was significantly reduced at week 2 p.i. and significantly increased at
213 weeks 6 and 8 p.i. in spleen tissue of *A. galli*-infected chickens (Table 2). Antimicrobial peptides
214 like defensins play an important role in innate immunity, and activity directed against bacteria,
215 fungi and viruses has been reported (Ganz, 2003). Interestingly, defensins may influence adaptive
216 immune responses as they can affect the maturation of dendritic cells as well as effector T cell
217 recruitment (Yang et al., 2002). In the chicken genome, 14 beta-defensin/gallinacin genes exist and
218 the nomenclature AvBD1-14 was suggested (Lynn et al., 2007). Local expression of several of the
219 AvBD genes and their antimicrobial activity against avian enteric pathogens have been described
220 (Evans and Harmon, 1995; Harmon, 1998; Hong et al., 2012; Sugiarto and Yu, 2004). However, the
221 role of AvBD in innate immunity towards helminth infections is not clear. In humans, some beta-
222 defensins are up regulated by pro inflammatory cytokines (McDermott et al., 2006; Scott and
223 Hancock, 2000). In the present study an increased expression of DEF β 1/AvBD1 coincided with an
224 increase in the expression of pro-inflammatory cytokine genes at week 6 p.i., but not at week 9 p.i.

225 3.3. Acute phase proteins

226 Mannose binding protein (MBL) and C-reactive protein (CRP) are soluble pattern recognition
227 receptors. Few reports exist on chicken CRP, but it appears that infections with *Eimeria* spp. and
228 *Histomonas meleagridis* induce high levels of CRP (Chamanza et al., 1999). In mammals, MBL binds
229 to microbial surface carbohydrates and mediates opsonophagocytosis directly or through
230 activation of the lectin complement pathway. A conserved function of MBL in the chicken was
231 suggested as cMBL in a heterologous in vitro assay was shown to enhance human complement
232 factor 4 (C4) deposition in a calcium dependent way (Norup and Juul-Madsen, 2007). As in
233 mammals, reduced levels of serum MBL in chickens may lead to increased disease susceptibility to

234 viral and bacterial infections (Juul-Madsen et al., 2007; Schou et al., 2010). Chicken MBL is mainly
235 produced in the liver, but constitutive and inducible local expression of the gene has also been
236 reported (Hogenkamp et al., 2006; Laursen et al., 1998; Nielsen et al., 1998). In this study, MBL
237 expression was significantly increased in spleen tissue of *A. galli*-infected chickens 6 weeks p.i.
238 (Table 2). An *in vivo* function of MBL in intestinal helminth infections has not yet been determined,
239 but preliminary results suggest that faecal shedding of *A. galli* eggs is reduced in infected inbred
240 chickens with high MBL serum levels (unpublished, Norup).

241 3.4. *Th signature cytokines*

242 In mammals, Th2 polarised cells drive responses to helminth infections. Also in the chicken a Th2
243 polarised cytokine response was reported in the jejunum and spleen of *A. galli*-infected chickens 2
244 weeks p.i. (Degen et al., 2005; Kaiser, 2007; Pleidrup et al., 2014; Schwarz et al., 2011). In
245 agreement with former studies, we observed an increased expression of the Th2 signature
246 cytokine IL-13 at 2 weeks p.i. in the spleen of *A. galli*-infected chickens, but not at later stages of
247 the infection (Table 2). This time-point corresponds to the mucosal phase of the infection which
248 co-incides with influx of both $\alpha\beta$ (including CD4+ve cells) and $\gamma\delta$ T cells in the jejunal mucosa as
249 reported by others (Schwarz et al., 2011). In the present study we observed a slightly decreased
250 expression of the Th1 signature cytokine IFN- γ at week 9 p.i. in spleen tissue of *A. galli*-infected
251 chickens. This observation is in contrast to earlier findings by Degen et al. (2005) who reported
252 decreased relative cytokine mRNA ratios (infected/non-infected) for IFN- γ as early as 2 weeks p.i.
253 Earlier reports do suggest that onset and length of the larvae mucosal phase depend on infection
254 dose which differed between the two experiments.

255 3.5. *Anti-inflammatory cytokines*

256 In human and murine infections the survival strategy of helminth parasites is largely based on
257 immunoregulation by excretory-secretory (ES) products through mechanisms involving regulatory
258 T cells (Taylor et al., 2012). No Foxp3 orthologue has been identified in the chicken, but thymic
259 CD4+CD25+ T cells were characterised as counterparts of mammalian natural Tregs by production
260 of IL-10 and TGF- β (Shanmugasundaram and Selvaraj, 2011). In the present study, an increased
261 expression of TGF- β 4 was observed 6 weeks p.i. in spleen tissue of *A. galli*- infected chickens
262 (Table 2). The chicken TGF- β gene-family includes: TGF- β 2, TGF- β 3 and TGF- β 4, of which the latter
263 is the chicken orthologue of mammalian TGF- β 1 acting as an anti-inflammatory cytokine
264 (Jakowlew et al., 1997; Pan and Halper, 2003). IL-10 has a conserved function in the chicken acting
265 as an anti-inflammatory cytokine (Rothwell et al., 2004). No increased expression of IL-10 was
266 observed in the present study; instead the expression was lower in the spleen tissue of *A. galli*-
267 infected chickens 6 and 9 weeks p.i. than in controls where expression increased by age (data not
268 shown). We have no explanation for this and further studies in other inbred chicken lines as well
269 as outbred lines need to be conducted in order to elucidate if this is a general response in *A. galli*
270 infections. Further studies of the expression of anti-inflammatory cytokines may also help us to
271 understand why *A. galli* infected chickens appear to have impaired vaccine responses towards
272 third party antigens (Pleidrup et al., 2014).

273 3.6. Conclusion

274 In summary, we have investigated the avian systemic immune response to *A. galli* infection by
275 expression analyses of immune genes in spleen. Interestingly, we observed only few differentially
276 expressed genes at week 2 p.i. which corresponds to the larvae mucosal phase. In contrast, by
277 week 6 p.i. where the larvae expectedly have matured and migrated back into the intestinal

278 lumen, we observed increased expression of pro-inflammatory cytokines and acute phase
279 proteins. It is yet to be determined if the observed pro-inflammatory response is caused by *A. galli*
280 specific pathogen-associated molecular pattern molecules (PAMPs), host specific damage-
281 associated molecular pattern molecules (DAMPs) released by tissue damage, DAMP homologues
282 in parasite secretions of even by opportunistic secondary infections.

283

284 **Conflict of interest statement**

285 The authors declare to have no conflicts of interest.

286 **Acknowledgements**

287 The authors wish to acknowledge financial support from The Danish Council for Strategic
288 Research, Aarhus University and the European Union Seventh Framework Network of Animal
289 Disease Infectiology Research Facilities (NADIR; reference number FP7-228394). Pete Kaiser and
290 Lisa Rothwell are thanked for fruitful comments and support, Karin Tarp, Lene Rosborg Dal and
291 Helle Handl for excellent technical assistance, and Karin V. Østergaard for proof reading of the
292 manuscript.

293

294 **References**

- 295 Balu, S., Kaiser, P., 2003. Avian interleukin-12 β (p40): cloning and characterization of the cDNA and
296 gene. *Journal of interferon & cytokine research : the official journal of the International Society for*
297 *Interferon and Cytokine Research* 23, 699-707.
- 298 Barker, K.A., Hampe, A., Stoeckle, M.Y., Hanafusa, H., 1993. Transformation-associated cytokine 9E3/CEF4 is
299 chemotactic for chicken peripheral blood mononuclear cells. *Journal of virology* 67, 3528-3533.
- 300 Brake, D.A., Fedor, C.H., Werner, B.W., Miller, T.J., Taylor, R.L., Jr., Clare, R.A., 1997. Characterization of
301 immune response to *Eimeria tenella* antigens in a natural immunity model with hosts which differ
302 serologically at the B locus of the major histocompatibility complex. *Infection and immunity* 65, 1204-1210.
- 303 Bustin, S.A., Benes, V., Garson, J.A., Hellemans, J., Huggett, J., Kubista, M., Mueller, R., Nolan, T., Pfaffl,
304 M.W., Shipley, G.L., Vandesompele, J., Wittwer, C.T., 2009. The MIQE guidelines: minimum information for
305 publication of quantitative real-time PCR experiments. *Clinical chemistry* 55, 611-622.
- 306 Chamanza, R., van Veenm, L., Tivapasi, M.T., Toussaint, M.J.M., 1999. Acute phase proteins in the domestic
307 fowl. *World's Poultry Science Journal* 55, 61-71.
- 308 Dahl, C., Permin, A., Christensen, J.P., Bisgaard, M., Muhairwa, A.P., Petersen, K.M., Poulsen, J.S., Jensen,
309 A.L., 2002. The effect of concurrent infections with *Pasteurella multocida* and *Ascaridia galli* on free range
310 chickens. *Veterinary microbiology* 86, 313-324.
- 311 Dalgaard, T.S., Vitved, L., Skjodt, K., Thomsen, B., Labouriau, R., Jensen, K.H., Juul-Madsen, H.R., 2005.
312 Molecular characterization of major histocompatibility complex class I (B-F) mRNA variants from chickens
313 differing in resistance to Marek's disease. *Scandinavian journal of immunology* 62, 259-270.
- 314 Darmawi, D., Balqis, U., Hambal, M., Tiuria, R., Frengki, F., Priosoeryanto, B.P., 2013. Mucosal Mast Cells
315 Response in the Jejunum of *Ascaridia galli*-Infected Laying Hens. *Media Peternakan* 36, 113-119.

316 Das, G., Kaufmann, F., Abel, H., Gauly, M., 2010. Effect of extra dietary lysine in *Ascaridia galli*-infected
 317 grower layers. *Veterinary parasitology* 170, 238-243.

318 Degen, W.G., Daal, N., Rothwell, L., Kaiser, P., Schijns, V.E., 2005. Th1/Th2 polarization by viral and helminth
 319 infection in birds. *Veterinary microbiology* 105, 163-167.

320 Eigaard, N.M., Schou, T.W., Permin, A., Christensen, J.P., Ekstrom, C.T., Ambrosini, F., Cianci, D., Bisgaard,
 321 M., 2006. Infection and excretion of *Salmonella* Enteritidis in two different chicken lines with concurrent
 322 *Ascaridia galli* infection. *Avian pathology : journal of the W.V.P.A* 35, 487-493.

323 Evans, E.W., Harmon, B.G., 1995. A review of antimicrobial peptides: defensins and related cationic
 324 peptides. *Veterinary clinical pathology / American Society for Veterinary Clinical Pathology* 24, 109-116.

325 Ferdushy, T., Luna-Olivares, L.A., Nejsun, P., Roepstorff, A.K., Thamsborg, S.M., Kyvsgaard, N.C., 2013.
 326 Population dynamics of *Ascaridia galli* following single infection in young chickens. *Parasitology* 140, 1078-
 327 1084.

328 Ferdushy, T., Schou, T.W., Norup, L.R., Dalgaard, T.S., Thamsborg, S.M., Nejsun, P., Permin, A., Juul-
 329 Madsen, H.R., Kyvsgaard, N.C., 2014. Acquisition of resistance after continuous infection with *Ascaridia galli*
 330 in chickens. *Parasitology*, 1-8.

331 Fioretti, D.P., Veronesi, F., Diaferia, M., Franciosini, M.P., Proietti, P.C., 2005. *Ascaridia galli*: a report of
 332 erratic migration. *Ital J Anim Sci* 4, 310-312.

333 Ganz, T., 2003. Defensins: antimicrobial peptides of innate immunity. *Nature reviews. Immunology* 3, 710-
 334 720.

335 Gauly, M., Bauer, C., Preisinger, R., Erhardt, G., 2002. Genetic differences of *Ascaridia galli* egg output in
 336 laying hens following a single dose infection. *Veterinary parasitology* 103, 99-107.

337 Gauly, M., Homann, T., Erhardt, G., 2005. Age-related differences of *Ascaridia galli* egg output and worm
338 burden in chickens following a single dose infection. *Veterinary parasitology* 128, 141-148.

339 Godot, V., Harraga, S., Podoprigora, G., Liance, M., Bardonnnet, K., Vuitton, D.A., 2003. IFN alpha-2a protects
340 mice against a helminth infection of the liver and modulates immune responses. *Gastroenterology* 124,
341 1441-1450.

342 Harmon, B.G., 1998. Avian heterophils in inflammation and disease resistance. *Poultry science* 77, 972-977.

343 Henriksen, S.A., Aagaard, K., 1976. [A simple flotation and McMaster method (author's transl)]. *Nordisk*
344 *veterinaermedicin* 28, 392-397.

345 Herd, R.P., McNaught, D.J., 1975. Arrested development and the histotropic phase of *Ascaridia galli* in the
346 chicken. *International journal for parasitology* 5, 401-406.

347 Hogenkamp, A., van Eijk, M., van Dijk, A., van Asten, A.J., Veldhuizen, E.J., Haagsman, H.P., 2006.
348 Characterization and expression sites of newly identified chicken collectins. *Molecular immunology* 43,
349 1604-1616.

350 Hong, Y.H., Song, W., Lee, S.H., Lillehoj, H.S., 2012. Differential gene expression profiles of beta-defensins in
351 the crop, intestine, and spleen using a necrotic enteritis model in 2 commercial broiler chicken lines.
352 *Poultry science* 91, 1081-1088.

353 Idi, A., Permin, A., Murrell, K.D., 2004. Host age only partially affects resistance to primary and secondary
354 infections with *Ascaridia galli* (Schrunk, 1788) in chickens. *Veterinary parasitology* 122, 221-231.

355 Jakowlew, S.B., Mathias, A., Lillehoj, H.S., 1997. Transforming growth factor-beta isoforms in the developing
356 chicken intestine and spleen: increase in transforming growth factor-beta 4 with coccidia infection.
357 *Veterinary immunology and immunopathology* 55, 321-339.

358 Jansson, D.S., Nyman, A., Vagsholm, I., Christensson, D., Goransson, M., Fossum, O., Hoglund, J., 2010.
 359 Ascarid infections in laying hens kept in different housing systems. *Avian pathology : journal of the W.V.P.A*
 360 39, 525-532.

361 Joyner, L.P., Norton, C.C., 1973. The immunity arising from continuous low-level infection with *Eimeria*
 362 *tenella*. *Parasitology* 67, 333-340.

363 Juul-Madsen, H.R., Norup, L.R., Handberg, K.J., Jorgensen, P.H., 2007. Mannan-binding lectin (MBL) serum
 364 concentration in relation to propagation of infectious bronchitis virus (IBV) in chickens. *Viral immunology*
 365 20, 562-570.

366 Kaiser, P., 2007. The avian immune genome--a glass half-full or half-empty? *Cytogenetic and genome*
 367 *research* 117, 221-230.

368 Katakam, K.K., Nejsun, P., Kyvsgaard, N.C., Jorgensen, C.B., Thamsborg, S.M., 2010. Molecular and
 369 parasitological tools for the study of *Ascaridia galli* population dynamics in chickens. *Avian pathology : journal of the W.V.P.A* 39, 81-85.

371 Kaufmann, F., Das, G., Preisinger, R., Schmutz, M., Konig, S., Gauly, M., 2011. Genetic resistance to natural
 372 helminth infections in two chicken layer lines. *Veterinary parasitology* 176, 250-257.

373 Kilpinen, O., Roepstorff, A., Permin, A., Norgaard-Nielsen, G., Lawson, L.G., Simonsen, H.B., 2005. Influence
 374 of *Dermanyssus gallinae* and *Ascaridia galli* infections on behaviour and health of laying hens (*Gallus gallus*
 375 *domesticus*). *British poultry science* 46, 26-34.

376 Laurent, F., Mancassola, R., Lacroix, S., Menezes, R., Naciri, M., 2001. Analysis of chicken mucosal immune
 377 response to *Eimeria tenella* and *Eimeria maxima* infection by quantitative reverse transcription-PCR.
 378 *Infection and immunity* 69, 2527-2534.

379 Laursen, S.B., Hedemand, J.E., Nielsen, O.L., Thiel, S., Koch, C., Jensenius, J.C., 1998. Serum levels, ontogeny
 380 and heritability of chicken mannan-binding lectin (MBL). *Immunology* 94, 587-593.

381 Lillehoj, H.S., Kim, C.H., Keeler, C.L., Jr., Zhang, S., 2007. Immunogenomic approaches to study host
 382 immunity to enteric pathogens. *Poultry science* 86, 1491-1500.

383 Luhken, G., Gauly, M., Kaufmann, F., Erhardt, G., 2011. Association study in naturally infected helminth
 384 layers shows evidence for influence of interferon-gamma gene variants on *Ascaridia galli* worm burden.
 385 *Veterinary research* 42, 84.

386 Luna-Olivares, L.A., Ferdushy, T., Kyvsgaard, N.C., Nejsun, P., Thamsborg, S.M., Roepstorff, A., Iburg, T.M.,
 387 2012. Localization of *Ascaridia galli* larvae in the jejunum of chickens 3 days post infection. *Veterinary*
 388 *parasitology* 185, 186-193.

389 Lynn, D.J., Higgs, R., Lloyd, A.T., O'Farrelly, C., Herve-Grepinet, V., Nys, Y., Brinkman, F.S., Yu, P.L., Soulier,
 390 A., Kaiser, P., Zhang, G., Lehrer, R.I., 2007. Avian beta-defensin nomenclature: a community proposed
 391 update. *Immunology letters* 110, 86-89.

392 Marcos-Atxutegi, C., Gandolfi, B., Aranguena, T., Sepulveda, R., Arevalo, M., Simon, F., 2009. Antibody and
 393 inflammatory responses in laying hens with experimental primary infections of *Ascaridia galli*. *Veterinary*
 394 *parasitology* 161, 69-75.

395 McConnell, S.K., Dawson, D.A., Wardle, A., Burke, T., 1999. The isolation and mapping of 19 tetranucleotide
 396 microsatellite markers in the chicken. *Animal genetics* 30, 183-189.

397 McDermott, A.M., Rich, D., Cullor, J., Mannis, M.J., Smith, W., Reid, T., Murphy, C.J., 2006. The in vitro
 398 activity of selected defensins against an isolate of *Pseudomonas* in the presence of human tears. *The British*
 399 *journal of ophthalmology* 90, 609-611.

400 Nielsen, O.L., Jorgensen, P.H., Hedemand, J., Jensenius, J.C., Koch, C., Laursen, S.B., 1998.
 401 Immunohistochemical investigation of the tissue distribution of mannan-binding lectin in non-infected and
 402 virus-infected chickens. *Immunology* 94, 122-128.

403 Norup, L.R., Dalgaard, T.S., Pleidrup, J., Permin, A., Schou, T.W., Jungersen, G., Fink, D.R., Juul-Madsen, H.R.,
 404 2013. Comparison of parasite-specific immunoglobulin levels in two chicken lines during sustained infection
 405 with *Ascaridia galli*. *Veterinary parasitology* 191, 187-190.

406 Norup, L.R., Juul-Madsen, H.R., 2007. An assay for measuring the mannan-binding lectin pathway of
 407 complement activation in chickens. *Poultry science* 86, 2322-2326.

408 Pan, H., Halper, J., 2003. Cloning, expression, and characterization of chicken transforming growth factor
 409 beta 4. *Biochemical and biophysical research communications* 303, 24-30.

410 Permin, A., Bisgaard, M., Frandsen, F., Pearman, M., Kold, J., Nansen, P., 1999. Prevalence of
 411 gastrointestinal helminths in different poultry production systems. *British poultry science* 40, 439-443.

412 Permin, A., Bojesen, M., Nansen, P., Bisgaard, M., Frandsen, F., Pearman, M., 1997. *Ascaridia galli*
 413 populations in chickens following single infections with different dose levels. *Parasitology research* 83, 614-
 414 617.

415 Permin, A., Christensen, J.P., Bisgaard, M., 2006. Consequences of concurrent *Ascaridia galli* and *Escherichia*
 416 *coli* infections in chickens. *Acta veterinaria Scandinavica* 47, 43-54.

417 Permin, A., Ranvig, H., 2001. Genetic resistance to *Ascaridia galli* infections in chickens. *Veterinary*
 418 *parasitology* 102, 101-111.

419 Pleidrup, J., Dalgaard, T.S., Norup, L.R., Permin, A., Schou, T.W., Skovgaard, K., Vadekaer, D.F., Jungersen,
 420 G., Sorensen, P., Juul-Madsen, H.R., 2014. *Ascaridia galli* infection influences the development of both
 421 humoral and cell-mediated immunity after Newcastle Disease vaccination in chickens. *Vaccine* 32, 383-392.

422 Reid, W.M., Mabon, J.L., Harshbarger, W.C., 1973. Detection of worm parasites in chicken eggs by candling.
 423 Poultry science 52, 2316-2324.

424 Rothwell, L., Young, J.R., Zoorob, R., Whittaker, C.A., Hesketh, P., Archer, A., Smith, A.L., Kaiser, P., 2004.
 425 Cloning and characterization of chicken IL-10 and its role in the immune response to *Eimeria maxima*.
 426 Journal of immunology 173, 2675-2682.

427 Saif, Y.M., Barnes, H.J., Glisson, J.R., Fadly, A.M., McDougald, L.R., Swayne, D.E., 2003. Diseases of Poultry.
 428 Wiley.

429 Sangster, N.C., 1999. Anthelmintic resistance: past, present and future. International journal for
 430 parasitology 29, 115-124; discussion 137-118.

431 Schneider, K., Puehler, F., Baeuerle, D., Elvers, S., Staeheli, P., Kaspers, B., Weining, K.C., 2000. cDNA cloning
 432 of biologically active chicken interleukin-18. Journal of interferon & cytokine research : the official journal
 433 of the International Society for Interferon and Cytokine Research 20, 879-883.

434 Schou, T., Permin, A., Roepstorff, A., Sorensen, P., Kjaer, J., 2003. Comparative genetic resistance to
 435 *Ascaridia galli* infections of 4 different commercial layer-lines. British poultry science 44, 182-185.

436 Schou, T.W., Permin, A., Christensen, J.P., Cu, H.P., Juul-Madsen, H.R., 2010. Mannan-binding lectin (MBL)
 437 in two chicken breeds and the correlation with experimental *Pasteurella multocida* infection. Comparative
 438 immunology, microbiology and infectious diseases 33, 183-195.

439 Schultz, U., Kaspers, B., Rinderle, C., Sekellick, M.J., Marcus, P.I., Staeheli, P., 1995. Recombinant chicken
 440 interferon: a potent antiviral agent that lacks intrinsic macrophage activating factor activity. European
 441 journal of immunology 25, 847-851.

442 Schwarz, A., Gauly, M., Abel, H., Das, G., Humburg, J., Rohn, K., Breves, G., Rautenschlein, S., 2011.
 443 Immunopathogenesis of *Ascaridia galli* infection in layer chicken. Developmental and comparative
 444 immunology 35, 774-784.

445 Scott, M.G., Hancock, R.E.W., 2000. Cationic Antimicrobial Peptides and Their Multifunctional Role in the
 446 Immune System. 20, 24.

447 Shanmugasundaram, R., Selvaraj, R.K., 2011. Regulatory T cell properties of chicken CD4+CD25+ cells.
 448 Journal of immunology 186, 1997-2002.

449 Sick, C., Schultz, U., Staeheli, P., 1996. A family of genes coding for two serologically distinct chicken
 450 interferons. The Journal of biological chemistry 271, 7635-7639.

451 Skallerup, P., Luna, L.A., Johansen, M.V., Kyvsgaard, N.C., 2005. The impact of natural helminth infections
 452 and supplementary protein on growth performance of free-range chickens on smallholder farms in El
 453 Sauce, Nicaragua. Preventive veterinary medicine 69, 229-244.

454 Skovgaard, K., Cirera, S., Vasby, D., Podolska, A., Breum, S.O., Durrwald, R., Schlegel, M., Heegaard, P.M.,
 455 2013. Expression of innate immune genes, proteins and microRNAs in lung tissue of pigs infected
 456 experimentally with influenza virus (H1N2). Innate immunity 19, 531-544.

457 Skovgaard, K., Mortensen, S., Boye, M., Hedegaard, J., Heegaard, P.M., 2010. Hepatic gene expression
 458 changes in pigs experimentally infected with the lung pathogen *Actinobacillus pleuropneumoniae* as
 459 analysed with an innate immunity focused microarray. Innate immunity 16, 343-353.

460 Staeheli, P., Puehler, F., Schneider, K., Gobel, T.W., Kaspers, B., 2001. Cytokines of birds: conserved
 461 functions--a largely different look. Journal of interferon & cytokine research : the official journal of the
 462 International Society for Interferon and Cytokine Research 21, 993-1010.

463 Sugiarto, H., Yu, P.L., 2004. Avian antimicrobial peptides: the defense role of beta-defensins. *Biochemical*
464 *and biophysical research communications* 323, 721-727.

465 Taylor, M.D., van der Werf, N., Maizels, R.M., 2012. T cells in helminth infection: the regulators and the
466 regulated. *Trends in immunology* 33, 181-189.

467 Tongson, M.S., McCraw, B.M., 1967. Experimental ascariasis: influence of chicken age and infective egg
468 dose on structure of *Ascaridia galli* populations. *Experimental parasitology* 21, 160-172.

469 Tugwell, R.L., Ackert, J.E., 1952. On the tissue phase of the life cycle of the fowl nematode *Ascaridia galli*
470 (Schrunk). *The Journal of parasitology* 38, 277-288.

471 Weining, K.C., Sick, C., Kaspers, B., Staeheli, P., 1998. A chicken homolog of mammalian interleukin-1 beta:
472 cDNA cloning and purification of active recombinant protein. *European journal of biochemistry / FEBS* 258,
473 994-1000.

474 Withanage, G.S., Kaiser, P., Wigley, P., Powers, C., Mastroeni, P., Brooks, H., Barrow, P., Smith, A., Maskell,
475 D., McConnell, I., 2004. Rapid expression of chemokines and proinflammatory cytokines in newly hatched
476 chickens infected with *Salmonella enterica* serovar typhimurium. *Infection and immunity* 72, 2152-2159.

477 Yang, D., Biragyn, A., Kwak, L.W., Oppenheim, J.J., 2002. Mammalian defensins in immunity: more than just
478 microbicidal. *Trends in immunology* 23, 291-296.

479

480