Comment on “Temporal Correlations of the Running Maximum of a Brownian Trajectory”

Mortensen, Kim; Pedersen, Jonas Nyvold

Published in:
Physical Review Letters

Link to article, DOI:
10.1103/PhysRevLett.117.248901

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

Comment on "Temporal Correlations of the Running Maximum of a Brownian Trajectory"

Bénichou *et al.* [1] use the running maximum (RM) position in a single experimental trajectory of a particle exhibiting 1D Brownian motion (BM) to estimate its diffusion coefficient. This is unreliable: While the estimator's precision (reproducibility) increases with the suggested parameter tuning, so does its *inaccuracy* (bias), as increasing emphasis is put on the RM's maximum value.

In the mathematical idealization for BM used in Ref. [1], B_t is the position of a particle diffusing with coefficient D. However, $B_t = \sqrt{2D}W_t$, where W_t is the Wiener process. In this model, BM is a scale-free process.

Experimentally, one samples positions $x_{i=1,\ldots,N}$ at time points $t_{i=1,\ldots,N}$ [1]. Typically, constant time lapse Δt is used, such that $t_i = i\Delta t$ and $T = N\Delta t$. For BM, measured positions relate as $x_{i+1} = x_i + \sqrt{2D}\eta_i$, where $\eta_i = W_{t_{i+1}} - W_{t_i}$ is a Gaussian white noise with $\langle \eta_i \rangle = 0$ and $\langle \eta_i \eta_j \rangle = \Delta t \delta_{i,j}$ for all i, j. Each of the $N-1$ displacements $\Delta x_i = x_{i+1} - x_i$ contains information about D; hence, variances of estimators in this discrete case are limited by N, not T, due to the scale invariance of BM.

A reasonable estimator $\hat{D}$ for D should (i) be unbiased, i.e., $\langle \hat{D} \rangle = D$, and (ii) have a variance that decreases as $1/N$, for sufficiently large but practically relevant N. The discretized version $\hat{D}^{(N)}_{\mathrm{msd}}$ of D_{msd} [1] with $\tau = \Delta t$, i.e., $\hat{D}^{(N)}_{\mathrm{msd}} = \sum_{i=1}^{N-1} (\Delta x_i)^2 / [2(N-1)\Delta t]$, complies with (i) and (ii) for $N \geq 2$ in the present case of instantaneous recording of positions and in the absence of measurement noise. It is even optimal: It achieves the Cramér-Rao lower bound [2,3] and thus has the lowest possible variance among unbiased estimators.

With discrete sampling, the RM is $M_i = \max_{j=1,\ldots,i} x_j$, and thus the RM-based estimator of Ref. [1] must read $\hat{D}^{(N,k)}_{\mathrm{es}} = [C(k) \sum_{i=1}^{N} M_i^k]^{2/k}$, with $C(k) \equiv ([\Delta t \sqrt{\pi}(k/2+1)] / \{2^k \Gamma[(k+1)/2] T^{k/2+1}\})$ and $k > 0$. As a function of N, the information available to $\hat{D}^{(N,k)}_{\mathrm{es}}$ increases so slowly that its variance approaches a constant value [1]. This is in conflict with (ii). The variance can be made arbitrarily small, however, by increasing k [1]; thus it is argued that $\hat{D}^{(N,k)}_{\mathrm{es}}$ is superior to $\hat{D}^{(N)}_{\mathrm{msd}}$ for small T [1].

Application of both estimators to Monte Carlo (MC) simulated BM shows, however, that the estimates of $\hat{D}^{(N)}_{\mathrm{msd}}$ scatter with a normal distribution around D, while the estimates of $\hat{D}^{(N,k)}_{\mathrm{es}}$ are skewed [Figs. 1(a) and 1(b)]. This results in a bias, $\langle \hat{D}^{(N,k)}_{\mathrm{es}} \rangle \neq D$, which is in conflict with (i). The bias becomes worse with increasing k [Fig. 1(c)], while the variance indeed decreases [Fig. 1(d)]. The bias of $\hat{D}^{(N,k)}_{\mathrm{es}}$ vanishes too slowly with N to ensure any practical relevance of $\hat{D}^{(N,k)}_{\mathrm{es}}$ relative to $\hat{D}^{(N)}_{\mathrm{msd}}$ [Figs. 1(c) and 1(d)].



FIG. 1. (a) Histograms of estimates obtained from application of, respectively, $\hat{D}^{(N)}_{\mathrm{msd}}$ (red) and $\hat{D}^{(N,k)}_{\mathrm{es}}$ with $k = 1$ (blue) to 10^5 MC simulated, discretely sampled BM trajectories using $D = 0.25$, $\Delta t = 1$, and $N = 10^4$. (b) The same as (a) for $k = 100$. (c) Mean values of estimates obtained as in (a) for various values of N. Results are shown for $\hat{D}^{(N)}_{\mathrm{msd}}$ (pluses) and $\hat{D}^{(N,k)}_{\mathrm{es}}$ with, respectively, k values of 1 (full circles), 10 (open circles), and 100 (crosses). (d) The same as (c) for the variances of the estimates. The theoretical variance $2D^2/(N-1)$ for $\hat{D}^{(N)}_{\mathrm{msd}}$, the Cramér-Rao lower bound, is indicated (full line).

In summary, the estimator suggested by Bénichou *et al.* [1] unfortunately yields biased values for the diffusion coefficient, while optimal, plug-and-play alternatives already exist [2,3].

The Danish Council for Strategic Research (Grant No. 10-092322/DSF) and the European Union's Horizon 2020 research and innovation programme (Grant No. 696656) are acknowledged.

Kim I. Mortensen* and Jonas N. Pedersen†
Department of Micro-and Nanotechnology
Technical University of Denmark
DK-2800 Kongens Lyngby, Denmark

Received 25 August 2016; published 6 December 2016
DOI: 10.1103/PhysRevLett.117.248901

*kim.mortensen@nanotech.dtu.dk
†jonas.pedersen@nanotech.dtu.dk

[1] O. Bénichou, P. L. Krapivsky, C. Mejía-Monasterio, and G. Oshanin, Phys. Rev. Lett. **117**, 080601 (2016).
[2] C. L. Vestergaard, P. C. Blainey, and H. Flyvbjerg, Phys. Rev. E **89**, 022726 (2014).
[3] C. L. Vestergaard, J. N. Pedersen, K. I. Mortensen, and H. Flyvbjerg, Eur. Phys. J. Spec. Top. **224**, 1151 (2015).

