Delivery of TLR7 agonist to monocytes and dendritic cells by DCIR targeted liposomes induces robust production of anti-cancer cytokines.

Klauber, Thomas Christopher Bogh; Laursen, Janne Marie; Zucker, Daniel; Pedersen, Susanne Brix; Jensen, Simon Skjøde; Andresen, Thomas Lars

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Klauber, T. C. B., Laursen, J. M., Zucker, D., Pedersen, S. B., Jensen, S. S., & Andresen, T. L. (2016). *Delivery of TLR7 agonist to monocytes and dendritic cells by DCIR targeted liposomes induces robust production of anti-cancer cytokines..* Poster session presented at The 43rd Annual Meeting & Exposition of the Controlled Release Society, Seattle, WA, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Delivery of TLR7 agonist to monocytes and dendritic cells by DCIR targeted liposomes induces robust production of anti-cancer cytokines

Thomas C. B. Klauber^{a*}, Janne M. Laursen^b, Daniel Zucker^b, Susanne Brix^a, Simon S. Jensen^{a,c}, Thomas L. Andresen^{a,c}

^aDepartment of Micro-and Nanotechnology, Center for Nanomedicine and Theranostics, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark; ^bDepartment of Systems Biology, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark; ^cMonTa Biosciences, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark, *Email: tcbk@nanotech.dtu.dk

Background

Tumor immune escape is today recognized as an important cancer hallmark and is therefore a major focus area in cancer therapy. Monocytes and dendritic cells (DCs) are central to creating a robust anti-tumor immune response and establishing an anti-tumorigenic microenvironment, but are directly targeted by the tumor escape mechanisms to develop immunosuppressive phenotypes. Providing activated monocytes and DCs to the tumor tissue is therefore an attractive way to break the tumor-derived immune suppression and reinstate cancer immune surveillance^{1,2}.

Aim

Our aim was to activate monocytes and DCs with high efficiency. To that end we investigated an immunotherapeutic Toll-Like Receptor (TLR) agonist delivery system comprising liposomes targeted to the dendritic cell immunoreceptor (DCIR) and examined the targeting properties of the liposomes as well as their immune activating potential in blood-derived monocytes, myeloid DCs (mDCs), and plasmacytoid DCs (pDCs)².

Methods

The Toll-like receptor 7 agonist TMX-202 was formulated in maleimide-functionalized liposomes prepared by the lyophilization method. All liposomes were Rhodamine B (RhB) labeled. Anti-DCIR mAbs were thiolated using Traut's reagent and conjugated to the maleimide-functionalized liposomes followed by purification by size exclusion chromatography (SEC). Size and zeta potential was determined by light scattering, ICP-MS and Bradford Assay were used to quantify lipid and Ab concentration and thus the amount of Ab per liposome. HPLC was used to quantify TMX-202 concentration.

Results

Liposomes were 150 - 180 nm of size and anionic. The conjugation procedure was optimized to provide 50-60 mAbs per liposome³. Flow cytometry showed that the anti-DCIR immunoliposomes showed strong preferential association to monocytes and myeloid dendritic cells (mDCs) over the combined population of T, B, and NK cells as well as some association to plasmacytoid DCs (pDCs) (fig. 2). Non-targeted liposomes showed little association to any cell subset (fig. 3), while a the IgG-control immunoliposomes showed some uptake by unspecific mechanisms.

Fig. 2. Anti-DCIR immunoliposomes associate preferentially to monocytes, mDCs, and pDCs (PBMCs). *** $p < 0.0001$, * $p < 0.005$, Mann-Whitney U test. $n = 9$ different donors. Error bars = S.E.M.

Fig. 3. Association of anti-DCIR immunoliposomes, control-IgG immunoliposomes, and m-liposomes to PBMC subsets. A. Association to monocytes; B. Association to mDCs; C. Association to pDCs; D. Association to B, T, and NK cells. *** $p < 0.0001$, ** $p < 0.0005$, * $p < 0.005$, Mann-Whitney U test. $n = 9$ different donors. Error bars = S.E.M.

Peripheral Blood Mononuclear Cells (PBMCs) were purified from buffy coats by gradient centrifugation and incubated with anti-DCIR agonist-immunoliposomes, non-targeted agonist-liposomes, agonist alone, or buffer and the association of the liposomes with PBMC subsets was analyzed by flow cytometry. Following a washing step, PBMCs were cultured for 5 days and on each day the secretion of cytokines was analyzed using Meso Scale Discovery (MSD) sandwich ELISAs.

MSD ELISAs showed that the targeted delivery greatly increased the potency of the TLR7 agonist compared to the free drug. Further, incubation with the anti-DCIR immunoliposomes resulted in potent secretion of key anti-cancer cytokines (IL-12p70, IFN- γ and IFN- α) as well as the pro-inflammatory cytokines TNF- α and IL-6. Importantly, due to gradually decreasing secretion of IL-10, the important IFN- γ :IL-10 ratio increased over the 5 day assay period.

Fig. 4. Cytokine secretion into the culture supernatants on day 1-5. A. IFN- α 2a, B. IL-12p70, C. IFN- γ , D. TNF- α , E. IL-10, F. IL-6. *** $p < 0.0001$, Kruskal-Wallis test. $n = 9$ different donors. Error bars = S.E.M.

Conclusions

Monocytes and mDCs were targeted with high specificity over lymphocytes, and we observed potent TLR7-specific secretion of the anti-cancer cytokines IL-12p70, IFN- α 2a, and IFN- γ . This delivery system could be a way to improve cancer treatment either in the form of a vaccine with co-formulated antigen or as an immunotherapeutic vector to boost monocyte and DC activity in combination with other treatment protocols such as chemotherapy or radiotherapy.

References

- Kalinski, P. *et al.* Natural killer-dendritic cell cross-talk in cancer immunotherapy *Expert Opin. Biol. Ther.* **5**, 1303–15 (2005).
- Monjazeb, A. M. *et al.* Immunoeediting and antigen loss: overcoming the achilles heel of immunotherapy with antigen non-specific therapies *Front. Oncol.* **3**: 1-10 (2013).
- Bates, E. E. *et al.* APCs express DCIR, a novel C-type lectin surface receptor containing an immunoreceptor tyrosine-based inhibitory motif *J. Immunol.* **163**, 1973–83 (1999).
- Kirpotin, D. *et al.* Sterically stabilized anti-HER2 immunoliposomes: design and targeting to human breast cancer cells in vitro *Biochemistry* **36**, 66–75 (1997)