



Stability of vitamin D in fish and mushrooms during different cooking procedures

Ložnjak, Petra; Jakobsen, Jette

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Ložnjak, P., & Jakobsen, J. (2016). *Stability of vitamin D in fish and mushrooms during different cooking procedures*. Poster session presented at 4th International Vitamin Conference, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Stability of vitamin D in fish and mushrooms during different cooking procedures

Petra Ložnjak*, Jette Jakobsen

Division of Food Technology, National Food Institute, Technical University of Denmark

*Corresponding author: petlon@food.dtu.dk



Introduction

Vitamin D is a group of fat-soluble sterols that come in several forms. The two major forms, which differ in their side chain construction (Fig.1), are cholecalciferol (vitamin D₃) found mainly in foodstuffs of animal origin, and ergocalciferol (vitamin D₂) which is found in certain fungi (mushrooms) and plants. There are very few studies which examine the changes in vitamin D content during cooking processing (1, 2). The information in food composition databases is primarily based on the content of vitamin D in raw food. Dietary intake of processed food needs to be corrected by the retention during the cooking process. Therefore, the purpose of the present study was to determine by HPLC-UV method the content of vitamin D₃ in rainbow trout and the content of vitamin D₂ in mushrooms after different cooking procedures. The retention of vitamin D during cooking procedures usually performed in households i.e. boiling, frying and baking was calculated by formula 1 (3).




Figure 1. Chemical structure of ergocalciferol (vitamin D₂) and cholecalciferol (vitamin D₃) with their natural sources

Material

- Rainbow trout (*Oncorhynchus mykiss*)
- Champignon mushrooms (*Agaricus bisporus*)

The types of heat treatments used in this study are listed for trout in Table 1 and mushrooms in Table 2.

Table 1. Processing performed on trout samples

Rainbow trout (<i>Oncorhynchus mykiss</i>)			
Cooking procedure	Abbreviation	Temperature (°C)	Time (min)
Boiling	C	88 ± 2	8
Boiling, pH 4,0	CL	88 ± 2	8
Steam cooked	S	87 ± 3	6
Microwave cooked	M	High power	2
Pan-fried	P	170 ± 4	8
Baked, oven (uncovered)	BA	110 ± 9	30
Baked, oven (uncovered)	BB	210 ± 12	10
Baked, oven (covered)	BC	190 ± 14	15

Table 2. Processing performed on mushrooms samples

Champignon mushroom (<i>Agaricus bisporus</i>)			
Cooking procedure	Abbreviation	Temperature (°C)	Time (min)
Boiling	C7	88 ± 2	20
Boiling, pH 3,5	C3,5	88 ± 2	20
Pan-fried	PS	Strong fire	5
Pan-fried	PW	Weak fire	20
Baked, oven (uncovered)	O200	200 ± 9	20
Baked, oven (uncovered)	O70	70 ± 6	90

Method

After heat treatment, the samples were homogenized, freeze-dried and saponified over-night. The vitamin D metabolites were extracted from the non-saponified matter followed by solid phase clean-up. Further clean-up was performed with semi-preparative HPLC where fractions of vitamin D₂ and D₃ were collected. The extracts were analyzed by HPLC-UV and quantified using vitamin D₂ and D₃ as internal standards.

Results

$$\% \text{ True retention} = \left(\frac{\mu\text{g vit D per 100 gram of cooked food} \cdot \text{amount of cooked food}}{\mu\text{g vit D per 100 gram of raw food} \cdot \text{amount of raw food}} \right) \cdot 100 \quad (1)$$




Figure 2. True retention in fish and mushrooms obtained after different cooking procedures

The preparation conditions affected the degree of retention ($p < 0.05$) both in trout and mushrooms samples.

- Trout samples – high true retention (85-114%)
- Champignon samples – lower true retention (62-89%)
 - Longer duration of heating procedure performed in champignon samples
 - Acidic pH environment enabled preservation of vitamin D in samples which were cooked in a boiling water

Conclusion

Cooking may cause the significant loss of vitamin D but it depends on the foodstuffs and the kind of heating procedure. Further research has to be done to optimize cooking procedure to enhance retention of vitamin D. Changes in vitamin D retention during heat treatment should be taken into consideration in future calculations of dietary intake of vitamin D.

References

1. J. Jakobsen and P. Knuthsen, "Stability of vitamin D in foodstuffs during cooking", *Food Chem.*, vol. 148, pp. 170–175, 2014.
2. P. Mattila, R. Ronkainen, K. Lehtikoinen, and V. Piironen, "Effect of Household Cooking on the Vitamin D content in Fish, Eggs, and Wild Mushrooms", *J. Food Compos. Anal.*, vol. 12, no. 3, pp. 153–160, 1999.
3. E. W. Murphy, P. E. Criner and B. C. Gray, "Comparisons of methods for calculating retentions of nutrients in cooked foods", *J. Agric. Food Chem.*, vol 23, pp. 1153-1157., 1975.