



## Monitoring of active layer refreezing in Greenlandic permafrost

**Doetsch , Joseph ; Christiansen, Anders Vest; Auken, Esben; Fiandaca, Gianluca; Ingeman-Nielsen, Thomas**

*Published in:*  
2nd International Workshop on Geoelectrical Monitoring

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Doetsch , J., Christiansen, A. V., Auken, E., Fiandaca, G., & Ingeman-Nielsen, T. (2013). Monitoring of active layer refreezing in Greenlandic permafrost. In *2nd International Workshop on Geoelectrical Monitoring* (pp. 28-28). Article 13

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## 13

### **Monitoring of active layer refreezing in Greenlandic permafrost**

Doetsch Joseph<sup>1</sup>, Christiansen Anders Vest<sup>1</sup>, Auken Esben<sup>1</sup>, Fiandaca Gianluca<sup>1</sup>, Ingeman-Nielsen Thomas<sup>2</sup>

<sup>1</sup>Aarhus University, Denmark

<sup>2</sup>Technical University of Denmark

The state of the high-latitude permafrost for the Earth's climate system has been widely acknowledged, but permafrost dynamics in a changing climate are poorly understood. Especially the annually freezing and thawing *active layer* is critical for climate dynamics, but it is difficult to monitor without disturbing the studied system.

We use minimally invasive surface geoelectrical monitoring to image active layer refreezing and its lateral variation at high temporal resolution. Our monitoring setup on Disko island, Greenland, allows acquisition of up to 12 data sets per day, using 54 electrodes on two parallel lines. Data is automatically acquired, backed up locally and uploaded to an online database.

We test different time-lapse inversion strategies on the large amount of data and correlate our inversion results with local temperature point measurements. The comparison of the time-lapse inversion results and the temperature and moisture content measurements aims at developing a petrophysical model for partially and fully frozen soils.