



Uplift and tilting of the Shackleton Range in East Antarctica driven by glacial erosion and normal faulting

Paxman, Guy J. G.; Jamieson, Stewart S. R.; Ferraccioli, Fausto; Bentley, Michael J. ; Forsberg, René; Ross, Neil; Watts, Anthony B.; Corr, Hugh F. J.; Jordan, Tom A.

Published in:
Journal of Geophysical Research

Link to article, DOI:
[10.1002/2016JB013841](https://doi.org/10.1002/2016JB013841)

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Paxman, G. J. G., Jamieson, S. S. R., Ferraccioli, F., Bentley, M. J., Forsberg, R., Ross, N., Watts, A. B., Corr, H. F. J., & Jordan, T. A. (2017). Uplift and tilting of the Shackleton Range in East Antarctica driven by glacial erosion and normal faulting. *Journal of Geophysical Research*, 122(3), 2390-2408.
<https://doi.org/10.1002/2016JB013841>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



RESEARCH ARTICLE

10.1002/2016JB013841

Key Points:

- Cenozoic glacial erosion and associated flexural responses have driven 40–50% percent of uplift in the Shackleton Range and Theron Mountains
- Mountain block tilting and a further 40–50% of the total uplift are attributed to normal faulting processes of Jurassic-Cretaceous age

Supporting Information:

- Supporting Information S1

Correspondence to:

G. J. G. Paxman,
guy.j.paxman@durham.ac.uk

Citation:

Paxman, G. J. G., S. S. R. Jamieson, F. Ferraccioli, M. J. Bentley, R. Forsberg, N. Ross, A. B. Watts, H. F. J. Corr, and T. A. Jordan (2017), Uplift and tilting of the Shackleton Range in East Antarctica driven by glacial erosion and normal faulting, *J. Geophys. Res. Solid Earth*, 122, 2390–2408, doi:10.1002/2016JB013841.

Received 7 DEC 2016

Accepted 22 FEB 2017

Accepted article online 27 FEB 2017

Published online 15 MAR 2017

©2017. The Authors.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Uplift and tilting of the Shackleton Range in East Antarctica driven by glacial erosion and normal faulting

Guy J. G. Paxman¹ , Stewart S. R. Jamieson¹ , Fausto Ferraccioli², Michael J. Bentley¹ , Rene Forsberg³ , Neil Ross⁴, Anthony B. Watts⁵ , Hugh F. J. Corr², and Tom A. Jordan²
¹Department of Geography, Durham University, Durham, UK, ²British Antarctic Survey, Cambridge, UK, ³National Space Institute, University of Denmark, Kongens Lyngby, Denmark, ⁴School of Geography, Politics and Sociology, Newcastle University, Newcastle upon Tyne, UK, ⁵Department of Earth Sciences, Oxford University, Oxford, UK

Abstract Unravelling the long-term evolution of the subglacial landscape of Antarctica is vital for understanding past ice sheet dynamics and stability, particularly in marine-based sectors of the ice sheet. Here we model the evolution of the bedrock topography beneath the Recovery catchment, a sector of the East Antarctic Ice Sheet characterized by fast-flowing ice streams that occupy overdeepened subglacial troughs. We use 3-D flexural models to quantify the effect of erosional unloading and mechanical unloading associated with motion on border faults in driving isostatic bedrock uplift of the Shackleton Range and Theron Mountains, which are flanked by the Recovery, Slessor, and Bailey ice streams. Inverse spectral (free-air admittance) and forward modeling of topography and gravity anomaly data allow us to constrain the effective elastic thickness of the lithosphere (T_e) in the Shackleton Range region to ~20 km. Our models indicate that glacial erosion, and the associated isostatic rebound, has driven 40–50% of total peak uplift in the Shackleton Range and Theron Mountains. A further 40–50% can be attributed to motion on normal fault systems of inferred Jurassic and Cretaceous age. Our results indicate that the flexural effects of glacial erosion play a key role in mountain uplift along the East Antarctic margin, augmenting previous findings in the Transantarctic Mountains. The results suggest that at 34 Ma, the mountains were lower and the bounding valley floors were close to sea level, which implies that the early ice sheet in this region may have been relatively stable.

1. Introduction

Antarctica's bedrock topography is an important boundary condition that influences the dynamics of the overlying ice sheet [Gasson *et al.*, 2015]. In particular, Antarctic ice sheet stability in regions proximal to the grounding line is heavily dependent on the local ice thickness and bedrock elevation and slope [Pollard *et al.*, 2015]. Near-coastal regions of Antarctica where the ice sheet is marine-based (i.e., the bed is below present-day sea level) are particularly susceptible to rapid grounding line retreat via marine ice sheet instability [Schoof, 2007] and calving mechanisms such as hydrofracturing and ice cliff failure [Pollard *et al.*, 2015].

The Recovery catchment in East Antarctica is located on the eastern margin of the Weddell Sea (Figure 1). It is one of the largest and yet least explored marine-based sectors of the East Antarctic Ice Sheet (EAIS) that may be susceptible to such instability mechanisms [Le Brocq *et al.*, 2008]. The area of the catchment is 1.5×10^6 km²; it drains ~10% of the EAIS and contains ~5 m of sea level equivalent, which is approximately equivalent to the entire West Antarctic Ice Sheet [Rignot *et al.*, 2008]. The regional ice velocity field shows that ice flow in the catchment is focused through three major outlet glaciers—Recovery, Slessor, and Bailey—where velocities reach almost 1000 m/yr at the grounding line [Rignot *et al.*, 2011]. These glaciers are the major arteries that drain the EAIS into the Filchner Ice Shelf (Figure 1).

Situated between these outlet glaciers are the Shackleton Range and Theron Mountains, over which ice velocities are less than 10 m/yr. The strong bimodality in the ice velocity field is reflected in, and caused by, the bedrock topography (Figure 1). The summits of the Shackleton Range and Theron Mountains protrude above the EAIS as nunataks at up to 1.8 km above sea level, while the bed at the floor of the Recovery, Slessor, and Bailey troughs is as deep as 2.5 km below sea level; the ice thickness in these glaciers exceeds 3 km [Fretwell *et al.*, 2013]. The troughs trend E-W and are 300–500 km long and 50–100 km wide. This fjord-like landscape renders the Recovery catchment particularly susceptible to ice sheet retreat in a warming world [DeConto and Pollard, 2016]. However, if the subglacial landscape has evolved significantly in the past




Figure 1. Regional setting of the Recovery catchment within Antarctica. (a) Ice sheet velocities [Rignot *et al.*, 2011]. EAIS = East Antarctic Ice Sheet. The inset denotes the major Antarctic ice divides [Rignot *et al.*, 2008], the area bounded by the red colored polygon is the Recovery catchment, and the black box denotes the area shown in the main figure. (b) Bedrock topography. The red lines show the major onshore basement faults that bound the Shackleton Range and Theron Mountains [Marsh, 1985]. The submarine Thiel trough is bounded by the Filchner Rift (magenta lines, ticks point to the downthrown side) [Jordan *et al.*, 2013, 2016]. The black dashed box marks our main study area.

34 Ma, the response of this sector to climatic and oceanic change in the past may have been very different compared to that of its modern configuration.

The timing and mechanism(s) responsible for the uplift of the Shackleton Range and Theron Mountains and the subsidence of the Recovery, Slessor, and Bailey troughs remain outstanding questions. Apatite fission track and (U-Th)/He dating indicate multiple phases of denudation and burial of the Shackleton Range in the Mesozoic before final uplift and formation of the present landscape since EAIS inception at 34 Ma [Krohne *et al.*, 2016]. Previous studies have suggested that the uplift of the mountain ranges and subsidence of the troughs are inherently coupled. Sugden *et al.* [2014] hypothesize that post-Eocene uplift of the Shackleton Range was driven by the regional isostatic response to glacial overdeepening and erosion within the Recovery and Slessor troughs. Furthermore, they speculate that the observed tilt of the Shackleton block, with the highest elevations along the southern escarpment (Figures 2 and 3) occurred because excavation of the larger Recovery trough caused more flank uplift than the smaller Slessor trough.

The Recovery, Slessor, and Bailey glaciers likely exploited preexisting fault systems that separate the metamorphic basement of the Shackleton Range [Tessensohn *et al.*, 1999b; Will *et al.*, 2009, 2010] from the Palaeozoic Beacon sediments and Jurassic dolerite sills exposed in the Theron Mountains and the isolated




Figure 2. (a) Perspective image of the bedrock of the Recovery catchment (vertical exaggeration (VE) = x50). The traces of inferred range-bounding faults [Marsh, 1985] are marked by the red dashed lines. The arrow marks the direction of grid north in the adopted polar stereographic projection. The inset shows the location of the study area within Antarctica. (b) Field photograph of a peneplanation surface exposed on Stephenson Bastion in the Shackleton Range (location marked by blue star in Figure 2a). (c) Profile X-Y across the Recovery catchment (VE = x50). Bedrock (black line) and ice surface (blue line) topography were assembled from three RES flight lines. Ice flow direction is out of the page. The green dashed lines highlight the tilting of the mountain blocks. The schematic red lines mark the position of the faults (the arrows show the inferred sense of dip-slip motion).

Whichaway Nunataks [Brook, 1972] (Figure 2). These fast-flowing glaciers are bounded by ice surface lineaments that reflect the trend of major subglacial fault systems (Figure 1) [Marsh, 1985]. These lineaments trend parallel to E-W trending approximately 500 Ma thrust faults in the Shackleton Range [Tessensohn et al., 1999b], regional aeromagnetic lineaments interpreted as reflecting major basement faults [Jordan et al., 2016], and inferred half-graben basins upstream of the Slessor Glacier [Bamber et al., 2006; Shepherd et al., 2006]. These observations support the hypothesis that the deep subglacial troughs are structurally controlled [Patton et al., 2016]. Jurassic extension and horst-and-graben formation have been recognized in the adjacent Weddell Sea Rift System [Jordan et al., 2016, and references therein] and also onshore, in particular in Dronning Maud Land where the Jurassic Jutulstraumen Rift has been imaged [Ferraccioli et al., 2005a, 2005b]. The Shackleton Range and Theron Mountains may therefore represent fault-bounded horst blocks that experienced tectonic uplift and tilting [Skidmore and Clarkson, 1972].

The relative roles of erosion-driven and tectonic uplift in driving Shackleton Range and Theron Mountains uplift have yet to be quantified, in contrast to the Gamburtsev Subglacial Mountains [Ferraccioli et al., 2011; Paxman et al., 2016] or the Transantarctic Mountains (TAM) [Stern et al., 2005], where the relative roles of these processes have been addressed with the aid of quantitative modeling. In this study, we use 3-D




Figure 3. (a) Bedrock topography DEM. The solid lines show the location of profiles 1–5 used to construct Figure 3c. The red lines show the location of the profiles used in our elastic thickness forward models (Figure 4). (b) Free-air gravity anomaly gridded from IceBridge and ICEGRAV line data. The red box shows the area of the grids used to compute the free-air admittance (Figure 4). Both grids are projected in Antarctic polar stereographic with true scale at 71°S. (c) Ensemble averaging of bedrock topography. Topographic profiles (colored lines) were constructed by sampling the DEM (Figure 3a) along five equally spaced (~20 km spacing), parallel lines. The profiles were isostatically rebounded to remove the effect of present-day ice sheet loading using an elastic plate model with $T_e = 20$ km, which corresponds to our regional T_e estimate (section 3.1). Only lines 1, 3, and 5 are shown for clarity. The black line is an ensemble average of profiles 1–5. The Shackleton Range and Theron Mountains are tilted (in the absence of ice loading) by ~1.2°N and ~0.8°S, respectively. BIS = Bailey Ice Stream, TM = Theron Mountains, SG = Slessor Glacier, SR = Shackleton Range, RG = Recovery Glacier.

flexural isostatic models to quantify for the first time both the mechanical unloading associated with normal border faults and erosional unloading associated with Cenozoic glacial incision in the Shackleton Range and Theron Mountains region. Our results have significant implications for understanding the evolution of paleotopography in this part of East Antarctica and for assessing how these changes in topography may have influenced the early history of the EAIS.

2. Geophysical Data Sets

This study utilizes bedrock elevation and free-air gravity data acquired during a number of recent airborne geophysical surveys over the previously unexplored Recovery catchment.

2.1. Bedrock Topography

We collated onshore ice thickness data from a series of recent airborne radio-echo sounding (RES) surveys over Coats Land and the Recovery catchment, including ICEGRAV (2013) [Ferraccioli *et al.*, 2014], Operation IceBridge (2009–2012) [Leuschen *et al.*, 2010, updated 2016], and a 2001/2002 survey of the upper reaches of the Bailey Ice Stream and Slessor Glacier [Rippin *et al.*, 2003; Bamber *et al.*, 2006; Shepherd *et al.*, 2006]. We also include direct ice thickness measurements that were previously incorporated into Bedmap2 [Fretwell *et al.*, 2013] (data coverage is shown in Figure S1 in the supporting information).

RES profiles reveal that the Theron Mountains exhibit a lightly dissected mesa-like topography, whereas the Shackleton Range is more heavily incised (Figure 2). The mesas in the Theron Mountains resemble those observed westward of the TAM, which are interpreted as the result of the Ferrar dolerite sills capping Beacon Supergroup sedimentary rocks [Ferraccioli *et al.*, 2001, 2009; Studinger *et al.*, 2004]. Both lithologies are also present in the Theron Mountains [Brook, 1972], hinting at a common mode of formation. A number of plateau surfaces are also exposed at up to 1.8 km above sea level in the Shackleton Range (Figure 2) [Skidmore and Clarkson, 1972; Kerr and Hermichen, 1999]. The plateau surfaces in the Shackleton Range cut different geological units; they do not reflect a stratigraphic dip slope but instead are surfaces that experienced erosion and were subsequently uplifted. These plateaux have been interpreted as remnant fragments of the Devonian Kukri Penepine, a flat, once-continuous undulating erosion surface which is extensively observed in the TAM [Stern and ten Brink, 1989; Fitzgerald, 1994; Tessensohn *et al.*, 1999a].

We gridded the new flight line data together with the existing direct ice thickness measurements from Bedmap2 [Fretwell *et al.*, 2013] using a 2 km grid mesh with a continuous curvature tensional spline algorithm [Wessel *et al.*, 2013]. The grid was masked to remove interpolated values more than 10 km from the nearest data point. These grid nodes were replaced by ice thickness values from the Bedmap2 compilation; while there are no direct ice thickness estimates in these areas in Bedmap2, approximate ice thicknesses have been computed using satellite-derived gravity field models [Fretwell *et al.*, 2013]. We then subtracted the ice thickness grid from the surface digital elevation model (DEM) [Fretwell *et al.*, 2013] to produce a bedrock DEM (Figure 3). Offshore bathymetry data were taken from Bedmap2.

We took the spectral average [see, e.g., Bassett and Watts, 2015] of an ensemble of five profiles crossing the mountain ranges. The ensemble average enhances the “common” features of the topographic profiles, such as the tilted plateau surface and the deep *U*-shaped glacial troughs, while at the same time suppresses the effects of the more localized dissection of the plateau surface by cirques and rivers. It can be seen that the Shackleton Range is on average tilted by 1.2° to the north and the Theron Mountains are tilted by 0.8° to the south (Figure 3).

2.2. Free-Air Gravity Anomaly

Our new free-air gravity anomaly (FAA) grid for the Recovery catchment (Figure 3) was generated from flight line data from the Operation IceBridge [Cochran and Bell, 2010, updated 2016] and ICEGRAV 2011 and 2013 surveys [Ferraccioli *et al.*, 2014]. Continuation to 500 m above the bedrock elevation, crossover analysis, and leveling of the lines was performed, and a satisfactory standard deviation of 1 mGal at crossovers between intersecting flight tracks was achieved. Gravity data were gridded at 2 km horizontal spacing using a continuous curvature tensional spline algorithm [Wessel *et al.*, 2013]. The grid was masked to remove interpolated values more than 10 km from the nearest data point. Uncertainties in the FAA grid were estimated to be ± 2 mGal. The FAA grid was used in conjunction with the bedrock topography grid to estimate the regional flexural rigidity of the lithosphere (section 3.1).

3. Methods

3.1. Effective Elastic Thickness Estimation

The isostatic response of the lithosphere to (un)loading may be computed by modeling the lithosphere as a flexed elastic plate overlying an inviscid (non-viscous) fluid [Watts, 2001]. The amplitude and wavelength of the isostatic response are determined by the effective elastic thickness (T_e), a proxy for the integrated strength of the lithosphere [Watts and Burov, 2003]. We employed two independent methods to determine the appropriate T_e for the Recovery catchment to see whether the T_e values converged.

3.1.1. The 2-D Forward Modeling

The bedrock topography along two RES flight lines exhibits intermediate-wavelength (100–500 km) warping characteristic of plate flexure in response to surface loading (Figures 4a and 4b). In these profiles, the Bailey trough is downwarped toward the elevated Theron Mountains mesa. We envisage that this topography is largely the product of regional erosion of material from within the Bailey trough and normal fault action. Unloading of the material within the trough is equivalent to the loading of a flat sheet by the mesa. The topography is therefore analogous to the loading of a seamount on the ocean floor, except the mesa displaces ice rather than water. We do not explicitly model the mechanism of loading; our 2-D forward model comprised a distributed load approximating the shape of the mountain range (with a topographic density of 2670 kg m^{-3}) (Figures 4a and 4b), which was applied to a thin elastic plate with a uniform T_e overlying an inviscid fluid mantle (with density 3330 kg m^{-3}) (equation (1)). The density of the material displaced by the load and infilling the flexure was that of ice (915 kg m^{-3}). We modeled the topography for a series of T_e values between 10 and 50 km. The wavelength of flexure is consistent with T_e values of 20–30 km. The best fitting T_e values (24 and 25 km for the two models) were determined using the root-mean-square misfit (Figures 4a and 4b).

3.1.2. The 3-D Inverse (Spectral) Modeling

The gravitational admittance is a transfer function that describes the relationship between the FAA and the bedrock topography for a range of load sizes (wavelengths) on an elastic plate with a given T_e . The observed admittance was computed by taking Fourier transforms of our newly compiled bedrock topography and FAA grids over a $900 \text{ km} \times 900 \text{ km}$ window (Figure 3) [following McKenzie and Fairhead, 1997; McKenzie, 2003]. Theoretical admittance functions for an elastic plate subject to surface loading were computed for a range




Figure 4. Effective elastic thickness modeling. (a and b) Forward modeling of observed bedrock topography (red lines) along two flight lines (A–A' and B–B') crossing the Bailey Ice Stream and Theron Mountains (locations are marked in Figure 3a). Comparison of predicted topography from elastic plate models (black lines) with the observed topography indicates a best fitting T_e of (Figure 4a) 25 and (Figure 4b) 24 km. (c) Comparison of the observed free-air gravitational admittance (red dots with standard error bars) with model curves for $T_e = 0, 5, 10, 20,$ and 40 km. The admittance recovers a best fitting T_e of 11 km, which is calibrated to 18 km (see Figure S2 in the supporting information).

of T_e values [Watts, 2001] and compared to the observed admittance (Figure 4c), providing an estimate of the average T_e value across the region. This method recovers a best fitting T_e value of 11 km. However, taking a Fourier transform of data sets with limited lateral extent causes spectral leakage into the result, downward biasing the recovered T_e [Kirby, 2014]. A calibration that accounts for the consequent underestimation of T_e [Kalinin and Watts, 2009] was used to correct the recovered T_e to 18 km (Figure S2).

Despite the uncertainties associated with the interpretation of the admittance for topography/gravity data sets of limited lateral extent, such as spectral leakage and the fact that the window-based spectral estimates reflect a wide range of spatial and temporal loads [Kirby, 2014], the corrected T_e value of 18 km is broadly consistent with the 24 and 25 km results from our 2-D forward models. In our subsequent flexure

calculations, we used a 3-D elastic plate model with a uniform T_e of 20 km, which is intermediate between our estimates, and tested the sensitivity of our model by running the calculations for T_e values between 5 and 50 km (Figure S3).

3.2. Calculation of Erosional Unloading

3.2.1. Spatial Distribution of Erosion

In order to determine the 3-D distribution of eroded material, we used a peak accordance method [Stern *et al.*, 2005; Champagnac *et al.*, 2007]. This approach involves the identification of peaks and flat-topped surfaces in the bedrock topography that are assumed to have not experienced any erosion and the interpolation of a smooth surface between them. The resulting “peak accordance surface” represents the restoration of the eroded material to the topography without accounting for the associated isostatic response. The difference between the accordance surface and the bedrock topography is the eroded material.

We identified over 80 flat-topped surfaces in the vicinity of the Shackleton Range and Theron Mountains in RES flight lines (Figure 2), Google Earth satellite imagery, and field photographs (Figure 2). We assumed that these now high-elevation plateau surfaces originally formed a contiguous, low elevation, and flat landscape prior to incision into it and thus have not experienced erosion since the onset of continental glaciation at 34 Ma. This assumption is supported by very low (0.10–0.35 m/Myr) long-term cosmogenic nuclide-derived erosion rates on the plateau surfaces in the Shackleton Range [Fogwill *et al.*, 2004; Sugden *et al.*, 2014] and also by a lack of glacial modification of these surfaces [Kerr and Hermichen, 1999]. If the peaks have been lowered since 34 Ma, the amount of erosion will be an underestimate. In addition, we used a spatial filter to identify local highs in the bedrock topography DEM within a circular moving window with a fixed radius of 15 km [following Champagnac *et al.*, 2007; Paxman *et al.*, 2016]. Peaks where the present-day ice velocity exceeds 10 m/yr, and have therefore likely experienced significant erosion, were discarded. The remainder were assumed to have experienced negligible erosion since 34 Ma; a smooth surface was interpolated between the peaks and flat-topped surfaces to produce a peak accordance surface that was assumed to exist just prior to the onset of glaciation at 34 Ma (Figure 5).

The accordance surface was constructed by (1) adjusting the DEM to account for the loading of the present-day ice sheet using our preferred elastic plate model with a T_e of 20 km (see section 3.2.3), (2) sampling the adjusted DEM at the location of each peak, and (3) interpolating between peaks using a continuous curvature tensional spline [Wessel *et al.*, 2013]. The eroded material was calculated by subtracting the ice-free bedrock topography from the peak accordance surface (Figure 5).

The assumption that the difference between the peak accordance surface and the bedrock topography is entirely due to removal of material by glacial (post-34 Ma) erosion is probably reasonable within the Theron Mountains and Shackleton Range themselves. However, this may not be the case over the large troughs, where some of the difference may also be attributable to tectonic subsidence due, for example, to movement on the border faults. For this reason there is uncertainty in the amount of material eroded from the troughs. We envisage two end-member scenarios for the amount of material that has been eroded from the troughs:

1. Minimum erosion scenario—tectonic subsidence has contributed to trough depth. In this scenario the peak accordance surface is dipped over troughs (Figure 5), representing a preexisting depression caused by mechanical subsidence on border faults (section 3.3). We dipped the surface such that when the contributions of erosional unloading and fault motion were summed (see section 3.3), the modeled trough depth matched the observed trough depth. It is therefore assumed that the tectonic subsidence was not infilled with sediment.
2. Maximum erosion scenario—subsidence of the trough floors below sea level is entirely attributable to glacial erosion. In this scenario, the peak accordance surface is stretched across the tops of the troughs (Figure 5) and the difference between the accordance surface and the bedrock topography is all glacially eroded material. Under this maximum erosion scenario, it is assumed that any fault movement predated glaciation and the resulting subsidence of the hanging wall blocks was completely infilled with sediment [Bamber *et al.*, 2006; Shepherd *et al.*, 2006].

Assuming an average eroded rock density of $2300\text{--}2700\text{ kg m}^{-3}$ (reflecting sedimentary and basement rock end-members), the minimum and maximum estimated mass of eroded material were $1.0 \times 10^{18}\text{ kg}$ and $1.5 \times 10^{18}\text{ kg}$, respectively.




Figure 5. Erosional unloading modeling. (a) Present-day bedrock topography adjusted for ice sheet loading ($T_e = 20$ km). The dashed lines show the traces of range-bounding faults. The symbols mark the points used to construct the peak accordance surface. Blue diamonds = subglacial flat-topped surfaces (mesas), green stars = plateau surfaces exposed in the Shackleton Range, yellow circles = local maxima within a fixed (15 km) radius. (b) Peak accordance surface. This represents the maximum erosion scenario (see text for description). (c) Distribution of eroded material (warm colors) and offshore sediment (cool colors). SR = Shackleton Range, TM = Theron Mountains. (d) Computed flexural response ($T_e = 20$ km) to unloading of eroded material and loading of sediment. (e) Maximum and (f) minimum erosion scenario along profile A-B (location marked in Figures 5a–5d). Black line = bedrock topography, magenta = peak accordance surface, yellow shaded region = eroded material, red = modeled flexure.

3.2.2. Offshore Sediment Estimates

The estimated mass of eroded material was compared to the mass of sediment located offshore on the continental shelf. Isopach maps for the Weddell Sea shelf north of the calving front (Figure 5) have been constructed by interpolating sediment package thicknesses measured from seismic reflection lines [Huang *et al.*, 2014]. Sediments are divided into preglacial (145–34 Ma), transitional (34–14 Ma), and full-glacial (14–0 Ma) sequences based on correlation of seismic stratigraphic facies across lines and age constraints from sediment cores [Huang *et al.*, 2014]. Because of the uncertainties associated with the volume, provenance, and postdepositional reworking of sediment, we determine a maximum and minimum total 34–0 Ma sediment volume under the following assumptions:

1. Material eroded from the Recovery catchment is now located on the southeastern Weddell Sea shelf (eastward of 50°W and south of 75°S), including the Crary Fan [Diekmann and Kuhn, 1999] (Figure 5). However, the Support Force Glacier and (during glacial periods) the Foundation Ice Stream also drain into the southeastern Weddell Sea via the Filchner Ice Shelf (Figure 1), so some fraction of the sediment will have been derived from this catchment. We assume that between 50 and 100% of the detrital sediment entered the Weddell Sea via the Recovery, Slessor, and Bailey glaciers, since they drain a larger area than the Support Force and Foundation glaciers.
2. Five to fifteen percent of the total offshore sediment is pelagic (biogenic) rather than detrital and therefore was not derived from onshore [Wilson *et al.*, 2012].
3. The average bulk density of offshore sediment is between

2100 and 2300 kg m⁻³ [Wilson *et al.*, 2012]. This accounts for uncertainties in the amount of pore space between the grains (i.e., the degree of mechanical compaction) and their lithology.

By computing the total volume of sediment and applying these assumptions, the mass of 34–0 Ma Recovery catchment-derived detrital sediment in the Weddell Sea basin was determined to be 0.66–1.6 × 10¹⁸ kg. The mass of rock eroded from onshore (1.0–1.5 × 10¹⁸ kg) is therefore within the range of uncertainty of the mass of offshore material. It might be expected that the mass of eroded material exceeds the mass of offshore sediment, since material has likely been lost from the shelf and reworked in the Weddell Gyre or by contourite currents, and some sediment may have instead been deposited within interior sedimentary basins [e.g., Shepherd *et al.*, 2006] during the early stages of EAIS development. Therefore, even our maximum erosion scenario does not obviously overestimate the amount of post-34 Ma erosion from the region. Due to a lack of constraints, we do not incorporate post-34 Ma sediment deposition onshore or beneath the Filchner Ice Shelf or the associated isostatic response in our models. Although this leads to an unrealistic gradient in sediment thickness/erosion at the calving front (Figure 5), sensitivity testing indicates that onshore flexural uplift is insensitive to the amount of offshore erosion/deposition (Figure S3).

3.2.3. Flexural Isostasy

The flexural isostatic adjustment ($w(x,y)$) to erosional unloading and sediment loading (Figure 5) was computed by solving the general equation for the (un)loading ($h(x,y)$) of an elastic plate overlying a nonviscous fluid [Turcotte and Schubert, 1982].

$$\nabla^2 [D(x,y) \nabla^2 w(x,y)] + (\rho_{\text{mantle}} - \rho_{\text{infill}})gw(x,y) = (\rho_{\text{load}} - \rho_{\text{displace}})gh(x,y) \quad (1)$$

where

$$D(x,y) = \frac{ET_e(x,y)^3}{12(1-\nu^2)} \quad (2)$$

is the flexural rigidity as a function of spatial dimensions x and y . Density terms represent the load (ρ_{load}), the material infilling the flexure (ρ_{infill}), the material displaced by the (un)loading (ρ_{displace}), and the mantle (ρ_{mantle}). We assumed values of 9.81 ms⁻² for the acceleration due to gravity (g), 100 GPa for Young's modulus (E), and 0.25 for the Poisson ratio (ν). By solving equation (1), we calculated the flexural uplift/subsidence due to the removal of the modern-day ice sheet (assuming an ice density of 915 kg m⁻³), the removal of the eroded material (assuming an average eroded material density of 2500 kg m⁻³), and the loading of offshore sediments (assuming an average sediment density of 2200 kg m⁻³).

The unloads were our updated ice thickness grid (section 2.1) and the grid of eroded material (section 3.2.1); the offshore sediment load was the postglacial (34–0 Ma) sediment isopach of Huang *et al.* [2014]. The modeled flexure is most sensitive to T_e , since this governs the amplitude and wavelength of the flexural response; sensitivity testing was carried out by computing the flexure for T_e values between 5 and 50 km, a typical range of values for the continental lithosphere [Watts, 2001]. Since crustal-scale faults may introduce a discontinuity in the plate where the flexural rigidity is effectively zero [Watts, 2001], we also tested a “broken plate” scenario, where T_e was decreased to zero at the plate break along one or more of the major faults bounding the Shackleton Range and Theron Mountains. We used a fast Fourier transform method [Watts, 2001] to solve equation (1) analytically for spatially uniform T_e scenarios and a numerical centered finite difference technique [e.g., Stewart and Watts, 1997] for spatially variable T_e scenarios.

3.3. Calculation of Mechanical Unloading

The bedrock topography of the Recovery catchment, with broad valleys bounded by faults and uplifted flanks, is typical of extensional terranes. Vening Meinesz [1950] proposed a model for the uplift of rift flanks as a consequence of failure of the lithosphere by normal faulting. In this case, slip on a normal fault causes unloading of the footwall block by removal of the hanging wall; the result is flexural isostatic rebound and uplift of the footwall. Concomitant replacement of footwall crustal rock by the mantle causes isostatic subsidence of the hanging wall block. Long-term normal fault displacement may therefore be modeled as the flexural isostatic adjustment to the rigid uplift/subsidence of the footwall/hanging wall blocks, assuming that the lithosphere retains a finite flexural rigidity during extension [Weissel and Karner, 1989]. The resulting topography resembles a half-graben, and the footwall is flexurally uplifted. Uplift on the shoulders of normal

faults is therefore the result of this so-called “mechanical unloading” of the lithosphere [Weissel and Karner, 1989; Watts, 2001].

To determine the contribution of mechanical unloading associated with the border faults to Shackleton Range and Theron Mountains uplift, we modeled the displacement across each fault as the flexural isostatic adjustment to the rigid uplift and subsidence of the footwall and hanging wall [following Weissel and Karner, 1989] (Figure 6). The amount of flexure depends on the elastic thickness, thickness of the faulted layer (the crust), material densities, and dip and heave of the faults. We used our preferred uniform T_e scenario (20 km) and a crustal thickness of 35 km [An *et al.*, 2015]. The assumed densities of the crust, infill (air), and mantle were 2670, 1, and 3330 kg m⁻³, respectively. For simplicity, we assumed that each fault is continuous, dips at 60° toward the downthrown side, and exhibits a constant vertical displacement (throw) along-strike. We tested the sensitivity of the results to the elastic and crustal thicknesses and the fault dip and found that only T_e strongly influences the distribution of flexure (Figure S4). The amount of extension (heave) was tuned so the modeled displacement matched the observed topography next to the fault(s).

We also incorporated the diffusion of the scarp due to mass-wasting processes, which is given by [Watts, 2001]:

$$h_t(k) = h_0(k) e^{-\left[1 - \left(\frac{\rho_c}{\rho_m}\right) \phi_e(k)\right] \kappa k^2 t} \quad (3)$$

where

$$\phi_e(k) = \left[\frac{Dk^4}{\rho_m g} + 1 \right]^{-1} \quad (4)$$

is the flexural response function, $h_t(k)$ is the topography after time t , $h_0(k)$ is the initial topography, κ is the “subducing coefficient,” and k is the wave number (the computation is carried out in the frequency domain). This equation assumes that erosion is a diffusive process that transports mass from the uplifted side of the fault to the subsided region (and the resulting flexural isostatic adjustment is computed). The result is to smooth the edge of the fault-generated topography so it is more exponential in form. Values of t and κ were chosen so the modeled scarp slope matched the observed slopes of the mountain ranges.

The (diffused) flexure was calculated in 2-D along a series of 1000 km long profiles (with 10 km horizontal spacing) trending orthogonal to the faults (Figure 6). These profiles were then gridded to produce a 3-D map of flexure driven by mechanical unloading (Figure 6). The displacement on the four major border faults was superimposed in various combinations to produce the total 3-D fault-driven displacement. We estimated the throw on the faults bounding the Recovery and Bailey troughs by measuring the difference in elevation of the bedrock on either side of the troughs. Elevation differences of 600–700 m provide first-order estimates of the cumulative long-term throw on the faults, assuming that the flexure associated with erosional unloading is approximately symmetrical either side of the troughs (Figure 5).

4. Results

4.1. Erosion-Driven Uplift

We calculated the contribution of erosional unloading to Shackleton Range and Theron Mountains uplift for our minimum and maximum erosion scenarios. For our preferred T_e scenario of 20 km, we find that erosion in the Recovery, Slessor, and Bailey troughs has driven on average between 600 m (minimum erosion scenario) and 800 m (maximum erosion scenario) of flexural uplift throughout the Shackleton Range and Theron Mountains (Figure 5). This represents ~40–50% of the total elevation of the mountain blocks. The greatest amount of flexural uplift (~1 km in the maximum erosion scenario) occurs along the southern flank of the Shackleton Range, which is bounded by the Recovery trough. The Recovery trough is deeper and wider than the Slessor trough, resulting in a larger-magnitude and longer-wavelength erosional unload [Sugden *et al.*, 2014]. However, this differential erosional unloading only confers a maximum northward tilt of 0.2° on the upper surface of the Shackleton Range, compared to the observed tilt of 1.2°N (Figures 5 and 7).

The flexure onshore is very insensitive to the amount and distribution of sediment offshore (Figure S4). Offshore sediment loading accounts for <3% of uplift/subsidence in the Shackleton Range, Theron Mountains, and bounding glacial troughs. We suggest that this is because the locus of sediment loading




Figure 6. Mechanical unloading modeling. (a) Location of profiles used to produce the 3-D flexural uplift distribution due to motion on a single fault (dashed line). The 2-D flexure profile (black line in Figure 6d) was sampled onto each 1000 km long profile at 1 km spacing. Each profile trends perpendicular to the fault trace. (b) Gridded flexural uplift due to mechanical unloading associated with dip-slip motion on the border fault (dashed line). (c) Flexural uplift due to mechanical unloading on four border faults, calculated by superimposing the individual displacements (e.g., as shown in Figure 6b). SR = Shackleton Range, TM = Theron Mountains. (d) Profile X-Y (location marked in Figure 6b). Dashed line = flexural uplift due to normal faulting [Weissel and Karner, 1989], solid line = diffused topography, red line = topography sampled from the grid (Figure 6b) along the same profile. Gridding causes a minor reduction in the amplitude of the topography but retains the distinct flexed pattern. (e) Profile A-B (location marked in Figure 6c). Locations of faults, with sense of motion, are shown schematically. Black line = bedrock topography, red line = modeled flexure due to mechanical unloading.

(the southeastern Weddell Sea) is too distal for significant isostatic uplift/subsidence to be transmitted to the inland fjord system, even if a flexurally rigid ($T_e = 50$ km) lithosphere is assumed.

We tested the sensitivity of the model to the assumed T_e scenario (Table 1 and Figure S3). However, we found that while the pattern of erosion-driven flexure is sensitive to the assumed T_e , no value between 5 and 50 km was able to produce a satisfactory agreement with the observed magnitude and wavelength of mountain uplift. Intermediate T_e values of 20–30 km give the best agreement with the observed wavelength of tilting, but the modeled tilt is only $\sim 0.2^\circ$. We also tested a scenario where the elastic plate was broken along faults bounding the Shackleton Range and Theron Mountains to investigate whether this could reproduce the observed tilting of the mountain ranges (Figure S3). However, the difference between continuous- and broken-plate flexure is relatively minor except for regions very close to the faults.




Figure 7. Contribution of erosional unloading and mechanical unloading to Shackleton Range and Theron Mountains uplift. (a) Rebounded bedrock topography. (b) Sum of glacial erosion and associated isostatic rebound (maximum erosion scenario). (c) Normal fault-driven uplift (mechanical unloading). (d) Total model uplift (sum of Figures 7b and 7c). A continuous elastic plate model with a T_e of 20 km was used. SR = Shackleton Range, TM = Theron Mountains. (e) Profile A-B across the Shackleton Range and Theron Mountains for the maximum erosion scenario. Under this scenario, fault-driven hanging wall subsidence was filled to sea level. The sum of the modeled erosion- and fault-driven uplift (blue line) compares well with the observed topography (black line). (f) Profile A-B for the minimum erosion scenario. Under this scenario, hanging wall subsidence was not infilled, meaning that the contribution of erosion was reduced. The match between observed and modeled topography is worse than the maximum erosion scenario, but the relative contributions of erosional and mechanical unloading remain similar.

Table 1. Misfit Between Observed and Modeled Bedrock Topography for Various Erosion and Flexural Uplift Scenarios^a

Flexural Uplift Scenario	Minimum Erosion Scenario RMS Misfit (m)	Maximum Erosion Scenario RMS Misfit (m)
Erosional unloading only; continuous elastic plate; $T_e = 20$ km	590	540
Erosional unloading and sediment loading; continuous elastic plate; $T_e = 20$ km	580	530
Erosional unloading and sediment loading; continuous elastic plate; $T_e = 5$ km	640	650
Erosional unloading and sediment loading; continuous elastic plate; $T_e = 50$ km	610	570
Erosional unloading and sediment loading; broken elastic plate; $T_e = 20$ km	570	520
Mechanical unloading only; faults bounding Recovery and Bailey only; $T_e = 20$ km	580	580
Mechanical unloading only; faults bounding Recovery, Bailey, and Slessor; $T_e = 20$ km	620	620
Erosional unloading, sediment loading, and mechanical unloading; faults bounding Recovery and Bailey only; $T_e = 20$ km	330	240

^aThe root-mean-square (RMS) misfit is the average misfit along five parallel and equally spaced 2-D profiles (the five ensemble profiles in Figure 3) crossing the Shackleton Range and Theron Mountains.

We find that irrespective of the assumed erosion and T_e scenario, erosion-driven flexure accounts for ~40–50% of the total elevation (and only ~0.2° of tilting) of the mountain blocks (Figure 7). The misfit between the modeled and observed topography is small on the flanks of the mountain ranges bounding the Slessor trough, but increases toward the flanks of the Recovery and Bailey troughs, where flexure underestimates the topography by up to 800 m (Figure 7).

4.2. Fault-Driven Uplift

Erosional unloading due to the removal of material (rock) from the troughs cannot account for the total observed elevations of the northern Theron Mountains and southern Shackleton Range or the observed tilt of the mountain surfaces (Figure 7). We therefore invoked mechanical unloading due to the unloading of the footwall by normal faults bounding the Bailey and Recovery troughs. For the maximum erosion scenario, the depressions created due to subsidence of the hanging wall blocks were assumed to be filled to sea level. For the minimum erosion scenario, the subsidence was not filled. The throw on the faults was estimated as 600–700 m (section 3.3). Incorporating mechanical unloading on these two major fault systems significantly improved the match between the observed and modeled flexural uplift and tilting of the Shackleton Range and Theron Mountains (Figure 7). Modeled tilts agree very well with the 1.2°N and 0.8°S tilting of the Shackleton Range and Theron Mountains, respectively (Figure 7). We find that erosional unloading accounts for 40–50% of the uplift of the mountains and mechanical unloading accounts for a further 40–50%. There is a small residual misfit; some of the topographic signature is likely the result of nonflexural processes, such as brittle deformation on faults. The maximum erosion scenario, where tectonic subsidence is infilled, produces a better overall fit with the observed topography than the minimum erosion scenario (Figure 7 and Table 1). This suggests fault activity and subsequent sedimentation predated glaciation (section 5.2).

In order to calculate a 34 Ma paleotopography, we restored the eroded material to the ice-rebounded topography and computed and subtracted the associated isostatic response (Figure 8). This calculation was based on the assumption that fault activity mostly predated the onset of Antarctic glaciation at 34 Ma (section 5.2) and has therefore not contributed to mountain uplift or trough subsidence since glacial inception. Since we determined a maximum and a minimum erosion scenario, which differ in their respective assumptions of how deep the troughs were prior to glaciation, we present a minimum and a maximum paleotopography (Figure 8).

Our key finding is that the model scenario that produces a best fit between process-oriented model topography and the observed modern topography requires major contributions from both erosion- and mechanically driven flexure (as well as slope diffusion, which is needed to explain the regrading of the fault scarps). Both processes, operating together, are necessary to achieve a satisfactory agreement with the observed elevation and tilt of the Shackleton Range and Theron Mountains. None of the processes alone can satisfactorily explain these observations. The model results are summarized in Table 1.

5. Discussion

5.1. Other Mechanisms for Shackleton Range and Theron Mountains Uplift

Glacial erosional unloading combined with mechanical unloading provides a simple and elegant model for the asymmetric uplift (tilting) of the Shackleton Range and Theron Mountains. Our results indicate that the




Figure 8. Reconstructions of preglacial topography in the Shackleton Range region. (a) Maximum paleotopography prior to the onset of continental glaciation at 34 Ma. This reconstruction corrects for glacial erosion and the resulting flexure assuming the minimum erosion scenario and no fault movement since 34 Ma. (b) Minimum 34 Ma paleotopography. This reconstruction corrects for glacial erosion and the resulting flexure assuming the maximum erosion scenario and no fault movement since 34 Ma. Bedrock elevations are relative to present-day sea level. The blue lines show the estimated pathways of preglacial river networks. RL = Recovery Lakes, SR = Shackleton Range, TM = Theron Mountains. (c) Profile A-B across the Shackleton Range and Theron Mountains. Black line = present-day (ice free) topography, red line = maximum 34 Ma paleotopography, blue line = minimum 34 Ma paleotopography.

flexural effects of glacial erosion have driven 40–50% of mountain uplift in this region near the Antarctic margin, which is similar to previous findings in the TAM [Stern *et al.*, 2005]. Ongoing glacial erosion of the Recovery, Slessor, and Bailey troughs and associated flexural isostatic uplift also provides a simple explanation for the emergence of the Shackleton Range from beneath the EAIS at 2.5 Ma [Sugden *et al.*, 2014]. Are there other processes that could account for the observed asymmetric pattern of uplift of the Shackleton Range and Theron Mountains (Figure 2)?

Bedrock uplift in the Recovery catchment could be linked to rift flank uplift on the margin of the Jurassic Weddell Sea Rift System. Such a mechanism has been suggested for the early Cenozoic uplift of the TAM on the flank of the West Antarctic Rift System [Stern and ten Brink, 1989; ten Brink *et al.*, 1997]. However, the TAM are very wide (~300 km) for a rift flank, which is in part attributed to a major inferred contrast in T_e across the lithospheric boundary between East ($T_e = 85$ km) and West Antarctica ($T_e = 5$ km) [ten Brink *et al.*, 1997]. In contrast, this study indicates that the Recovery catchment is characterized by T_e values of ~20 km. Moreover, the highest elevations of the Shackleton Range are >300 km from the Filchner Rift, which marks the easternmost extent of the Weddell Sea Rift System [Jordan *et al.*, 2013, 2016] (Figure 1). In addition, the trends of the faults inferred flanking the Shackleton Range are approximately orthogonal to the Weddell Sea Rift System. Together, this suggests that although Jurassic rift flank uplift and passive margin

development can explain the uplift of the Antarctic margin (e.g., in Coats Land north of the Bailey Ice Stream; Figure 1), it is unlikely that they can explain the observed patterns and extent of uplift farther inland in the Recovery catchment.

Another option is that the observed asymmetry in the topography is the result of spatially variable erosion rather than spatially variable uplift. However, this is unlikely to be the case, since the tilt is observed in the mesa/plateau surfaces (Figure 2), which have experienced negligible incision and cut different geological units [Kerr and Hermichen, 1999; Sugden *et al.*, 2014; Krohne *et al.*, 2016]. The presence of surfaces that all tilt away from the region of unloading and have slopes that are not the same as geological dip slopes is strong evidence for flexural tilting [Watts *et al.*, 2000].

Krohne *et al.* [2016] have proposed a model in which thick (up to 3.4 km) sedimentary basins formed in the region following the opening of the Weddell Sea. Post-Jurassic sediments are not observed in the regional outcrops, suggesting that these sequences have been eroded. Could erosion of this overburden, which is not considered in our models, have driven isostatic uplift of the Shackleton Range and Theron Mountains? Erosion of sediments would indeed drive isostatic uplift of the underlying bedrock. However, removal of a spatially uniform overburden is not capable of driving spatially variable (asymmetric) bedrock uplift as is observed. Moreover, if the top of the sedimentary basin was close to sea level [Krohne *et al.*, 2016], sediment erosion could not uplift the top of the bedrock (i.e., the basin floor) to above sea level, because the unload (the sediment) is less dense than the material it displaces (the mantle).

5.2. Timing of Fault Activity

While the timing of glacial incision is well constrained to the last 34 Ma [Coxall *et al.*, 2005; Thomson *et al.*, 2013; Krohne *et al.*, 2016], the timing of fault activity remains a source of uncertainty. The inferred faults that bound the Shackleton Range and Theron Mountains lie approximately parallel to major Pan-African age thrust faults and proposed crustal-scale transpressional shear zones recognized within the Shackleton Range itself and in western Dronning Maud Land [e.g., Jacobs *et al.*, 2015]. Recent thermochronology studies indicate a period of significant exhumation in the Shackleton Range area at approximately 190–180 Ma, which is attributed to the onset of crustal extension in the Weddell Sea Rift System [Jordan *et al.*, 2013, 2016] and widespread mafic magmatism associated with Ferrar Large Igneous Province [Krohne *et al.*, 2016]. A renewed period of exhumation at approximately 120–100 Ma is attributed to a change in spreading direction in the oceanic crust north of the Weddell Sea Rift System, which may have triggered oblique trans-tension onshore [Krohne *et al.*, 2016].

Because 120–100 Ma is the most recent episode of exhumation prior to glaciation at 34 Ma [Krohne *et al.*, 2016], it is the most likely time at which the faults inferred to bound the Shackleton Range and Theron Mountains were last active. The time between the conclusion of fault activity and the onset of glaciation was likely relatively short in order to preserve the (asymmetric) topography associated with faulting (Figure 2). Although the faults may have been moving since 34 Ma, as has been inferred in other regions of East Antarctica [Cianfarra and Salvini, 2016], there is no geological evidence for this in the Shackleton Range region. Furthermore, the presence of fluvial valley slopes flowing toward the Slessor trough close to sea level would appear to rule out significant post-34 Ma tectonic uplift [Sugden *et al.*, 2014]. However, valley incision can have a strong positive feedback on the growth and life span of major range-bounding normal faults in extensional systems [Olive *et al.*, 2014]. The offsets on the range-bounding faults in the Shackleton Range region may therefore, in part, be the result of glacial or preglacial (fluvial) erosion within the troughs. We speculate that the ongoing process of erosion-driven isostatic uplift is accommodated on these faults, as has been suggested for the Lambert Glacier region [Phillips and Läufer, 2009]. The resultant unloading of the footwall by the hanging wall would also contribute to the total flexural uplift, highlighting that the faults were likely the cause and effect of uplift.

5.3. Landscape Evolution

The landscape evolution of the Shackleton Range region since Gondwana breakup was likely dominated initially by rifting in the Weddell Sea (commencing at approximately 180 Ma [Jordan *et al.*, 2016]), uplift of the passive continental margin, and dissection of the landscape by continental river systems [Sugden *et al.*, 2014; Krohne *et al.*, 2016]. With the locus of uplift along the continental margin, it is likely that Jurassic-Cretaceous river systems initially flowed eastward. At some stage, the passive margin was breached at the




Figure 9. Cartoon showing the proposed landscape evolution of the Shackleton Range region. The geometry of the region has been simplified to two elongate, subparallel mountain ranges bounded by three subparallel troughs. (a) Fault activity in the Jurassic/Cretaceous uplifted the Shackleton Range (SR) and Theron Mountains (TM) blocks. The troughs were filled with sediment. Large river networks drained the continental interior, flowing westward toward the Jurassic age passive margin. (b) After 34 Ma, the region was covered by the early East Antarctic Ice Sheet (EAIS). Valley floors subsided due to the loading effect of the ice sheet. (c) By the Quaternary, the EAIS had grown to continental scale, and three large ice streams had excavated large overdeepened troughs. The location of these troughs was controlled by the preexisting fault structure and river networks. Ice flow is from east to west. Erosional unloading in the troughs drove isostatic bedrock uplift of the Shackleton Range and Theron Mountains, causing the peaks to emerge from beneath the EAIS as nunataks [Sugden *et al.*, 2014]. As a result of ice sheet loading and glacial erosion, the floors of the glacial troughs now lie up to 2.5 km below sea level.

location of the present-day confluence of the Recovery, Slessor, and Bailey glaciers; this could have occurred prior to or after glaciation. The modern ice streams exploit this breach today—it is the point through which the entire catchment drains into the Filchner Ice Shelf (Figure 1).

Plate reorganization at approximately 120–100 Ma triggered activity on the faults inferred to flank the Shackleton Range and Theron Mountains and drove exhumation of the region [Krohne *et al.*, 2016]. Our models indicate that the fault systems that drove the majority of mountain uplift were those bounding the Recovery and Bailey troughs. However, magnetic modeling indicates that further upstream the Slessor Glacier is underlain by a sediment-filled half-graben [Shepherd *et al.*, 2006]. Because the topography prior to faulting is unconstrained, our models cannot be used to estimate the total amount of uplift on the faults. Our models suggest that the amount of uplift driven by the Recovery and Bailey faults was ~600–700 m greater than by the Slessor faults. As well as following the location of the preexisting fault systems (see below), the

location of the Slessor Glacier was likely controlled by the flexural downwarping induced by mechanical unloading on the faults bounding the Recovery and Bailey troughs (Figure 7).

Assuming that fault activity had ceased by the Late Cretaceous, significant (500–1000 m) topography must have existed in the Shackleton Range region prior to glaciation (Figures 8 and 9). During or shortly after faulting, the grabens bounded by the faults were likely infilled with sediment [Krohne *et al.*, 2016] and rivers exploited the structurally controlled topography and cut the valley floors to base level [Sugden *et al.*, 2014] (matching our “maximum erosion scenario”; Figure 5). These river networks (Figures 8 and 9) would have flowed westward if the passive margin had been breached by this time; near the head of the Recovery trough, rivers may have drained east into the Recovery Lakes [Bell *et al.*, 2007] (Figure 8).

From 34 Ma onward, the landscape has been shaped significantly by glaciation. The modern landscape of the Recovery catchment bears the hallmarks of selective linear erosion [Sugden and John, 1976] by the EAIS. For selective linear erosion to occur, an existing (lower amplitude) topographic feature must have existed prior to glaciation [Sugden and John, 1976; Wilson *et al.*, 2012]. The Recovery, Bailey, and Slessor glaciers therefore likely exploited preexisting depressions controlled by the faults flanking by the mountains and occupied by rivers prior to glaciation (Figure 9). The focusing of ice through the preexisting troughs initiated a strong positive feedback, whereby the troughs were rapidly overdeepened by fast-flowing, warm-based erosive ice, while the peaks of the neighboring mountain blocks were protected (and isostatically uplifted) beneath slow-moving, cold-based nonerosive ice [Kessler *et al.*, 2008]. We estimate that ~2 km of rock has been eroded from the Recovery, Slessor, and Bailey troughs. This implies long-term average vertical erosion rates of ~0.06 mm/yr, which is consistent with observed erosion rates beneath modern polar glaciers [Koppes *et al.*, 2015].

5.4. Implications for Past Ice Sheet Dynamics

The evolution of the bedrock topography of the Recovery catchment has significant implications for the dynamics and stability of past Antarctic ice sheets. With more subdued topographic relief at 34 Ma (Figure 8), topographic steering of the ice sheet would have been less effective during the early stages of glaciation. Early ice sheets may therefore have simply overridden the mountains and troughs. As the bed within the troughs was progressively overdeepened, topographic steering will have become more effective, allowing ice and subglacial erosion to be focused through the troughs [Kessler *et al.*, 2008]. Gradually, ice thicknesses and flow velocities will have increased in the troughs and decreased over the mountain blocks [Sugden *et al.*, 2014].

By correcting for erosion and erosion-driven uplift, we have shown that the Shackleton Range and Theron Mountains were ~700 m lower at the time of EAIS inception at the Eocene-Oligocene climate transition (34 Ma) than today (Figure 8). Furthermore, the Bailey, Slessor, and Recovery trough floors were likely close to sea level at this time (Figure 8), compared to almost 2.5 km below sea level today. This paleotopography therefore provides a new input for models of early ice sheet initiation and evolution. Crucially, the implication of the reconstructed topography is that the early ice sheets were less responsive to climate and ocean forcing, because the bed was not significantly overdeepened below sea level.

6. Conclusions

In this study, we have presented new compilations of radar and gravity data over the previously largely unexplored Recovery catchment and used the data sets to quantify for the first time the roles of erosion-driven and tectonic uplift. The 2-D forward and 3-D inverse (spectral) modeling indicates that the Recovery catchment is characterized by T_e values of around 20 km. Our 3-D flexural models show that erosion-driven uplift has driven a substantial amount (~700 m) of post-Eocene uplift of the Shackleton Range and Theron Mountains, augmenting the previous study of Sugden *et al.* [2014]. However, the model results show that erosion alone cannot account for the elevation or the tilt of the Shackleton Range and the Theron Mountains. We propose that the Recovery, Slessor, and Bailey glaciers are structurally controlled. The glacially overdeepened troughs were superimposed on preexisting fault-bounded half-grabens that may have been active during Jurassic rifting and Cretaceous intraplate faulting as proposed from independent recent thermochronology studies [Krohne *et al.*, 2016]. Overall, our results indicate that the Shackleton Range and Theron Mountains were likely ~700 m lower and the bounding valley floors were close to sea level at the Eocene-Oligocene climate transition at 34 Ma. This has important implications for developing more robust models of the dynamics and stability of the early EAIS.

Acknowledgments

G.J.G.P. was supported by Natural Environment Research Council UK studentship NE/L002590/1. We thank Xiaoxia Huang and Karsten Gohl for providing their Weddell Sea isopach grids and some fruitful discussions and Nicole Krohne and Frank Lisker for sharing an early version of their thermochronology manuscript. We are grateful for the work carried out by the Institute for Geophysics, University of Texas, who planned and flew the ICEGRAV 2011 survey. Owen King is thanked for his work in picking the bed for the ICEGRAV 2013 survey. We also wish to thank Vincent Strak and an anonymous reviewer for their helpful and constructive comments. Operation IceBridge data products used in this study are available through the IceBridge data portal at the National Snow and Ice Data Center (<http://nsidc.org/icebridge/portal/>). Grids and figures were produced using the Generic Mapping Tools software package [Wessel *et al.*, 2013].

References

- An, M., D. A. Weins, Y. Zhao, M. Feng, A. A. Nyblade, M. Kanao, Y. Li, A. Maggi, and J.-J. L  v  que (2015), S-velocity model and inferred Moho topography beneath the Antarctic plate from Rayleigh waves, *J. Geophys. Res. Solid Earth*, **120**, 359–383.
- Bamber, J. L., F. Ferraccioli, I. Joughin, T. Shepherd, D. M. Rippin, M. J. Siegert, and D. G. Vaughan (2006), East Antarctic ice stream tributary underlain by major sedimentary basin, *Geology*, **34**(1), 33–36.
- Bassett, D., and A. B. Watts (2015), Gravity anomalies, crustal structure, and seismicity at subduction zones: 1. Seafloor roughness and subducting relief, *Geochem. Geophys. Geosyst.*, **16**, 1508–1540, doi:10.1002/2014GC005684.
- Bell, R. E., M. Studinger, C. A. Shuman, M. A. Fahnestock, and I. Joughin (2007), Large subglacial lakes in East Antarctica at the onset of fast-flowing ice streams, *Nature*, **445**, 904–907.
- Brook, D. (1972), Stratigraphy of the Theron Mountains, *Brit. Antarct. Surv. Bull.*, **29**, 67–89.
- Champagnac, J. D., P. Molnar, R. S. Anderson, C. Sue, and B. Delacou (2007), Quaternary erosion-induced isostatic rebound in the western Alps, *Geology*, **35**(3), 195–198.
- Cianfarra, P., and F. Salvini (2016), Origin of the Adventure Subglacial Trench linked to Cenozoic extension in the East Antarctic Craton, *Tectonophysics*, **670**, 30–37.
- Cochran, J. R., and R. E. Bell (2010, updated 2016), *IceBridge Sander AIRGrav L1B Geolocated Free Air Gravity Anomalies, Version 1*, NASA National Snow and Ice Data Center Distributed Active Archive Center, Boulder, Colo.
- Coxall, H. K., P. A. Wilson, H. P  like, C. H. Lear, and J. Backman (2005), Rapid stepwise onset of Antarctic glaciation and deeper calcite compensation in the Pacific Ocean, *Nature*, **433**, 53–57.
- DeConto, R. M., and D. Pollard (2016), Contribution of Antarctica to past and future sea-level rise, *Nature*, **531**, 591–597.
- Diekmann, B., and G. Kuhn (1999), Provenance and dispersal of glacial-marine surface sediments in the Weddell Sea and adjoining areas, Antarctica: Ice-rafting versus current transport, *Mar. Geol.*, **158**(1–4), 209–231.
- Ferraccioli, F., F. Coren, E. Bozzo, C. Zanolla, S. Gandolfi, I. Tabacco, and M. Frezzotti (2001), Rifted(?) crust at the East Antarctic Craton margin: Gravity and magnetic interpretation along a traverse across the Wilkes Subglacial Basin region, *Earth Planet. Sci. Lett.*, **192**, 407–421.
- Ferraccioli, F., P. C. Jones, M. L. Curtis, P. T. Leat, and T. R. Riley (2005a), Tectonic and magmatic patterns in the Jutulstraumen rift (?) region, East Antarctica, as imaged by high-resolution aeromagnetic data, *Earth Planets Space*, **57**, 767–780.
- Ferraccioli, F., P. C. Jones, M. L. Curtis, and P. T. Leat (2005b), Subglacial imprints of early Gondwana break-up as identified from high resolution aerogeophysical data over western Dronning Maud Land, East Antarctica, *Terra Nova*, **17**, 573–579.
- Ferraccioli, F., E. Armadillo, T. A. Jordan, E. Bozzo, and H. F. J. Corr (2009), Aeromagnetic exploration over the East Antarctic Ice Sheet: A new view of the Wilkes Subglacial Basin, *Tectonophysics*, **478**, 62–77.
- Ferraccioli, F., C. A. Finn, T. A. Jordan, R. E. Bell, L. M. Anderson, and D. Damaske (2011), East Antarctic rifting triggers uplift of the Gamburtsev Mountains, *Nature*, **479**, 388–392.
- Ferraccioli, F., R. Forsberg, T. A. Jordan, K. Matsuoka, A. V. Olsen, O. King, and M. E. Ghidella (2014), New aerogeophysical views of crustal architecture in the Recovery frontier of East Antarctica, *Geophys. Res. Abstr.*, **16**, EGU2014–4374.
- Fitzgerald, P. G. (1994), Thermochronologic constraints on post-Paleozoic tectonic evolution of the central Transantarctic Mountains, Antarctica, *Tectonics*, **13**(4), 818–836, doi:10.1029/94TC00595.
- Fogwill, C. J., M. J. Bentley, D. E. Sugden, A. R. Kerr, and P. W. Kubik (2004), Cosmogenic nuclides ¹⁰Be and ²⁶Al imply limited Antarctic Ice Sheet thickening and low erosion in the Shackleton Range for >1 m.y., *Geology*, **32**(3), 265–268.
- Fretwell, P., H. D. Pritchard, D. G. Vaughan, et al. (2013), Bedmap2: Improved ice bed, surface and thickness datasets for Antarctica, *The Cryosphere*, **7**(1), 375–393.322.
- Gasson, E., R. M. DeConto, and D. Pollard (2015), Antarctic bedrock topography uncertainty and ice sheet stability, *Geophys. Res. Lett.*, **42**, 5372–5377, doi:10.1002/2015GL064322.
- Huang, X., K. Gohl, and W. Jokat (2014), Variability in Cenozoic sedimentation and paleo-water depths of the Weddell Sea basin related to pre-glacial and glacial conditions of Antarctica, *Global Planet. Change*, **118**, 25–41.
- Jacobs, J., M. Elburg, A. L. L  ufer, I. C. Kleinmanns, F. Henkes-Kunst, S. Estrada, A. S. Ruppel, D. Damaske, P. Montero, and F. Bea (2015), Two distinct Late Mesoproterozoic/Early Neoproterozoic basement provinces in central/eastern Dronning Maud Land, East Antarctica: The missing link, 15–21  E, *Precambrian Res.*, **265**, 249–272.
- Jordan, T. A., F. Ferraccioli, N. Ross, H. F. J. Corr, P. T. Leat, R. G. Bingham, D. M. Rippin, A. le Brocq, and M. J. Siegert (2013), Inland extent of the Weddell Sea Rift imaged by new aerogeophysical data, *Tectonophysics*, **585**, 137–160.
- Jordan, T. A., F. Ferraccioli, and P. T. Leat (2016), New geophysical compilations link crustal block motion to Jurassic extension and strike-slip faulting in the Weddell Sea Rift System of West Antarctica, *Gondw. Res.*, **42**, 29–48.
- Kalnins, L. M., and A. B. Watts (2009), Spatial variations in effective elastic thickness in the Western Pacific Ocean and their implications for Mesozoic volcanism, *Earth Planet. Sci. Lett.*, **286**(1–2), 89–100.
- Kerr, A. R., and W. D. Hermichen (1999), Glacial modification of the Shackleton Range, Antarctica, *Terra Antarct.*, **6**(3), 353–360.
- Kessler, M. A., R. S. Anderson, and J. P. Briner (2008), Fjord insertion into continental margins driven by topographic steering of ice, *Nat. Geosci.*, **1**(6), 365–369.
- Kirby, J. F. (2014), Estimation of the effective elastic thickness of the lithosphere using inverse spectral methods: The state of the art, *Tectonophysics*, **631**, 87–116.
- Koppes, M., B. Hallet, E. Rignot, J. Mouginot, J. S. Wellner, and K. Boldt (2015), Observed latitudinal variations in erosion as a function of glacier dynamics, *Nature*, **526**, 100–103.
- Krohne, N., F. Lisker, G. Kleinschmidt, A. Kl  gel, A. L. L  ufer, S. Estrada, and C. Spiegel (2016), The Shackleton Range (East Antarctica): An alien block at the rim of Gondwana, *Geol. Mag.*, **1–24**.
- Le Brocq, A. M., A. Hubbard, M. J. Bentley, and J. L. Bamber (2008), Subglacial topography inferred from ice surface terrain analysis reveals a large un-surveyed basin below sea level in East Antarctica, *Geophys. Res. Lett.*, **35**, L16503, doi:10.1029/2008GL034728.
- Leuschen, C., P. Gogineni, F. Rodr  guez-Morales, J. Paden, and C. Allen (2010, updated 2016), *IceBridge MCoRDS L2 Ice Thickness, Version 1*, NASA National Snow and Ice Data Center Distributed Active Archive Center, Boulder, Colo.
- Marsh, P. D. (1985), Ice surface and bedrock topography in Coats Land and part of Dronning Maud Land, Antarctica, from satellite imagery, *Brit. Antarct. Surv. Bull.*, **68**, 19–36.
- McKenzie, D. (2003), Estimating T_e in the presence of internal loads, *J. Geophys. Res.*, **108**(B9), 1–21, doi:10.1029/2002JB001766.
- McKenzie, D., and D. Fairhead (1997), Estimates of the effective elastic thickness of the continental lithosphere from Bouguer and free air gravity anomalies, *J. Geophys. Res.*, **102**(B12), 27,523–27,552, doi:10.1029/97JB02481.

- Olive, J.-A., M. D. Behn, and L. C. Malatesta (2014), Modes of extensional faulting controlled by surface processes, *Geophys. Res. Lett.*, *41*, 6725–6733, doi:10.1002/2014GL061507.
- Patton, H., D. A. Swift, C. D. Clark, S. J. Livingstone, and S. J. Cook (2016), Distribution and characteristics of overdeepenings beneath the Greenland and Antarctic ice sheets: Implications for overdeepening origin and evolution, *Quat. Sci. Rev.*, *148*, 128–145.
- Paxman, G. J. G., A. B. Watts, F. Ferraccioli, T. A. Jordan, R. E. Bell, S. S. R. Jamieson, and C. A. Finn (2016), Erosion-driven uplift in the Gamburtsev Subglacial Mountains of East Antarctica, *Earth Planet. Sci. Lett.*, *452*, 1–14.
- Phillips, G., and A. L. Läufer (2009), Brittle deformation relating to the Carboniferous-Cretaceous evolution of the Lambert Graben, East Antarctica: A precursor for Cenozoic relief development in an intraplate and glaciated region, *Tectonophysics*, *471*(3–4), 216–224.
- Pollard, D., R. M. DeConto, and R. B. Alley (2015), Potential Antarctic Ice Sheet retreat driven by hydrofracturing and ice cliff failure, *Earth Planet. Sci. Lett.*, *412*, 112–121.
- Rignot, E., J. L. Bamber, M. R. van den Broeke, C. Davis, Y. Li, W. J. van de Berg, and E. van Meijgaard (2008), Recent Antarctic ice mass loss from radar interferometry and regional climate modeling, *Nat. Geosci.*, *1*(2), 106–110.
- Rignot, E., J. Mouginot, and B. Scheuchl (2011), Ice flow of the Antarctic Ice Sheet, *Science*, *333*(6048), 1427–1430.
- Rippin, D. M., J. L. Bamber, M. J. Siegert, D. G. Vaughan, and H. F. J. Corr (2003), Basal topography and ice flow in the Bailey/Slessor region of East Antarctica, *J. Geophys. Res.*, *108*(F1), 6008, doi:10.1029/2003JF000039.
- Schoof, C. (2007), Ice sheet grounding line dynamics: Steady states, stability, and hysteresis, *J. Geophys. Res.*, *112*, F03S28, doi:10.1029/2006JF000664.
- Shepherd, T., J. L. Bamber, and F. Ferraccioli (2006), Subglacial geology in Coats Land, East Antarctica, revealed by airborne magnetics and radar sounding, *Earth Planet. Sci. Lett.*, *244*(1–2), 323–335.
- Skidmore, M. J., and P. D. Clarkson (1972), Physiography and glacial geomorphology of the Shackleton Range, *Brit. Antarct. Surv. Bull.*, *30*, 69–80.
- Stern, T. A., and U. S. ten Brink (1989), Flexural uplift of the Transantarctic Mountains, *J. Geophys. Res.*, *94*(B8), 10,315–10,330, doi:10.1029/JB094iB08p10315.
- Stern, T. A., A. Baxter, and P. Barrett (2005), Isostatic rebound due to glacial erosion within the Transantarctic Mountains, *Geology*, *33*(3), 221–224.
- Stewart, J., and A. B. Watts (1997), Gravity anomalies and spatial variations of flexural rigidity at mountain ranges, *J. Geophys. Res.*, *102*(B3), 5327–5352, doi:10.1029/96JB03664.
- Studinger, M., R. E. Bell, W. R. Buck, G. D. Karner, and D. D. Blankenship (2004), Sub-ice geology inland of the Transantarctic Mountains in light of new aerogeophysical data, *Earth Planet. Sci. Lett.*, *220*, 391–408.
- Sugden, D. E., and B. S. John (1976), *Glaciers and Landscape*, Edward Arnold, London, U. K.
- Sugden, D. E., C. J. Fogwill, A. S. Hein, F. M. Stuart, A. R. Kerr, and P. W. Kubik (2014), Emergence of the Shackleton Range from beneath the Antarctic Ice Sheet due to glacial erosion, *Geomorphology*, *208*, 190–199.
- ten Brink, U., R. I. Hackney, S. Bannister, T. A. Stern, Y. Makovsky (1997), Uplift of the Transantarctic Mountains and the bedrock beneath the East Antarctic ice sheet, *J. Geophys. Res.*, *102*(B12), 27,603–27,621.
- Tessensohn, F., G. Kleinschmidt, and W. Buggisch (1999a), Permo-Carboniferous glacial beds in the Shackleton Range, *Terra Antarct.*, *6*(3), 337–344.
- Tessensohn, F., G. Kleinschmidt, F. Talarico, W. Buggisch, A. Brommer, F. Henjes-Kunst, U. Kroner, I. L. Millar, and A. Zeh (1999b), Ross-age amalgamation of East and West Gondwana: Evidence from the Shackleton Range, *Terra Antarct.*, *6*(3), 317–325.
- Thomson, S. N., P. W. Reiners, S. R. Hemming, and G. E. Gehrels (2013), The contribution of glacial erosion to shaping the hidden landscape of East Antarctica, *Nat. Geosci.*, *6*(3), 203–207.
- Turcotte, D. L., and G. Schubert (1982), *Geodynamics: Applications of Continuum Physics to Geological Problems*, John Wiley, New York.
- Vening Meinesz, F. A. (1950), Les Grabens Africains resultant de compression ou de tension dans la croûte terrestre?, *Bull. Inst. R. Colon. Belge*, *21*, 539–552.
- Watts, A. B. (2001), *Isostasy and Flexure of the Lithosphere*, Cambridge Univ. Press, Cambridge.
- Watts, A. B., and E. B. Burov (2003), Lithospheric strength and its relationship to the elastic and seismogenic layer thickness, *Earth Planet. Sci. Lett.*, *213*(1–2), 113–131.
- Watts, A. B., W. S. McKerrow, and E. Fielding (2000), Lithospheric flexure, uplift, and landscape evolution in south-central England, *J. Geol. Soc. London*, *157*(6), 1169–1177.
- Weissel, J. K., and G. D. Karner (1989), Flexural uplift of rift flanks due to mechanical unloading of the lithosphere during extension, *J. Geophys. Res.*, *94*(B10), 13,919–13,950, doi:10.1029/JB094iB10p13919.
- Wessel, P., W. H. F. Smith, R. Scharroo, J. Luis, and F. Wobbe (2013), Generic Mapping Tools: Improved version released, *Eos Trans. AGU*, *94*(45), 409–410, doi:10.1002/2013EO450001.
- Will, T. M., A. Zeh, A. Gerdes, H. E. Frimmel, I. L. Millar, and E. Schmädicke (2009), Palaeoproterozoic to Palaeozoic magmatic and metamorphic events in the Shackleton Range, East Antarctica: Constraints from zircon and monazite dating, and implications for the amalgamation of Gondwana, *Precambrian Res.*, *172*, 25–45.
- Will, T. M., H. E. Frimmel, A. Zeh, P. Le Roux, and E. Schmädicke (2010), Geochemical and isotopic constraints on the tectonic and crustal evolution of the Shackleton Range, East Antarctica, and correlation with other Gondwana crustal segments, *Precambrian Res.*, *180*, 85–112.
- Wilson, D. S., S. S. R. Jamieson, P. J. Barrett, G. Leitchenkov, K. Gohl, and R. D. Larter (2012), Antarctic topography at the Eocene-Oligocene boundary, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, *335*–336, 24–34.