Polarization extinction ratio and polarization dependent intensity noise in long-pulse supercontinuum generation (Conference Presentation)

Chin, Catherine; Engelsholm, Rasmus Dybbro; Moselund, Peter Morten; Feuchter, Thomas; Leick, Lasse; Podoleanu, Adrian; Bang, Ole

Published in:
Proceedings of SPIE

Link to article, DOI:
[10.1117/12.2253478](https://doi.org/10.1117/12.2253478)

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Chin, C., Engelsholm, R. D., Moselund, P. M., Feuchter, T., Leick, L., Podoleanu, A., & Bang, O. (2017). Polarization extinction ratio and polarization dependent intensity noise in long-pulse supercontinuum generation (Conference Presentation). In *Proceedings of SPIE* (Vol. 10089). [100890L] SPIE - International Society for Optical Engineering. Proceedings of SPIE - The International Society for Optical Engineering
<https://doi.org/10.1117/12.2253478>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Polarization extinction ratio and polarization dependent intensity noise in long-pulse supercontinuum generation (Conference Presentation)

Catherine Chin, NKT Photonics A/S (Denmark); **Rasmus Dybbro Engelsholm**, DTU Fotonik (Denmark); **Peter M. Moselund**, **Thomas Feuchter**, **Lasse Leick**, NKT Photonics A/S (Denmark); **Adrian Podoleanu**, Univ. of Kent (United Kingdom); **Ole Bang**, DTU Fotonik (Denmark)

ABSTRACT

We investigate the polarization of supercontinuum generated in nominally non-birefringent silica photonic crystal fibers over the entire spectrum of the source (450-2400 nm). We demonstrate that the degree of polarization varies over the spectrum but that some parts of the spectrum show stable polarization extinction ratios (PER) of over 10 dB. We experimentally demonstrate how the spectrally resolved polarization develops with increasing power and along the length of the nonlinear fiber. The experimental results are compared to numerical simulations of coupled polarization states mimicking the experimental conditions. Subsequently, a single-shot pulse-to-pulse polarization dependent relative intensity noise (PD-RIN) was measured and the noise characteristics were analyzed using long-tailed and rogue wave statistics. To do this, we used a range of 10 nm narrow bandpass filters (BPF) between 550 nm to 2200 nm, and fast photo detectors, to record 800 consecutive pulses. Peaks from these pulses are first extracted, then distribution of their pulse height histogram (PHH) is constructed. Analysis using higher-order moments about the mean (variance, skewness and kurtosis) showed that: (1) around the pump wavelength of 1064nm, the PD-RIN is lowest, PHH exhibits a Gaussian distribution, and higher order moments are zero, (2) further away from pump, PD-RIN increases in parabolic fashion, PHH follows a left-skewed long-tailed Gamma distribution, and higher-order moments increase. Spectrally, the difference of the PD-RIN in the two orthogonal axes increases with PER.

View presentation recording on the SPIE Digital Library:
<http://dx.doi.org/10.1117/12.2253478.5387835751001>