



LCIA framework and cross-cutting issues guidance within the UNEP-SETAC Life Cycle Initiative

Verones, Francesca ; Bare, Jane; Bulle, Cécile ; Frischknecht, Rolf; Hauschild, Michael Zwicky; Hellweg, Stefanie; Henderson, Andrew ; Jolliet, Olivier ; Laurent, Alexis; Liao, Xun

Total number of authors:
23

Published in:
Journal of Cleaner Production

Link to article, DOI:
[10.1016/j.jclepro.2017.05.206](https://doi.org/10.1016/j.jclepro.2017.05.206)

Publication date:
2017

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Verones, F., Bare, J., Bulle, C., Frischknecht, R., Hauschild, M. Z., Hellweg, S., Henderson, A., Jolliet, O., Laurent, A., Liao, X., Lindner, J. P., Maia de Souza, D., Michelsen, O., Patouillard, L., Pfister, S., Posthuma, L., Prado-Lopez, V., Ridoutt, B., Rosenbaum, R. K., ... Fantke, P. (2017). LCIA framework and cross-cutting issues guidance within the UNEP-SETAC Life Cycle Initiative. *Journal of Cleaner Production*, 161, 957-967. <https://doi.org/10.1016/j.jclepro.2017.05.206>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 10'974 words including all affiliations, captions, tables and references

2 **LCIA framework and cross-cutting issues guidance within the UNEP-SETAC Life Cycle**

3 **Initiative**

4 Francesca **Verones**^{1*}, Jane **Bare**², Cécile **Bulle**³, Rolf **Frischknecht**⁴, Michael **Hauschild**⁵,
5 Stefanie **Hellweg**⁶, Andrew **Henderson**⁷, Olivier **Jolliet**⁸, Alexis **Laurent**⁵, Xun **Liao**⁹, Jan Paul
6 **Lindner**¹⁰, Danielle **Maia de Souza**¹¹, Ottar **Michelsen**¹², Laure **Patouillard**¹³, Stephan
7 **Pfister**⁶, Leo **Posthuma**^{14,15}, Valentina **Prado**¹⁶, Brad **Ridoutt**^{17,18}, Ralph K. **Rosenbaum**¹⁹,
8 Serenella **Sala**²⁰, Cassia **Ugaya**²¹, Marisa **Vieira**²², Peter **Fantke**⁵

9 * corresponding author: *francesca.verones@ntnu.no*

10 ¹ Industrial Ecology Programme, Department of Energy and Process Engineering, Norwegian
11 University of Science and Technology (NTNU), NO-7491 Trondheim, Norway

12 ² US EPA, Office of Research and Development, National Risk Management Research
13 Laboratory, 26 West MLK Dr., Cincinnati, OH 45268

14 ³ CIRAI, Ecole des Sciences de la Gestion, Université du Québec À Montréal, 315, rue
15 Sainte-Catherine Est, Montréal, QC, Canada

16 ⁴ treeze Ltd, Kanzleistrasse 4, Uster, Switzerland

17 ⁵ Division for Quantitative Sustainability Assessment, Department of Management
18 Engineering, Technical University of Denmark, Bygningstorvet 116B, 2800 Kgs. Lyngby,
19 Denmark

20 ⁶ ETH Zurich, Institute of Environmental Engineering, 8093 Zürich, Switzerland

21 ⁷ Noblis, 16414 San Pedro Ave, San Antonio, TX 78232

22 ⁸ School of Public Health, University of Michigan, Ann Arbor, MI, United States

23 ⁹ Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

24 ¹⁰ Fraunhofer Institute for Building Physics, Stuttgart, Germany

25 ¹¹ University of Alberta, Department of Agricultural, Food and Nutritional Science, T6G 2P5,
26 Edmonton, Alberta

27 ¹² NTNU Sustainability, Norwegian University of Science and Technology (NTNU), NO-7491
28 Trondheim, Norway

29 ¹³ CIRAI, École Polytechnique de Montréal, P.O. Box 6079, Montréal, Québec H3C 3A7,
30 Canada

31 ¹⁴ RIVM (Dutch National Institute for Public Health and the Environment), Centre for
32 Sustainability, Environment and Health, P.O. Box 1, 3720 BA Bilthoven, the Netherlands

33 ¹⁵ Radboud University Nijmegen, Department of Environmental Science, Institute for Water
34 and Wetland Research, Heyendaalseweg 135, 6525 AJ Nijmegen, The Netherlands

35 ¹⁶ Institute of Environmental Sciences CML, Leiden University, Einsteinweg 2, 2333 CC,
36 Leiden

37 ¹⁷ Commonwealth Scientific and Industrial Research Organisation (CSIRO) Agriculture and
38 Food, Private Bag 10, Clayton South, Victoria 3169, Australia

39 ¹⁸ University of the Free State, Department of Agricultural Economics, Bloemfontein 9300,
40 South Africa

41 ¹⁹ IRSTEA, UMR ITAP, ELSA-PACT – Industrial Chair for Environmental and Social Sustainability
42 Assessment, 361 rue Jean-François Breton, BP 5095, 34196 Montpellier, France

43 ²⁰ European Commission, Joint Research Centre, Directorate D: Sustainable Resource,
44 Bioeconomy unit, Via E. Fermi 2749, Ispra VA, Italy

45 ²¹ Federal University of Technology, Avenida Sete de Setembro, Rebouças Curitiba, Paraná,
46 Brazil

47 ²² PRé Consultants B.V., Stationsplein 121, 3818 LE Amersfoort, Netherlands

48

49 **Abstract**

50 Increasing needs for decision support and advances in scientific knowledge within life cycle
51 assessment (LCA) led to substantial efforts to provide global guidance on environmental life
52 cycle impact assessment (LCIA) indicators under the auspices of the UNEP-SETAC Life Cycle
53 Initiative. As part of these efforts, a dedicated task force focused on addressing several LCIA
54 cross-cutting issues as aspects spanning several impact categories, including spatiotemporal
55 aspects, reference states, normalization and weighting, and uncertainty assessment. Here,
56 findings of the cross-cutting issues task force are presented along with an update of the
57 existing UNEP-SETAC LCIA emission-to-damage framework. Specific recommendations are
58 provided with respect to metrics for human health (Disability Adjusted Life Years, DALY) and
59 ecosystem quality (Potentially Disappeared Fraction of species, PDF). Additionally, we stress
60 the importance of transparent reporting of characterization models, reference states, and
61 assumptions, in order to facilitate cross-comparison between chosen methods and
62 indicators. We recommend developing spatially regionalized characterization models,
63 whenever the nature of impacts shows spatial variability and related spatial data are
64 available. Standard formats should be used for reporting spatially differentiated models, and
65 choices regarding spatiotemporal scales should be clearly communicated. For normalization,
66 we recommend using external normalization references. Over the next two years, the task
67 force will continue its effort with a focus on providing guidance for LCA practitioners on how
68 to use the UNEP-SETAC LCIA framework as well as for method developers on how to
69 consistently extend and further improve this framework.

70

71 **Keywords.** life cycle impact assessment, characterization framework, uncertainty
72 assessment, human health, ecosystem quality, natural resources

73 **Highlights**

- 74 • The existing UNEP-SETAC LCIA framework was updated.
 - 75 • Recommendations were formulated for several LCIA cross-cutting issues.
 - 76 • Recommendations were provided for specific areas of protection.
 - 77 • Continuous efforts will focus on further harmonizing cross-cutting issues in LCIA.
- 78

79 **1. Introduction**

80 Life Cycle Assessment (LCA) is a method for environmental assessment and management,
81 which has evolved to provide decision support. LCA is used for quantifying potential
82 environmental impacts of products, processes, or services. The adverse impacts are usually
83 assessed for several impact categories, such as acidification, eutrophication, and climate
84 change. LCA is often used for comparative studies to support the selection of
85 environmentally preferable alternatives, for eco-design purposes, and for identification of
86 the potentially largest environmental impacts and trade-offs in a product life cycle (Hellweg
87 et al. 2014). The LCA approach has also recently been extended to assessments of
88 organizations (ISO/TS 14072 2014; UNEP et al. 2015), thereby increasing its range of
89 applications and its reach to high-level decision- and policy-makers. Consequently, LCA-
90 based decisions have become more and more relevant for recognizing and reducing
91 environmental impacts of products and processes.

92 Triggered by the increasing needs for reliable decision support and by ongoing advances in
93 scientific knowledge, the UNEP-SETAC Life Cycle Initiative (LC Initiative) has been initiated to
94 improve the science and practices in the field of life cycle thinking (UNEP-SETAC 2016). The
95 LC Initiative has established several task forces, aimed at 1) harmonizing current approaches,
96 2) furthering the development of life cycle impact assessment (LCIA), and 3) providing
97 guidance on recommended models and methods for calculating environmental indicators so
98 that their application provides the best possible transparency, reproducibility, and validity,
99 as well as the best possible support for decision-making.

100 One of these UNEP-SETAC task forces has been addressing LCIA cross-cutting issues, i.e.
101 topics that are relevant across several, or all, of the existing impact categories. The activities
102 of this task force concentrated on the improvement and harmonization of the LCIA

103 characterization framework, and on aspects such as furthering consensus regarding
104 normalization and weighting, spatial differentiation, uncertainty assessment, endpoint
105 indicators for human health, ecosystem quality, and natural resources, as well as the
106 identification of representative reference states.

107 In 2004, the LC Initiative published a recommendation for an LCIA framework, embracing an
108 overview of existing impact categories, and the status of their development (Jolliet et al.
109 2004). Since then, there has been substantial progress in LCIA methods, as well as underlying
110 models and data, both in terms of covered impact pathways, spatial differentiation and
111 resolution, novelties in endpoint indicators, and normalization procedures. It is therefore
112 time to review and evaluate these developments and innovations in a structured way,
113 especially for the damage (endpoint) level, while midpoints are kept as they were described
114 in the 2004 framework. It is the aim of the cross-cutting issues task force to improve the
115 applicability and operationalization of LCIA methods and to integrate scientific advances into
116 the LCIA framework in a compatible and consistent way.

117 In January 2016, a Pellston workshop (i.e. a workshop hosted by the Society for
118 Environmental Toxicology and Chemistry (SETAC) on critical and urgent topics) was
119 conducted in Valencia, Spain, uniting efforts of the cross-cutting issues and other, topical,
120 task forces, which worked on impacts derived from land and water use, exposure to fine
121 particulate matter, and climate change (Frischknecht et al. 2016a). The workshop
122 participants discussed several cross-cutting issues, such as the need to revise the LCIA
123 framework, in order to include recent advances in LCIA science and achieve a more
124 comprehensive coverage of indicators. In addition, recommendations for harmonization of
125 reference states, spatial differentiation, normalization and weighting, uncertainty
126 assessment across impact categories, as well as specific issues for individual areas of
127 protection (e.g. aggregated metrics for damages on human health and on ecosystem quality)
128 were discussed. This paper provides an overview of the current state of development of the
129 previously mentioned cross-cutting issues, and presents expert recommendations. We
130 deliver recommendations that are currently ready for consideration (section 3), and give an
131 outlook where further research and harmonization are needed (section 4).

132 **2. Approach**

133 The task force on cross-cutting issues was established in January 2015, when it started to
134 work on different issues in individual subtasks, as mentioned in the introduction. In late
135 autumn 2015, all active members of the cross-cutting issues task force consolidated findings
136 from the different subtasks into an internal white paper, which served as starting point for
137 proposing recommendations during the Pellston workshop, to which several members of the
138 cross-cutting issues task force but also members from all other guidance project tasks forces
139 were invited along with various sector experts. Discussions between the workshop
140 participants led to the formulation of recommendations, which were presented and
141 discussed in a workshop plenary session, then finalized and agreed upon, and finally
142 published in the official Pellston workshop report in early 2017, complemented with the
143 main content of the initial cross-cutting issues white paper (Frischknecht et al. 2016b).

144 For some of the cross-cutting issues subtasks, participants produced and published final
145 recommendations, while for other subtasks it was decided to collate further analytical
146 reports on the current state-of-the-art, as a foundation for ongoing discussions. In the
147 following, a status is given for each of the subtasks in the cross-cutting issues theme,
148 followed by the outlook. The supporting information (SI, Tables S1 to S3) and Table 2 contain
149 case study results for different production and consumption scenarios of 1kg rice, based on
150 Frischknecht et al. (2016a), to exemplify the compliance of the topical indicators to and
151 relevance of recommendations made for cross-cutting issues.

152 **3. Results and recommendations**

153 The discussions on the cross-cutting issues yielded various results, which are summarized
154 below under separate subjects.

155 3.1. Update to the LCIA framework and damage categories

156 Currently, LCIA analyses result in outputs for three areas of protection for damages on:
157 human health, ecosystem quality and natural resources. The definition of these areas aims
158 to safeguard the values that are considered important to society (Table 1). For instance, the
159 area of protection “human health” uses aggregated morbidity and mortality impacts as an
160 indicator for measuring damages on human health.

161 Various methodological developments over the last decade indicate the need for an update
162 of the existing LCIA framework and the harmonization of the different impact categories

163 within and across areas of protection. There are, for example, damage methods published
164 without midpoint indicators because of the lack of linear relationships between these
165 midpoints and elementary flows, as well as between midpoints and observed damages. Also,
166 for some impact categories no good suggestion for midpoints does currently exist (e.g. land
167 use). This makes it necessary to allow for possibilities beyond modeling the impact pathway
168 via midpoints to damages only (e.g. (Chaudhary et al. 2015; Verones et al. 2016b)).
169 Moreover, research is progressing to include other environmental issues, such as ecosystem
170 services, into LCIA (e.g. (Koellner et al. 2013; Cao et al. 2015; Othoniel et al. 2016)). After the
171 scoping phase of the LC Initiative, ecosystem services appeared as a joint area of protection
172 with natural resources (Jolliet et al. 2014). Thus, after analyzing recent developments, we
173 propose to distinguish between two overarching systems (1: natural systems and, 2: humans
174 and man-made systems) with three different types of values, in order to distinguish the
175 reasons for identifying the different areas of protection more clearly. This leads in total to
176 the identification of six potential areas of protection for consideration in LCIA (Table 1).
177 Natural systems are broadly defined and go beyond the concept of ecosystems, including
178 also immaterial assets, such as natural heritage, whereas humans and man-made systems
179 are defined to only relate to anthropocentric values. “Values” in this context refer to aspects
180 society deems worth protecting and are independent of the terms “values” and “value
181 choices” as used in weighting.

182 The first set of values refers to intrinsic values, i.e. values given for the sake of the existence
183 in itself. For instance, the damage categories human health and ecosystem quality
184 encompass intrinsic values. It is generally recognized that human beings have a right to life
185 on their own, and that non-human species have a value in their existence, i.e., value that
186 would be lost if the species did not exist. A second set of values refers to instrumental
187 values. These encompass values that have a clear utility to humans and are defined from an
188 anthropocentric standpoint. They include, for example, any kind of resource, ecosystem
189 service, or built infrastructure (socio-economic assets) exploitable or otherwise usable by
190 humans. The third set are cultural values. These are again set from a human point of view
191 and refer to spiritual, aesthetic, or recreational dimensions, including cultural and natural
192 heritage. An example is a cultural heritage site (a damage will occur if this site is flooded for

193 a hydropower dam, such as in Turkey, where the damming of the Tigris river risks flooding
194 the ancient city of Hasankeyf (Berkun 2010)).

195 The cross-cutting issues task force is aware that additional work is required (see section 4 on
196 outlook) to further refine the LCIA framework regarding the consideration of damage
197 categories that have not yet sufficiently been addressed in LCA, such as those addressing
198 ecosystem services and cultural and natural heritage. The inclusion of the latter two borders
199 on social LCA. Recommendations on how to avoid potential double-counting of these values
200 will need to be established (Zimdars et al. 2017) when combining environmental and social
201 life cycle indicators (e.g. also considering the loss of an aesthetically-valued species), once
202 methods for assessing impacts on these values have been developed and are operational.
203 Ecosystem services may also contain cultural values (Millennium Ecosystem Assessment
204 2005) and therefore also need to be addressed in a way to avoid double-counting. This is a
205 subject for further discussions.

206 *Table 1: Overview of the human societal values and how damages on these values are measured and the respective links to*
207 *humans/man-made and natural systems.*

	Intrinsic values	Instrumental values	Cultural values
Humans and man-made systems	Human health (measured as damages on humans from morbidity & mortality)	Socio-economic assets (measured as damages on man-made environment such as built infrastructure, loss of cash crops, etc.)	Cultural heritage (measured as damages on buildings, historic monuments, artwork, landscapes, etc.)
Natural systems	Ecosystem quality (measured as damages on ecosystems, i.e. biodiversity loss, by means of species richness & vulnerability)	Natural resources & Ecosystem services (measured as damages on resources, such as exhaustion of mineral primary resources, loss of availability of crops, wood, loss of water flow regulation potentials, etc.)	Natural heritage (measured as damages on flora, fauna, geological elements, etc.)

208

209 In the original UNEP-SETAC LCIA framework (Jolliet et al. 2004) two modeling options are
210 distinguished: 1) modeling up to midpoint impact indicators only, 2) modeling up to damage


211 categories *via* midpoint impact indicators. The direct link between life cycle inventory (LCI)
212 and damage category was not foreseen. A midpoint impact indicator was defined as an
213 indicator “*located on the impact pathway at an intermediate position between the LCI results*
214 *and the ultimate environmental damage*” (Jolliet et al. 2004). However, since then numerous
215 methods, dealing with various impact categories, have been developed that do not contain
216 midpoint impact indicators, but are instead modelled straight to a damage level (e.g. (Souza
217 et al. 2013; Chaudhary et al. 2015; Verones et al. 2016b; Vieira et al. 2016). This is often the
218 case when it is difficult and/or not informative to identify a separately quantifiable midpoint
219 impact indicator for some impact pathways, such as for land use impacts, where in some
220 cases only the area of land being occupied or transformed is provided (inventory parameter)
221 (Vidal-Legaz et al. 2016).

222 It has been common to provide the linkage between combined impact categories at
223 midpoint level and impact categories at damage level with one constant conversion factor
224 for the whole world. However, since 2004, several impact categories have been developed
225 that take spatial differentiation into account (e.g. land use, water use, and freshwater
226 eutrophication). The consideration of spatial differentiation makes it difficult - or even
227 impossible - to apply constant conversion factors, since the cause-effect model from
228 midpoint impact indicator to damage indicator might vary spatially as well, depending on the
229 impact category.

230 Even though midpoint impact indicators may be desirable in some circumstances, they are
231 not required for an impact assessment model, nor are damage level indicators necessary.
232 Models stopping at midpoint level, or models going directly to damage, or models
233 encompassing both, are equally appropriate. As mentioned, traditionally, midpoint impact
234 indicators have been converted to damage indicators via constant conversion factors. We
235 assert explicitly that this is not a fixed requirement, but that instead spatially explicit
236 conversion matrices can be used to improve validity, if the impact category in question
237 contains a relevant spatial aspect. This has, for example, been explained for water impacts,
238 where it is acknowledged that differences between regions matter substantially when
239 considering this indicator (e.g. Pfister et al. (2009)). We are aware that non-globally uniform
240 conversion factors may potentially be leading to different conclusions at the midpoint
241 impact versus the damage level due to the introduction of additional information

242 (variability). The discrepancy reflects that modelling beyond the midpoint introduces
243 relevant additional information and hence that the midpoint result is less environmentally
244 relevant than the damage result. We accept, though do not encourage, that, for the case
245 that no relevant midpoint impact indicator can be identified along the impact pathway,
246 proxy indicators can be designed, which are not defined along an impact pathway itself, such
247 as for example water scarcity indicators (Boulay et al. 2016; Boulay et al. in review). These
248 proxies need to be justified, labelled, and documented to avoid confusion. All in all, the
249 proposed extensions to the LCIA framework as triggered by developments in science and
250 societal concerns leads to an increased comprehensiveness, but also potentially more
251 flexibility in the characterization framework (Figure 1). This has the implication that there is
252 an even greater need than before to transparently report which impact pathway has been
253 modelled up to what level, specifying whether (proxy) midpoint levels have been in- or
254 excluded and providing, if possible, a documentation of their uncertainty.

255 During the Pellston workshop, the topical task forces proposed specific recommendations
256 for indicators and characterization models for land stress, water stress, fine particulate
257 matter formation, and climate change (Frischknecht et al. 2016b). All of these
258 recommendations consistently fit into the recommended updated LCIA framework (Table 1
259 and Figure 1) and highlight the breadth of options and the need for a more flexible
260 framework. Factors for climate change are recommended for a midpoint level only. While
261 this indicator is on the impact pathway for potentially both human health and ecosystem
262 quality, this is not the case for the recommended water scarcity indicator, which is defined
263 as a proxy midpoint. Impacts from exposure to fine particulate matter on human health are
264 defined at both midpoint and damage level, while water use impacts on human health and
265 land stress impacts on ecosystems are defined on a damage level only. For land stress, no
266 operational midpoint indicator is currently available.



267

268 *Figure 1: Updated LCIA framework. The lists of impact categories (on midpoint and damage level) are not complete and are*
 269 *meant to be indicative. Impact characterization models can link the Life Cycle Inventory (LCI) to midpoint impact level*
 270 *(column 2, black dashed lines) and stop there or continue to damage level (column 3, solid black lines), or they can go*
 271 *directly from the life cycle inventory (LCI) to damage level (column 3, grey, dotted line). Similar to midpoint modeling,*
 272 *damage modeling is based on natural science and involves assumptions and choices but is not a weighting step. Note that*
 273 *damage categories are available on a disaggregated level (e.g. climate change, land impacts), or they can be aggregated*
 274 *into overarching categories (column 4, colored lines for existing areas of protection, grey lines for not yet operational ones),*
 275 *if wished. Areas of protection that are operational are indicated with colors, those that are not yet fully operational are*
 276 *shown in the grey box. Weighting of damage category scores may include normalization and is an optional step (in grey)*
 277 *distinct from the damage modeling. Normalization and weighting can also be performed on midpoint impact indicator level.*

278 3.2. Specific recommendation for areas of protection

279 Within each area of protection (aggregated impact categories at damage level), several
 280 different impacts may be combined (such as impacts on human health from toxicity, climate
 281 change and photochemical ozone formation, *i.e.* aggregation over items in the two left hand
 282 side columns in Figure 1). To aggregate, units and metrics need to be consistent among the
 283 categories that are aggregated. Thus, our focus here is on recommendations for the damage
 284 level, in order to make sure that consistent comparisons within areas of protection are
 285 possible. Aggregation into single scores per area of protection may ease the decision-making
 286 process and the communication of the results (fewer indicators have to be communicated),
 287 but may at the same time decrease transparency with respect to uncertainties and trade-
 288 offs among impact categories. Aggregation is a procedure that is commonly applied in LCA
 289 practice, and we include it for the sake of completeness, without advocating that
 290 assessments at damage level need to be aggregated, as this depends on the goal and scope

291 of the study. Whenever aggregated damage level results are used, comparability of metrics
292 used and values addressed by the different areas of protection needs to be ensured, which is
293 therefore an important part of the normalization and weighting subtask. Generally, we want
294 to stress that calculating results at a damage level does not necessarily need to entail an
295 aggregation into a single score per area of protection (note that aggregation across areas of
296 protection relates to normalization and weighting processes, addressed in Section 3.5).

297 In the previous section, we described a potential broadening of areas of protection to
298 consider in environmental decision-making. However, since some of them do not yet exist or
299 are not yet fully evaluated, we will not give recommendations for these at this stage.
300 Instead, we focus on improving the three main established categories, human health,
301 ecosystem quality, as well as natural resources (in color in Figure 1).

302 Human health: Human health is an area of protection that deals with the intrinsic values of
303 human health, addressing both mortality and morbidity. Several impact categories
304 contribute to damages on human health, covering a wide variety of potential impacts. These
305 range from toxic impacts from exposure to substances (e.g., increasing the incidence of
306 cancer) to malnutrition (e.g., water shortages leading to crop shortages leading to
307 malnutrition) to heat stress-related impacts (cardiovascular diseases) associated with
308 greenhouse gas emissions. To compare impacts of these different categories at a damage
309 level (i.e. the net damages on human health), it is crucial to have a common metric. In this
310 respect, human health impact categories generally build on a well-established and widely
311 adopted metric, which is the disability-adjusted life year (DALY) (Murray et al. 1996; Lopez
312 2005; Forouzanfar et al. 2015). We recommend to continue using DALYs in LCIA for human
313 health, as proposed and motivated by Fantke et al. (2015). Topical indicators recommended
314 at the damage level by the LC Initiative follow this recommendation (fine particulate matter,
315 impacts of water use on human health; see illustrative rice case study in SI and Table 2).
316 However, it is recommended that methods use the most recent severity weights originating
317 from the Global Burden of Disease (GBD) study series (Salomon et al. 2012; Salomon et al.
318 2015). This is noteworthy, since the DALYs from the GBD 2010 study (Murray et al. 2012) do
319 not embed age weighting and discounting in their base case anymore (for transparency
320 reasons), which is compatible with the LCIA context. In line with enhancing and moving
321 towards more transparent reporting, we also recommend to document the different

322 components of a DALY separately (e.g., the years of life lost (YLL), the years lived disabled
323 (YLD), and disability weighting).

324 Table 2 illustrates the usage of DALY in a case study on rice produced in different countries.
325 It brings on the same common DALY scale potential impacts of malnutrition due to water use
326 and impacts due to exposure to primary and secondary fine particulate matter. For India,
327 these impacts per kg cooked rice are of similar order of magnitude, with 2.1×10^{-5} to 3.6×10^{-5}
328 DALY/kg_{rice} for water use impacts, and 1.3×10^{-5} DALY/kg_{rice} for PM_{2.5} related impacts, but are
329 lower than the potential reduction in malnutrition impacts of 1.4×10^{-4} DALY/kg_{rice} associated
330 with the production of one kg rice.

331 *Table 2: Results for the human health impact of the functional unit (FU) of 1 kg of white, cooked rice (cooked at home in*
332 *rural India, urban China, or Switzerland). The impact is shown at damage level. Further detail of the case study definition can*
333 *be found in Frischknecht et al. (2016a).*

Impact category	Spatial region/Archetype			
Water use impacts		Inventory [m³/FU]	CF [DALY/m³]	Damage[DALY/FU]
	Average India		4.59E-05	3.58E-05
Rural India	Ganges	0.78	3.80E-05	2.96E-05
	Godavari		2.70E-05	2.11E-05
	Average China		7.31E-05	3.36E-05
Urban China	Yellow River	0.46	1.20E-04	5.38E-05
	Pearl River		4.50E-06	2.07E-06
	Average US		5.63E-05	4.51E-06
US/Switzerland	Red River	0.08	1.30E-06	1.01E-07
	Arkansas River		6.70E-05	5.36E-06
Particulate matter formation (marginal)		Inventory [kg/FU]	CF [DALY/kg]	Damage[DALY/FU]
	Indoor, primary PM _{2.5}	1.71E-03	5.13E-03	8.80E-06
	Rural Outdoor, primary PM _{2.5}	4.36E-04	9.65E-05	4.21E-08
Rural India	Urban Outdoor, primary PM _{2.5}	-	-	-
	NH ₃	6.07E-03	5.04E-04	3.06E-06
	Outdoor, secondary PM _{2.5} : SO ₂	3.32E-03	2.34E-04	7.77E-07
	NO _x	3.49E-03	5.04E-05	1.76E-07
	Indoor, primary PM _{2.5}	-	-	-
	Rural Outdoor, primary PM _{2.5}	3.89E-04	9.65E-05	3.76E-08
Urban China	Urban Outdoor, primary PM _{2.5}	2.25E-04	3.74E-03	8.41E-07
	NH ₃	6.07E-03	5.04E-04	3.06E-06
	Outdoor, secondary PM _{2.5} : SO ₂	3.52E-03	2.34E-04	8.24E-07
	NO _x	3.38E-03	5.04E-05	1.70E-07
	Indoor, primary PM _{2.5}	2.13E-06	1.69E+00	3.60E-06
	Rural Outdoor, primary PM _{2.5}	2.64E-04	9.65E-05	2.54E-08
US/Switzerland	Urban Outdoor, primary PM _{2.5}	1.46E-05	3.74E-03	5.46E-08
	NH ₃	1.50E-03	5.04E-04	7.56E-07
	Outdoor, secondary PM _{2.5} : SO ₂	3.43E-03	2.34E-04	8.04E-07
	NO _x	3.59E-03	5.04E-05	1.81E-07
Particulate matter formation (average)		Inventory [kg/FU]	CF [DALY/kg]	Damage[DALY/FU]
	Indoor, primary PM _{2.5}	1.71E-03	1.66E-02	2.85E-05
Rural India	Rural Outdoor, primary PM _{2.5}	4.36E-04	2.31E-04	1.01E-07

	Urban Outdoor, primary PM _{2.5}	-	-	-
	Outdoor, secondary PM _{2.5} : NH ₃	6.07E-03	5.04E-04	3.06E-06
	SO ₂	3.32E-03	2.34E-04	7.77E-07
	NO _x	3.49E-03	5.04E-05	1.76E-07
	Indoor, primary PM _{2.5}	-	-	-
	Rural Outdoor, primary PM _{2.5}	3.89E-04	2.31E-04	8.97E-08
Urban China	Urban Outdoor, primary PM _{2.5}	2.25E-04	5.29E-03	1.19E-06
	Outdoor, secondary PM _{2.5} : NH ₃	6.07E-03	5.04E-04	3.06E-06
	SO ₂	3.52E-03	2.34E-04	8.24E-07
	NO _x	3.38E-03	5.04E-05	1.70E-07
	Indoor, primary PM _{2.5}	2.13E-06	2.32E+00	4.93E-06
	Rural Outdoor, primary PM _{2.5}	2.64E-04	2.31E-04	6.08E-08
US/Switzerland	Urban Outdoor, primary PM _{2.5}	1.46E-05	5.29E-03	7.72E-08
	Outdoor, secondary PM _{2.5} : NH ₃	1.50E-03	5.04E-04	7.56E-07
	SO ₂	3.43E-03	2.34E-04	8.04E-07
	NO _x	3.59E-03	5.04E-05	1.81E-07

334

335 Ecosystem quality: The area of protection “Ecosystem Quality” deals with damages on the
 336 intrinsic value of natural ecosystems; to date, most models focus on compositional
 337 attributes of biodiversity only, such as species richness (e.g. Goedkoop et al. (2009); (Curran
 338 et al. 2016; Teixeira et al. 2016)). This area of protection encompasses diverse drivers and
 339 pathways of impacts (e.g., water stress, emissions of chemicals leading to eutrophication or
 340 acidification or ecotoxicity). Building consistency across the diverse models in this field is as
 341 important as it is challenging (Curran et al. 2011). However, we stress here that further
 342 research and developments should by no means be stifled by recommendations based on
 343 this paper.

344 Due to the prevalence of indicators for loss of species richness, we currently recommend the
 345 use of potentially disappeared fraction of species (PDF) as a common endpoint metric.
 346 However, the currently-used PDFs only seemingly represent a single metric, while
 347 representing sometimes (widely) different meanings, e.g., when they have been derived
 348 from models based on data from different scales (local, regional, global) or from effects data
 349 on different species groups for different stressors (discussed in Curran et al. (2011)). For
 350 instance, the action of building a parking lot may lead to a very high local loss of species on
 351 the plot occupied (local-scale PDF), but if only regionally and globally abundant species are
 352 lost, the regional-scale and global-scale PDF of the same intervention would be negligible.
 353 This example illustrates that PDFs of different scales should under no circumstances be
 354 mixed without a proper conversion. Also, impacts using different species groups are not to
 355 be mixed without proper consideration (first: recognizing possible differences) or conversion
 356 (second: handling the difference between groups). If other metrics than PDF are used, we

357 recommend providing (preferably validated) conversion factors to PDF. Transparent
358 reporting is also crucial to document the development of PDFs (e.g., which taxonomic groups
359 or spatial locations were considered). Additionally, we recommend that the model
360 developers report PDFs in a disaggregated way (i.e. separately for freshwater, marine and
361 terrestrial ecosystems), and, if applicable, for specific taxonomic groups (i.e., specifically for
362 plants, or invertebrates, when those were used to define a PDF). If possible, to facilitate
363 application, aggregation procedures across taxonomic groups and ecosystems to one final
364 set of values should be made available. First approaches for this exist (e.g. Verones et al.
365 (2015)), but we recommend putting further efforts into researching options for this
366 aggregation. Until consistent aggregation across taxonomic groups is possible, we
367 recommend developing impact indicators for different taxonomic groups separately. The
368 choice of taxonomic groups and modelling approaches should be documented clearly and
369 transparently to facilitate the understanding by practitioners. Impacts on ecosystems, both
370 at regional and global scales, should be reported whenever possible (global levels reporting
371 on irreversible extinction, regional levels being important for preserving ecosystem functions
372 in places where endemism is low) (see also section 3.3). The indicator recommended for
373 land stress is fully aligned with these recommendations (Chaudhary et al. 2015; Frischknecht
374 et al. 2016b). This PDF indicator quantifies both regional losses and global losses, and clearly
375 does so for a set of taxonomic groups, while, for the ease of application, also providing taxa-
376 aggregated characterization factors. Table S1 (SI) illustrates how this indicator applies to the
377 rice case study for the global PDF impacts of land occupation, showing that three types of
378 land occupation dominate the impact of species, i.e., the production (cultivation) of the rice
379 as could be expected, the intensive forest production of wood for cooking in the India
380 scenario and the use of urban area in the US production/Swiss consumption scenario. Other
381 improvements of this indicator (e.g. regarding intensities of land use) are recommended by
382 the land use task force (Milà i Canals et al. 2016), but do not affect the recommendations
383 related to cross-cutting issues.

384 Natural resources and ecosystem services: To date, many impact assessment methods (e.g.
385 (Goedkoop et al. 1999; Jolliet et al. 2003; Goedkoop et al. 2009)) consider a third damage
386 category focusing on resources. This is the only category that so far focuses on “instrumental
387 values” (Table 1). We recommend refining the scope of this damage category to “natural

388 resources” (Sonderegger et al. accepted). As of now there are several different definitions of
389 what should be in- or excluded in such an area of protection (see e.g. the discussion in
390 Dewulf et al. (2015)).

391 Ecosystem services have an instrumental value for humans, and are defined as “*the benefits*
392 *people obtain from ecosystems*” (Millennium Ecosystem Assessment 2005). Thus, ecosystem
393 services can also be seen as a part of the natural resources, but are seldom operationalized
394 in LCIA models at this time. However, the LCIA research community has made first steps
395 towards their inclusion (e.g. (Zhang et al. 2010a; Zhang et al. 2010b; Saad et al. 2013)),
396 including the identification of challenges of doing so (Zhang et al. 2010a; Zhang et al. 2010b;
397 Bare 2011; Othoniel et al. 2016), but further efforts are needed to adequately include the
398 different types of ecosystem services (provisioning, regulating, supporting and cultural) in
399 models with global coverage (models covering only a small spatial unit, such as an individual
400 country or part of an ecoregion are often not applicable in other world regions due to
401 differences in present services and environmental conditions. Therefore, models are
402 required that can deliver individual factors for different world regions).

403 3.3. Guidance on temporal and spatial modelling issues

404 It is becoming increasingly clear that, in various instances, spatial and temporal issues are of
405 utmost relevance in LCIA (Hauschild 2006). For instance, when evaluating water use impacts,
406 the sensitivity of receiving ecosystems towards impacts can vary significantly, and can
407 therefore lead to spatially different characterization factors (CF) (Boulay et al. 2015). Taking
408 global CFs (averages) may lead to over- or underestimations of impacts. Therefore,
409 introducing spatial differentiation (or regionalization) in LCIA models can help improve the
410 accuracy of LCA results (Mutel et al. 2009). The same is true for aggregation of temporal data
411 in the case of water consumption (e.g. Pfister et al. (2014)) and also for photochemical
412 ozone (Shah and Ries 2009; Huijbregts 1998).

413 Spatially differentiated LCIA models and CFs are available in various existing LCIA methods,
414 such as LC-Impact (Verones et al. 2016a), TRACI (Bare 2002), IMPACT World+ (Bulle et al.
415 2012), Ecological Scarcity (Frischknecht et al. 2013), or EDIP (Potting et al. 2004) for either
416 multiple impact categories or single indicators (e.g. water use impacts, eutrophication, land
417 use impacts, toxicity, acidification).

418 For all recommended impact categories except climate change, some kind of spatial
419 differentiation is included, either through the use of spatial archetypes for capturing at the
420 global level relevant variabilities across various urban and rural areas for particulate matter
421 formation or via full inclusion of spatial details on an ecoregion (land stress) or watershed
422 (water scarcity and water consumption impacts) level. Although these spatial aspects are all
423 clearly reported, the data format of characterization factors is often not consistent. The
424 importance of including spatial differentiation in relation to water stress – the impact
425 category with the largest spatial variation in characterization factors - is highlighted in Table
426 S3 (SI) for the illustrative rice case study: Between the Yellow and Pearl watersheds in urban
427 China, there is almost a factor of 200 difference in terms of how scarce water is, and impacts
428 from water consumption on human health vary more than a factor 25. Using a Chinese or
429 global average would underestimate the impact greatly in one case (Yellow river), while
430 overestimating it in the other case (Pearl River). Moving towards including spatial detail is
431 therefore a crucial recommendation for improving environmental assessments. Still, for the
432 ease of application, all topical indicators recommended in the guidance process provided
433 aggregated CFs (country level, for instance) in addition to regionalized ones to also allow for
434 impact characterization when e.g. emission regions are unknown.

435 Spatial variation is also high for human impacts from exposure to fine particulate matter due
436 to variation in population density around the locations of emission or the more than 100
437 times difference in intake fractions between indoor and outdoor releases as function of
438 location. Accounting for such spatial variation based on exact location of emission would
439 require to know the exact emission location and to model the dispersion at a 10 km or
440 higher resolution, which is usually not practical for LCA applications. Table 2 illustrates for
441 the rice case study how such spatial variation can be handled via the definition of
442 characterization factors differentiated by indoor, rural outdoor and urban outdoor
443 archetypes, which can then be linked to present life cycle inventory databases, such as
444 ecoinvent. The exact parameterization of the indoor archetypes can be further customized
445 to the country or continental region of production and consumption, the CFs of Table 2
446 accounting for regional person density and building tightness in each region. In the case of
447 human health impacts of fine particulate matter exposure, archetypes need to not only
448 reflect spatial variation in population density, but also the level of exposure, since the

449 considered dose-response is non-linear and depends on background exposure of the
450 considered individuals.

451 If spatial differentiation is meaningful to the nature of the impact category covered, and if
452 data are available, we recommend developing spatial characterization factors for midpoint
453 and damage impact categories. Spatial differentiation is meaningful, if the potentially
454 “impacted entity” shows clear differences in spatial distribution, such as water scarcity or
455 biodiversity. The geographical resolution should ideally reflect the spatial characteristics of
456 the impacted entity (e.g. watersheds for water consumption impacts, ecoregions for land-
457 use impacts, or population density for human toxicity). The recommended topical indicators
458 fulfill these recommendations (Frischknecht et al. 2016b), as shown in the case study results
459 presented in the SI.

460 In order to facilitate the use of regionalized CF and the interpretation of final LCA results,
461 LCIA method developers should use a standardized format for reporting regionalized CFs.
462 Standards from the Open Geospatial Consortium (OGC 2016) are recommended as a good
463 starting point. For instance, they recommend using the GeoTIFF format for raster data and
464 the GeoPackage Vector format for vector data.

465 Transparent reporting urges a clear specification of all assumptions related to the inclusion
466 of regionalization in LCIA models (e.g., the level of spatial differentiation of input LCIA
467 parameters, the choice for the resulting spatial resolution for spatially differentiated LCIA
468 methods and the way spatially aggregated CFs have been calculated). This is imperative,
469 even if the chosen model has global resolution without regionalized CFs.

470 3.4 Reference states

471 Most impact categories require a baseline scenario, which is commonly referred to as the
472 “reference state.” This can be either a historical situation, a (hypothetical) future state of the
473 environment, a situation in absence of human interventions, a political target situation, or
474 the current situation. A reference state, thus, refers to both time and space. Choices in the
475 reference state may influence the outcome of the characterization factors. However, many
476 LCIA methods do not mention explicitly which reference state they use, which makes it hard
477 for researchers and practitioners to judge whether these models are compatible (referring to
478 the same reference state) or not. We therefore recommend that the choice of reference

479 state be reported transparently and explicitly. Table S4 in the SI summarizes the chosen
480 reference states for all topical indicators recommended. Except for land use, all indicators
481 are using current, fixed situations (e.g. a fixed reference year), and represent a pragmatic
482 approach (i.e. constrained by data availability). Land use defines a “natural” situation as
483 baseline and represents a normative approach (i.e. based on desirability).

484 Regarding modeling procedures, there are also different possibilities, such as modelling
485 marginal or average impacts. Marginal approaches depart from the current situation (i.e.
486 influencing also the choice of reference state) and assess the impact of one additional unit of
487 emission/resource use. Average assessments focus on the difference between the current
488 situation and the background concentration (historical or zero). This also has an implication
489 for the characterization factors and should, for the sake of transparency and user-
490 friendliness for practitioners, be explicitly reported by model developers. Especially
491 regarding emission-based impact categories, we recommend model developers provide both
492 marginal and average characterization factors. The former are useful for practitioners in the
493 case of small changes being assessed (e.g. individual products), while the latter are useful for
494 assessing larger changes in an economy or longer time frames (Huijbregts et al. 2011). The
495 provided CFs for land use and fine particulate matter follow this recommendation, providing
496 both marginal and average CFs. Table 2 compares the marginal and average characterization
497 factors applied in the illustrative rice case study for human health impacts of fine particulate
498 matter exposure. The difference is especially important in the case of indoor emissions from
499 solid fuel combustion with a factor 3 higher average CF than the marginal CF due to the non-
500 linear dose-response with decreasing slope at higher exposure levels. In this particular case
501 of indoor cooking, the average dose-response may be more adequate for LCA decision
502 contexts, since switching to another type of cooking or to low emission cook stoves would
503 reduce exposure by one or several orders of magnitude, which does not correspond any
504 more to a marginal change.

505 3.5. Normalization and weighting

506 To date, there is no recommendation for which normalization or weighting approach should
507 be used. According to the ISO standard 14044 both normalization and weighting are optional
508 steps in LCA (ISO 2006). Normalization has three main purposes, namely 1) checking the
509 plausibility of LCA results (i.e. their magnitude of results), 2) setting the results into

510 perspective by comparing the magnitude of every individual impact category, and,
511 optionally, 3) preparing the results for further weighting by translating them into a common
512 unit. The main purpose of weighting is to facilitate aggregation of indicators and to reflect
513 the preferences of decision-maker(s) and stakeholders in the assessment. Weighting factors
514 can be elicited a number of ways: from direct elicitation of preferences to weighting
515 methods based on policy targets (Huppes et al., 2012). In the end, weighting is typically
516 applied to obtain a single score for the assessment. Normalization and weighting may
517 sometimes also be useful when reporting footprints that cover more than one impact
518 pathway (Ridoutt et al. 2015).

519 A review of the normalization and weighting approaches, including an assessment of their
520 strengths and weaknesses as well as recommendations for their applications and further
521 developments, can be found in Pizzol et al. (2016). Following the outcome of the Pellston
522 workshop, the current recommendation is to favor external normalization approaches in
523 studies that apply normalization, i.e. approaches in which the reference system is
524 independent from or not directly related to the alternatives assessed in the study (e.g.
525 society's background load within a given region or the world). Compared to internal
526 normalization approaches, where the reference system is a function of the assessed
527 alternatives, external approaches are the only ones capable of meeting all three
528 aforementioned purposes. As a subsequent recommendation, wherever possible, LCA
529 practitioners should opt for global instead of regional or national normalization references
530 to avoid the risk of inconsistency between the geographical scopes of the LCI results of the
531 study and that of the inventory behind the normalization references. In a globalized market,
532 LCA studies are typically associated with a geographical scope – and hence LCI results –
533 spread over the entire world. In practice, it is important to note that there are data gaps in
534 current external normalization references, which may lead to biases in the impact results
535 and which the LCA practitioners should be aware of (Heijungs et al. 2006; Laurent et al.
536 2015; Pizzol et al. 2016; Cucurachi et al. 2017). In all cases, a sensitivity analysis should be
537 performed to test the influence of different weighting and normalization approaches, and
538 sources of uncertainties should be clearly identified, described, and discussed by
539 practitioners.

540 3.6. Handling of uncertainties

541 The models underlying each LCIA come with uncertainties, and neglecting these
542 uncertainties may lead to incorrect LCIA interpretations and thus biased decision support.
543 This can be circumvented and made transparent by uncertainty analysis. A complete and
544 fully quantitative uncertainty analysis makes it clear whether predicted median differences
545 for an impact reflect real differences or only reflect a slight (or no) difference (due to
546 overlapping confidence intervals of the items being compared).

547 In the models and data underlying LCA, there are different types of uncertainty, such as
548 parameter uncertainty, model uncertainty, or value choices (Huijbregts 1998; Hertwich et al.
549 2001a; Hertwich et al. 2001b). Although it is clear that uncertainties in models and data
550 exist, LCIA methods rarely report uncertainties for their characterization factors. However,
551 first attempts have been made to quantify chemical-specific uncertainty for characterization
552 results related to certain impact pathways, (e.g. Fantke et al. (2016)), or to provide a generic,
553 quantitative uncertainty estimate for characterization results across chemicals, e.g.
554 Rosenbaum et al. (2008), to propagate parameter uncertainty using a Monte Carlo approach
555 (Roy et al. 2014), or to combine model and parameter uncertainty (Henderson et al. 2017).
556 Because of lack of uncertainty information on CFs, uncertainty of LCIA results is rarely
557 included in LCA reports and publications. If sound and transparent decisions are to be
558 supported, reporting of uncertainties should become a routine practice to avoid over-
559 interpretation and biased decisions. Identifying, qualitatively or even quantitatively
560 describing, and finally documenting uncertainties would also allow highlighting assumptions,
561 data and model components for model developers that need special attention to further
562 improve the LCIA methods. We recommend that model developers and practitioners alike
563 report uncertainties at least in a qualitative way (if a quantitative approach is not possible).
564 This advice is followed by the topical indicators who all discuss uncertainty at least in a
565 qualitative way (Frischknecht et al. 2016b). Explicit 95% confidence intervals are given for
566 the land stress impacts, while others, such as the water scarcity indicator reports results of
567 sensitivity analyses or spatial variability (water consumption impacts on human health,
568 particulate matter related impacts).

569 **4. Outlook**

570 Apart from the issues discussed here, there are still multiple cross-cutting issues that need
571 future research and more comprehensive discussion within the UNEP-SETAC cross-cutting

572 issues task force and with external experts and stakeholders. The task force calls for further
573 discussion and development on issues across all areas of protection (especially those not yet
574 developed, see Figure 1), as well as spatial and temporal issues and uncertainty assessment.
575 Below, we discuss some specific, concrete suggestions, without the ambition to be
576 comprehensive, but as a way to stimulate and suggest priority items for research.

577 Ecosystem quality is an area of protection with a large need for further development.
578 Scientific analyses suggest that a multitude of approaches can be chosen to quantify
579 ecological impacts (e.g., McGill et al. (2015)), warranting close attention to models, metrics
580 and underlying data to define ecological impacts within and across the various impact
581 categories. Apart from completing and improving the coverage of impact pathways, there is
582 a need for increasing the harmonization across impact categories. This includes, for example,
583 thoughts about whether vulnerability measures should be considered. Such measures could
584 include that there are species or ecosystems that are more vulnerable to certain types of
585 interventions than others and that there may be large differences in the importance of
586 different species for the functioning of ecosystems. Impact assessment models that account
587 for several taxonomic groups (e.g. plants, birds and mammals) need to take care to include
588 the differences in species numbers between the groups. Species-rich taxonomic groups tend
589 to dominate the impact assessment, even though they may not be the taxon that is
590 potentially losing the largest fraction of species. Taxonomic groups should not be weighted
591 based on their species richness alone, as this may lead to underestimating impacts on
592 smaller taxonomic groups, whose species may be more threatened. In terms of which
593 species should be used for constructing impact assessment models, we argue that species
594 should be taken into account that are representative for an ecosystem, and its functions and
595 niches, reflecting different levels of threats and endemism.

596 Damage categories related to natural resources and ecosystem services are in need of
597 further development too. However, there is little consensus on how to model impacts and
598 which endpoint indicators to aspire to. Due to the challenges associated with the damage
599 category of natural resources, from definitions to harmonization and coherence in
600 modelling, a dedicated task force will be in place in the next phase (2016-2017) of the UNEP-
601 SETAC flagship project for guidance on LCIA indicators.

602 Further research and development is also needed on how temporally and spatially
603 differentiated LCIA methods can be integrated into LCA approaches and how aggregations
604 across different temporal and spatial scales should take place. Uncertainty related to
605 temporal and spatial variability should be reported for temporally and spatially aggregated
606 CFs. Also, future efforts will focus on developing guidance on which uncertainties should and
607 could be reported quantitatively in LCIA. It is suggested to consider the possibility of
608 assigning a generic uncertainty factor to impact assessment methods that do not provide
609 uncertainty values. Such a generic factor is usually much higher than truly quantified
610 uncertainty values to motivate practitioners and developers to report uncertainty values. If
611 such values can be provided (quantitatively or qualitatively, for example through a Pedigree
612 matrix (Weidema et al. 1996; Fantke et al. 2012), this generic factor will be reduced.

613 For normalization two topics are of interest for further investigation: (i) the Planetary
614 Boundary concept and its integration in LCIA, and (ii) the incorporation of Multi Criteria
615 Decision Analysis (MCDA) methods. The former has recently gained important momentum in
616 environmental assessment and management as it paves the way for developing approaches
617 and tools allowing to benchmark impacts from an analyzed system with absolute thresholds,
618 which should not be exceeded to keep earth systems functioning (Rockstrom et al. 2009).
619 Some early studies have discussed ways of integrating it as part of the characterization, the
620 normalization, or the weighting steps (Fang et al. 2015; Sandin et al. 2015; Bjørn et al. 2016).
621 No consensus currently exists on this aspect and further research that clearly identify the
622 implications of such integration (e.g. uncertainties, applicability to diverse case studies, etc.)
623 are needed before recommendations can be formulated. With respect to Multi Criteria
624 Decision Analysis (MCDA), some methods aiming at improving decision support in
625 comparative LCAs have also been proposed (Benoit et al. 2003; Prado et al. 2012). These
626 methods are typically applied after characterization and require uncertainty information
627 which may not be available to practitioners.

628 **5. Conclusions**

629 The UNEP-SETAC task force on cross-cutting issues in LCIA evaluated an update of the LCIA
630 framework, and worked on harmonizing several other issues, such as regionalization. The
631 evaluations showed latitude for improving LCIA-practices for existing and future indicators.
632 Recommendations are presented with possible improvements on the short and longer term.

633 The improvements will help increase the comprehensiveness as well as the meaningfulness
634 of LCIA outputs for decision-support. The activities of the task force are still ongoing and will
635 focus on further progress towards harmonizing several cross-cutting issues in LCIA.
636 Recommendations made here were followed partly by the topical task forces present at the
637 Pellston workshop (land use, water use, fine particulate matter, climate change) in
638 establishing the consensual indicators. For the LCIA research community our
639 recommendations have three main implications: 1) the call for increased comprehensiveness
640 on the coverage of areas of protection, 2) the call for an improved transparency in model
641 documentation to ease the identification of compatibility among models and indicator
642 results, and 3) an enhanced recognition of the importance of aligning different cross-cutting
643 aspects, such as standards for spatial differentiation and/or how uncertainty is addressed.
644 Recommendations are targeted towards the LCA community in an effort to contribute to
645 improved decision making through the transparent use of LCIA methods.

646 **Disclaimer**

647 The views expressed in this article are those of the authors and do not necessarily represent
648 the views or policies of the organizations to which they belong. The designations employed
649 and the presentation of the material in this publication do not imply the expression of any
650 opinion whatsoever on the part of the UNEP-SETAC Life Cycle Initiative concerning the legal
651 status of any country, territory, city or area or of its authorities, or concerning delimitation of
652 its frontiers or boundaries. Moreover, the views expressed do not necessarily represent the
653 decision or the state policy of the UNEP-SETAC Life Cycle Initiative, nor does citing of trade
654 names or commercial processes constitute endorsement.

655 **Acknowledgements**

656 The authors acknowledge the contribution from the UNEP-SETAC Life Cycle Initiative for
657 funding this activity.

658 **5. References**

- 659 Bare, J. (2011). "Recommendation for land use impact assessment: first steps into framework,
660 theory, and implementation." Clean Technologies and Environmental Policy **13**(1): 7-18.
661 Bare, J. C. (2002). "Traci: The Tool for the Reduction and Assessment of Chemical and Other
662 Environmental Impacts." Journal of Industrial Ecology **6**(3-4): 49-78.

- 663 Benoit, V. and Rousseaux, P. (2003). "Aid for aggregating the impacts in Life Cycle assessment." The
664 International Journal of Life Cycle Assessment **8**(2): 74-82.
- 665 Berkun, M. (2010). "Hydroelectric potential and environmental effects of multidam hydropower
666 projects in Turkey." Energy for Sustainable Development **14**(4): 320-329.
- 667 Bjørn, A., Margni, M., Roy, P.-O., Bulle, C. and Hauschild, M. Z. (2016). "A proposal to measure
668 absolute environmental sustainability in life cycle assessment." Ecological Indicators **63**: 1-13.
- 669 Boulay, A.-M., Bare, J., Benini, L., Berger, M., Lathuillière, M., Manzardo, A., Margni, M., Motoshita,
670 M., Núñez, M., Pastor, A. V., Ridoutt, B., Oki, T., Worbe, S. and Pfister, S. (in review). "The
671 WULCA consensus characterization model for water scarcity footprints: Assessing impacts of
672 water consumption based on available water remaining (AWARE)." Environ. Sci. Technol.
- 673 Boulay, A.-M., Motoshita, M., Pfister, S., Bulle, C., Muñoz, I., Franceschini, H. and Margni, M. (2015).
674 "Analysis of water use impact assessment methods (part A): evaluation of modeling choices
675 based on a quantitative comparison of scarcity and human health indicators." The
676 International Journal of Life Cycle Assessment **20**(1): 139-160.
- 677 Boulay, A.-M., Pfister, S., Motoshita, M., Schencker, U., Benini, L., Gheewala, S., Brito de Figueiredo,
678 M. C. and Harding, K. (2016). Chapter 5: Water use related impacts: Water scarcity and
679 human health effects. Part 1: Water Scarcity. Global Guidance for Life Cycle Impact
680 Assessment Indicators, Volume 1. R. Frischknecht and O. Jolliet. France, Paris, UNEP.
- 681 Bulle, C., Jolliet, O., Humbert, S., Rosenbaum, R. and Margni, M. (2012). "IMPACT World +: A new
682 global regionalized life cycle impact assessment method. ." Proceedings of the International
683 Conference on Ecobalance , Yokohama, Japan.
- 684 Cao, V., Margni, M., Favis, B. D. and Deschênes, L. (2015). "Aggregated indicator to assess land use
685 impacts in life cycle assessment (LCA) based on the economic value of ecosystem services." Journal of Cleaner Production **94**: 56-66.
- 687 Chaudhary, A., Verones, F., De Baan, L. and Hellweg, S. (2015). "Quantifying Land Use Impacts on
688 Biodiversity: Combining Species-Area Models and Vulnerability Indicators." Environ. Sci.
689 Technol. **49**(16): 9987-9995.
- 690 Cucurachi, S., Seager, T. P. and Prado, V. (2017). "Normalization in Comparative Life Cycle
691 Assessment to Support Environmental Decision Making." Journal of Industrial Ecology: n/a-
692 n/a.
- 693 Curran, M., De Baan, L., De Schryver, A., Van Zelm, R., Hellweg, S., Koellner, T., Sonnemann, G. and
694 Huijbregts, M. A. J. (2011). "Toward Meaningful End Points of Biodiversity in Life Cycle
695 Assessment." Environ. Sci. Technol. **45**(1): 70-79.
- 696 Curran, M., De Souza, D. M., Antón, A., Teixeira, R., Michelsen, O., Vidal-Legaz, B., Sala, S. and Milà i
697 Canals, L. (2016). "How well does LCA model land use impacts on biodiversity?—A
698 comparison with approaches from ecology and conservation " Environ. Sci. Technol. **50**:
699 2782-2795.
- 700 Dewulf, J., Benini, L., Mancini, L., Sala, S., Blengini, G. A., Ardente, F., Recchioni, M., Maes, J., Pant, R.
701 and Pennington, D. (2015). "Rethinking the Area of Protection "Natural Resources" in Life
702 Cycle Assessment." Environmental Science & Technology **49**(9): 5310-5317.
- 703 Fang, K., Heijungs, R. and De Snoo, G. R. (2015). "Understanding the complementary linkages
704 between environmental footprints and planetary boundaries in a footprint–boundary
705 environmental sustainability assessment framework." Ecological Economics **114**: 218-226.
- 706 Fantke, P. and Jolliet, O. (2016). "Life cycle human health impacts of 875 pesticides." The
707 International Journal of Life Cycle Assessment **21**(5): 722-733.
- 708 Fantke, P., Jolliet, O., Evans, J. S., Apte, J. S., Cohen, A. J., Hänninen, O. O., Hurley, F., Jantunen, M. J.,
709 Jerrett, M., Levy, J. I., Loh, M. M., Marshall, J. D., Miller, B. G., Preiss, P., Spadaro, J. V., Tainio,
710 M., Tuomisto, J. T., Weschler, C. J. and McKone, T. E. (2015). "Health effects of fine
711 particulate matter in life cycle impact assessment: findings from the Basel Guidance
712 Workshop." The International Journal of Life Cycle Assessment **20**(2): 276-288.

- 713 Fantke, P., Wieland, P., Juraske, R., Shaddick, G., Itoiz, E. S., Friedrich, R. and Jolliet, O. (2012).
714 "Parameterization Models for Pesticide Exposure via Crop Consumption." Environmental
715 Science & Technology **46**(23): 12864-12872.
- 716 Forouzanfar, M. H., Alexander, L., Anderson, H. R., Bachman, V. F., Biryukov, S., et al. (2015). "Global,
717 regional, and national comparative risk assessment of 79 behavioural, environmental and
718 occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a
719 systematic analysis for the Global Burden of Disease Study 2013." The Lancet **386**(10010):
720 2287-2323.
- 721 Frischknecht, R. and Büsser Knöpfel, S. (2013). Swiss Eco-Factors 2013 according to the Ecological
722 Scarcity Method. Methodological fundamentals and their application in Switzerland.
723 Environmental studies no. 1330. , Federal Office for the Environment, Bern, Switzerland.
- 724 Frischknecht, R., Fantke, P., Tschümperlin, L., Niero, M., Antón, A., et al. (2016a). "Global guidance on
725 environmental life cycle impact assessment indicators: progress and case study." The
726 International Journal of Life Cycle Assessment: 1-14.
- 727 Frischknecht, R. and Jolliet, O. (2016b). Global Guidance for Life Cycle Impact Assessment Indicators,
728 Volume 1. Paris, France, United Nations Environment Programme.
- 729 Goedkoop, M., Heijungs, R., Huijbregts, M. A. J., De Schryver, A., Struijs, J. and van Zelm, R. (2009).
730 ReCiPe 2008: A life cycle impact assessment method which comprises harmonised category
731 indicators at the midpoint and endpoint levels. First edition. Report i: Characterization. The
732 Netherlands, Ruimte en Milieu, Ministerie van Volkshuisvesting, Ruimtelijke Ordening en
733 Milieubeheer.
- 734 Goedkoop, M. and Spriensma, R. (1999). The Eco-Indicator 99. A damage oriented method for life
735 cycle impact assessment. Methodology report and Annex, Pré Consultants, Amersfoort, The
736 Netherlands. <http://www.pre.nl/eco-indicator99/>.
- 737 Hauschild, M. (2006). "Spatial Differentiation in Life Cycle Impact Assessment: A decade of method
738 development to increase the environmental realism of LCIA." The International Journal of
739 Life Cycle Assessment **11**(1): 11-13.
- 740 Heijungs, R., Guinée, J., Kleijn, R. and Rovers, V. (2006). "Bias in normalization: Causes, consequences,
741 detection and remedies." The International Journal of Life Cycle Assessment **12**(4): 211.
- 742 Hellweg, S. and Milà i Canals, L. (2014). "Emerging approaches, challenges and opportunities in life
743 cycle assessment." Science **344**(6188): 1109-1113.
- 744 Henderson, A. D., Asselin-Balençon, A. C., Heller, M., Lessard, L., Vionnet, S. and Jolliet, O. (2017).
745 "Spatial Variability and Uncertainty of Water Use Impacts from U.S. Feed and Milk
746 Production." Environmental Science & Technology **51**(4): 2382-2391.
- 747 Hertwich, E. and Hammitt, J. (2001a). "A decision-analytic framework for impact assessment part I:
748 LCA and decision analysis." The International Journal of Life Cycle Assessment **6**(1): 5-12.
- 749 Hertwich, E. G. and Hammitt, J. K. (2001b). "A decision-analytic framework for impact assessment."
750 The International Journal of Life Cycle Assessment **6**(5): 265.
- 751 Huijbregts, M. A. J. (1998). "Application of uncertainty and variability in LCA." The International
752 Journal of Life Cycle Assessment **3**(5): 273-280.
- 753 Huijbregts, M. A. J., Hellweg, S. and Hertwich, E. (2011). "Do We Need a Paradigm Shift in Life Cycle
754 Assessment?" Environ. Sci. Technol. **45**: 3833-3834.
- 755 ISO (2006). Environmental management - Life Cycle Assessment - Requirements and guidelines.
756 International Standard ISO 14044, International Organisation for Standardisation. Geneva,
757 Switzerland.
- 758 ISO/TS 14072 (2014). Environmental management -- Life cycle assessment -- Requirements and
759 guidelines for organizational life cycle assessment. International Organisation for
760 Standardisation. Geneva, Switzerland.
- 761 Jolliet, O., Frischknecht, R., Bare, J., Boulay, A.-M., Bulle, C., et al. (2014). "Global guidance on
762 environmental life cycle impact assessment indicators: findings of the scoping phase." The
763 International Journal of Life Cycle Assessment **19**(4): 962-967.

- 764 Jolliet, O., Margni, M., Charles, R., Humbert, S., Payet, J., Rebitzer, G. and Rosenbaum, R. (2003).
 765 "IMPACT 2002+: A new life cycle impact assessment methodology." The International Journal
 766 of Life Cycle Assessment **8**(6): 324-330.
- 767 Jolliet, O., Müller-Wenk, R., Bare, J., Brent, A., Goedkoop, M., Heijungs, R., Itsubo, N., Peña, C.,
 768 Pennington, D., Potting, J., Rebitzer, G., Stewart, M., de Haes, H. and Weidema, B. (2004).
 769 "The LCIA midpoint-damage framework of the UNEP/SETAC life cycle initiative." The
 770 International Journal of Life Cycle Assessment **9**(6): 394-404.
- 771 Koellner, T., De Baan, L., Beck, T., Brandão, M., Civit, B., Margni, M., Milà i Canals, L., Saad, R., de
 772 Souza, D. M. and Müller-Wenk, R. (2013). "UNEP-SETAC guideline on global land use impact
 773 assessment on biodiversity and ecosystem services in LCA." International Journal of LCA
 774 **18**(6): 1188-1202.
- 775 Laurent, A. and Hauschild, M. Z. (2015). Normalisation in LCA. Life Cycle Impact Assessment. M. Z.
 776 Hauschild and M. A. J. Huijbregts. Springer, Dordrecht NL.
- 777 Lopez, A. D. (2005). "The evolution of the Global Burden of Disease framework for disease, injury and
 778 risk factor quantification: developing the evidence base for national, regional and global
 779 public health action." Globalization and Health **1**(1): 1-8.
- 780 McGill, B. J., Dornelas, M., Gotelli, N. J. and Magurran, A. E. (2015). "Fifteen forms of biodiversity
 781 trend in the Anthropocene." Trends in Ecology & Evolution **30**(2): 104-113.
- 782 Milà i Canals, L., Antón, A., Bauer, C., De Camillis, C., Freiermuth, R., Grant, T., Michelsen, O. and
 783 Stevenson, M. (2016). Chapter 6: Land use related impacts on biodiversity. Global Guidance
 784 for Life Cycle Impact Assessment Indicators, Volume 1. R. Frischknecht and O. Jolliet. Paris,
 785 France, UNEP.
- 786 Millennium Ecosystem Assessment (2005). Ecosystems and Human Well-being. Synthesis. Island
 787 Press, Washington, DC.
- 788 Murray, C. J. L. and Lopez, A. D. (1996). "Evidence-Based Health Policy--Lessons from the Global
 789 Burden of Disease Study." Science **274**(5288): 740-743.
- 790 Mutel, C. L. and Hellweg, S. (2009). "Regionalized Life Cycle Assessment: Computational Methodology
 791 and Application to Inventory Databases." Environ. Sci. Technol. **43**: 5797-5803.
- 792 OGC, (Open Geospatial Consortium)). (2016). "OGC® Standards and Supporting Documents."
 793 Retrieved 27 January, 2016, from <http://www.opengeospatial.org/standards>.
- 794 Othoniel, B., Rugani, B., Heijungs, R., Benetto, E. and Withagen, C. (2016). "Assessment of Life Cycle
 795 Impacts on Ecosystem Services: Promise, Problems, and Prospects." Environmental Science &
 796 Technology **50**(3): 1077-1092.
- 797 Pfister, S. and Bayer, P. (2014). "Monthly water stress: spatially and temporally explicit consumptive
 798 water footprint of global crop production." Journal of Cleaner Production **73**: 52-62.
- 799 Pfister, S., Koehler, A. and Hellweg, S. (2009). "Assessing the Environmental Impacts of Freshwater
 800 Consumption in LCA." Environ. Sci. Technol. **43**(11): 4098-4104.
- 801 Pizzol, M., Laurent, A., Sala, S., Weidema, B., Verones, F. and Koffler, C. (2016). "Normalisation and
 802 weighting in life cycle assessment: quo vadis?" The International Journal of Life Cycle
 803 Assessment: 1-14.
- 804 Potting, J. and Hauschild, M. Z. (2004). "Background for spatial differentiation in life cycle impact
 805 assessment. The EDIP2003 methodology."
- 806 Prado, V., Rogers, K. and Seager, T. P. (2012). Integration of MCDA Tools in Valuation of Comparative
 807 Life Cycle Assessment. Life Cycle Assessment Handbook, John Wiley & Sons, Inc.: 413-431.
- 808 Ridoutt, B., Pfister, S., Manzardo, A., Bare, J., Boulay, A.-M., et al. (2015). "Area of concern: a new
 809 paradigm in life cycle assessment for the development of footprint metrics." The
 810 International Journal of Life Cycle Assessment: 1-5.
- 811 Rockstrom, J., Steffen, W., Noone, K., Persson, A., Chapin, F. S., et al. (2009). "A safe operating space
 812 for humanity." Nature **461**(7263): 472-475.
- 813 Rosenbaum, R. K., Bachmann, T. M., Gold, L. S., Huijbregts, M. A. J., Jolliet, O., Juraske, R., Koehler, A.,
 814 Larsen, H. F., MacLeod, M., Margni, M., McKone, T. E., Payet, J., Schuhmacher, M., van de
 815 Meent, D. and Hauschild, M. Z. (2008). "USEtox - the UNEP-SETAC toxicity model:

- 816 recommended characterisation factors for human toxicity and freshwater ecotoxicity in life
817 cycle impact assessment." *Int. J. Life Cycle Assess.* **13**: 532-546.
- 818 Roy, P.-O., Deschênes, L. and Margni, M. (2014). "Uncertainty and spatial variability in
819 characterization factors for aquatic acidification at the global scale." *Int. J. Life Cycle Assess.*
820 **19**: 882-890.
- 821 Saad, R., Koellner, T. and Margni, M. (2013). "Land use impacts on freshwater regulation, erosion
822 regulation, and water purification: a spatial approach for a global scale level." *The
823 International Journal of Life Cycle Assessment* **18**(6): 1253-1264.
- 824 Salomon, J. A., Haagsma, J. A., Davis, A., de Noordhout, C. M., Polinder, S., Havelaar, A. H., Cassini, A.,
825 Devleeschauwer, B., Kretzschmar, M., Speybroeck, N., Murray, C. J. L. and Vos, T. (2015).
826 "Disability weights for the Global Burden of Disease 2013 study." *The Lancet Global Health*
827 **3**(11): e712-e723.
- 828 Salomon, J. A., Vos, T., Hogan, D. R., Gagnon, M., Naghavi, M., et al. (2012). "Common values in
829 assessing health outcomes from disease and injury: disability weights measurement study for
830 the Global Burden of Disease Study 2010." *The Lancet* **380**(9859): 2129-2143.
- 831 Sandin, G., Peters, G. M. and Svanström, M. (2015). "Using the planetary boundaries framework for
832 setting impact-reduction targets in LCA contexts." *The International Journal of Life Cycle
833 Assessment* **20**(12): 1684-1700.
- 834 Sonderegger, T., Dewulf, J., Drielsma, J., Fantke, P., Maia de Souza, D., Pfister, S., Stoessel, F.,
835 Verones, F., Vieira, M., Weidema, B. P. and Hellweg, S. (accepted). "Towards harmonizing
836 natural resources as an area of protection in Life Cycle Impact Assessment " *International
837 Journal of LCA*(DOI:10.1007/s11367-017-1297-8).
- 838 Souza, D. M., Flynn, D., Rosenbaum, R. K., de Melo Lisboa, H. and Koellner, T. (2013). "Land use
839 impacts on biodiversity: proposal of characterization factors based on functional diversity." *int. J. Life Cycle Assess.* **18**(6): 1231-1242.
- 841 Teixeira, R. F. M., Maia de Souza, D., Curran, M. P., Antón, A., Michelsen, O. and Milà i Canals, L.
842 (2016). "Towards consensus on land use impacts on biodiversity in LCA: UNEP/SETAC Life
843 Cycle Initiative preliminary recommendations based on expert contributions." *Journal of
844 Cleaner Production* **112, Part 5**: 4283-4287.
- 845 UNEP-SETAC. (2016). "Life Cycle Initiative." Retrieved 21 April, 2016, from
846 <http://www.lifecycleinitiative.org/>.
- 847 UNEP, (United Nations Environment Programme) and SETAC, (Society for Environmental Toxicology
848 and Chemistry) (2015). Guidance on Organizational Life Cycle Assessment, UNEP.
- 849 Verones, F., Hellweg, S., Azevedo, L. B., Chaudhary, A., Cosme, N., Fantke, P., Goedkoop, M.,
850 Hauschild, M. Z., Laurent, A., Mutel, C. L., Pfister, S., Ponsioen, T., Steinmann, Z., Van Zelm,
851 R., Verones, F., Vieira, M. and Huijbregts, M. A. J. (2016a). "LC-IMPACT Version 0.5- A
852 spatially differentiated life cycle impact assessment approach " Retrieved 28 April, 2017,
853 from <http://www.lc-impact.eu/>.
- 854 Verones, F., Huijbregts, M. A. J., Chaudhary, A., De Baan, L., Koellner, T. and Hellweg, S. (2015).
855 "Harmonizing the Assessment of Biodiversity Effects from Land and Water Use within LCA."
856 *Environ. Sci. Technol.* **49**(6): 3584-3592.
- 857 Verones, F., Pfister, S., van Zelm, R. and Hellweg, S. (2016b). "Biodiversity impacts from water
858 consumption on a global scale for use in life cycle assessment." *The International Journal of
859 Life Cycle Assessment*: 1-10.
- 860 Vidal-Legaz, B., Sala, S., Antón, A., Maia de Souza, D., Nocita, M., Putman, B. and Texeira, R. F. M.
861 (2016). Land-use related environmental indicators for Life Cycle Assessment: Analysis of key
862 aspects in land use modelling. *JRC Technical Report*. Luxemburg: Publications Office of the
863 European Union.
- 864 Vieira, M. D. M., Ponsioen, T. C., Goedkoop, M. J. and Huijbregts, M. A. J. (2016). "Surplus Ore
865 Potential as a Scarcity Indicator for Resource Extraction." *Journal of Industrial Ecology*.
- 866 Weidema, B. P. and Wesnæs, M. S. (1996). "Data quality management for life cycle inventories—an
867 example of using data quality indicators." *Journal of Cleaner Production* **4**(3-4): 167-174.

- 868 Zhang, Y., Baral, A. and Bakshi, B. R. (2010a). "Accounting for Ecosystem Services in Life Cycle
869 Assessment, Part II: Toward an Ecologically-Based LCA." Environ. Sci. Technol. **44**(7): 2624-
870 2631.
- 871 Zhang, Y., Singh, S. and Bakshi, B. R. (2010b). "Accounting for Ecosystem Services in Life Cycle
872 Assessment, Part I: A Critical Review." Environmental Science & Technology **44**(7): 2232-
873 2242.
- 874 Zimdars, C., Haas, A. and Pfister, S. (2017). "Enhancing comprehensive measures of social impacts in
875 S-LCA by including environmental and economic aspects." International Journal of LCA: DOI:
876 10.1007/s11367-11017-11305-z.
- 877