Anthelmintic effects of forage chicory (*Cichorium intybus*) against free-living and parasitic stages of *Cooperia oncophora*

Pena-Espinoza, Miguel Angel; Williams, Andrew R. ; Thamsborg, Stig M.; Simonsen, Henrik Toft; Enemark, Heidi L.

Published in:
Veterinary Parasitology

Link to article, DOI:
[10.1016/j.vetpar.2017.07.008](https://doi.org/10.1016/j.vetpar.2017.07.008)

Publication date:
2017

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Pena-Espinoza, M. A., Williams, A. R., Thamsborg, S. M., Simonsen, H. T., & Enemark, H. L. (2017). Anthelmintic effects of forage chicory (*Cichorium intybus*) against free-living and parasitic stages of *Cooperia oncophora*. *Veterinary Parasitology*, 243, 204-207. <https://doi.org/10.1016/j.vetpar.2017.07.008>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Anthelmintic effects of forage chicory (*Cichorium intybus*) against free-living and parasitic stages of *Cooperia oncophora*

Miguel Peña-Espinoza^{a,e*}, Andrew R. Williams^b, Stig M. Thamsborg^b, Henrik T. Simonsen^c, Heidi L. Enemark^d

^aNational Veterinary Institute, Technical University of Denmark, Bülowsvej 27, 1870 Frederiksberg C, Denmark

^bDepartment of Veterinary Disease Biology, Faculty of Health and Medical Sciences, University of Copenhagen, Dyrlægevej 100, 1870 Frederiksberg C, Denmark

^cDepartment of Biotechnology and Biomedicine, Technical University of Denmark, Søtofts Plads, 2800 Kongens Lyngby, Denmark

^dNorwegian Veterinary Institute, Ullevålsveien 68, P.O. Box 750 Sentrum, N-0106, Oslo, Norway

^eInstituto de Farmacología y Morfofisiología, Facultad de Ciencias Veterinarias, Universidad Austral de Chile, Valdivia, Chile

*Corresponding author. E-mail: miguel.pena@uach.cl; miguelpenaesinoza@gmail.com

Graphical abstract

Highlights

- Chicory can exert direct anthelmintic effects against *Cooperia oncophora*
- Chicory extracts inhibited the *in vitro* hatching of *C. oncophora* eggs
- Chicory exerted a potent and dose-dependent paralysis of *C. oncophora* adults *ex vivo*
- *In vivo* inactivity of chicory on *C. oncophora* may relate to host-related factors

Abstract

Chicory shows great promise as an anthelmintic forage for grazing ruminants that can reduce reliance on anti-parasitic drugs. Recently, we reported potent anthelmintic effects of chicory-based diets in infected cattle with significant reductions in worm burdens of the abomasal nematode *Ostertagia ostertagi*, whilst no apparent activity was observed against the small intestinal parasite *Cooperia oncophora*. To explore this discrepancy, we investigated direct anthelmintic effects of forage chicory against *C. oncophora* *in vitro*. Chicory leaves (cultivar ‘Spadona’) were extracted with methanol in a Soxhlet apparatus and the resulting extract was purified by solid-phase extraction to concentrate bioactive phytochemicals such as sesquiterpene lactones. *C. oncophora* eggs and adult worms from mono-infected donor calves were exposed to decreasing concentrations of the chicory extract. In an egg hatch assay, the chicory extract induced a marked and dose-dependent inhibition of egg hatching, with 95% inhibition at 2500 µg extract/mL (EC₅₀ = 619 [95% CI: 530 – 722] µg extract/mL). In the adult motility inhibition assays, the chicory extract induced a potent and dose-dependent worm paralysis. At 12 h of incubation, worms exposed to chicory showed a total paralysis at ≥ 500 µg extract/mL, while after 48 h of incubation a complete inhibition of worm motility was observed at ≥ 250 µg extract/mL (EC₅₀ = 80 [95% CI: 67 – 95] µg extract/mL). We have demonstrated that forage chicory can induce potent inhibitory effects on the egg hatching and exert direct anthelmintic activity against parasitic stages of *C. oncophora*. These results suggest that the previously reported absence of *in vivo* effects of chicory towards *C. oncophora* in infected animals may be related with host-mediated factors and/or inhibitory digestive conditions, rather than an inherent inactivity of chicory and its bioactive phytochemicals.

Keywords: Forage chicory; anthelmintic; nematode; *Cooperia oncophora*; cattle

1. Introduction

In the search for novel control strategies of livestock nematodes in a context of expanding parasite drug-resistance, a promising and increasingly explored approach is the inclusion of forages with anthelmintic activity in animal diets (Sandoval-Castro et al., 2012; Hoste et al., 2015). One bioactive forage under investigation is chicory (*Cichorium intybus* L., Asteraceae), well known for its content of several biologically active compounds, most notably the sesquiterpene lactones (SL) (Rees and Harborne, 1985; Simonsen et al., 2013). *In vivo*, chicory feeding has been shown to reduce the burdens of abomasal nematodes in sheep (Scales et al., 1995; Tzamaloukas et al., 2005). Recently, we reported a potent anti-parasitic activity of chicory-based diets against the abomasal cattle nematode *Ostertagia ostertagi*, the most pathogenic bovine parasite in temperate regions (Peña-Espinoza et al., 2016). However, in the same study, the small intestinal nematode *Cooperia oncophora* was not affected by dietary chicory. It is unclear whether this lack of effect is due to an inherent inactivity of chicory towards *C. oncophora*, or if other factors such as the host digestive physiology may affect the availability and/or activity of phytochemicals along the gastrointestinal tract, as recently substantiated for condensed tannins in cattle (Desrues et al., 2017). Previously, we reported that forage chicory extracts have direct and dose-dependent activity against *O. ostertagi* *in vitro* (Peña-Espinoza et al., 2015), but corresponding evidence with *C. oncophora* or other small intestinal nematodes of ruminants is absent. Consequently, the objective of the present study was to investigate whether forage chicory can exert direct *in vitro* anthelmintic effects against free-living and parasitic stages of *C. oncophora*.

2. Materials and methods

2.1 Plant material and extraction procedure

Forage chicory extracts were prepared with chicory leaves collected from an organic dairy farm in Årre, Denmark (55°32'50''N, 8°35'49''E) in early December 2012. Chicory leaves were hand-picked from a permanent pasture sown in spring 2010 containing forage chicory (cultivar [cv.]

Spadona; <10% dry matter in the field), ryegrass and white clover. At collection, all chicory plants were at the vegetative stage. Chicory leaves were stored at -20°C in the dark until extraction. Extracts were prepared following the method described by Foster et al. (2011), with minor modifications. This method enhances the isolation of SL, which are likely the main anthelmintic compounds in chicory (Foster et al., 2011; Peña-Espinoza et al., 2015). Briefly, dried and ground leaf tissue (10 g) was extracted with methanol in a Soxhlet apparatus. The resulting crude extract was evaporated to dryness, dissolved in 48 mL of ethanol/Mili-Q-H₂O/methanol (1/1/1; v/v/v) and loaded into 8×6 mL solid-phase extraction (SPE) tubes (EmporeTM SPE Cartridges, 66873–U Supelco) set in a SPE vacuum manifold. Each SPE tube was further washed with 2×6 mL of the ethanol/Mili-Q-H₂O/methanol solution and the three resulting eluates were pooled. The obtained extract was concentrated to dryness, resuspended in 48 mL Mili-Q-H₂O and loaded into new 6×6 mL SPE tubes. After SPE, all tubes were washed with 2×6 mL Mili-Q-H₂O and SL were eluted with 3×6 mL methanol. The obtained eluates were pooled and dried under reduced pressure. The resulting purified extract was highly viscous and poorly soluble in Mili-Q-H₂O. Therefore, the extract was dissolved in 100% dimethyl sulfoxide (DMSO) at a concentration of 250 mg dry extract/mL (stock solution) and kept at -20°C until use for *in vitro* assays.

2.2 Parasite material

Parasite material for the *in vitro* assays was collected from two nematode-naïve donor calves (4 months-old) mono-infected with 20,000 third-stage larvae of an anthelmintic-susceptible, laboratory propagated strain of *C. oncophora* (kindly provided by Prof. Dr. Janina Demeler, Freie Universität Berlin, Germany). The experiment was approved by the Animal Experiments Inspectorate of the Danish Ministry of Environment and Food (License j. No. 2013-15-2934-00763). Nematode eggs were detected in faeces from both calves after 14 days post-infection. Unembryonated and clean *C. oncophora* eggs were immediately recovered from fresh faeces following the procedure described by Demeler et al. (2012), and then suspended in Mili-Q-H₂O for use in the egg hatch assay. The two

donor calves were euthanised 28 and 29 days post-infection, respectively, to obtain live *C. oncophora* adults for worm motility inhibition assays. Calves were stunned by captive bolt, bled and the digestive tract was immediately removed. The first 5 m of the small intestine were recovered and motile *C. oncophora* adults were isolated from the intestinal content using a modification of the agar-migration method (Christensen et al., 1995) as described in Peña-Espinoza et al. (2015). Isolated nematodes were washed twice in a sterile incubation medium (RPMI 1640 with L-glutamine, Gibco) containing 200 µg/mL streptomycin, 200 U/mL penicillin and 2 µg/mL amphotericin B at 37 °C. After washing, nematodes were kept in Petri dishes with warm incubation medium and used immediately for the *in vitro* assays.

2.3 *In vitro* assays

First, we investigated the effect of chicory on the hatching of *C. oncophora* eggs by the egg hatch assay (EHA). The assay was conducted according to von Samson-Himmelstjerna et al. (2009), with some modifications. The stock solution of chicory extract was serially diluted in Mili-Q-H₂O and five two-fold decreasing concentrations (in triplicates) were prepared in a 24 well-plate. The final test concentrations were: 2500, 1250, 625, 313 and 156 µg dry chicory extract/mL (final concentration 2% DMSO). Wells containing thiabendazole (Sigma-Aldrich, 1 mg/mL, 2% DMSO) and 2% DMSO diluted in Mili-Q-H₂O were also run in triplicates as positive and negative controls, respectively. Immediately after recovery, ~100 unembryonated *C. oncophora* eggs in Mili-Q-H₂O were added to each well. The 24-well plate was incubated at 25 °C for 48 h, and after incubation 100 µL of Lugol iodine was added to each well. The plate was observed under an inverted microscope and a minimum of 100 eggs and first-stage larvae (L1) were counted per well. Egg hatching percentages were calculated for each well as: % egg hatching = $100 \times [(\text{number of L1 per well}) / (\text{total number of eggs} + \text{L1 per well})]$.

Next, we investigated the activity of chicory on the motility of *C. oncophora* adults by the adult motility inhibition assay (AMIA). The assays were performed according to Peña-Espinoza et al.

(2015), with minor changes. Two independent AMIAs were conducted using decreasing concentrations of the chicory extract with *C. oncophora* adults from each calf (i.e. one assay per animal). The chicory extract stock solution was serially diluted in 100% DMSO to obtain six test concentrations. For each assay, 24 well-plates were prepared with the six extract concentrations diluted in the warm sterile incubation media described above. Final tested concentrations in well were: 1000, 500, 250, 125, 60 and 30 µg dry chicory extract/mL media (final concentration 2% DMSO), run in triplicates for each AMIA. Wells containing ivermectin (Sigma-Aldrich, 1 mg/mL, 2% DMSO) and 2% DMSO diluted in the incubation media were run in triplicates as positive and negative controls, respectively. Immediately after recovery, three to four motile *C. oncophora* adults (males and females) were gently added to each well and the plates were incubated at 37 °C. Worm motility was examined at 12, 24 and 48 h after incubation using an inverted microscope and the number of motile and non-motile (no movement detected in 10 s) nematodes per well was recorded. At 24 h of incubation, the incubation media were replaced by fresh media (37 °C) with the same final concentrations run at the start of the assay. At every time point, the number of motile or non-motile worms in the triplicates exposed to the same extract concentration in each AMIA were merged. This resulted in two final replicates per concentration (one replicate per AMIA = per calf) with a larger number of exposed worms to the chicory extract (n= 10 – 12 worms per replicate and concentration) for analysis of motility inhibition. For each AMIA (i.e. replicate) and time point, worm motility percentages per concentration were calculated as: % worm motility = $100 \times [(\text{number of motile worms in all wells of the same concentration})/(\text{total number of worms in all wells of the same concentration})]$.

2.4 Statistical analyses

The tested chicory extract concentrations were log transformed and the hatching/motility percentages were analysed as described by Peña-Espinoza et al. (2015), with bottom values fixed as 0% and top values set as the mean hatching/motility percentages observed in the negative controls.

Dose-response curves and the effective extract concentrations needed to inhibit the hatching/motility in 50% of incubated eggs/adults (EC_{50}) were calculated, including the 95% confidence interval (CI) and R squared measure of goodness of fit (R^2) for each dose-response curve. All analyses were performed in GraphPad Prism® version 6.05.

3. Results and discussion

In the EHA, the mean egg hatching rate in negative control wells after 48 h was 90%. In contrast, the chicory extract displayed a marked and dose-dependent inhibition of egg hatching (Fig. 1a), with 95% of eggs unhatched at the highest chicory concentration and an $EC_{50} = 619$ (95% CI: 530 – 722) μg dry extract/mL ($R^2 = 0.97$). In addition, considerable damage was observed in dead L1 inside unhatched eggs incubated with chicory, in comparison with unexposed eggs (Fig. 1b–d). Our results showed a higher anthelmintic potency of the investigated chicory (cv. Spadona) than in a previous EHA testing the effects of two different chicory varieties (cv. Grasslands Puna and cv. Forage Feast) on the hatching of a mainly *Haemonchus contortus* egg population, with reported EC_{50} values between 2600 – 6400 μg extract/mL (Foster et al., 2011). While the same extraction method was used in the cited work and in our present investigation, the variation in anthelmintic potencies between the studies is likely explained by differences in the tested chicory cultivars, cultivation conditions and nematode species.

In the AMIA, *C. oncophora* adults in negative control wells showed 100% motility until 48 h of incubation. Contrastingly, the chicory extract induced a potent and dose-dependent inhibition of worm motility at 12, 24 and 48 h of incubation (Fig. 2), with the highest extract concentration achieving total worm paralysis at all time points in both AMIA. Furthermore, a time-dependent inhibitory effect of the extract was noticed, illustrated by decreasing EC_{50} values as the incubation time progressed. At 12 h of incubation, worms exposed to chicory showed a total motility inhibition at concentrations ≥ 500 μg dry extract/mL with an $EC_{50} = 189$ (95% CI: 184 – 194) μg dry extract/mL ($R^2 = 0.99$), whereas at 24 h of incubation an $EC_{50} = 150$ (95% CI: 147 – 152) μg dry

extract/mL ($R^2 = 0.99$) was observed. After 48 h of incubation, a complete worm paralysis was observed at concentrations ≥ 250 μg dry extract/mL with an $\text{EC}_{50} = 80$ (95% CI: 67 – 95) μg dry extract/mL ($R^2 = 0.98$). This direct and potent *in vitro* effect of chicory against *C. oncophora* adults contrasts with the lack of any anthelmintic activity on *C. oncophora* in chicory-fed cattle (~74% of chicory dry matter in the diet; Peña-Espinoza et al., 2016). Previous studies have also shown no *in vivo* activity of chicory feeding on small intestinal nematode burdens in sheep (Marley et al., 2003; Athanasiadou et al., 2005; Heckendorn et al., 2007). Our present *in vitro* results suggest that this apparent absence of *in vivo* effects against *C. oncophora*, and perhaps towards other small intestinal worms, may not be explained by an intrinsic inactivity of chicory and its bioactive phytochemicals against this particular nematode per se. Instead, the observed *in vivo* inactivity of chicory against small intestinal nematodes could be related with the digestion processes and/or the intestinal physico-chemical environment in the host (i.e. affecting the concentration and/or composition of the active compounds). Recent studies from our group have reported that dietary condensed tannins ingested by cattle can be present at much higher concentrations in the abomasum than in the small intestine (Desrues et al., 2017), which could explain the anthelmintic effects of a tannin-rich diet against *O. ostertagi* but not towards *C. oncophora* (Desrues et al., 2016). To date, the pharmacokinetics of SL or other chicory bioactive compounds in the ruminant gastrointestinal tract are poorly understood, but previous studies have suggested that SL are more stable at low pH (Ferreira and Gonzalez, 2008; Saroglou et al., 2008). Certainly, further research is needed to unveil the nematocidal compounds of chicory and to elucidate how these molecules can be affected by the host metabolism in different gut compartments.

Our previous studies with chicory cv. Spadona confirmed the presence of several SL and their dihydro-derivatives in tested leaf extracts (Peña-Espinoza et al., 2015; Williams et al., 2016).

However, high-performance liquid chromatography analysis of the current chicory extract did not allow the characterisation of individual compounds (data not shown). Nevertheless, the potent and

dose-dependent *in vitro* activity of the current chicory (cv. Spadona) extract against *C. oncophora* correlates with our previous results with *O. ostertagi* (Peña-Espinoza et al., 2015), and also with recent findings of a direct activity of chicory (cv. Spadona) against swine nematodes (Williams et al., 2016), indicating that chicory has a broad nematocidal effect *in vitro* that warrants the further elucidation of the major anthelmintic compound(s).

In conclusion, we have demonstrated that forage chicory has direct and dose-dependent inhibitory effects on the egg hatching and adult worm motility in free-living and parasitic stages of *C. oncophora*, respectively. These potent *in vitro* anthelmintic effects suggest that the absence of *in vivo* effects of chicory feeding towards *C. oncophora* may be associated with host-mediated factors and/or inhibitory digestive conditions, rather than an inherent inactivity of chicory and its bioactive phytochemicals.

Conflicts of interest statement

The authors declare that they have none conflicts of interest.

Acknowledgements

The EMIDA ERA-NET project ‘Coping with anthelmintic resistance in ruminants’ (CARES; 3405-11-0430/32), CONICYT Chile (Becas Chile Scholarship and FONDECYT Postdoctorado #3170875) and the Danish Council for Independent Research (project no. DFF-6111-00394) are acknowledged for financial support.

References

- Athanasiadou, S., Tzamaloukas, O., Kyriazakis, I., Jackson, F., Coop, R.L., 2005. Testing for direct anthelmintic effects of bioactive forages against *Trichostrongylus colubriformis* in grazing sheep. *Vet. Parasitol.* 127, 233–243.
- Christensen, C., Barnes, E., Nansen, P., Roepstorff, A., Slotved, H., 1995. Experimental *Oesophagostomum dentatum* infection in the pig: worm populations resulting from single infections with three doses of larvae. *Int. J. Parasitol.* 25, 1491–1498.

- Demeler, J., Kleinschmidt, N., Küttler, U., Koopmann, R., von Samson-Himmelstjerna, G., 2012. Evaluation of the egg hatch assay and the larval migration inhibition assay to detect anthelmintic resistance in cattle parasitic nematodes on farms. *Parasitol. Int.* 61, 614–618.
- Desrues, O., Peña-Espinoza, M., Hansen, T.V.A., Enemark, H.L., Thamsborg, S.M., 2016. Anti-parasitic activity of pelleted sainfoin (*Onobrychis viciifolia*) against *Ostertagia ostertagi* and *Cooperia oncophora* in calves. *Parasit. Vectors* 9, 329. doi:10.1186/s13071-016-1617-z
- Desrues, O., Mueller-Harvey, I., Pellikaan, W.F., Enemark, H.L., Thamsborg, S.M., 2017. Condensed tannins in the gastrointestinal tract of cattle after sainfoin (*Onobrychis viciifolia*) intake and their possible relationship with anthelmintic effects. *J. Agric. Food Chem.* 65, 1420–1427.
- Ferreira, J., Gonzalez, J., 2008. Chemical and biological stability of artemisinin in bovine rumen fluid and its kinetics in goats (*Capra hircus*). *Brazil J. Vet. Parasitol.* 17, 103–109.
- Foster, J., Cassida, K., Turner, K., 2011. In vitro analysis of the anthelmintic activity of forage chicory (*Cichorium intybus* L.) sesquiterpene lactones against a predominantly *Haemonchus contortus* egg population. *Vet. Parasitol.* 180, 298–306.
- Heckendorn, F., Häring, D.A., Maurer, V., Senn, M., Hertzberg, H., 2007. Individual administration of three tanniferous forage plants to lambs artificially infected with *Haemonchus contortus* and *Cooperia curticei*. *Vet. Parasitol.* 146, 123–134.
- Hoste, H., Torres-Acosta, J.F.J., Sandoval-Castro, C., Mueller-Harvey, I., Sotiraki, S., Louvandini, H., Thamsborg, S.M., Terrill, T.H., 2015. Tannin containing legumes as a model for nutraceuticals against digestive parasites in livestock. *Vet. Parasitol.* 212, 5–17.
- Marley, C.L., Cook, R., Keatinge, R., Barrett, J., Lampkin, N., 2003. The effect of birdsfoot trefoil (*Lotus corniculatus*) and chicory (*Cichorium intybus*) on parasite intensities and performance of lambs naturally infected with helminth parasites. *Vet. Parasitol.* 112, 147–155.
- Peña-Espinoza, M., Boas, U., Williams, A.R., Thamsborg, S.M., Simonsen, H.T., Enemark, H.L.,

2015. Sesquiterpene lactone containing extracts from two cultivars of forage chicory (*Cichorium intybus*) show distinctive chemical profiles and in vitro activity against *Ostertagia ostertagi*. *Int. J. Parasitol. Drugs Drug Resist.* 5, 191–200.
- Peña-Espinoza, M., Thamsborg, S.M., Desrués, O., Hansen, T.V.A., Enemark, H.L., 2016. Anthelmintic effects of forage chicory (*Cichorium intybus*) against gastrointestinal nematode parasites in experimentally infected cattle. *Parasitology* 143, 1279–1293.
- Rees, S., Harborne, J., 1985. The role of sesquiterpene lactones and phenolics in the chemical defence of the chicory plant. *Phytochemistry* 24, 2225–2231.
- Sandoval-Castro, C.A., Torres-Acosta, J.F.J., Hoste, H., Salem, A.Z.M., Chan-Pérez, J.I., 2012. Using plant bioactive materials to control gastrointestinal tract helminths in livestock. *Anim. Feed Sci. Technol.* 176, 192–201.
- Saroglou, V., Demetzos, C., Skaltsa, H., 2008. Effects of pH and temperature on the stability of sesquiterpene lactones. *Planta Med* 74–PC102. doi:10.1055/s-0028-1084620
- Scales, G.H., Knight, T.L., Saville, D.J., 1995. Effect of herbage species and feeding level on internal parasites and production performance of grazing lambs. *New Zeal. J. Agric. Res.* 38, 237–247.
- Simonsen, H., Weitzel, C., Christensen, S., 2013. Guaianolide sesquiterpenoids: pharmacology and biosynthesis, in: Ramawat, K., Mérillon, J. (Eds.), *Natural Products*. Springer-Verlag Berlin Heidelberg, pp. 3069–3098.
- Tzamaloukas, O., Athanasiadou, S., Kyriazakis, I., Jackson, F., Coop, R.L., 2005. The consequences of short-term grazing of bioactive forages on established adult and incoming larvae populations of *Teladorsagia circumcincta* in lambs. *Int. J. Parasitol.* 35, 329–335.
- von Samson-Himmelstjerna, G., Coles, G.C., Jackson, F., Bauer, C., Borgsteede, F., Cirak, V.Y., Demeler, J., Donnan, A., Dorny, P., Epe, C., Harder, A., Höglund, J., Kaminsky, R., Kerboeuf, D., Küttler, U., Papadopoulos, E., Posedi, J., Small, J., Várady, M., Vercruysse, J., Wirtherle,

N., 2009. Standardization of the egg hatch test for the detection of benzimidazole resistance in parasitic nematodes. *Parasitol. Res.* 105, 825–834.

Williams, A.R., Peña-Espinoza, M., Boas, U., Simonsen, H.T., Enemark, H.L., Thamsborg, S.M., 2016. Anthelmintic activity of chicory (*Cichorium intybus*): in vitro effects on swine nematodes and relationship to sesquiterpene lactone composition. *Parasitology* 143, 770–777.

Figure captions

Fig. 1 (a) Dose–response curve obtained in an egg hatch assay (EHA) with *Cooperia oncophora* eggs incubated for 48 h at different concentrations of chicory extract ([log] μg dry extract/mL). Error bars represent S.D. between replicates. Data points with no error bars indicate that the variation among values was 0 (or close to 0); (b) Unembryonated *Cooperia oncophora* egg before exposure to chicory extract; (c) Newly hatched first stage larvae (L1) after 48 h of incubation (25 °C) in 2% DMSO (negative control); (d) Unhatched embryonated egg with a dead L1 after 48 h of incubation (25 °C) with chicory extract at 625 μg dry extract/mL

Fig. 2. Dose–response curves obtained in adult motility inhibition assays (AMIA) with *Cooperia oncophora* at 12, 24 and 48 h after incubation with different concentrations of chicory extract ([log] μg dry extract/mL). Each data point represents the mean worm motility rates of 2 replicates (one replicate per AMIA; $n = 10 - 12$ worms per replicate) for each concentration (see text for further information). Error bars represent S.D. between replicates. Data points with no error bars indicate that the variation among values was 0 (or close to 0)

Fig. 1

Fig. 2