Normal and Inverse Diffusive Isotope Fractionation of Deuterated Toluene and Benzene in Aqueous Systems

Rolle, Massimo; Jin, Biao

Published in:
Environmental Science & Technology Letters

Link to article, DOI:
[10.1021/acs.estlett.7b00159](https://doi.org/10.1021/acs.estlett.7b00159)

Publication date:
2017

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Rolle, M., & Jin, B. (2017). Normal and Inverse Diffusive Isotope Fractionation of Deuterated Toluene and Benzene in Aqueous Systems. *Environmental Science & Technology Letters*, 4(7), 298-304.
<https://doi.org/10.1021/acs.estlett.7b00159>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

This is a Post Print of the article published on line 31st May 2017 in Environmental Science & Technology Letters. The publishers' version is available at the permanent link: doi: [10.1021/acs.estlett.7b00159](https://doi.org/10.1021/acs.estlett.7b00159)

Normal and Inverse Diffusive Isotope Fractionation of Deuterated Toluene and Benzene in Aqueous Systems

Massimo Rolle^{*,1} and Biao Jin¹

¹Department of Environmental Engineering, Technical University of Denmark, Bygningstorvet

Building 115, DK-2800 Kgs. Lyngby, Denmark

Corresponding author phone: +45 45251566; e-mail: masro@env.dtu.dk

1 **Abstract**

2 Diffusive isotope fractionation of organic contaminants in aqueous solution is difficult to quantify
3 and only a few experimental datasets are available for compounds of environmental interest. In this
4 study, we investigate diffusive fractionation of perdeuterated and non-deuterated benzene and
5 toluene. Multitracer experiments were carried out in 1-D gel dissection tubes and in a quasi 2-D
6 flow-through porous medium. The experiments allowed us to simultaneously and directly compare
7 the diffusive and dispersive behavior of benzene and toluene. We observed an unexpected, opposite
8 behavior of the two monoaromatic hydrocarbons. Toluene showed a normal diffusive isotope effect
9 ($D_{C_7D_8}/D_{C_7H_8}=0.96$) with enrichment of the non-deuterated isotopologue in the direction of the
10 diffusive and transverse dispersive fluxes. Conversely, the measured trends for benzene indicate
11 inverse diffusive fractionation ($D_{C_6D_6}/D_{C_6H_6}=1.02$), with a remarkably faster diffusion rate of the
12 perdeuterated isotopologue that was enriched in the downgradient portion of the diffusion tubes and
13 at the fringes of the contaminant plumes in the flow-through setup. These outcomes can neither be
14 interpreted as mass-dependent fractionation nor be described as purely hydrodynamic (i.e., mass
15 independent) effects. The results of this study are relevant for the use of labeled/non-labeled
16 mixtures of organic compounds as conservative and (bio)reactive tracers in environmental
17 applications.

18

19 **Introduction**

20 Aromatic hydrocarbons are widespread contaminants, frequently found in soils and aquatic
21 environments. Among these chemicals BTEX (i.e., benzene, toluene, ethylbenzene and xylene)
22 compounds are of particular concern due to their relatively high solubility, mobility and toxicity.¹
23 Mixtures of labeled and nonlabeled compounds have been often applied as a diagnostic tool to
24 understand and quantify contaminant transport and transformation mechanisms. For instance, in the
25 field of groundwater contamination, deuterium-labeled BTEX compounds have been used in tracer
26 tests, push-pull tests and in situ microcosms to evaluate contaminant retardation and in situ rates of
27 transformation during groundwater bioremediation and natural attenuation.²⁻⁶ One important aspect
28 that has not (yet) been evaluated in detail is the extent of diffusive isotope fractionation of BTEX
29 compounds in aqueous solution and its implications in environmental systems. The evaluation of
30 such effect is important since recent investigation of subsurface solute transport has highlighted the
31 key controlling role of aqueous diffusion for groundwater contaminant transport at different
32 scales.⁷⁻¹⁰ Despite the increased recognition of the quantitative importance and of the macroscopic
33 impact of small scale diffusive processes on large scale transport of organic contaminants, only a
34 few experimental and modeling studies have attempted to quantify diffusive isotope fractionation
35 for organic compounds.¹¹⁻¹⁶ The lack of data and mechanistic understanding of organic chemicals'
36 diffusive isotope fractionation becomes apparent when compared with the advances in the related
37 field of inorganic isotope geochemistry, in which numerous studies have been carried out to
38 investigate diffusive isotope effects of major cations, anions and dissolved gases in both aqueous
39 solutions¹⁷⁻²⁹ and non-aqueous systems.³⁰⁻³³

40 In this work we focus on diffusive transport of perdeuterated and non-deuterated benzene and
41 toluene. The diffusive behavior of these chemicals (particularly of benzene) was investigated in
42 early studies,³⁴⁻³⁶ mostly as self-diffusion or as tracer diffusion in organic solvents. However, to the

best of our knowledge, no study has reported a comparison of diffusive isotope fractionation of these contaminants in aqueous solution. The purpose of this Letter is to report the unexpected, contrasting fractionation behavior of toluene and benzene that we have observed in a series of experiments performed in different setups under purely diffusive conditions (1-D gel dissection tubes), as well as in flow-through systems (2-D flow-through chamber). A key feature of the experiments was the simultaneous presence of non-deuterated and perdeuterated mixtures of toluene and benzene undergoing diffusion and lateral dispersion. In this way, in each experiment the 4 tracers (C_6H_6 , C_6D_6 , C_7H_8 , C_7D_8) were all transported under the same conditions, which facilitates a direct comparison of the effects of diffusion and transverse dispersion on the concentration of non-deuterated and perdeuterated benzene and toluene and, more importantly, on their diffusive isotope fractionation.

Materials and Methods

Chemicals and Analytical Methods. High-purity organic compounds (99.5%, Sigma-Aldrich, Germany) were used in the experiments. Measurements were carried out with a 7890A gas chromatograph (GC) with a capillary column (30m $\times$ 250 μ m, 1.0 μ m film thickness; Agilent, USA) coupled to a 5975C tri-axis mass selective detector (MSD) (Agilent, USA). Headspace samples were injected for analysis using a COMBIPAL multi-purpose autosampler system.

1-D Diffusion Experiment. Gel dissection experiments were performed in cylindrical glass tubes (1.1 cm diameter and 20 cm length) using agarose gel prepared with a minimum amount (1% w/w) of phyto agar (Duchefa, Netherlands). Agar solutions containing mixtures of dissolved toluene, perdeuterated toluene, benzene and perdeuterated benzene (1:1:1:1 volume proportion) were prepared and filled in the first 10 cm of the diffusion tubes. This zone acted as contaminant source during the experiments in which the compounds diffused towards the remaining portion of the tubes

that was filled with pure gel medium. The tubes were kept horizontally at a constant temperature of 20 °C. After 9 and 15 days, the tubes were sampled by cutting the gel into 1 cm slices with a scalpel. The slices were immediately sealed in 10 ml glass vials with screw caps and sent to GC analysis.

2-D Flow-through Experiment. Flow-through experiments were performed in a quasi two-dimensional flow-through chamber (inner dimensions: 80cm×18cm×1cm, L×H×W), equipped with 10 equally-spaced (1 cm spacing) ports both at the inlet and at the outlet. The flow-through system was filled with homogeneous quartz sand (Euroquarz, Germany) with grain diameter of 1.0-1.5 mm. The sand was washed in an acidic solution and dried for 12 hours in an oven at 120 °C before filling the flow-through chamber. The sand was filled with a wet-packing procedure using ultra-pure Milli-Q water (EvoquaWater, USA) to avoid air entrapment in the porous medium.³⁷ The inlet and outlet ports were connected to two high-precision multi-channel peristaltic pumps (IPC-N24, Ismatec, Switzerland). Sampling was performed with a 10-channel syringe pump (KD Scientific, USA). The system was operated at a seepage velocity of 0.8 m/d. An aqueous solution containing the four isotopologues was continuously injected from the two lowermost ports at the inlet of the flow-through chamber. After establishing a steady-state plume (i.e., exchanging at least two pore volumes), samples were taken at the outlet ports and analyzed for the concentrations of deuterated and non-deuterated compounds.

Modeling Approach. The governing equations for contaminant transport in the two experimental setups are the 1-D Fick's second law of diffusion³⁸ in the gel dissection tubes, and the 2-D steady state advection dispersion equation in the 2-D flow-through setup.³⁹⁻⁴⁰ The models used to quantitatively interpret the experimental results are based, respectively, on a numerical and an analytical solution of these governing equations. The key parameters controlling transport of the different toluene and benzene isotopologues are their diffusion coefficients and, in the flow-through setup, their transverse hydrodynamic dispersion coefficients. The governing equations and their

boundary conditions are summarized in the Supporting Information together with concentration maps of the steady-state plumes in the 2-D flow-through setup.

Results and Discussion

Two examples of benzene and toluene spatial profiles measured in the 1-D setup after 9 and 15 days of diffusion are illustrated in Figure 1. The plots show typical diffusion curves with small differences between deuterated and non-deuterated compounds that are difficult to appreciate in the concentration plots. The differences become apparent by plotting the ratios C_6D_6/C_6H_6 and C_7D_8/C_7H_8 . These ratios surprisingly show an opposite trend with a decreasing pattern for toluene, and a reverse, increasing trend for benzene. To quantify these observations we used a simple inverse power law model,^{21,41} which relates the diffusion coefficients of the deuterated and non-deuterated isotopologues to their molecular masses:

$$\frac{D_{C_nD_m}}{D_{C_nH_m}} = \left(\frac{m_{C_nH_m}}{m_{C_nD_m}} \right)^\beta \quad (1)$$

103

104 Figure 1. Spatial profiles of concentration and isotopologue ratio for benzene and toluene along the
 105 length of the diffusion tubes in the 9-day experiment A (panels a-c) and in the 15-day experiment C
 106 (panels d-f). The symbols (squares for the non-deuterated, triangles for the perdeuterated
 107 isotopologues, circles for the ratios) represent the measured data, whereas the lines are the outcomes
 108 of the simulations.

109

110 Fitting the experimental data allowed estimating the values of the diffusion coefficients for the non-
 111 deuterated and D-labeled isotopologues and, thus, the value of the exponent β expressing the mass
 112 dependence of the isotopologues diffusion coefficients. The fitting procedure was carried out using
 113 the function *lsqnonlin* implemented in MATLAB. Table 1 summarizes the results obtained for the
 114 different tube experiments and reports the observed diffusive isotope fractionation. Both the
 115 graphical representation of the experimental data (Figure 1) and the parameters reported in Table 1

show a normal isotope effect for toluene and an inverse isotope effect for benzene. In the case of toluene, the deuterated isotopologue has a slightly lower diffusion coefficient and results in decreasing isotope ratios as the toluene species diffuse towards the pure gel medium. For benzene, instead, the data show an inverse diffusive isotope effect: the molecules of the deuterated isotopologue diffuse at a slightly faster rate and become enriched in the initially pure gel medium, as benzene diffuses from the contamination source. This behavior was consistently observed in all experiments. The β (0.444-0.490) values for toluene are consistent with those of previous experiments,¹⁵ whereas the negative values for benzene clearly indicate an inverse isotope effect. Average values characterizing normal and inverse isotope fractionation observed for toluene and benzene in these diffusion experiments are: $D_{C_7D_8}/D_{C_7H_8}=0.962\pm0.002$ and $D_{C_6D_6}/D_{C_6H_6}=1.019\pm0.002$.

127

Table 1. Summary of the results for the multitracer 1-D diffusion experiments.

Experiment	Time [Days]	$D_{C_nH_m}$ [$\times 10^{-9} m^2 s^{-1}$]	$D_{C_nD_m}$ [$\times 10^{-9} m^2 s^{-1}$]	β [-]	$D_{C_nD_m}/D_{C_nH_m}$ [-]
Benzene					
A	9	0.958 $\pm$ 0.040	0.976 $\pm$ 0.017	-0.251	1.019
B	9	0.958 $\pm$ 0.008	0.978 $\pm$ 0.007	-0.279	1.021
C	15	0.956 $\pm$ 0.021	0.972 $\pm$ 0.020	-0.218	1.016
D	15	0.957 $\pm$ 0.025	0.974 $\pm$ 0.024	-0.238	1.018
Toluene					
A	9	0.800 $\pm$ 0.010	0.768 $\pm$ 0.009	0.490	0.960
B	9	0.811 $\pm$ 0.015	0.780 $\pm$ 0.012	0.460	0.962
C	15	0.798 $\pm$ 0.012	0.769 $\pm$ 0.010	0.444	0.964
D	15	0.810 $\pm$ 0.019	0.780 $\pm$ 0.016	0.453	0.963

129

Figure 2 shows vertical profiles of concentration and isotope ratio observed for the simultaneous transport of deuterated and non-deuterated benzene and toluene in the flow-through chamber. The concentration trends show a typical transverse dispersion profile for plumes continuously injected in a homogeneous porous medium. The isotope ratio measured at the different ports confirms the

normal diffusive fractionation for toluene and the inverse fractionation for benzene. In fact, the ratio C_7D_8/C_7H_8 decreases from the core towards the outer fringe of the plume, whereas the ratio C_6D_6/C_6H_6 increases towards the plume fringe. No fitting procedure was used to evaluate the data from the flow-through experiment. Instead, the experimental results were evaluated with pure forward modeling of solute transport based on the average diffusion coefficients of the 4 compounds determined in the tube diffusion setups. The key parameter in this setup is the transverse hydrodynamic dispersion coefficients, which is compound (and isotopologue) specific and was described with the following parameterization:

$$D_T = D_P + D_{aq} \left(\frac{Pe^2}{Pe + 2 + 4\delta^2} \right)^{0.5} \quad (2)$$

where D_P [L^2/T] denotes the velocity-independent pore diffusion coefficient; $Pe = vd/D_{aq}$ [-] is the grain Péclet number, in which d [L] is the average grain size diameter and v [L/T] is the seepage velocity; δ [-] is the ratio between the length of a pore channel and its hydraulic radius; a value of $\delta=5.37$ was determined in previous tracer experiments performed in a range of porous media, grain sizes and seepage velocities comprising the conditions of the current flow-through setup.⁴² This parameterization of transverse dispersion has been tested for solute transport in similar quasi 2-D systems⁴³⁻⁴⁵ and verified in pore-scale studies⁴⁶⁻⁴⁷ and fully three-dimensional flow-through experiments.³⁸ The essential feature of Eq. 2 is that it acknowledges the importance of aqueous diffusion also in the non-linear, velocity-dependent mechanical dispersion term. The good agreement of the predictive, purely forward simulations for both measured concentrations and ratios (Figure 2) shows the capability of the model to capture the diffusive fractionation effects in the flow-through system and also the accuracy of the experimentally determined values of aqueous diffusion for the deuterated and non-deuterated tracers obtained from the 1-D tube experiments. The latter compare very well with the values of aqueous diffusion coefficients computed with classical

empirical correlations for organic compounds (see Table S2 in the Supporting Information); this also indicates that sorption of toluene and benzene was not important in our experiments. The values of the transverse dispersion coefficients, determined according to Eq. 2, and used in the profiles and isotope ratio maps shown in Figure 2 are: $D_{T,C7H8}=1.563\times10^{-9}$, $D_{T,C7D8}=1.547\times10^{-9}$, $D_{T,C6H6}=1.641\times10^{-9}$ and $D_{T,C6D6}=1.649\times10^{-9}$ m²/s.

Figure 2. Observed and simulated vertical profiles of concentrations and ratios at the outlet of the flow-through setup (a-b, d-e) and spatial maps of the isotopologue ratio for benzene (c) and toluene (f). The concentration data at the outlet ports (squares for the non-deuterated, triangles for the perdeuterated isotopologues) are average of triplicate measurements of two sampling events of the steady-state plumes, carried out after flushing 2 and 4 pore volumes (i.e., 48 and 96 hours, respectively).

Diffusive isotope fractionation in condensed systems, such as aqueous solutions, is challenging to investigate and to conceptualize in a solid theoretical framework. Different factors that may affect diffusive isotope fractionation include solute mass, volume, shape, molecular structure, polarity, temperature and solute-solvent interactions. Mass and volume dependencies have dominated the discussion about diffusive fractionation in condensed systems and the interpretation of diffusion experiments. Models that have been proposed to capture the effects of diffusive fractionation include the so-called hydrodynamic description (e.g., Stokes-Einstein relation), as well as the extension to the condensed phase of formulations derived from Chapman-Enskog kinetic theory of gas diffusion.³⁸ The hydrodynamic model does not include mass dependence and the key parameter is the radius and, thus, the molar volume of the solute, whereas the kinetic theory predicts an inverse square root relation with the reduced mass.⁴¹ Most of the experimental studies for different aqueous solutes have reported a mass dependence that is typically weaker than the inverse square root relation from kinetic theory. A simple power law expression (Eq. 1), in which the exponent is derived by fitting the experimental data, is the model that is most often used to relate the diffusive isotope effects to the mass of the diffusing solutes in theoretical, computational and experimental studies.^{21,24-26,41} It is worth pointing out that in this simple model the exponent β is an empirical coefficient which also lumps other factors such as steric effects and solute-solvent interactions that are likely to play an important role on diffusive isotope fractionation. Indeed, the data presented in this study show normal and inverse isotope effects that can neither be interpreted on the basis of mass dependencies nor be ascribed as purely hydrodynamic (i.e., mass independent) effects. In fact, based on the sole mass difference the deuterated benzene should have shown lower diffusion rates than the non-labeled isotopologue. Instead, an inverse effect was observed both in the tube diffusion experiments and in the flow-through setup. Our experiments also allow excluding a purely hydrodynamic interpretation of the results. Molar volumes of deuterated benzene and toluene have

193 been shown to be smaller than those of the corresponding non-labeled isotopologues; for instance,
194 Bartell and Roskos⁴⁸ report the following values at 20 °C: $V_{C_6D_6}$ =88.62 mL/mol, $V_{C_6H_6}$ =88.86
195 mL/mol, $V_{C_7D_8}$ =105.98 mL/mol, and $V_{C_7H_8}$ =106.28 mL/mol. These differences are commonly
196 attributed to the smaller effective radius of the C-D bonds compared to the C-H bonds.⁴⁸⁻⁴⁹ However,
197 a pure hydrodynamic interpretation, based on a simple Stokes-Einstein relation and considering the
198 molar volumes reported above, would predict very weak inverse isotope effects
199 (D_{CnDm}/D_{CnHm} =1.0009) for both compounds. This was not observed in this study, in which stronger
200 inverse and normal diffusive isotope fractionation effects were observed for benzene and toluene,
201 respectively. We hypothesize that the geometry of the hydration shells surrounding the aromatic
202 molecules and the solvation dynamics play a major role for diffusive isotope fractionation of
203 deuterated and non-deuterated benzene and toluene in aqueous solutions. For instance, a recent
204 computational study of benzene hydration⁵⁰ reports that the hydration shell of a benzene molecule
205 has an average number of 31 water molecules and this number can change and diminish
206 substantially upon increase of temperature and decrease of density. To explain the unexpected
207 inverse fractionation observed for benzene, one could hypothesize a similar effect on benzene
208 hydration due to the deuterium substitutions. Processes connected to the hydrophobic effect of
209 aromatic molecules in solution and associated changes of density and orientation of water
210 molecules surrounding the solute,⁵¹ as well as the structure and patterns of solute hydration⁵² appear
211 to be more important than the effects due to mass and molar volume differences. This hypothesis
212 will require future substantiation that might be provided by molecular dynamic simulations of
213 aqueous diffusion of deuterated and non-deuterated species. Such simulations have been performed
214 for instance for charged species^{22,53-54} and have contributed to elucidate the importance of the
215 hydration shell and of the interaction between solute and water molecules for the observed isotope
216 fractionation of different ions in aqueous solution. The different interaction of deuterated and non-

217 deuterated benzene and toluene molecules with water molecules appears to be an important factor to
218 explain the experimental observations of this study. This can be deduced also by comparing the
219 results of this study with earlier experiments. For instance, benzene self-diffusion⁵⁵ and tracer
220 diffusion in chlorobenzene⁵⁶ showed only very minor or no isotope effects during diffusion of C₆H₆
221 and C₆D₆ in these organic liquids, which contrasts with the rather large fractionation consistently
222 observed in the different aqueous systems considered in our study.

223 We think that the normal and inverse diffusive fractionation effects observed for deuterated and
224 non-deuterated BTEX will stimulate further research to develop a mechanistic understanding of
225 transport and transformation of these contaminants and their labeled mixtures in different
226 environments. Despite fractionation of D/H substituted aromatic compounds due to biodegradation
227 can be large and may be dominant for many transformation pathways,⁵⁷ our study shows that
228 diffusive fractionation is also significant and should be considered when these chemicals are used as
229 tracers and diagnostic tools in environmental systems in which transport and transformation
230 processes are typically coupled. On a broader perspective, the results of this study, showing
231 different diffusive isotope fractionation of very similar compounds, also have implications for the
232 fast developing field of compound specific stable isotope analysis (CSIA) techniques⁵⁸⁻⁵⁹, which
233 will greatly benefit from data for different environmental contaminants. The path towards
234 developing a comprehensive database of organic pollutant diffusive isotope fractionation and a
235 sound theory capturing the key mechanisms causing diffusive fractionation is inherently complex
236 but rich of intriguing future challenges.

237

238 **Supporting Information**

239 Description of the flow-through system and of the modeling approach used to quantitatively
240 interpret the results in the different experimental setups.

241 The authors declare no competing financial interests.

242

243 Acknowledgments

244 The authors acknowledge the support of the Deutsche Forschungsgemeinschaft (Grant RO 4169/2-1)
245 and the help of Chrysanthi-Elisabeth Nika, Hanne Bøggild and Mikael Emil Olsson for assistance in
246 the experimental work.

247 References

- 248 1. Wiedemeier, T. H.; Rifai, H. S.; Newell, C. H.; Wilson, T. H., Natural attenuation of fuels and
249 chlorinated solvents in the subsurface. John Wiley, New York, **1999**, 617 p.
- 250 2. Thierrin, J.; Davis, G. B.; Barber, C., A Groundwater Tracer Test with Deuterated Compounds for
251 Monitoring In-Situ Biodegradation and Retardation of Aromatic Hydrocarbons. *Ground Water* **1995**, *33*,
252 469-475.
- 253 3. Reusser, D. E.; Istok, J. D.; Beller, H. R.; Field, J. A., In situ transformation of deuterated toluene and
254 xylene to benzylsuccinic acid analogues in BTEX-contaminated aquifers. *Environ. Sci. Technol.* **2002**,
255 *36*, 4127-4134.
- 256 4. Fisher, A.; Bauer, J.; Meckenstock, R. U.; Stichler, W.; Griebler, C.; Maloszewski, P.; Kastner, M.;
257 Richnow, H. H., A multitracer test proving the reliability of Rayleigh equation-based approach for
258 assessing biodegradation in a BTEX contaminated aquifer. *Environ. Sci. Technol.* **2006**, *40*, 4245-4252.
- 259 5. Gieg, L. M.; Alumbaugh, R. E.; Field, J.; Jones, J.; Istok, J. D.; Suflita, J. M., Assessing in situ rates of
260 anaerobic hydrocarbon bioremediation. *Microb. Biotechnol.* **2009**, *2*, 222-233.
- 261 6. Cozzarelli, I. M.; Bekins, B. A.; Eganhouse, R. P.; Warren, E.; Essaid, H. I., In situ measurements of
262 volatile aromatic hydrocarbon biodegradation rates in groundwater. *J. Contam. Hydrol.* **2010**, *111*, 48-64.
- 263 7. Parker, B. L.; Chapman, S. W.; Guilbeault, M. A. Plume persistence caused by back diffusion from thin
264 clay layers in a sand aquifer following TCE source-zone hydraulic isolation. *J. Contam. Hydrol.* **2008**,
265 *102*, 86-104.
- 266 8. Carrera, J.; Sanchez-Vila, X.; Benet, I.; Medina, A.; Galarza, G.; Guimera, J., On matrix diffusion:
267 formulations, solution methods and qualitative effects. *Hydrogeol. J.* **1998**, *6*, 178-90.
- 268 9. Rolle, M.; Chiogna, G.; Hochstetler, D. L.; Kitanidis, P. K., On the importance of diffusion and
269 compound-specific mixing for groundwater transport: An investigation from pore to field scale, *J.*
270 *Contam. Hydrol.* **2013**, *153*, 51-68.
- 271 10. Rolle, M.; Kitanidis, P.K., Effects of compound-specific dilution on transient transport and solute
272 breakthrough: A pore-scale analysis. *Adv. Water Resour.* **2014**, *71*, 186-199.
- 273 11. LaBolle, E. M.; Fogg, G. E.; Eweis, J. B.; Gravner, J.; Leaist, D. G., Isotopic fractionation by
274 diffusion in groundwater. *Water Resour. Res.* **2008**, *44*.
- 275 12. Rolle, M.; Chiogna, G.; Bauer, R.; Griebler, C.; Grathwohl, P., Isotopic Fractionation by Transverse
276 Dispersion: Flow-through Microcosms and Reactive Transport Modeling Study. *Environ. Sci. Technol.*
277 **2010**, *44*, 6167-6173.
- 278 13. Eckert, D.; Rolle, M.; Cirpka, O. A., Numerical simulation of isotope fractionation in steady-state
279 bioreactive transport controlled by transverse mixing. *J. Contam. Hydrol.* **2012**, *140*, 95-106.
- 280 14. Van Breukelen, B. M.; Rolle, M., Transverse Hydrodynamic Dispersion Effects on Isotope Signals in
281 Groundwater Chlorinated Solvents' Plumes. *Environ. Sci. Technol.* **2012**, *46*, 7700-7708.

- 282 15. Jin, B.; Rolle, M.; Li, T.; Haderlein, S.B., Diffusive fractionation of BTEX and chlorinated ethenes in
283 aqueous solution: quantification of spatial isotope gradients. *Environ. Sci. Technol.* **2014**, 48, 6141-6150.
- 284 16. Wanner, P.; Hunkeler, D., Carbon and chlorine isotopologue fractionation of chlorinated hydrocarbons
285 during diffusion in water and low permeability sediments. *Geochim. et Cosmochim. Acta* **2015**, 157,
286 198-212.
- 287 17. Jähne, B.; Heinz, G.; Dietrich, W., Measurement of the diffusion coefficients of sparingly soluble gases
288 in water. *Journal of Geophysical Research: Oceans* **1987**, 92, 10767-10776.
- 289 18. Eggenkamp, H. G. M.; Middelburg, J. J.; Kreulen, R., Preferential diffusion of ³⁵Cl relative to ³⁷Cl in
290 sediments of Kau Bay, Halmahera, Indonesia. *Chem. Geol.* **1994**, 116, 317-325.
- 291 19. Prinzhofer, A.; Pernaton, E.; Isotopically light methane in natural gas: bacterial imprint or diffusive
292 fractionation? *Chem. Geol.* **1997**, 142, 193-200.
- 293 20. Schloemer, S.; Krooss, B. M., Molecular transport of methane, ethane and nitrogen and the influence of
294 diffusion on the chemical and isotopic composition of natural gas accumulations. *Geofluids* **2004**, 4, 81-
295 108.
- 296 21. Richter, F. M.; Mendybaev, R. A.; Christensen, J. N.; Hutcheon, I. D.; Williams, R. W.; Sturchio, N. C.;
297 Beloso, A. D., Kinetic isotopic fractionation during diffusion of ionic species in water. *Geochim.*
298 *Cosmochim. Acta* **2006**, 70, 277-289.
- 299 22. Bourg, I. C.; Sposito, G., Molecular dynamics simulations of kinetic isotope fractionation during the
300 diffusion of ionic species in liquid water. *Geochim. Cosmochim. Acta* **2007**, 71, 5583-5589.
- 301 23. Donahue, M. A.; Werne, J. P.; Meile, C.; Lyons, T. W., Modeling sulfur isotope fractionation and
302 differential diffusion during sulfate reduction in sediments of the Cariaco Basin. *Geochim. Cosmochim.*
303 *Acta* **2008**, 72, 2287-2297.
- 304 24. Bourg, I. C.; Sposito, G., Isotopic fractionation of noble gases by diffusion in liquid water: Molecular
305 dynamics simulations and hydrologic applications. *Geochim. Cosmochim. Acta* **2008**, 72, 2237-2247.
- 306 25. Eggenkamp, H. G. M.; Coleman, M. L., The effect of aqueous diffusion on the fractionation of chlorine
307 and bromine stable isotopes. *Geochim. Cosmochim. Acta* **2009**, 73, 3539-3548.
- 308 26. Bourg, I. C.; Richter, F. M.; Christensen, J. N.; Sposito, G., Isotopic mass dependence of metal cation
309 diffusion coefficients in liquid water. *Geochim. Cosmochim. Acta* **2010**, 74, 2249-2256.
- 310 27. Beekman, H. E.; Eggenkamp, H. G. M.; Appelo, C. A. J., An integrated modelling approach to
311 reconstruct complex solute transport mechanisms - Cl and delta Cl-37 in pore water of sediments from a
312 former brackish lagoon in The Netherlands. *Appl. Geochem.* **2011**, 26, 257-268.
- 313 28. Wortmann, U. G.; Chrnayavsky, B. M., The significance of isotope specific diffusion coefficients for
314 reaction-transport models of sulfate reduction in marine sediments. *Geochim. Cosmochim. Acta* **2011**, 75,
315 3046-3056.
- 316 29. Tempest, K.; Emerson, S., Kinetic isotope fractionation of argon and neon in during air–water transfer.
317 *Marin. Chem.* **2013**, 153, 39–47.
- 318 30. Richter, F. M.; Liang, Y.; Davis, A. M.; Isotope fractionation by diffusion in molten oxides. *Geochim.*
319 *Cosmochim. Acta* **1999**, 63, 2853-2861.
- 320 31. Richter, F. M.; Davis, A. M.; DePaolo, D. J.; Watson, E. B.; Isotope fractionation by chemical diffusion
321 between molten basalt and rhyolite. *Geochim. Cosmochim. Acta* **2003**, 67, 3905-3923.
- 322 32. Watkins, M. W.; DePaolo, D. J.; Ryerson, F. J.; Peterson, B. T., Influence of liquid structure on diffusive
323 isotope separation in molten silicates and aqueous solutions. *Geochim. Cosmochim. Acta* **2011**, 75, 3103-
324 3118.
- 325 33. Watkins, M. W.; Liang, Y.; Richter, F. M.; Ryerson, F. J.; DePaolo, D. J., Diffusion of multi-isotopic
326 chemical species in molten silicates. *Geochim. Cosmochim. Acta* **2014**, 139, 313-326.

- 327 34. Mills, R., Search for isotope effects in the self-diffusion of benzene and cyclohexane at 25°. *J. Phys.*
328 *Chem.* **1975**, 79, 852-853.
- 329 35. Mills, R.; Harris, K. R., The effect of isotopic substitution on diffusion in liquids. *Chem. Soc. Rev.* **1976**,
330 5, 215-231.
- 331 36. Freer, R.; Sherwood, J. N., Diffusion in organic liquids. *J.C.S. Faraday I* **1980**, 76, 1030-1037.
- 332 37. Haberer, C. M.; Rolle, M.; Cirpka, O.A.; Grathwohl P., Oxygen transfer in a fluctuating capillary fringe.
333 *Vadose Zone J.* **2012**, 11.
- 334 38. Cussler, E.L.; Diffusion – mass transfer in fluid systems. Cambridge University Press (3rd Edition), **2009**,
335 631p.
- 336 39. Cirpka, O.A.; Rolle, M.; Chiogna, G.; de Barros, F.P.J.; Nowak, W, Stochastic evaluation of mixing-
337 controlled steady-state plume lengths in two-dimensional heterogeneous domains. *J. Contam.Hydrol.*
338 **2012**, 138-139, 22-39.
- 339 40. Muniruzzaman, M.; Haberer, C.M.; Grathwohl, P.; Rolle, M., Multicomponent ionic dispersion during
340 transport of electrolytes in heterogeneous porous media: Experiments and model-based interpretation.
341 *Geochim. Cosmochim. Acta* **2014**, 656-669.
- 342 41. Bhattacharyya, S.; Bagchi, B., Power law mass dependence of diffusion: A mode coupling theory
343 analysis. *Phys. Rev. E* **2000**, 61, 3850-3856.
- 344 42. Ye, Y.; Chiogna, G.; Cirpka, O. A.; Grathwohl, P.; Rolle M., Experimental investigation of compound-
345 specific dilution of solute plumes in saturated porous media: 2-D vs. 3-D flow-through systems. *J.*
346 *Contam. Hydrol.* **2015**, 172, 33–47.
- 347 43. Chiogna, G.; Eberhardt, C.; Grathwohl, P; Cirpka, O. A.; Rolle, M., Evidence of compound-dependent
348 hydrodynamic and mechanical transverse dispersion by multitracer laboratory experiments. *Environ. Sci.*
349 *Technol.* **2010**, 44, 688–693.
- 350 44. Rolle, M.; Muniruzzaman, M.; Haberer, C. M.; Grathwohl, P., Coulombic effects in advection-
351 dominated transport of electrolytes in porous media: Multicomponent ionic dispersion. *Geochim.*
352 *Cosmochim. Acta* **2013**, 120, 195-205.
- 353 45. Muniruzzaman, M.; Rolle, M., Impact of multicomponent ionic transport on pH fronts propagation in
354 saturated porous media. *Water Resour. Res.* **2015**, 51, 6739–6755.
- 355 46. Rolle, M.; Hochstetler, D. L.; Chiogna, G.; Kitanidis, P. K.; Grathwohl P., Experimental investigation
356 and pore-scale modeling interpretation of compound-specific transverse dispersion in porous media.
357 *Transp. Porous Med.* **2012**, 93, 347–362.
- 358 47. Hochstetler, D. L.; Rolle, M.; Chiogna, G.; Haberer, C. M.; Grathwohl, P.; Kitanidis, P. K., Effects of
359 compound-specific transverse mixing on steady-state reactive plumes: Insights from pore-scale
360 simulations and Darcy-scale experiments. *Adv. Water Resour.* **2013**, 54, 1-10.
- 361 48. Bartell, L. S.; Roskos, R. R.; Isotope effects on molar volume and surface tension: simple theoretical
362 model and experimental data for hydrocarbons. *J. Phys. Chem.* **1966**, 44, 457-463.
- 363 49. Dunitz, J.; Ibberson, R. M., Is deuterium always smaller than protium? *Angewandte Chemie-*
364 *International Edition* **2008**, 120, 4276-4278.
- 365 50. Choudhary, A.; Chandra, A., Spatial and orientational structure of the hydration shell of benzene in sub-
366 and supercritical water. *J. Phys. Chem. B* **2015**, 119, 8600-8612.
- 367 51. Raschke, T.; Levitt, M., Nonpolar solutes enhance water structure within hydration shells while reducing
368 interactions between them. *PNAS* **2005**, 102, 6777-6782.
- 369 52. Schravendijk, P.; van der Vegt, N.F.A., From hydrophobic to hydrophilic solvation: An application to
370 hydration of benzene. *J. Chem. Theory Comput.* **2005**, 1, 643-652.
- 371 53. Moller, K. B.; Rey, R.; Masia, M.; Hynes, J. T., On the coupling between molecular diffusion and
372 solvation shell exchange. *J. Chem. Phys.* **2005**, 122.

- 373 54. Hofmann, A. E.; Bourg, I. C.; DePaolo, D. J., Ion desolvation as a mechanism for kinetic isotope
374 fractionation in aqueous systems. *PNAS* **2012**, 109, 18689-18694.
- 375 55. Mills, R., Diffusion relationships in the binary system benzene-perdeuteriobenzene at 25 C. *J. Phys.*
376 *Chem.* **1976**, 80, 888-890.
- 377 56. Shankland, I. R.; Dunlop, P. J.; Barr, L. W., Isotope effect in liquids as shown by tracer diffusion at 25 C
378 of several labeled benzenes in benzene, chlorobenzene, n-octane, and cyclohexane, and the relationship
379 to the isotope effect in solids. *Phys. Rev. B* 1975, 12, 2249-2252.
- 380 57. Morasch, B; Richnow, H-H.; Schink, B.; Meckenstock, R.U., Stable hydrogen and carbon isotope
381 fractionation during microbial toluene degradation: mechanistic and environmental aspects. *Appl.*
382 *Environ. Microbiol.* **2001**, 67, 4842-4849.
- 383 58. Thullner, M.; Centler, F.; Richnow, H. H.; Fischer, A., Quantification of organic pollutant degradation in
384 contaminated aquifers using compound specific stable isotope analysis - Review of recent developments.
385 *Org. Geochem.* **2012**, 42, 1440-1460.
- 386 59. Hatzinger, P. B.; Bohlke, J.; Sturchio, N. C., Application of stable isotope ratio analysis for
387 biodegradation monitoring in groundwater. *Current opinion in biotechnology* **2013**, 24, 542-549.