



Trophic impact of Atlantic bluefin tuna migrations in the North Sea

Mariani, Patrizio; Andersen, Ken Haste; Lindegren, Martin ; MacKenzie, Brian

Published in:
ICES Journal of Marine Science

Link to article, DOI:
[10.1093/icesjms/fsx027](https://doi.org/10.1093/icesjms/fsx027)

Publication date:
2017

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Mariani, P., Andersen, K. H., Lindegren, M., & MacKenzie, B. (2017). Trophic impact of Atlantic bluefin tuna migrations in the North Sea. *ICES Journal of Marine Science*, 74(6), 1552-1560.
<https://doi.org/10.1093/icesjms/fsx027>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Trophic impact of Atlantic bluefin tuna migrations in the North Sea**

2

3 Patrizio Mariani^{1*}, Ken H Andersen¹, Martin Lindegren¹, Brian R MacKenzie¹

4

5 ¹ Centre for Ocean Life, National Institute for Aquatic Resources, Technical University of Denmark,
6 Charlottenlund, Denmark

7

8 * Corresponding author:

9 Patrizio Mariani

10 Kavalergården 6, 2920 Charlottenlund

11 Denmark

12 phone: +45 35 88 33 53

13 fax: +45 35 88 33 11

14 pat@aqua.dtu.dk

15

16 **Abstract**

17 Large highly migratory predators can have major impacts on local marine ecosystems
18 by reducing prey populations and leading to trophic cascades that affect the entire fish
19 community. These trophic interactions are typically non-linear and can alter both the
20 migratory behavior of the predator and the stability of the fish community. The
21 impact of a migrating top-predator is investigated here for Atlantic bluefin tuna in the
22 North Sea. Bluefin tuna has been absent from the region for half-century, but recent
23 years have seen recovery of migrations and a return of bluefin tuna in the area. We
24 use a size spectrum model to analyse the trophic impact of the returning tuna on the
25 entire fish community, under scenarios with varying levels of tuna consumption and
26 fishing mortality on the prey. We show that with high level of prey fishing mortality
27 in the North Sea, the effect of a tuna re-colonization results in only limited trophic
28 cascades. However, high tuna consumption or changes in fishing mortality may result
29 in a sudden recruitment failure of small-pelagic fish due to cascading effects on the
30 fish community. In present-day conditions, the level of tuna consumption that triggers
31 recruitment failure is lower at increasing fishing mortalities on their prey, providing
32 indications for the future sustainable management of both small-pelagics and bluefin
33 tuna in the area.

34

35 *Keywords: fish community, long distance migration, size spectrum model, trophic*
36 *cascade*

37

38 **Introduction**

39 Trophic relationships form the backbone of the functioning and biodiversity of
40 ecosystems (Paine 1966, Worm and Duffy 2003). Several factors can affect the
41 magnitude and importance of trophic interactions driving non-linear processes and
42 complex dynamics in food webs (Levin 1998). For example, changes in the
43 abundance of individuals in one trophic level can elicit direct and indirect changes on
44 other trophic levels, a process known as a trophic cascade (Paine 1980, Pace et al.,
45 1999, Polis et al., 2000). Evidences of trophic cascades have been reported both in
46 terrestrial and in aquatic ecosystems (Pace et al., 1999, Shurin et al., 2002, Schmitz et
47 al., 2004, Casini et al., 2008, Frank et al., 2005) supporting the hypothesis of a
48 widespread process in ecosystem dynamics. Ecologists have often debated about how
49 ubiquitous trophic cascades are in ecosystems (Polis 1994, Polis et al., 2000) partly
50 because several compensatory mechanisms can dampen or eliminate the effects of
51 trophic cascades (Pace et al., 1999, Cury et al., 2003, Andersen and Pedersen, 2010,
52 Heath et al., 2014, Lindegren et al., 2016). It has been shown that in marine food
53 webs a high degree of connectivity, presence of omnivory, ontogenic diet shifts and
54 fishing can all prevent or dampen trophic cascades (Baum and Worm 2009, Andersen
55 and Pedersen, 2010). Nonetheless, high fishing pressure has been suggested to trigger
56 regime shifts in large marine ecosystems (Daskalov et al., 2007, Möllmann et al.,
57 2009) and are likely responsible for recent changes in the fish community structure in
58 coastal ecosystems (Jackson et al., 2001).

59
60 Generally, cascades are initiated by any external perturbation that can lower the
61 abundance of some part of the community. Thus apart from fishing, another example
62 of an external perturbation is the appearance of migrating larger predators that
63 consume substantial amounts of prey over a relative narrow time period (Polis, 1994,
64 Scheffer et al. 2005). Casini et al. (2012) showed that when high abundances of Baltic
65 cod *Gadus morhua* migrated into unoccupied habitats in the Baltic Sea, their
66 predatory impact induced four-level trophic cascades in the forage fish, zooplankton
67 and phytoplankton communities of the region.

68
69 Atlantic bluefin tuna (*Thunnus thynnus*) is one of the largest apex predators in some
70 north Atlantic marine ecosystems and have declined due mainly to overexploitation
71 (Fromentin and Powers, 2005). This species used to migrate into the North Sea region
72 for many years in the early-mid 1900s but ceased to do so in the mid-1960s and has
73 been rare or absent during most years ever since (Mather et al., 1995; MacKenzie and
74 Myers 2007; Fromentin and Restrepo 2008), [but the last 2-3 years have provided](#)
75 [evidence of their reappearance \(Waage 2016, Bursell 2016; Thuesen Bleeg 2016;](#)
76 [Christiansen and Gyldenkræne 2016\)](#). Bluefin tuna migrated mainly from the
77 Mediterranean Sea spawning ground and remained in the North Sea region for 2-4
78 months before migrating southward in the autumn (Tiews 1978). While present in the
79 North Sea, tuna consumption of prey (mainly mackerel and herring) has been
80 estimated to be large (~300 kton per year) with a likely large impact on the
81 distribution of the local fish community (Tiews 1978). [This species has recently](#)
82 [reappeared in the northern North Sea and Norwegian Sea. Norway has recently been](#)
83 [allocated a commercial fishing quota from ICCAT increasing from about 36 Mt to 52](#)
84 [Mt in the period 2015 - 2017 \(ICCAT 2014a\), and its entire quota for 2016 was](#)
85 [captured in a single haul with 190 tunas \(Waage 2016\)](#). The presence of tunas has
86 been also reported in the inner North Sea with observations of both vagrant
87 [individuals and schools in the Skagerrak and Kattegat in 2015 and 2016 \(Bursell](#)

88 2016; Thuesen Bleeg 2016; Christiansen and Gyldenkræne 2016). Additionally
89 Atlantic bluefin tuna has recently appeared in other northern areas of the north
90 Atlantic such as Denmark Strait and east Greenland (MacKenzie et al., 2014).

91
92 In this paper, our objective is to analyse the trophic effects induced by the
93 reappearance of Atlantic bluefin tuna (henceforth tuna) on the northern North Sea
94 food web. We investigate whether predation by tuna could have impacts on trophic
95 levels farther down the food web through trophic cascades and evaluate the effects of
96 interactions between tuna consumption and fishing mortality on the prey. We use a
97 size- and trait-based model for the fish community that is able to account for the
98 changes in trophic levels during ontogeny and resolve fishing mortality on larger sizes
99 (Andersen and Pedersen 2010). The model is used to reconstruct the North Sea fish
100 community size spectra under different scenarios of tuna migration and fishing
101 pressure and allows estimating trophic cascades induced by predation from this top
102 predator.

103 104 105 **Methods**

106 *Model formulation*

107 We use a previously published size-spectrum model to represent the baseline fish
108 community (Andersen and Pedersen, 2010; see appendix for concise description
109 including equations and parameters). The model is a trait-based size spectrum model
110 (Andersen et al., 2016) based on a description of the energy budget of individuals. All
111 rates and processes are parameterized using the size of individuals and the asymptotic
112 size of species (Hartvig et al., 2011). The result of the model is the distribution of
113 individual abundances $n(w, W_\infty, t)$ as function of size w , asymptotic size W_∞ and
114 variation over time, t . The central process in the model is predation of small
115 individuals by large individuals. The food obtained from predation fuels growth and
116 reproduction. Reproduction is limited by a Beverton-Holt stock-recruitment
117 relationship to ensure coexistence of a continuum of asymptotic sizes, by introducing
118 additional intra-specific density-dependence to population dynamics not represented
119 by other processes in the model. Fishing mortality is imposed as a function of
120 individual size and asymptotic size on all species (excluding tuna) and assuming a
121 logistic size selectivity function, calibrated to emulate the fishing pattern in the North
122 Sea (Pope et al., 2006). The total biomass of fish in the North Sea has been estimated
123 to be between 8,600 kton and 13,000 kton (Sparholt 1990), which is consistent with
124 estimates from multi-species fish stock assessment models providing 8,000 kton for
125 the entire North Sea (ICES 2016). To simulate the northern North Sea fish community
126 the total biomass of the ecosystem is set in the model as a fraction of what has been
127 estimated and we use a reference biomass of 5,000 kton.

128
129 The impact of predation by tuna on the resident fish community is represented in the
130 model as an extra predation mortality μ_T inflicted on the size within the tunas' prey
131 size preference $\phi_T(w_p)$ where w_p is the weight of prey. Bluefin tuna are opportunistic
132 predators and feed on a diverse diet, both in terms of species and sizes, and their diet
133 composition more likely reflects available prey distributions and abundances in the
134 surrounding environment than strong preferences for particular species or sizes (Goñi,
135 and Arrizabalaga, 2010; Olafsdottir et al., 2016). We assume that the prey range of
136 tuna is constrained within a minimum value ($w_S = 100$ gr) and some large size
137 ($w_L = 2.5$ kg), i.e., $w_S \leq w_p \leq w_L$. This size range approximates the size range of

138 the dominant fish species known to be consumed by tuna in the North Sea during the
 139 1950s-1960s (Tiews 1978) and by tunas in other northern shelf regions (e.g., Chase
 140 2002, Overholtz 2005, Butler et al., 2010). However we also analyse the results at
 141 different values of w_S to test the sensitivity of the model to this parameter.

142

143 The total consumption of tuna S_T can be calculated as (biomass per time):

144

$$145 \quad S_T = \mu_T \int w_p N(w_p) \phi_T(w_p) dw_p \quad (1)$$

146

147 where $N(w)$ is the community spectrum which is the integral over all asymptotic
 148 sizes: $N(w) = \int n(w, W_\infty) w dW_\infty$. Hence $w_p N(w_p)$ is equal to the total biomass in
 149 the prey range on which tuna can impose a predation mortality μ_T with size
 150 preference $\phi_T(w_p)$. The equation for S_T can be rewritten to isolate the mortality:

$$151 \quad \mu_T = S_T \frac{1}{\int w_p N(w_p) \phi_T(w_p) dw_p}. \quad (2)$$

152 which is then used as an extra mortality term in the integration of the model
 153 (Appendix 1).

154

155 We focus our analyses on four scenarios (Table 1): unperturbed ecosystems with no
 156 fishing or tuna consumption, i.e., fishing mortality $F = 0$, consumption by tuna $S_T = 0$
 157 (N_{00}); an unexploited ecosystem with no fishing but with tuna consumption, i.e., $F =$
 158 0 , $S_T = 150$ kton year⁻¹ (N_{01}); an exploited ecosystem with high fishing but no tunas,
 159 $F = 0.7$ year⁻¹, $S_T = 0$ (N_{10}); an exploited ecosystem with tunas, i.e., $F = 0.7$ year⁻¹, S_T
 160 $= 150$ kton/year (N_{11}). In addition, we analyse the trophic impact of fishing and
 161 migrations for a larger range of fishing mortalities ($F \in [0 - 1.5]$ year⁻¹) and tuna
 162 consumptions ($S_T \in [0 - 800]$ kton/year).

163

164 Given a scenario with either fishing or tuna migration ($\{N_{10}, N_{01}, N_{11}\} \in \tilde{N}$), we can
 165 calculate the direct impact of tuna predation as the difference between \tilde{N} and the
 166 unperturbed case: $\int w_p \tilde{N} - w_p N_{00} dw_p$. This difference in biomass estimates the
 167 direct trophic impact on the tuna prey range relative to an unperturbed scenario.
 168 Moreover, in order to estimate the trophic cascade initiated by tuna consumption we
 169 develop an index that expresses the total magnitude (regardless of sign) of the change
 170 in biomass relative to a reference size spectrum. The index (Δ) is defined as the
 171 integral of the size-spectra relative to the case of no tuna predation:

172

$$173 \quad \Delta = - \int_{w_0}^{w_L} \left| \log \frac{N_{\alpha 1}(w)}{N_{\alpha 0}(w)} \right| d(\log w) \quad (3)$$

174

175 where N_α indicates the specific fishing scenario considered (unexploited or exploited
 176 at different levels). This index is always positive and measures the change in size
 177 structure driven by tuna consumption for all sizes smaller than the maximum tuna's
 178 prey size (w_L) but larger than the minimum size included in the model (i.e., fish egg
 179 size, $w_0 = 0.5$ mg). We exclude from the computation effects on sizes larger than the
 180 maximum prey size (w_L) as the abundance of those individuals is relatively low
 181 compared to smaller size classes, hence although perturbation can propagate upwards
 182 on the higher trophic levels and lead to a large change relative to the unexploited
 183 biomass, they have negligible effects on the absolute biomasses.

184

185 *Consumption by tuna in the northern North Sea*

186 Estimating the total prey consumption by tuna requires knowledge of their abundance,
 187 their body size, and their daily ration. No abundance estimates are available for this
 188 time period. However, by combining commercial catch data (ICCAT 2012) and
 189 estimates of instantaneous fishing mortality rates (Fromentin and Restrepo 2008) for
 190 the most important fishery in the region (i.e., the Norwegian fishery), it is possible to
 191 derive approximate estimates of biomass. Most of the Norwegian catches were made
 192 in ICES Subdivision IVa (Tiews 1978). We therefore assume that most of the
 193 foraging by this tuna biomass occurred in this region. Our tuna biomass and
 194 consumption estimates will therefore underestimate the total consumption because
 195 they exclude the catches by other countries; however non-Norwegian landings were
 196 relatively small (ca. 27% of all tuna landings in ICES Areas II-VII; Huse et al.,
 197 2015). We therefore provide sensitivity analyses of the model results to changes in S_T .

198
 199 We expressed the instantaneous fishing mortality rates on an annual basis, which
 200 allows estimation of the percentage of stock biomass removed annually. This removal
 201 rate can be considered as an annual exploitation rate, U (Dick and MacCall 2011).
 202 Consequently, given the reported catches (C) and the estimated U , biomass in the
 203 region can be calculated (Worm et al., 2009; Rosenberg et al., 2014) as $B = C/U$,
 204 where B is the total biomass. Note that U is the fraction of the biomass removed by
 205 the fishery on an annual basis (i.e., obtained by converting the instantaneous fishing
 206 mortality rate to annual removal rates). The instantaneous fishing mortality rates were
 207 estimated to be $F = 0.3$, $F = 0.2$ and $F = 0.1$ per year, for the years 1950s, 1960s and
 208 1970s respectively (Fromentin and Restrepo 2008). These fishing mortality rates
 209 correspond approximately to annual removal rates of 26%, 18% and 10% respectively
 210 for the three time periods. As most of the individuals captured in this fishery were
 211 adults (Tiews 1978, Fromentin and Restrepo 2008), the biomass would correspond
 212 only to a spawning stock biomass (i.e., excluding juveniles) in this region. Indeed the
 213 reported weights in the catches (Tiews 1978) were mainly in the range 150 – 400 kg.
 214 We used two reference tuna weights, $w_1 = 200$ kg and $w_2 = 300$ kg, to derive the
 215 total number of fish in the area ($n_1 = B/w_1$; $n_2 = B/w_2$) and the average daily
 216 consumptions of the population (S_T) using the allometric relationship (Innes et al.,
 217 1987, Overholtz 2005):

$$218$$

$$219 \quad S_T = 0.5 k (w_1^{0.8} n_1 + w_2^{0.8} n_2) \quad (4)$$

$$220$$

221 where $k = 0.123$ and S_T has units of kg per day. Based on the relationship above we
 222 obtain an average daily individual consumption rate of 4% body weight. The daily
 223 ration of tuna in the North Sea and during the 1950s-60s has been previously assumed
 224 to be 3-6% day⁻¹ (Tiews 1978) and 1-4.7% day⁻¹ for tuna in the Gulf of Maine-
 225 Georges Bank during 1970-2002 (Overholtz 2005), which are then consistent with the
 226 estimates obtained above.

227

228 **Results**

229 *Consumption by tuna in the northern North Sea*

230 Norwegian landings of tuna in the region varied between 15 kton in 1952 to 0 kton
 231 after 1987. This corresponds to an estimated maximum biomass of about $B = 60$ kton
 232 ($B = 14$ kton, as average 1950 – 1987). We can then estimate the total annual
 233 consumption by tuna assuming a residence period of 100 days (Figure 1). This
 234 consumption is estimated to be between 100 - 200 kton/year with a maximum value
 235 of 232 kton/year in 1952. Moreover, as the catches declined, the tuna consumption

236 decreases to around 30 kton/year after 1963 and then to 0 in most recent periods.
237 Those values compare well with previous estimates of tuna consumption in the area
238 (Tiews 1978) (Figure 1).

239

240 *Community response to tuna migration and fishing*

241 At equilibrium and under unperturbed conditions (N_{00} ; $F = 0$, $S_T = 0$) the simulated
242 ecological community in the northern North Sea distributes according to a size spectra
243 (slope of -1.47), which provides the assumed total biomass of about 5,000 kton
244 (Figure 2).

245

246 Assuming migration of tuna in this unperturbed system (Figure 3a), we find that the
247 tuna consumption can substantially reduce the abundance of the larger prey in the
248 feeding range ($w > 1$ kg) and have positive indirect impacts on the smaller prey size
249 classes ($0.1 \text{ kg} < w < 0.3 \text{ kg}$). This is mainly because smaller sizes outside the tuna
250 prey range experience a reduction of both predation pressure and competition from
251 the larger sizes, thus contributing to more fish growing into the smallest tuna prey
252 range. The perturbation generated by tuna consumption in the preferred prey range
253 cascades down to smaller size ranges with gradually weaker effects (oscillations). The
254 perturbation is indeed damped while moving from larger to smaller individuals and it
255 becomes very small in the region dominated by zooplankton biomass ($w < 0.5 \text{ g}$).

256

257 The structure of this type of ecosystem is qualitatively different from the fish
258 community simulated when fishing pressure is introduced (N_{10}). In case of a fishing
259 mortality $F = 0.7 \text{ year}^{-1}$ and no tuna consumption ($S_T = 0$) the model results in a 20%
260 increase in fish biomass within the range $0.1 \text{ kg} < w < 1 \text{ kg}$. As above the increase in
261 biomass in a certain range of sizes has cascading effects in the community size
262 spectra. Hence, while negligible effects are predicted on the smaller size classes in the
263 model, $w < 10^{-2} \text{ g}$, biomass increase and decreases are alternated throughout the fish
264 community. Small perturbations to this general trophic cascade are introduced when
265 tuna consumption is considered in this exploited scenario (N_{11} , Figure 3a). The main
266 effect is the slight shifts of the peaks of the perturbation towards smaller sizes as a
267 larger reduction of biomass occur in the larger tuna prey range. At equilibrium the
268 highest biomass in the range $0.1 \text{ kg} < w < 2.5 \text{ kg}$ is obtained in the exploited case with
269 no tuna (N_{10}) while the opposite conditions (no fishing, but tuna present, N_{01})
270 provides the lowest (Table 1). Trophic impact is then largest when tuna is introduced
271 in an exploited scenario, while it becomes much smaller when the ecosystem is
272 exposed to fishery (index Δ in Table 1).

273

274 The sensitivity of the fish community to tuna consumption in exploited or unexploited
275 ecosystems is smaller when a larger prey size range is considered in the model
276 (Figure 3b). Assuming tuna able to feed on prey as small as $w_s = 1 \text{ g}$ the relative
277 change for both ecosystems is always very close to 1 with a trophic impact $\Delta \approx 0.1$ in
278 both cases (Figure 3c). In general an increase in the prey size range has damping
279 effects on trophic cascades. Reducing the prey size range, by increasing w_s , larger
280 effects of tuna consumption are simulated in the community structure and those are
281 generally much stronger for unexploited ecosystems ($\Delta \approx 0.4$, Figure 3c). Indeed the
282 similarities between exploited and unexploited ecosystems rapidly disappear as the
283 smallest prey item is increased to values larger than 20 g.

284

285 *Model sensitivity*

286 In case of no fishing the introduction of tuna in the ecosystem generates a lower
287 equilibrium biomass in the prey range (Figure 4a). The higher the tuna consumption,
288 the larger the reduction in prey biomass. A sharp transition towards low values of
289 biomass (around 500 kton) is shown by the model at $S_T \approx 400$ kton/year and no
290 fishing ($F = 0$). At this level of consumption, the prey biomass reaches a critical
291 minimum spawning stock size that cannot support recruitment. With a low
292 recruitment of the larger species in the prey range, the smaller prey can increase their
293 biomass because of reduced predation. Hence, at this critical level, the temporal
294 dynamic of the size spectrum model shows an oscillatory behaviour of the abundance
295 of tunas' prey. The coefficient of variation (CV, i.e., standard deviation relative to the
296 mean) in the abundance of prey fish within the tuna size range calculated over the last
297 realizations of the model (i.e., last year) is generally small ($CV < 0.05$), while it
298 suddenly increases when the model approaches the critical transition ($CV \approx 0.6$).
299 However, those oscillations start occurring when the fish biomass is low, hence
300 further increases in tuna consumption do not affect the outcome.

301

302 Introducing fishing will generally have positive effects on biomass in the range of the
303 tuna prey. This occurs because the reduction of large fish (such as cod) reduces the
304 predation pressure on small fish, which then increase in abundance. Indeed, at a
305 moderate level of fishing ($0.5 \leq F \leq 1 \text{ year}^{-1}$) and no tuna migration, there is a
306 maximum increase of 500 kton of tuna prey biomass produced (Figure 4a). Further
307 increases in fishing mortality can however reduce smaller fish including the tuna prey
308 size. The same conclusions can be drawn when both tuna migration and fishing are
309 present in the model, resulting in a general non-linear relation between the two
310 processes (Figure 4). Interestingly, at relatively low levels of fishing mortality ($F =$
311 0.5 year^{-1}) up to 500 kton/year of tuna consumption can be present in the model
312 without largely altering the prey biomass compared to an unperturbed scenario.

313

314 The direct impact of tuna on the entire fish community size structure can be evaluated
315 at different values of consumption using the perturbation index Δ (Figure 4b). Since Δ
316 is a measure relative to the absence of tuna migration (Eq. 3), the index start from
317 $\Delta = 0$ at all levels of fishing mortality and can be generally described as the area
318 below a perturbation curve (Figure 3). Tuna can significantly affect the size structure
319 of the community, $\Delta > 0.5$, at high level of consumption or at low level of fishing
320 mortality (Figure 4b). If F is high the effect of tuna consumption on the size-structure
321 is negligible. While at lower levels of fishing ($F \leq 0.6$) the larger the tuna
322 consumption, the greater is the effect on the food web. This pattern is evident up to
323 the critical transition described above, where the collapse of the tuna prey range will
324 largely impact the size-structure and any further increase in S_T will not have any
325 effects.

326

327 Discussion

328 Although it is recognized that predation in marine ecosystems is an important factor
329 explaining food webs dynamics and community structure (Verity and Smetacek 1996,
330 Heithaus et al., 2008, Baum and Worm 2009, Mariani et al., 2013), effects of top
331 predators on the marine food web are more complex because of their cascading
332 impact on the entire community (Cury et al., 2003, Ferretti et al., 2012, Blenckner et
333 al., 2015). Our results show how effects of fishing and consumption of top predators
334 can produce large changes in the small pelagic fish communities with cascading
335 effects that are stronger at lower fishing mortalities. In the northern North Sea, with

336 high level of fishing mortality, the biomass of small pelagic fish is only marginally
337 controlled by predation of large top predators (e.g., cod, tunas, etc.). On the other
338 hand at lower fishing mortality, consumption by top predators can significantly
339 reduce the biomass of the small pelagic having cascading effects on other components
340 of the food web.

341

342 Thus, with an intermediate exploitation rate ($F \approx 0.3 \text{ year}^{-1}$), fishing may have
343 beneficial effects for increasing small pelagic fish biomass, which can then potentially
344 sustain consumption by migratory species such as tunas. Above these intermediate
345 values ($F > 0.5 \text{ year}^{-1}$), small pelagic fish biomass will also be reduced. Moreover,
346 the model shows a critical threshold for tuna consumption above which both trophic
347 impact on the fish community and small pelagic biomass are independent from
348 predation (Figure 4). This can be explained by a strong decline of fish biomass in the
349 preferred prey size range of tuna and the consequent failure in their recruitment.

350 Assuming that the feeding ground in the northern North Sea has a total fish biomass
351 of 5,000 kton and that tuna consumption could be in the range 100 - 200 kton/year,
352 then 2 – 4% of the entire fish community could be consumed by tunas in about 100
353 days. However, the consumption is concentrated in the preferred size range
354 ($w_S \leq w_P \leq w_L$) and at certain levels can become critical for the recruitment of
355 species in this range. Presently this critical level is at high values of tuna consumption
356 for the prey size range $0.1 \leq w_P \leq 1.5 \text{ kg}$. Indeed, when consumption by migrating
357 predators is at 650 kton (13% of the entire fish community) and $F = 0.7$ per year, the
358 recruitment of some prey species (Eq. M10 in the Appendix) oscillates around values
359 of $R_0 \approx 1$, hence the biomass of fish within the tuna preferred size range remains at a
360 minimum level with large impact on the trophic cascade.

361

362 Overall tuna biomass in the northeast Atlantic and Mediterranean Sea has been
363 increasing during the past 5-10 years (ICCAT 2014b) which should also stimulate
364 tuna to explore new (or former) foraging areas. There is ample evidence that
365 migratory species can explore the environment and discover new areas for feeding
366 (Dingle and Drake 2007, Alerstam et al., 2003, Aidley 1981, Hays et al., 2014).
367 Recently, migrations of mackerel and bluefin tuna have been observed in waters east
368 of Greenland (Denmark Strait), an area that was not previously occupied by either
369 species (MacKenzie et al., 2014; Jansen et al., 2016). Those changes are likely driven
370 by improved conditions of the habitats in northern areas (e.g., increase in temperature
371 and food) and can now have cascading effects on the local fish community. Indeed, a
372 re-appearance of tuna in the northern North Sea would presently have negligible
373 effects on both size-structure and pelagic biomass, since the fishing pressure in the
374 area is much greater than the potential impact of tuna feeding. *Despite the presently
375 negligible food web effects of tuna consumption in the northern North Sea a future
376 increase in tuna biomass could lead to pronounced changes in both size-structure and
377 pelagic biomass acting across larger spatial scales. Although a higher tuna biomass
378 would inevitably lead to calls for tuna fishing quotas by several Countries
379 surrounding the North Sea, the historical lesson learned from the past stock collapse
380 and disappearance of tuna would arguably motivate exploitation levels to be kept at a
381 low and sustainable level. Whether these levels would promote a tuna biomass at a
382 level at which consumption critically affects the prey population and the ecosystem as
383 a whole remains to be seen. However any fishing-related removal of tunas would
384 reduce the trophic impacts by reducing the tuna consumption within a range
385 consistent with that considered in the present study.*

386 In addition, it may be suggested that not only tuna but also other large fish predators
387 with similar size and partially overlapping diet preferences could invade or recolonize
388 the North Sea under future climate change. One such competing species could be the
389 warm-water predator swordfish (*Xiphias gladius*) (Stillwell and Kohler 1985), which at
390 least during the Stone Age used to inhabit these waters and could potentially recolonize
391 given future increases in temperature (Enghoff et al. 2007). Recent occurrences in
392 Norwegian waters suggest that such a scenario is indeed very likely (Sundby et al.,
393 2013). Given that our model approach is not species-specific but generalizable to all fish
394 across a great range of sizes, a potential increase in such ecologically similar predators
395 would simply serve to increase the consumption of prey fish within the size range of
396 tuna and support our findings on potential future changes in the North Sea food with
397 respect to its fish prey biomass and size structure.

398
399 Evidences of rapid trophic cascades driven by seasonal migration of top predators
400 have been suggested in the Baltic Sea where periodic migrations of cod and its
401 consumption on herring led to cascading effects across multiple trophic levels in the
402 Gulf of Riga (Casini et al., 2012). We note that fishing mortality on herring in the area
403 is about $F = 0.4 \text{ yr}^{-1}$ (ICES 2014) which in our model would allow a significant
404 trophic impact also at low values of predator consumption (Fig 4b). Moreover, the
405 patterns in trophic cascades observed in the Gulf of Riga are consistent with our tuna
406 results because they show larger impact on the prey (herring) and weaker impact on
407 the other components of the ecosystem (plankton) (Fig 3).

408
409 Recent estimates of the total amount of fish in the entire North Sea are ca. 8,000 kton,
410 (ICES 2016) thus our assumption of 5,000 kton could be reasonable for the northern
411 North Sea. However, changes in prey size range or total fish biomass can affect our
412 results. A change in the total fish biomass in the model is equivalent to a change in
413 the consumption levels of tunas, hence similar effects are produced both in terms of
414 biomass available to tuna feeding or in terms of trophic impact (Figure 4). Indeed, if
415 the fish community is assumed to have a lower total biomass, the critical transition
416 point is shifted towards smaller consumption levels. Beyond this threshold the
417 decrease in prey biomass becomes insensitive to further increases in tuna
418 consumption or, similarly, to reduction in total fish biomass. A wider prey size range
419 for tuna can significantly reduce the trophic impact of consumption. This is because
420 the amount of biomass available to tuna is larger and the feeding impact evenly
421 distributed over a wider size-spectra. On the contrary, a narrower prey size range can
422 induce stronger trophic-cascades illustrated by larger values of the perturbation index
423 Δ .

424
425 Our findings illustrate interactions between migration behaviour and trophic impacts
426 of a migratory predator on a food web. As such, they provide an additional ecosystem
427 context for the management of both forage species in the North Sea and for tuna
428 itself. This knowledge could be useful for understanding the conditions that could
429 allow tuna to re-establish a migration path that was lost in the 1960s, as well as the
430 food web consequences of such a feeding migration. The sudden interruption of fish
431 migration towards certain feeding areas has been explained by a loss of group
432 collective memory driven by the breaking of the social transmission of the
433 information (Petitgas et al., 2010; De Luca et al., 2014). Removing those individuals
434 that had some information and preference for specific feeding areas has likely
435 produced a rapid change in the migratory behaviour of the rest of the group that has
436 likely moved towards other areas (De Luca et al., 2014). Hence, given that food

437 resources are now available for tuna, the recent re-appearance in the region might
438 represent the beginnings of a re-establishment of the migratory path towards the
439 northern North Sea. This re-establishment will proceed if some of the individuals that
440 were able to rediscover this historical feeding area will survive to restore this
441 information in the tuna population.

442

443

444

445 **Acknowledgements**

446 Part of the research leading to these results has received support from the EU-FP7
447 project EURO-BASIN (grant agreement no. 264933).

448

449

450 **References**

- 451 Aidley, D. J. (Ed.). 1981. *Animal migration* (Vol. 13). CUP Archive.
- 452
- 453 Alerstam, T., Hedenström, A., & Åkesson, S. 2003. Long-distance migration: evolution and
454 determinants. *Oikos*, 103(2), 247-260.
- 455
- 456 Andersen, K. H., Jacobsen, N. S. and Farnsworth, K.D. 2016. The theoretical foundations for size
457 spectrum models of fish communities. *Canadian Journal of Fisheries and Aquatic Science*. 73(4): 575-
458 588. [doi:10.1139/cjfas-2015-0230](https://doi.org/10.1139/cjfas-2015-0230)
- 459
- 460 Andersen, K. H., and Pedersen, M. 2010. Damped trophic cascades driven by fishing in model marine
461 ecosystems. *Proceedings of the Royal Society B: Biological Sciences*, 277(1682), 795-802.
- 462
- 463 Baum, J. K., & Worm, B. 2009. Cascading top-down effects of changing oceanic predator
464 abundances. *Journal of Animal Ecology*, 78(4), 699-714.
- 465
- 466 Blenckner T, Llope M, Möllmann C, Voss R, Quaas MF, Casini M, Lindegren M, Folke C, Chr.
467 Stenseth N. 2015. Climate and fishing steer ecosystem regeneration to uncertain economic futures.
468 *Proceedings of the Royal Society B: Biological Sciences*, 282: 20142809.
469 <http://dx.doi.org/10.1098/rspb.2014.2809>
- 470
- 471 Britten, G.L., Dowd, M., Minto, C., Ferretti, F., Boero, F. and Lotze, H.K. 2014. Predator decline leads
472 to decreased stability in a coastal fish community. *Ecology letters*, 17(12), pp.1518-1525.
- 473
- 474 Bursel, J. 2016. 400 kilo + tun genudsat I dansk forvand! In Fisk og Fri, September 2016,
475 <http://www.fiskogfri.dk/400-kilo-tun-genudsat-i-dansk-farvand/> (Accessed: 29th January 2017).
- 476
- 477 Butler, C. M., Rudershausen, P. J., and Buckel, J. A. 2010. Feeding Ecology of Atlantic Bluefin Tuna
478 (*Thunnus Thynnus*) in North Carolina: Diet, Daily Ration, and Consumption of Atlantic Menhaden
479 (*Brevoortia Tyrannus*). *Fish. Bull.* 108, 56–69.
- 480
- 481 Casini, M., Lövgren, J., Hjelm, J., Cardinale, M., Molinero, J. C., & Kornilovs, G. 2008. Multi-level
482 trophic cascades in a heavily exploited open marine ecosystem. *Proceedings of the Royal Society B:*
483 *Biological Sciences*, 275(1644), 1793-1801.
- 484
- 485 Casini, M., Blenckner, T., Möllmann, C., Gårdmark, A., Lindegren, M., Llope, M., Kornilovs, G.,
486 Plikshs, M. and Stenseth, N.C. 2012. Predator transitory spillover induces trophic cascades in
487 ecological sinks. *Proceedings of the National Academy of Sciences*, 109(21), pp.8185-8189.
- 488
- 489 Chase, B. C. 2002. Differences in diet of Atlantic bluefin tuna (*Thunnus thynnus*) at five seasonal
490 feeding grounds on the New England continental shelf. *Fish. Bull.-Natl. Ocean. ATMOSPHERIC Adm.*
491 100, 168–180.
- 492
- 493 Chen, H., Hagerty, S., Crotty, S.M. and Bertness, M.D. 2016. Direct and indirect trophic effects of
494 predator depletion on basal trophic levels. *Ecology*, 97(2), pp.338-346.
- 495
- 496 Christiansen, M. & Gyldenkræne, P. Nordjysk lystfisker jubler: der er blåfannede tun overalt.
497 Danmarks Radio Nyheder (2016), [https://www.dr.dk/nyheder/indland/nordjysk-lystfisker-jubler-der-er-
498 blaaannede-tun-overalt](https://www.dr.dk/nyheder/indland/nordjysk-lystfisker-jubler-der-er-blaafannede-tun-overalt) (Accessed: 29th January 2017).
- 499
- 500 Cury, P., Shannon, L., & Shin, Y. J. 2003. The functioning of marine ecosystems: a fisheries
501 perspective. *Responsible fisheries in the marine ecosystem*, 103-123.
- 502
- 503 De Luca, G., Mariani, P., MacKenzie, B. R., & Marsili, M. 2014. Fishing out collective memory of
504 migratory schools. *Journal of The Royal Society Interface*, 11(95), 20140043.
- 505
- 506 de Roos, A.M., Persson, L. 2001. Physiologically structured models - from versatile technique to
507 ecological theory. *Oikos* 94, 51–71.
- 508

509 Daskalov, G.M., Grishin, A.N., Rodionov, S. and Mihneva, V., 2007. Trophic cascades triggered by
510 overfishing reveal possible mechanisms of ecosystem regime shifts. *Proceedings of the National*
511 *Academy of Sciences*, 104(25), pp.10518-10523.

512

513 De Luca, G., Mariani, P., MacKenzie, B.R. and Marsili, M., 2014. Fishing out collective memory of
514 migratory schools. *Journal of The Royal Society Interface*, 11, p.20140043.
515 DOI:10.1098/rsif.2014.0043

516

517 Dick, E. J., and MacCall, A. D. 2011. Depletion-Based Stock Reduction Analysis: A catch-based
518 method for determining sustainable yields for data-poor fish stocks. *Fisheries Research*, 110: 331–341.
519 <http://www.sciencedirect.com/science/article/pii/S0165783611001962> (Accessed 16 August 2016).

520

521 Dingle, H., & Drake, V. A. 2007. What is migration? *Bioscience*, 57(2), 113-121.

522

523 Doughty, C.E., Roman, J., Faurby, S., Wolf, A., Haque, A., Bakker, E.S., Malhi, Y., Dunning, J.B. and
524 Svenning, J.C. 2015. Global nutrient transport in a world of giants. *Proceedings of the National*
525 *Academy of Sciences*, p.201502549.

526

527 Enghoff, I.B., MacKenzie, B.R. and Nielsen, E.E., 2007. The Danish fish fauna during the warm
528 Atlantic period (ca. 7000–3900bc): Forerunner of future changes? *Fisheries Research*, 87(2), pp.167-
529 180.

530

531 Ferretti, F., Osio, G. C., Jenkins, C. J., Rosenberg, A. A., & Lotze, H. K. 2013. Long-term change in a
532 meso-predator community in response to prolonged and heterogeneous human impact. *Scientific*
533 *reports*, 3.

534

535 Fiskeritidende 2016. Større bestand af tun på vej. p. 5 (in Danish), n. 23, 9th July 2016

536

537 Frank, K. T., Petrie, B., Choi, J. S., & Leggett, W. C. 2005. Trophic cascades in a formerly cod-
538 dominated ecosystem. *Science*, 308(5728), 1621-1623.

539

540 Fromentin, J.M. and Powers, J.E. 2005. Atlantic bluefin tuna: population dynamics, ecology, fisheries
541 and management. *Fish and Fisheries*, 6(4), pp.281-306.

542

543 Fromentin, J. M., & Restrepo, V. 2008. A year-class curve analysis to estimate mortality of Atlantic
544 bluefin tuna caught by the Norwegian fishery from 1956 to 1979. *Collective Volume of Scientific*
545 *Papers*, 64(2), 480-490.

546

547 Fromentin, J.M., Reygondeau, G., Bonhommeau, S. and Beaugrand, G. 2014. Oceanographic changes
548 and exploitation drive the spatio-temporal dynamics of Atlantic bluefin tuna (*Thunnus thynnus*).
549 *Fisheries Oceanography*, 23(2), pp.147-156.

550

551 Goñi, N. and Arrizabalaga, H. 2010. Seasonal and interannual variability of fat content of juvenile
552 albacore (*Thunnus alalunga*) and bluefin (*Thunnus thynnus*) tunas during their feeding migration to the
553 Bay of Biscay. *Progress in Oceanography*, 86(1), pp.115-123.

554

555 Hartvig, M., Andersen, K. H., & Beyer, J. E. 2011. Food web framework for size-structured
556 populations. *Journal of Theoretical Biology*, 272(1), 113-122.

557

558 Hays, G. C., Christensen, A., Fossette, S., Schofield, G., Talbot, J., & Mariani, P. 2014. Route
559 optimisation and solving Zermelo's navigation problem during long distance migration in cross flows.
560 *Ecology letters*, 17(2), 137-143.

561

562 Heath, M.I R., D. C. Speirs, and J. H. Steele 2014. "Understanding patterns and processes in models of
563 trophic cascades." *Ecology letters* 17.1: 101-114.

564

565 Heithaus, M. R., Frid, A., Wirsing, A. J., & Worm, B. 2008. Predicting ecological consequences of
566 marine top predator declines. *Trends in Ecology & Evolution*, 23(4), 202-210.

567

568 Huse, G., MacKenzie, B.R., Trenkel, V., Doray, M., Nøttestad, L. and Oskarsson, G. 2015. Spatially
569 explicit estimates of stock sizes, structure and biomass of herring and blue whiting, and catch data of
570 bluefin tuna. *Earth System Science Data*, 7(1), pp.35-46.

571
572 ICCAT 2012. Report of the 2012 Atlantic Bluefin Tuna Stock Assessment Session. Doc. No. SCI-033 /
573 2012, p. 124.

574
575 ICCAT 2014a. Recommendation by ICCAT amending the recommendation 13-07 by ICCAT to
576 establish a multi-annual recovery plan for bluefin tuna in the eastern Atlantic and Mediterranean.
577 Recommendation 14-04, pp. 35.

578
579 ICCAT 2014b. Report of the 2014 Atlantic Bluefin Tuna Stock Assessment Session (Madrid, Spain –
580 September 22 to 27, 2014). SCRS Plenary Sessions 9 to 21. Pages 1-178.

581
582 ICES 2014. Report of the Baltic Fisheries Assessment Working Group (WGBFAS), 3–10 April 2014.
583 ICES Headquarters, Copenhagen ICES CM 2014/ACOM:10

584
585 ICES 2016. Report of the Working Group on Multispecies Assessment Methods (WGSAM), 9–13
586 November 2015, Woods Hole, USA. ICES CM 2015/SSGEPI:20. 206 pp.

587
588 Innes, S., Lavigne, D. M., Earle, W. M., & Kovacs, K. M. 1987. Feeding rates of seals and whales. *The*
589 *Journal of Animal Ecology*, 115-130.

590
591 Jackson, J.B., Kirby, M.X., Berger, W.H., Bjorndal, K.A., Botsford, L.W., Bourque, B.J., Bradbury,
592 R.H., Cooke, R., Erlandson, J., Estes, J.A. and Hughes, T.P., 2001. Historical overfishing and the
593 recent collapse of coastal ecosystems. *science*, 293(5530), pp.629-637.

594
595 Jansen, T., Post, S., Kristiansen, T., Óskarsson, G. J., Boje, J., MacKenzie, B. R., Broberg, M. and
596 Siegstad, H. 2016. Ocean warming expands habitat of a rich natural resource and benefits a national
597 economy. *Ecological Applications* 26: 2021-2032. DOI: 10.1002/eap.1384

598

599 Jennings, S., Greenstreet, S., & Reynolds, J. 1999. Structural change in an exploited fish community: a
600 consequence of differential fishing effects on species with contrasting life histories. *Journal of Animal*
601 *Ecology*, 68(3), 617-627.

602
603 Jennings, S. and Blanchard, J.L. 2004. Fish abundance with no fishing: predictions based on
604 macroecological theory. *Journal of Animal Ecology*, 73(4), pp.632-642.

605
606 Levin, S. A. 1998. Ecosystems and the biosphere as complex adaptive systems. *Ecosystems*, 1(5), 431-
607 436.

608
609 Lindegren, M., Checkley, D.M., Jr., Ohman, M.D., Koslow, A., Goericke, R. 2016. Resilience and
610 Stability of a Pelagic Marine Ecosystem. *Proceedings of the Royal Society B-Biological Sciences*, 283:
611 20151931

612
613 Lotze, H.K. and McClenachan, L. 2014. Marine historical ecology: informing the future by learning
614 from the past. *Marine Community Ecology and Conservation*, pp.165-200.

615
616 MacKenzie, B. R., Payne, M., Boje, J., Højer, J. L., Siegstad, H. 2014. A cascade of warming impacts
617 brings bluefin tuna to Greenland waters. *Global Change Biology* 20: 2484–2491, doi:
618 10.1111/gcb.12597

619
620 MacKenzie, B.R. and Myers, R.A., 2007. The development of the northern European fishery for north
621 Atlantic bluefin tuna *Thunnus thynnus* during 1900–1950. *Fisheries Research*, 87(2), pp.229-239.

622
623 Mariani, P., & Visser, A. W. 2010. Optimization and emergence in marine ecosystem models. *Progress*
624 *in Oceanography*, 84(1), 89-92.

625
626 Mather, F.J., Mason, J.M. and Jones, A.C., 1995. *Historical document: life history and fisheries of*
627 *Atlantic bluefin tuna*. NOAA Technical Memorandum NMFS-SEFSC 370: 1–165.
628
629 McCauley, D.J., Pinsky, M.L., Palumbi, S.R., Estes, J.A., Joyce, F.H. and Warner, R.R. 2015. Marine
630 defaunation: Animal loss in the global ocean. *Science*, 347(6219), p.1255641.
631
632 Myers, R. A., Baum, J. K., Shepherd, T. D., Powers, S. P., & Peterson, C. H. 2007. Cascading effects
633 of the loss of apex predatory sharks from a coastal ocean. *Science*, 315(5820), 1846-1850.
634
635 Möllmann, C., Diekmann, R., Müller-Karulis, B., Kornilovs, G., Plikshs, M. and Axe, P., 2009.
636 Reorganization of a large marine ecosystem due to atmospheric and anthropogenic pressure: a
637 discontinuous regime shift in the Central Baltic Sea. *Global Change Biology*, 15(6), pp.1377-1393.
638
639 Olafsdottir, D., MacKenzie, B. R., Chosson-P, V., Ingimundardottir, T. 2016. Dietary evidence of
640 mesopelagic and pelagic foraging by Atlantic bluefin tuna (*Thunnus thynnus* L.) during autumn
641 migrations to the Iceland Basin. *Frontiers in Marine Science* 3: Article 106, June 2016; 23 pp.
642 <http://dx.doi.org/10.3389/fmars.2016.00108>.
643
644 Overholtz, W. J. 2006. Estimates of consumption of Atlantic herring (*Clupea harengus*) by bluefin tuna
645 (*Thunnus thynnus*) during 1970–2002: an approach incorporating uncertainty. *J. Northwest Atl. Fish.*
646 *Sci*, 36, 55-63.
647
648 Pace, M. L., Cole, J. J., Carpenter, S. R., & Kitchell, J. F. 1999. Trophic cascades revealed in diverse
649 ecosystems. *Trends in ecology & evolution*, 14(12), 483-488.
650
651 Paine, R. T. 1966. Food web complexity and species diversity. *American Naturalist*, 65-75.
652
653 Paine, R. T. 1980. Food webs: linkage, interaction strength and community infrastructure. *Journal of*
654 *Animal Ecology*, 49(3), 667-685.
655
656 Petitgas, P., Secor, D.H., McQuinn, I., Huse, G. and Lo, N. 2010. Stock collapses and their recovery:
657 mechanisms that establish and maintain life-cycle closure in space and time. *ICES Journal of Marine*
658 *Science: Journal du Conseil*, 67(9), pp.1841-1848.
659
660 Pinnegar, J. K., Polunin, N. V. C., Francour, P., Badalamenti, F., Chemello, R., Harmelin-Vivien, M.
661 L., Hereu, B., Milazzo, M., Zabala, M., D’Anna, G., & Pipitone, C. 2000. Trophic cascades in benthic
662 marine ecosystems: lessons for fisheries and protected-area management. *Environmental*
663 *Conservation*, 27(02), 179-200.
664
665 Polis, G. A. 1994. Food webs, trophic cascades and community structure. *Australian Journal of*
666 *Ecology*, 19(2), 121-136.
667
668 Polis, G. A., Holt, R. D., Menge, B. A., & Winemiller, K. O. 1996. Time, space, and life history:
669 influences on food webs. In *Food Webs* (pp. 435-460). Springer US.
670
671 Polis, G. A., Sears, A. L., Huxel, G. R., Strong, D. R., & Maron, J. 2000. When is a trophic cascade a
672 trophic cascade?. *Trends in Ecology & Evolution*, 15(11), 473-475.
673
674 Pope, J. G., Rice, J. C., Daan, N., Jennings, S. & Gislason, H. 2006 Modelling an exploited marine fish
675 community with 15 parameters—results from a simple size-based model. *ICES J. Mar. Sci.* 63, 1029–
676 1044.
677
678 Rosenberg, A. A., Fogarty, M. J., Cooper, A. B., Dickey-Collas, M., Fulton, E. A., Gutierrez, N. L.,
679 Hyde, K. J. W., et al. 2014. Developing new approaches to global stock status assessment and fishery
680 production potential of the seas. *FAO Fisheries and Aquaculture Circular* (Rome), 1086: 1–175.
681
682 Scheffer, M., Carpenter, S., & Young, B. D. 2005. Cascading effects of overfishing marine
683 systems. *Trends in Ecology & Evolution*, 20(11), 579-581.
684

685 Schmitz, O. J., Krivan, V., & Ovadia, O. 2004. Trophic cascades: the primacy of trait-mediated
686 indirect interactions. *Ecology Letters*, 7(2), 153-163.
687
688 Shurin, J.B., Borer, E.T., Seabloom, E.W., Anderson, K., Blanchette, C.A., Broitman, B., Cooper, S.D.
689 and Halpern, B.S. 2002. A cross-ecosystem comparison of the strength of trophic cascades. *Ecology*
690 *letters*, 5(6), pp.785-791.
691
692 Sparholt, H., 1990. An estimate of the total biomass of fish in the North Sea. *Journal du Conseil: ICES*
693 *Journal of Marine Science*, 46(2), pp.200-210.
694
695 Steneck, R., & Sala, E. 2005. *Large marine carnivores: trophic cascades and top-down controls in*
696 *coastal ecosystems past and present* (pp. 110-37). University of Maine Sea Grant College Program.
697
698 Stillwell, C.E. and Kohler, N.E., 1985. Food and feeding ecology of the swordfish *Xiphias gladius* in
699 the western North Atlantic Ocean with estimates of daily ration. *Marine ecology progress series*, 22(3),
700 pp.239-247.
701
702 Sundby, S., Nøttestad, L., Myklevoll, S. and Tangen, Ø., 2013. Swordfish towards the Arctic Atlantic
703 in climate change. *COLLECT. VOL. SCI. PAP. ICCAT*, 69(3), pp.1296-1303.
704
705 Tiews, K. 1978. On the disappearance of bluefin tuna in the North Sea and its ecological implications
706 for herring and mackerel. *Rapports et Proces-Verbaux des Reunions Int. Explor. Mer.*, 172: 301-309.
707
708 Thuesen Bleeg, M. 2016. Her får danske fiskere den forbudte gigantfangst på 450 kilo krogen: - Ikke
709 for små drenge, [http://livsstil.tv2.dk/2016-09-24-her-faar-danske-fiskere-den-forbudte-gigantfangst-](http://livsstil.tv2.dk/2016-09-24-her-faar-danske-fiskere-den-forbudte-gigantfangst-paa-450-kilo-krogen-ikke-for-smaa)
710 [paa-450-kilo-krogen-ikke-for-smaa](http://livsstil.tv2.dk/2016-09-24-her-faar-danske-fiskere-den-forbudte-gigantfangst-paa-450-kilo-krogen-ikke-for-smaa) (Accessed 29th January 2017).
711
712 Verity, P. G., & Smetacek, V. 1996. Organism life cycles, predation, and the structure of marine
713 pelagic ecosystems. *Marine ecology progress series. Oldendorf*, 130(1), 277-293.
714
715 Werner, E. E., & Peacor, S. D. 2003. A review of trait-mediated indirect interactions in ecological
716 communities. *Ecology*, 84(5), 1083-1100.
717
718 Waage, N. 2016. Fishermen and oceanographers rejoice over major bluefin tuna catch. In Norway
719 Today, September 2016, <http://norwaytoday.info/finance/cheers-large-catches-bluefin-tuna/> (Accessed:
720 29th January 2017).
721
722 Worm, B., & Duffy, J. E. 2003. Biodiversity, productivity and stability in real food webs. *Trends in*
723 *Ecology & Evolution*, 18(12), 628-632.
724
725 Worm, B., Hilborn, R., Baum, J. K., Branch, T. A., Collie, J. S., Costello, C., Fogarty, M. J., et al.
726 2009. Rebuilding global fisheries. *Science*, 325: 578-585.
727
728 Worm, B., & Tittensor, D. P. 2011. Range contraction in large pelagic predators. *Proceedings of the*
729 *National Academy of Sciences*, 108(29), 11942-11947.
730
731
732
733

734 Tables and Figures


735 Table 1. Description of case studies with indication of fishing mortality (F) and tuna consumption (S_T).
 736 Fishing mortality does not include tuna population. Also shown calculated values of fish biomass in the
 737 tuna's preferred size range and the trophic impact index (Δ).
 738

Case	F year ⁻¹	S_T kton year ⁻¹	Prey biomass kton year ⁻¹	Δ
N_{00}	0	0	775	-
N_{01}	0	150	633	0.33
N_{10}	0.7	0	1286	-
N_{11}	0.7	150	1181	0.14

739

740


741



742
743
744
745
746
747
748
749




Figure 1. Total prey biomass consumed per year by tuna migrating into the northern North Sea between 1950 and 2009. Our values (black line) are compared with a series of estimates from Tiews (1978) (cfr. Table 150, grey lines). Catch data of the Norwegian fleet have been transformed to total biomass and then into daily consumption. The total consumed biomass is calculated assuming 100 days of feeding in the area and is used to define the parameter S_T .

750




751
752
753
754
755
756

Figure 2. Abundance size spectra of 25 asymptotic size classes in an unperturbed ecosystem (thin grey lines) and community spectrum (black line) and the zooplankton resource spectrum (thick grey line).



757
758
759
760
761
762
763
764

Figure 3. Impact of tuna in the northern North Sea using the ratio N_{01}/N_{00} (black) and N_{11}/N_{10} (blue) to evaluate the effects of tuna consumption on an unexploited and exploited ($F = 0.7 \text{ year}^{-1}$) ecosystem respectively. Tuna consumption ($S_T = 150 \text{ kton year}^{-1}$) is applied in the prey size range $w_S \leq w_P \leq w_L$ (grey areas) with $w_L = 2.5 \text{ Kg}$ and (a) $w_S = 100 \text{ g}$ (b) $w_S = 1 \text{ g}$. The index Δ Eq. 3 is graphically shown as the area under the relative change curve (blue shaded area). In (c) the value of Δ is shown at different values of w_S for unexploited (black) and exploited (blue) ecosystem.



766
 767
 768
 769
 770
 771
 772
 773
 774
 775
 776

Figure 4. Direct impact of fishing (F) and consumption by tuna (S_T) on (a) prey biomass (a) and (b) size-structure index Δ (Eq. 3). The tuna prey biomass is calculated as the difference between the equilibrium prey biomass, in a specific fishing and consumption scenario, and the unperturbed case, when $F = 0$ and $S_T = 0$. White contour lines are for negative values, black solid lines for negative biomass values and the zero contour line is shown with a dashed line. The prey interval is $w_S \leq w_P \leq w_L$, with $w_S = 100$ gr and $w_L = 2.5$ kg. The index Δ is defined in Eq. 3 as the changes in size structure for $w \leq w_L$.

Encounter and consumption

Prey size selection
$$\phi\left(\frac{w_{prey}}{w}\right) = \exp\left[-\left(\ln\left(\frac{\beta w_{prey}}{w}\right)\right)^2 / (2\sigma^2)\right]$$
 M1

Volumetric search rate
$$V(w) = \gamma w^q; \gamma = \frac{f_0 h \beta^{n-q}}{(1-f_0)\sqrt{2\pi}\kappa_r\sigma}$$
 M2

Encountered food
$$E(w) = V(w) \sum_j \int_0^\infty \phi\left(\frac{w_{prey}}{w}\right) N_j(w_{prey}) w_{prey} dw_{prey}$$
 M3

Maximum consumption rate
$$I_{max} = hw^n$$
 M4

Feeding level
$$f_i(w) = \frac{E_i(w)}{E_i(w) + I_{max,i}}$$
 M5

Growth and reproduction

Maturation function
$$\psi(w) = \left[1 + \left(\frac{w}{w_{mat,i}}\right)^{-10}\right]^{-1} \left(\frac{w}{W_i}\right)^{1-n}$$
 M6

Somatic growth
$$g_i(w) = (\alpha f_i(w) I_{max,i} - k_i w^p)(1 - \psi(w))$$
 M7

Egg production
$$g_r(w) = (\alpha f_i(w) I_{max,i} - k_i w^p)\psi(w)$$
 M8

Recruitment

Population egg production
$$R_{p,i} = \epsilon / (2w_0 N_i(w_0) g(w_0)) \int_{w_0}^{W_i} \psi(w) N_i(w) g_r(w) dw$$
 M9

Recruitment
$$R_i = R_{max,i} \frac{R_{p,i}}{R_{max,i} + R_{p,i}} \xi(t)$$
 M10

Mortality

Background mortality
$$\mu_0 = Z_0 W_i^z$$
 M11

Predation mortality
$$\mu_{p,i}(w_{prey}) = \sum_j \int_{w_0}^\infty \phi\left(\frac{w_{prey}}{w}\right) (1 - f_j(w)) V_j(w) \theta_{ij} N_j(w) dw$$
 M12

Fishing selectivity
$$S_i(w) = 1 / (1 + e^{(s1 - s2 w)})$$
 M13

Fishing mortality
$$\bar{F}_i(w) = S_i(w) F_i$$
 M14

Resource spectrum

Growth rate
$$\frac{\partial N_r(w)}{\partial t} = r_0 w^{n-1} (\kappa(w) - N_r(w)) - \mu_{p,r}(w) N_r(w)$$
 M15

Carrying capacity
$$\kappa(w) = \kappa_r w^{-\lambda}$$
 M16

778

779

780