Xylitol production by *Debaryomyces hansenii* and *Candida guilliermondii* from rapeseed straw hemicellulosic hydrolysate

López-Linares, Juan Carlos; Romero, Inmaculada; Cara, Cristobal; Castro, Eulogio; Mussatto, Solange I.

Published in:
Bioresource Technology

Link to article, DOI:
[10.1016/j.biortech.2017.09.139](https://doi.org/10.1016/j.biortech.2017.09.139)

Publication date:
2018

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
López-Linares, J. C., Romero, I., Cara, C., Castro, E., & Mussatto, S. I. (2018). Xylitol production by *Debaryomyces hansenii* and *Candida guilliermondii* from rapeseed straw hemicellulosic hydrolysate. *Bioresource Technology*, 247, 736-743. <https://doi.org/10.1016/j.biortech.2017.09.139>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Xylitol production by *Debaryomyces hansenii* and *Candida guilliermondii* from rapeseed straw hemicellulosic hydrolysate

Juan Carlos López-Linares^a, Inmaculada Romero^a, Cristobal Cara^a, Eulogio Castro^a, Solange I. Mussatto^{b,*}

^a Center for Advanced Studies in Energy and Environment (CEAEMA), University of Jaén, Campus Las Lagunillas, 23071, Jaén, Spain

^b Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Kemitorvet, Building 220, 2800, Kongens Lyngby, Denmark

* Corresponding author. Tel.: +45 93 51 18 91.

E-mail address: smussatto@biosustain.dtu.dk, solangemussatto@hotmail.com (S.I. Mussatto).

Abstract

This study evaluated the possibility of using rapeseed straw hemicellulosic hydrolysate as a fermentation medium for xylitol production. Two yeast strains, namely *Debaryomyces hansenii* and *Candida guilliermondii*, were used for this bioconversion process and their performance to convert xylose into xylitol was compared. Additionally, different strategies were evaluated for the hydrolysate detoxification before its use as a fermentation medium. Assays in semi-defined media were also performed to verify the influence of hexose sugars on xylose metabolism by the yeasts. *C. guilliermondii* exhibited higher tolerance to toxic compounds than *D. hansenii*. Not only the toxic compounds present in the hydrolysate affected the yeast's performance, but glucose also had a negative impact on their

performance. It was not necessary to completely eliminate the toxic compounds to obtain an efficient conversion of xylose into xylitol, mainly by *C. guilliermondii* ($Y_{P/S} = 0.55$ g/g and 0.45 g/g for *C. guilliermondii* and *D. hansenii*, respectively).

Keywords: rapeseed straw; hemicellulosic hydrolysate; detoxification; semi-defined media; xylitol

1. Introduction

Xylitol ($C_5H_{12}O_5$) is a natural food sweetener that has been used as a sucrose substitute due to the similar sweetening power and lower caloric value. In addition, several benefits to the health have been associated to the ingestion of xylitol. One of the most important is in oral health, reducing and/or preventing tooth decay. However, xylitol can also be used for the treatment of several diseases such as

diabetes, hemolytic anemia, inflammatory processes, colon diseases and parenteral and renal injuries, as well as to prevent respiratory infections, acute otitis media and osteoporosis. Due to these properties, the xylitol market, including applications in food and pharmaceutical industries, has significantly increased in the last decades (Mussatto, 2012; Mussatto and Roberto, 2002).

Xylitol is traditionally produced by chemical synthesis (xylose hydrogenation using chemical catalysts), but it can also be produced by fermentation using microorganisms with ability to convert xylose into xylitol, mainly yeasts. Xylose-to-xylitol bioconversion by yeasts is considered an attractive alternative to the currently large-scale commercial chemical process since the biological route has lower energy requirements (milder conditions of pressure and temperature are needed), is more sustainable and environmentally-friendly (Cortez et al., 2016; Mussatto, 2012).

Different feedstocks have been tested as carbon source for xylitol production by fermentation; the efficiency of the bioconversion process is clearly dependent on the feedstock and pretreatment applied to release xylose, as well as on the microorganism and cultivation conditions. Rapeseed straw is an agricultural residue with a high content of cellulosic and hemicellulosic sugars - greater than 55% (Choi et al., 2013); however, with no relevant application currently. The generation of this residue has significantly increased in the last years due to the growing interest on rape oil for biodiesel production (Pinkowska et al., 2013). In 2014, for example, 9.1 million and 36.1 million hectares of rapeseed were cultivated in Europe and worldwide, respectively (FAOSTAT, 2017).

Different types of pretreatment can be used to release sugars from lignocellulose biomass, among of which, diluted acid pretreatment is one of the most efficient to remove hemicellulose sugars (Mussatto, 2016). However, the generation of toxic compounds such as furan derivatives (furfural and hydroxymethylfurfural), acetic acid, and phenolic compounds, may also occur during this process, and such compounds affect the cell metabolism during fermentation. For this reason, the generation of these compounds must be avoided as much as possible during biomass pretreatment (Mussatto and Dragone, 2016). Some detoxification strategies can also be used in order to reduce the concentration of toxic compounds to levels that do not affect the microbial metabolism, which include vacuum evaporation, overliming, adsorption in activated charcoal or in ion-exchange resins, and extraction with organic solvent, among others (Mussatto and Roberto, 2004a).

The present study evaluated the possibility of producing xylitol from rapeseed straw hemicellulosic hydrolysate. The performance of two yeasts, namely *Debaryomyces hansenii* and *Candida guilliermondii*, to convert xylose into xylitol from hydrolysate medium was evaluated and compared. Such yeasts have been reported as being the most efficient yeasts for bioconversion of xylose to xylitol (Domínguez et al., 1999) and therefore they were both selected for use in the present study. In addition, different detoxification strategies were tested to reduce the negative effect of toxic compounds on fermentation performance. To the best of our knowledge, this is the first study on the use of rapeseed straw as a feedstock for xylitol production.

2. Materials and methods

2.1. *Raw material*

Rapeseed straw was collected from the fields of the province of Seville, Spain, after seed harvesting. As soon as collected, the feedstock was air dried, milled to particle size smaller than 0.4 cm using a laboratory hammer mill (Retsch, SM-100, Haan, Germany), homogenized and stored in a dry place until use. The feedstock (94% w/w dry matter) presented the following composition (% w/w): glucan 31.5 ± 0.3 , hemicellulose 17.4 ± 0.1 (xylan 13.2 ± 0.1 , galactan 1.9 ± 0.1 , arabinan 1.2 ± 0.02 , mannan 1.2 ± 0.04), acid insoluble lignin 16.2 ± 0.5 , acid soluble lignin 1.6 ± 0.05 , acetyl groups 3.4 ± 0.07 , ashes 6.7 ± 0.26 , and extractives 15.4 ± 1.3 (López-Linares et al., 2015).

2.2. *Hemicellulosic hydrolysate*

Rapeseed straw hemicellulosic hydrolysate (RH, raw hydrolysate) was produced by dilute acid pretreatment of the feedstock (10% w/v feedstock load) with 2% (w/v) sulfuric acid at 130 °C for 1 h. After pretreatment, the liquid fraction (hemicellulosic hydrolysate) was recovered by vacuum filtration and the contents of sugars (glucose, xylose, arabinose, mannose, and galactose) and toxic compounds (furfural, hydroxymethylfurfural (HMF), acetic acid, formic acid, and phenolic compounds) were determined.

2.3. *Hydrolysate detoxification*

Different strategies were used for detoxification of the rapeseed straw hemicellulosic hydrolysate in order to decrease the concentration of toxic compounds present (**Fig. 1**). Initially, the raw hydrolysate (RH) was treated with activated charcoal (100 mesh particle size - Sigma-Aldrich) in a 3.5% (w/v) ratio and the suspension was incubated in a rotary shaker (Certomat-R, B-Braun, Germany) at 45 °C, 150 rpm for 1 h (Mussatto and Roberto, 2004b). The liquid fraction was recovered by vacuum filtration in Whatman no. 1 filter paper (ACH, activated charcoal-treated hydrolysate) and 4.5-fold concentrated by evaporation at 60 °C under vacuum conditions. The concentrated liquid fraction thus obtained (CH, ACH 4.5-fold concentrated) had the pH adjusted to 6.0 by addition of solid KOH. Then, an ion-exchange resin treatment was carried out by passing the liquor through a copper column (1.2 cm wide × 50 cm length) filled with the resin Microionex MB200 (Rohm Haas, Copenhagen, Denmark) in a ratio of 0.2 g resin per mL liquor. At the end of this process, the pH of the liquor was corrected to 6 using solid KOH. The hydrolysate obtained after this procedure was denominated DH (detoxified hydrolysate). An aliquot of each hydrolysate sample was collected and used for sugars and toxic compounds determination.

To decrease even more the concentration of toxic compounds present in DH, some additional strategies were applied. 1) A portion of the DH was two-fold diluted to decrease the level of inhibitory compounds and then the diluted hydrolysate was supplemented with sugars up to the original level present in DH. The hydrolysate obtained after this step was denominated DDH (diluted detoxified hydrolysate). 2)

A portion of DH was submitted to an additional detoxification step using organic solvents. In this case, the hydrolysate had the pH adjusted to 3 by addition of H_2SO_4 (98% w/v) and was then mixed with ethyl acetate or chloroform in a 1:3 (v/v) ratio. The mixture was maintained in a rotary shaker at 200 rpm, 30 °C for 45 min. Finally, the residual organic solvent was eliminated by evaporation. The hydrolysates obtained after this step were denominated EA-DH (DH further treated with ethyl acetate) and CL-DH (DH further treated with chloroform).

2.4. Microorganisms and inocula

Debaryomyces hansenii (NRRL Y-7426) and *Candida guilliermondii* FTI 20037 (ATCC 201 935) were the yeast strains used in the experiments. The strains were maintained at 4 °C on malt extract agar slants containing (g/l): malt extract 30, peptone 5, and agar 15.

The inocula were grown in 250-mL Erlenmeyer flasks containing 100 ml of medium composed of (g/l): xylose 30, malt extract 3, yeast extract 3, and peptone 5. For preparation of this medium, a concentrated solution containing all the nutrients was prepared and sterilized at 121 °C for 20 min, while the xylose solution was autoclaved at 112 °C for 15 min. The cells were grown in a rotary shaker at 30 °C and 200 rpm for 24 h, being subsequently recovered by centrifugation (3000 g, 15 min) and resuspended in the fermentation medium.

2.5. Fermentation media and conditions

2.5.1. Hydrolysate-based fermentation media

The detoxified hydrolysate (DH) and its variations (DDH, EA-DH, and CL-DH) were used as fermentation medium for xylitol production. To be used as fermentation medium, the hydrolysates had the pH adjusted to 5.5 by addition of solid KOH and were subsequently autoclaved at 112 °C for 15 min.

Fermentation assays were performed in 100-mL Erlenmeyer flasks containing 50 ml of hydrolysate inoculated with an initial cell concentration of 3 g/l. The flasks were maintained in a rotary shaker at 200 rpm, 30 °C for 144 h. During the experiments, samples were withdrawn each 24 h, centrifuged at 8,000 g for 10 min, and used to determine cell, sugars, xylitol and ethanol concentration. All the experiments were carried out at least in duplicate.

2.5.2. *Semi-defined fermentation media*

A semi-defined medium with the same composition of sugars present in DH (g/l: glucose 11.7, xylose 40.6, galactose 8.7, arabinose 6.8, and mannose 2.2), was prepared and used for xylitol production. Such medium was denominated DM1. Additionally, fermentation media with the same composition of DM1 but without glucose (DM2), without galactose (DM3), and without glucose and galactose (DM4) were also formulated and used for fermentation. All the semi-defined media were supplemented with the same nutrients used for inoculum preparation. The nutrients solution was sterilized at 121 °C for 20 min, while the sugars solutions were autoclaved at 112 °C for 15 min. The pH of the semi-defined media was adjusted to 5.5 by addition of 2M KOH solution.

Fermentation assays were carried out in 100-mL Erlenmeyer flasks containing 50 ml of medium inoculated with 3 g/l of cells concentration, and were maintained under the same conditions used for the assays in hydrolysate-based medium (200 rpm, 30 °C for 72 h). Samples were periodically withdrawn for sugars, xylitol, ethanol, and cell growth determinations. All the experiments were carried out at least in duplicate.

2.6. Analytical methods

Glucose, xylose, arabinose, mannose and galactose concentrations in the hydrolysates were determined by high performance liquid chromatography (HPLC) in a Waters 2695 liquid chromatograph (Mildford, MA, EEUU) equipped with a refractive index detector (Waters 2414). A Transgenomic CHO-782 column operating at 70 °C with ultrapure water as a mobile-phase (0.6 ml/min) was used. Acetic acid, formic acid, furfural and HMF concentrations were also determined by HPLC using a refractive index detector, but, in these cases, a Bio-Rad HPX-87H column operating at 65 °C with 5 mM H₂SO₄ as mobile phase at a flow rate of 0.6 ml/min, was used. Phenolic compounds were estimated by the Folin-Cicalteau method (Singleton and Rossi, 1965) using gallic acid as standard.

Sugars, xylitol and ethanol concentrations from fermentation samples were determined by HPLC using a Bio-Rad HPX-87H column under the same conditions described for acetic acid determination. Cell concentration was estimated by dry weight. For analysis, fermentation samples were filtered through cellulose nitrate filter with 0.2 µm pore size (Sartorius Stedim Biotech, Germany), which were

previously dried to constant weight. The cell concentration was calculated as the ratio between the mass of dried biomass and filtered sample volume.

All analytical determinations were performed in triplicate and average results are shown. Relative standard deviations were below 3%.

2.7. Fermentation parameters

Cell yield ($Y_{X/S}$, g/g) was defined as the ratio between formed cells and total sugars consumed. Xylitol yield ($Y_{PX/S}$, g/g) was calculated as the ratio between xylitol produced and xylose consumed, while the ethanol yield ($Y_{PE/S}$, g/g) was considered as the ratio between the ethanol produced and total sugars consumed during the fermentation. Xylitol and ethanol productivities (Q_{PX} and Q_{PE} respectively, g/l.h) were determined as the ratio between the xylitol or ethanol concentration (g/l) and the fermentation time (h), respectively.

3. Results and discussion

3.1. Hydrolysate composition and detoxification strategies

Rapeseed straw pretreatment with dilute sulfuric acid was very efficient for solubilization of the feedstock's hemicellulose fraction. In total, 72.3% of the hemicellulosic sugars were recovered in the hydrolysate, xylose being the most representative corresponding to 58% of the total sugars present in the hydrolysate (**Table 1**, RH). Acetic acid was the most abundant among the inhibitor compounds. This organic acid is commonly found in hemicellulosic hydrolysates since it is also released from the hemicellulose structure during pretreatment (Mussatto and Roberto, 2004a). Besides acetic acid, other inhibitor compounds including formic acid, furfural and HMF (sugar's degradation products), and phenolic compounds (from the partial breakdown of lignin) were also present in the hydrolysate. The low tolerance of xylitol-producing microorganisms to the toxicity of furans, acetic acid and phenolic compounds is well-known (Carvalho et al., 2005; Felipe et al., 1995; Morita and Silva, 2000; Mussatto and Roberto, 2004b). For this reason, a detoxification step is required to at least partially remove the toxic compounds present and generate a hydrolysate more suitable for use as fermentation medium. In the specific case of this study, attention was mainly given to the removal of acetic acid and phenolic compounds, which were the two main toxic compounds present in RH (**Table 1**).

Unfortunately, there is not a specific method of detoxification able to efficiently remove all the toxic compounds and a combination of strategies is usually required. In the present study, the original hydrolysate (RH) was initially treated with activated charcoal as this method has been shown to be efficient to remove furans and phenolic compounds from hydrolysates produced from different feedstocks (Carvalho et al., 2005; Lee et al., 2011; Mussatto and Roberto, 2001). In agreement with that, the use of activated charcoal for

detoxification of rapeseed straw hemicellulosic hydrolysate resulted in an almost complete removal of furans and 68% removal in the content of phenolic compounds (**Table 1**, ACH). These results are very interesting, especially for furans, as according to Wannawilai et al. (2017) low concentrations of furfural (up to 0.3 g/l) enhanced the production of xylitol from xylose by *Candida magnoliae* TISTR 5663. On the other hand, the efficiency of activated charcoal to remove formic and acetic acids was very limited (only 27% and 14%, respectively).

Vacuum evaporation process has been suggested as an alternative to remove weak organic acids like acetic and formic acids from hydrolysate as such compounds are partially volatilized under these conditions (Converti et al., 2000; Mussatto and Roberto, 2004a). Besides removing organic acids, this method is also convenient since it promotes an increase in the concentration of sugars in the medium, which, in turn, is beneficial to obtain higher xylitol yield by fermentation (Mussatto and Roberto, 2003). As can be seen in **Table 1** (CH), the hydrolysate concentration step promoted an increase in the content of total sugars to 75.5 g/l (58% xylose, 17% glucose), which was proportional to the concentration factor employed (4.5-fold). Formic and acetic acids were partially volatilized during this process (33% and 56%, respectively), as their concentration increased less than 4.5-fold. On the other hand, the vacuum evaporation process did not remove phenolic compounds, which is in agreement with other authors (Chandel et al., 2011; Parajó et al., 1997).

Phenolic compounds are usually very toxic to microorganisms even when present in low concentrations in the fermentation medium (Mussatto and Roberto, 2004a). These compounds affect cell growth and metabolism decreasing the product formation. For this reason,

additional efforts were done to decrease even more the concentration of phenolic compounds in rapeseed straw hemicellulosic hydrolysate, by submitting this liquor to a treatment with ion-exchange resins, which has been suggested to be effective for phenolic compounds removal (Taherzadeh and Karimi, 2011). When applied for the detoxification of CH, this method resulted in phenolic compounds and formic acid reductions of 41 % and 53%, respectively, while the concentration of acetic acid was only slightly reduced, 18% (**Table 1**, DH).

Finally, the detoxified hydrolysate (DH) was submitted to extraction with organic solvents as a final strategy to reduce the concentration of toxic compounds, mainly acetic acid. The idea behind the extraction with solvents is to transfer the acetic acid from the hydrolysate medium to the solvent phase, which occurs based on concentration gradient and distribution coefficients. Two different organic solvents, ethyl acetate and chloroform, were used in this step. Acetic acid was removed by 62% when ethyl acetate was used and only 12% when chloroform was used. However, these treatments resulted in an increase in the level of phenolic compounds probably due to their concentration during the distillation step to eliminate the residual organic solvent. Overall, extraction with ethyl acetate was more efficient than with chloroform, generating a hydrolysate (EA-DH) with lower level of toxic compounds (**Table 1**).

3.2. Xylitol production from rapeseed straw hemicellulosic hydrolysates

Rapeseed straw detoxified hydrolysate (DH) was used as fermentation medium for xylitol production by *D. hansenii* and *C. guilliermondii*. Fermentation assays were also performed using the hydrolysate after extraction with ethyl acetate (EA-DH) and with

chloroform (CL-DH). Additionally, fermentation assays were carried out using the DH diluted with water (1:1) and supplemented with sugars up to the original level present in DH, in order to have another medium with a different composition in terms of toxic compounds (DDH). Besides evaluating and comparing the performance of the yeasts in the different hydrolysate-based media, these assays made also possible to estimate the level of toxic compounds that the yeasts are able to tolerate without affecting the efficiency of xylose conversion into xylitol. The differences in the composition of the hydrolysates used as fermentation medium can be seen in **Table 1**.

Both yeasts, *D. hansenii* and *C. guilliermondii* had a similar behavior when cultivated in DH (**Fig. 2**). Likewise, the time course of the fermentation from DDH was also very similar for both yeasts. However, when comparing the fermentation of DH and DDH, it is evident the faster consumption of xylose by both yeasts when cultivated in the medium containing lower concentration of toxic compounds (mainly acetic acid and phenolic compounds), DDH (**Figs. 2a,c**). From DDH, xylose was almost exhausted after 72 h, while from DH, more than 13 g/l of xylose remained in the medium at the same time and about 5 g/l after 144 h. Xylitol production was also higher and faster from DDH (**Figs. 2b,d**), resulting in productivities four-fold higher when compared to DH. Concerning cell growth, no difference was observed in the fermentation of both hydrolysates with *C. guilliermondii*. However, *D. hansenii* cell yield was three times higher when the cells were cultivated in DDH than in DH, achieving, in this case, a yield similar to those obtained with *C. guilliermondii* (0.06–0.07 g/g, **Table 2**).

It is worth mentioning that a gradual consumption of acetic acid was observed during the fermentation of DDH by both yeasts, being this acid completely exhausted after 144 h (data not shown). Previous studies also reported consumption of acetic acid during hydrolysate fermentation by *C. guilliermondii* (Morita and Silva, 2000) and during semi-defined medium fermentation by *D. hansenii* (Duarte et al., 2006). According to Felipe et al. (1995), acetic acid can be potentially toxic for *C. guilliermondii* in concentrations above 3 g/l, while lower concentrations of this acid may have a positive effect on xylose uptake, xylitol yield and cell growth. Under these conditions, yeasts are able to assimilate and use acetic acid as an additional source of nutrients for growth (Camargo et al., 2015). Wannawilai and Sirisansaneeyakul (2015) also reported a clear inhibition of acetic acid (3.5 g/l) on the cell growth and xylitol production during the fermentation of sugarcane bagasse hydrolysate by *Candida magnoliae* TISTR 5663. In the present study, DDH hydrolysate contained 2.65 g/l of acetic acid, while the concentration of this acid in DH was double (**Table 1**). This fact would explain the consumption of acetic acid during the fermentation of DDH, but not during the fermentation of DH.

When comparing the hydrolysates further treated with solvents (EA-DH and CL-DH) with DH, it is evident that the extraction with organic solvents improved the fermentability of the hydrolysate. Similar concentration of residual xylose was observed in the media after 144 h of fermentation (**Figs. 2a,c**). However, higher xylitol concentration was obtained in the fermentation of extracted hydrolysates by both yeasts, mainly by *C. guilliermondii* (**Figs. 2b,d**) probably due to their lower level of acetic acid. Fermentation of EA-DH and CL-DH by *C. guilliermondii* surpassed by more than 25% the xylitol yield obtained in DH medium (**Table 2**). Similar improvement in xylitol yield

was not observed by *D. hansenii*. Since the distillation after DH extraction with organic solvents resulted in an increase in the concentration of phenolic compounds in these media and the content of acetic acid in the extracted media was even lower than in DH (**Table 1**), it can be concluded that phenolic compounds in concentrations above 3 g/l were toxic for *D. hansenii*. Moreover, a synergistic effect between acetic acid and phenolic compounds on the growth of *D. hansenii* has been previously reported (Domínguez et al., 1999).

When comparing the performance of the yeasts in the extracted hydrolysates, the best results in terms of xylitol yield and productivity were obtained with *C. guilliermondii* (**Table 2**) independently of the solvent used for extraction. Xylitol yield, in particular (0.53-0.54 g/g) was very close to the value obtained in DDH fermentation (0.57 g/g). However, an important difference was observed in the productivity values, since maximum xylitol production by *C. guilliermondii* in EA-DH and CL-DH media occurred at 144 h, while for DDH medium it occurred at 48 h only. Phenolic compounds could have been the main responsible for such differences since they were present in much lower amount in DDH than in EA-DH and CL-DH, while the concentration of acetic acid was in the same range of values observed for EA-DH (**Table 1**).

Similar to DDH, a gradual consumption of acetic acid was also observed during the fermentation of the EA-DH by both yeasts, reaching approx. 0.2 g/l after 144 h (data not shown). It should be noted that the assimilation of acetic acid by *C. guilliermondii* and *D. hansenii* was only observed during the fermentation of DDH and EA-DH, which contained less than 3 g/l of this acid in the composition, and not during the fermentation of DH and CL-DH (acetic acid concentration > 3 g/l). This fact is in agreement with the studies reported by

Felipe et al. (1995) and Camargo et al. (2015) in terms of threshold level for acetic acid to act as a toxic compound or as a nutritional source for the microorganism. Moreover, both media where acetic acid was assimilated by the yeasts resulted also in the highest productions of xylitol.

In brief, the assays in hydrolysate media revealed that rapeseed straw hemicellulosic hydrolysate can be used as fermentation medium for xylitol production. However, a detoxification step is required to reduce the concentration of toxic compounds and improve the xylitol yield and productivity. The best results of xylitol production ($Y_{PX/S} = 0.54$ g/g and $Q_{PX} = 0.14$ g/l.h) were achieved when using the detoxified hydrolysate submitted to extraction with solvents as fermentation medium, and *C. guilliermondii* as yeast strain. These results, mainly the productivity, can be further improved by decreasing the concentration of toxic compounds present, especially phenolic compounds, as demonstrated through the assays in DDH.

3.3. Xylitol production from semi-defined media vs hydrolysate media

It is well known that other sugars present in the composition of hemicellulosic hydrolysates, mainly hexoses, may affect the xylose conversion to xylitol by yeast (Mussatto et al., 2006). In order to evaluate the influence of the other sugars present on rapeseed straw hemicellulosic hydrolysate composition on xylose metabolism by *D. hansenii* and *C. guilliermondii*, different semi-defined media (DM) were formulated containing the same concentration of sugars present in DH, but in different combinations. DM1 contained all the five

sugars identified in DH (xylose, glucose, arabinose, mannose and galactose), while DM2 was absent in glucose, DM3 was absent in galactose, and DM4 was absent in glucose and galactose. These assays made possible to identify the influence of glucose and galactose on xylose metabolism by both yeasts and revealed also the maximum values of xylitol yield and productivity able to be achieved in a medium containing the mixture of sugars present in DH but without toxic compounds.

As can be seen in **Figs. 3a** and **c**, both yeasts were able to assimilate all the xylose present in the different media after 48 h of fermentation; only a minimal amount ($< 10\%$) was still present at that time in DM3 and DM4 cultivated by *C. guilliermondii* (**Fig. 3c**). Besides xylose, all the other sugars present in the different media were also completely assimilated by both yeasts after 48 h. On the other hand, xylitol production was favored in the absence of glucose, DM2, achieving 27.5 g/l with *D. hansenii* (**Fig. 3b**) and 24.7 g/l with *C. guilliermondii* (**Fig. 3d**). These results suggest inhibition of glucose on xylose metabolism by both yeasts. Such effect is probably associated to a catabolic repression caused by glucose on the enzymes responsible for xylitol production. According to Tochampa et al. (2005), glucose/xylose ratios lower than 10% can even improve the activity of xylose-reductase increasing the xylitol volumetric productivity. However, higher glucose/xylose ratios inhibit the xylose transport into the cell and repress induction of relevant enzymes. In the present study, the glucose/xylose ratio present in rapeseed straw hemicellulosic hydrolysate (as well as in the semi-defined media) was above 30%. Therefore, it can be concluded that not only the toxic compounds present in the hydrolysate medium affected the yeast's performance to convert xylose into xylitol, but glucose also had a negative impact on the bioconversion process.

When compared to DM1, which contained all the sugars present in DH, xylitol production by both yeasts was also improved in the absence of glucose and galactose (DM4) (**Figs. 3b,d**). However, in this case the maximum xylitol production was lower than in DM2 (absent in glucose only), probably because in DM4 part of the xylose was used for cell growth and not for xylitol production. As can be seen in **Table 2**, cell yield was practically the same in DM2 and DM4, by both yeasts, supporting the theory that the yeasts used part of the xylose to keep the same growth level during the cultivations. Assays in absence of galactose only (DM3) revealed that this hexose did not have a negative impact on xylose metabolism by both yeasts, since the maximum xylitol production in DM3 was practically the same as that achieved in DM1 by *D. hansenii*, and even a bit lower than that achieved in DM1 by *C. guilliermondii* (**Figs. 3b,d**). These assays confirm that glucose was the hexose with the most significant negative impact on xylose metabolism by both yeasts.

Assays in DM1 revealed also the maximum production of xylitol able to be achieved by both yeasts when cultivated in a medium with the same composition of sugars present in DH but without toxic compounds. Such values corresponded to xylitol yield and productivity of 0.57 g/g and 0.80 g/l.h, respectively by *D. hansenii*, and 0.55 g/g and 0.43 g/l.h respectively, by *C. guilliermondii* (**Table 2**). These values were in close agreement with those achieved during the fermentation of DDH by both yeasts, with exception only for xylitol productivity by *D. hansenii*, which was better in fermentation from semi-defined medium. These results show that it is not necessary to completely eliminate the toxic compounds from the hydrolysate medium to obtain an efficient conversion of xylose into xylitol. As can

be seen in **Table 1**, DDH still contained acetic acid (2.65 g/l), formic acid (0.60 g/l), and phenolic compounds (0.85 g/l) in its composition and resulted in xylitol yield and productivity similar to DM1 (**Table 2**).

When comparing the yeasts, *C. guilliermondii* showed higher tolerance to the toxic compounds present in DDH than *D. hansenii*, since its performance to convert xylose into xylitol was very similar in DDH and DM1 ($Y_{PX/S} = 0.57$ and 0.55 g/g respectively, and $Q_{PX} = 0.41$ and 0.43 g/l.h, respectively), while xylitol productivity by *D. hansenii* in DDH was only half the value achieved in DM1 (**Table 2**). On the other hand, *D. hansenii* was able to produce xylitol faster than *C. guilliermondii* from semi-defined medium. However, to achieve this productivity from hydrolysate medium, additional detoxification steps would be required to reduce even more the level of toxic compounds present in DDH.

The strategy used in the present study for hydrolysate detoxification followed by extraction with solvents (EA-DH and CL-DH) was not enough to decrease the concentration of toxic compounds to the same level present in DDH (**Table 1**). Even so, xylitol yield was significantly improved (~ 30%) by *C. guilliermondii* when compared to the DH fermentation (independent of the solvent used for extraction), achieving values similar to those obtained from DM1 and DDH (0.55 - 0.57 g/g) (**Table 2**). Similar improvement in xylitol yield was not observed for *D. hansenii* cultivated in EA-DH and CL-DH. However, xylitol productivity was not improved from EA-DH and CL-DH by any of the strains, confirming that both require a lower level of toxic compounds in the medium for a faster conversion of xylose into xylitol.

3.4. By-product formation

Ethanol was detected as a by-product in all the fermentations in hydrolysate and semi-defined media. The formation of this compound is related to the metabolism of hexoses by yeasts. For this reason, ethanol production was lower in DM4 when compared to the other cultivation media (**Table 2**), as mannose was the only hexose sugar present in this medium. Maximum concentration of ethanol was achieved after 24 h of fermentation in all the media containing glucose in the composition. During this time, all the glucose present in the media was consumed by the yeasts, while galactose and mannose were only partially consumed at that time. However, assays in semi-defined media DM2 and DM4 revealed that galactose and mannose were also used for ethanol production by both yeasts (**Table 2**).

Another interesting fact observed during the cultivations was a simultaneous consumption of xylose and glucose by both yeasts with simultaneous formation of xylitol and ethanol, although xylitol was indeed the product produced in higher amount. On the other hand, arabinose was only slightly consumed by the yeasts after exhaustion of xylose from the media, being not observed simultaneous consumption of pentose sugars. Xylitol and ethanol were also partially consumed by the yeasts after exhaustion of the main carbon sources (xylose and glucose), being used for the cell's growth. Similar behavior was reported by Gírio et al. (2000) during the cultivation of *D. hansenii* in mixed substrates. According to these authors, cell growth increased during the fermentation and was also associated to the consumption of ethanol and xylitol by the yeast when xylose concentration in the medium was lower than 3 g/l.

4. Conclusions

Rapeseed straw hemicellulosic hydrolysate can be used as fermentation medium for xylitol production by *D. hansenii* or *C. guilliermondii*. However, *C. guilliermondii* exhibited higher tolerance to the toxic compounds present in the hydrolysate being therefore more attractive for use in this bioconversion process since it would require less detoxification steps. As the detoxification steps result in more expenses, other alternatives including adaptive laboratory evolution of the strains could be an interesting option to overcome the hydrolysate toxicity as well as the negative impact of glucose on xylose metabolism. These results open up new possibilities for the valorization of rapeseed straw.

Acknowledgements

JC López-Linares gratefully acknowledges the research grant received from University of Jaén (Plan de Apoyo). Support from the Novo Nordisk Foundation (Denmark) is also acknowledged.

References

1. Camargo, D., Sene, L., Variz, D.I.L.S., Felipe, M.G.A., 2015. Xylitol bioproduction in hemicellulosic hydrolysate obtained from sorghum forage biomass. *Appl. Biochem. Biotechnol.* 175, 3628–3642.
2. Carvalheiro, F., Duarte, L.C., Lopes, S., Parajó, J.C., Pereira, H., Gírio, F.M., 2005. Evaluation of the detoxification of brewery's spent grain hydrolysate for xylitol production by *Debaryomyces hansenii* CCM1 941. *Process Biochem.* 40, 1215–1223.
3. Chandel, A.K., Silva, S.S., Singh, O.V., 2011. Detoxification of lignocellulosic hydrolysates for improved bioethanol production, in: Bernardes, M.A.S. (Ed.), *Biofuel production-recent developments and prospects*. InTech, Rijeka, Croatia, pp. 225–246.
4. Choi, C.H., Um, B.H., Kim, Y.S., Oh, K.K., 2013. Improved enzyme efficiency of rapeseed straw through the two-stage fractionation process using sodium hydroxide and sulfuric acid. *Appl. Energy*, 102, 640–646.
5. Converti, A., Dominguez, J.M., Perego, P., Silva, S.S., Zilli, M., 2000. Wood hydrolysis and hydrolyzate detoxification for subsequent xylitol production. *Chem. Eng. Technol.* 23, 1013–1020.
6. Cortez DV, Mussatto SI, Roberto IC., 2016. Improvement on d-xylose to xylitol biotransformation by *Candida guilliermondii* using cells permeabilized with Triton X-100 and selected process conditions. *Appl. Biochem. Biotechnol.* 180, 969–979.
7. Domínguez, J.M., Cruz, J.M., Roca, E., Domínguez, H., Parajó, J.C., 1999. Xylitol production from wood hydrolyzates by entrapped *Debaryomyces hansenii* and *Candida guilliermondii* cells. *Appl. Biochem. Biotechnol.* 81, 119–130.

8. Duarte, L.C., Carvalheiro, F., Tadeu, J., Gírio, F.M., 2006. The combined effects of acetic acid, formic acid, and hydroquinone on *Debaryomyces hansenii* physiology. Appl. Biochem. Biotechnol. 130, 461–475.
9. FAOSTAT (2017). Food and Agriculture Organization of the United Nations. <http://faostat3.fao.org/>. Accessed 29 June 2017.
10. Felipe, M.G.A., Vieira, D.C., Vitolo, M., Silva, S.S., Roberto, I.C., Mancilha, I.M., 1995. Effect of acetic acid on xylose fermentation to xylitol by *Candida guilliermondii*. J. Basic Microbiol. 35, 171–177.
11. Gírio, F.M., Amaro, C., Azinheira, H., Pelica, F., Amaral-Collaco, M.T., 2000. Polyols production during single and mixed substrate fermentations in *Debaryomyces hansenii*. Bioresour. Technol. 71, 245–251.
12. Lee, J.M., Venditti, R.A., Jameel, H., Kenealy, W.R., 2011. Detoxification of woody hydrolyzates with activated carbon for bioconversion to ethanol by the thermophilic anaerobic bacterium *Thermoanaerobacterium saccharolyticum*. Biomass Bioenergy 35, 626–636.
13. López-Linares, J.C., Ballesteros, I., Tourán, J., Cara, C., Castro, E., Ballesteros, M., Romero, I., 2015. Optimization of uncatalyzed steam explosion pretreatment of rapeseed straw for biofuel production. Bioresour. Technol. 190, 97–105.
14. Morita, T.A., Silva, S.S., 2000. Inhibition of microbial xylitol production by acetic acid and its relation with fermentative parameters. Appl. Biochem. Biotechnol. 84, 801–808.

15. Mussatto, S.I., 2012. Application of xylitol in food formulations and benefits for health, in: Silva, S.S., Chandel, A.K. (Eds.), D-Xylitol. Springer-Verlag Berlin Heidelberg, Berlin, Germany, pp. 309–323.
16. Mussatto, S.I., 2016. Biomass pretreatment with acids, in: Mussatto, S.I. (Ed.), Biomass fractionation technologies for a lignocellulosic feedstock based biorefinery. Elsevier Inc., Waltham, MA, pp. 169–185.
17. Mussatto, S.I., Dragone, G.M., 2016. Biomass pretreatment, biorefineries and potential products for a bioeconomy development, in: Mussatto, S.I. (Ed.), Biomass fractionation technologies for a lignocellulosic feedstock based biorefinery. Elsevier Inc., Waltham, MA, pp. 1–22.
18. Mussatto, S.I., Roberto, I.C., 2001. Hydrolysate detoxification with activated charcoal for xylitol production by *Candida guilliermondii*. Biotechnol. Lett. 23, 1681–1684.
19. Mussatto, S.I., Roberto, I.C., 2002. Xylitol: a sweetener with benefits for human health. Braz. J. Pharm. Sci. 38, 401–413.
20. Mussatto, S.I., Roberto, I.C., 2003. Xylitol production from high xylose concentration: evaluation of the fermentation in bioreactor under different stirring rates. J. Appl. Microbiol. 95, 331–337.
21. Mussatto, S.I., Roberto, I.C., 2004a. Alternatives for detoxification of diluted-acid lignocellulosic hydrolyzates for use in fermentative processes - a review. Bioresour. Technol. 93, 1–10.

22. Mussatto, S.I., Roberto, I.C., 2004b. Optimal experimental condition for hemicellulosic hydrolyzate treatment with activated charcoal for xylitol production. *Biotechnol. Prog.* 20, 134–139.
23. Mussatto, S.I., Silva, C.J.S.M., Roberto, I.C., 2006. Fermentation performance of *Candida guilliermondii* for xylitol production on single and mixed substrate media. *Appl. Microbiol. Biotechnol.* 72, 681–686.
24. Parajó, J.C., Domínguez, H., Domínguez, J.M., 1997. Improved xylitol production with *Debaryomyces hansenii* Y-7426 from raw or detoxified wood hydrolysates. *Enzyme Microb. Technol.* 21, 18–24.
25. Pinkowska, H., Wolak, P., Oliveros, E., 2013. Production of xylose and glucose from rapeseed straw in subcritical water - Use of Doehlert design for optimizing the reaction conditions. *Biomass Bioenerg.* 58, 188–197.
26. Singleton, V.L., Rossi, S.A., 1965. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am. J. Enol. Vitic.* 16, 144–158.
27. Taherzadeh, M.J., Karimi, K., 2011. Fermentation inhibitors in ethanol processes and different strategies to reduce their effects, in: Pandey, A., Larroche, C., Ricke, S.C., Dussap, C.-G., Gnansounou, E. (Eds.), *Biofuels*. Elsevier, pp. 287–311.
28. Tochampa, W., Sirisansaneeyakul, S., Vanichsriratana, W., Srinophakun, P., Bakker, H.H.C., Chisti, Y., 2005. A model of xylitol production by the yeast *Candida mogii*. *Bioprocess Biosyst. Eng.* 28, 175–183.

29. Wannawilai, S., Sirisansaneeyakul, S., 2015. Economical production of xylitol from *Candida magnolia* TISTR 5663 using sugarcane bagasse hydrolysate. Kasetsart J. (Nat. Sci.) 49, 583-596.
30. Wannawilai, S., Lee, W-C., Chisti, Y., Sirisansaneeyakul S., 2017. Furfural and glucose can enhance conversion of xylose to xylitol by *Candida magnoliae* TISTR 5663. J. Biotechnol. 241, 147-157.

Figure Captions

Figure 1. Schematic representation of the steps performed for the production and detoxification of rapeseed straw hemicellulosic hydrolysate. RH: raw hydrolysate (obtained by dilute acid pretreatment of rapeseed straw); ACH: activated charcoal-treated hydrolysate; CH: ACH 4.5-fold concentrated; DH: detoxified hydrolysate (CH further treated with ion-exchange resin); DDDH: DH diluted with water (1:1) and supplemented with sugars up to the same concentration present in DH; EA-DH: DH further treated with ethyl acetate; CL-DH: DH further treated with chloroform.

Figure 2. Xylose consumption (A) and xylitol production (B) by *Debaryomyces hansenii* and xylose consumption (C) and xylitol production (D) by *Candida guilliermondii*. Fermentation assays in rapeseed straw hemicellulosic hydrolysate based media. DH: detoxified hydrolysate; DDH: DH diluted with water (1:1) and supplemented with sugars up to the same concentration present in DH; EA-DH: DH further treated with ethyl acetate; CL-DH: DH further treated with chloroform.

Figure 3. Xylose consumption (A) and xylitol production (B) by *Debaryomyces hansenii* and xylose consumption (C) and xylitol production (D) by *Candida guilliermondii*. Fermentation assays in semi-defined media. DM1: medium formulated with the same concentration of sugars present in the detoxified hydrolysate (DH); DM2: same composition of DM1 but without glucose; DM3: same composition of DM1 but without galactose; DM4: same composition of DM1 but without glucose and galactose.

Figure 1

Figure 2

Figure 3

Table 1. Composition of rapeseed straw hemicellulosic hydrolysate (g/l) in the original form and after each detoxification step.

Sugars				Inhibitor compounds			
Hydrolysate	Glucose	Xylose	Galactose	Arabinose	Mannose	Formic acid	Phenolic compounds
RH	2.96 ± 0.04	10.30 ± 0.07	2.20 ± 0.04	1.72 ± 0.03	0.66 ± 0.03	1.20 ± 0.02	1.93 ± 0.27
ACH	2.81 ± 0.12	9.91 ± 0.38	2.09 ± 0.11	1.66 ± 0.06	0.62 ± 0.03	0.88 ± 0.11	0.61 ± 0.09
CH	12.49 ± 0.19	43.82 ± 0.79	9.13 ± 0.19	7.31 ± 0.08	2.76 ± 0.06	2.65 ± 0.03	2.94 ± 0.13
DH	11.68 ± 0.09	40.59 ± 0.21	8.67 ± 0.12	6.79 ± 0.09	2.22 ± 0.06	1.25 ± 0.02	1.74 ± 0.01
EA-DH	14.83 ± 0.41	49.66 ± 0.90	10.88 ± 0.40	8.41 ± 0.37	2.79 ± 0.43	1.22 ± 0.11	3.74 ± 0.12
CL-DH	13.04 ± 0.25	43.62 ± 0.49	9.57 ± 0.11	7.64 ± 0.14	2.45 ± 0.12	1.49 ± 0.18	3.43 ± 0.17
DDH	11.82 ± 0.09	41.51 ± 0.10	8.79 ± 0.13	6.83 ± 0.08	2.25 ± 0.05	0.60 ± 0.02	0.85 ± 0.01

RH: raw hydrolysate (obtained by dilute acid pretreatment of rapeseed straw);

ACH: activated charcoal-treated hydrolysate

CH: ACH 4.5-fold concentrated

DH: detoxified hydrolysate (CH further treated with ion-exchange resin)

EA-DH: DH further treated with ethyl acetate

CL-DH: DH further treated with chloroform

DDH: DH diluted with water (1:1) and supplemented with sugars up to the same concentration present in DH

Table 2. Fermentation parameters obtained during the production of xylitol from rapeseed straw detoxified hydrolysates (DH) and semi-defined media (DM) by *Debaryomyces hansenii* and *Candida guilliermondii*.

Fermentation medium	Yeast strain	Time (h) *	Xylitol yield (Y _{PXS} , g/g)	Xylitol productivity (Q _{PX} , g/l.h)	Cell yield (Y _{X/S} , g/g)	Ethanol yield (Y _{PE/S} , g/g)	Ethanol productivity (Q _{PE} , g/l.h)
DH	D	72	0.42 ± 0.01	0.16 ± 0.00	0.02 ± 0.00	0.18 ± 0.00	0.13 ± 0.00
	C	144	0.42 ± 0.01	0.10 ± 0.00	0.06 ± 0.00	0.12 ± 0.01	0.05 ± 0.00
DDH	D	48	0.55 ± 0.01	0.42 ± 0.00	0.07 ± 0.00	0.19 ± 0.01	0.24 ± 0.01
	C	48	0.57 ± 0.01	0.41 ± 0.06	0.06 ± 0.00	0.19 ± 0.00	0.23 ± 0.03
EA-DH	D	144	0.45 ± 0.01	0.12 ± 0.01	0.13 ± 0.01	0.10 ± 0.00	0.04 ± 0.01
	C	144	0.54 ± 0.01	0.14 ± 0.00	0.12 ± 0.01	0.13 ± 0.00	0.06 ± 0.01
CL-DH	D	144	0.38 ± 0.00	0.09 ± 0.00	0.04 ± 0.00	0.11 ± 0.00	0.05 ± 0.00
	C	144	0.53 ± 0.03	0.13 ± 0.01	0.05 ± 0.00	0.13 ± 0.01	0.06 ± 0.00
DM1	D	24	0.57 ± 0.01	0.80 ± 0.04	0.08 ± 0.00	0.11 ± 0.00	0.23 ± 0.04
	C	48	0.55 ± 0.01	0.43 ± 0.01	0.06 ± 0.00	0.21 ± 0.00	0.25 ± 0.00
DM2	D	24	0.78 ± 0.01	1.15 ± 0.01	0.08 ± 0.01	0.13 ± 0.02	0.23 ± 0.03
	C	48	0.64 ± 0.01	0.51 ± 0.02	0.08 ± 0.00	0.15 ± 0.02	0.15 ± 0.01
DM3	D	48	0.57 ± 0.03	0.41 ± 0.00	0.14 ± 0.01	0.20 ± 0.01	0.17 ± 0.00
	C	48	0.47 ± 0.00	0.35 ± 0.00	0.09 ± 0.00	0.22 ± 0.01	0.21 ± 0.01
DM4	D	48	0.65 ± 0.02	0.50 ± 0.01	0.10 ± 0.00	0.06 ± 0.00	0.05 ± 0.00
	C	48	0.67 ± 0.01	0.49 ± 0.02	0.09 ± 0.01	0.06 ± 0.01	0.05 ± 0.00

* Time at which maximum xylitol production was achieved.

D: *Debaryomyces hansenii*; C: *Candida guilliermondii*; Fermentation time: time of maximum xylitol production;

DH: detoxified hydrolysate; DDH: DH diluted with water (1:1) and supplemented with sugars up to the same concentration present in DH;

EA-DH: DH further treated with ethyl acetate; CL-DH: DH further treated with chloroform.

DM1: semi-defined medium formulated with the same concentration of sugars present in DH; DM2: same composition of DM1 but without glucose;

DM3: same composition of DM1 but without galactose; DM4: same composition of DM1 but without glucose and galactose.

HIGHLIGHTS

Rapeseed straw hemicellulosic hydrolysate was used for xylitol production

Different strategies were evaluated for the hydrolysate detoxification

C. guilliermondii exhibited higher tolerance to the toxic compounds than *D. hansenii*

Toxic compounds and glucose concentration affected the yeast's performance

Partial removal of toxic compounds was enough to achieve an efficient bioconversion