Organophosphate esters in dust samples collected from Danish homes and daycare centers

Langer, Sarka; Fredricsson, Malin; Weschler, Charles J.; Bekö, Gabriel; Strandberg, Bo; Remberger, Mikael; Toftum, Jørn; Clausen, Geo

Published in:
Chemosphere

Link to article, DOI:
[10.1016/j.chemosphere.2016.04.016](https://doi.org/10.1016/j.chemosphere.2016.04.016)

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Langer, S., Fredricsson, M., Weschler, C. J., Bekö, G., Strandberg, B., Remberger, M., Toftum, J., & Clausen, G. (2016). Organophosphate esters in dust samples collected from Danish homes and daycare centers. *Chemosphere*, 154, 559-566. <https://doi.org/10.1016/j.chemosphere.2016.04.016>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Organophosphate esters in dust samples collected from
Danish homes and daycare centers**

Sarka Langer^{a,*}, Malin Fredricsson^a, Charles J. Weschler^{b,c}, Gabriel Bekö^c,
Bo Strandberg^d, Mikael Remberger^e, Jørn Toftum^b, Geo Clausen^b

^aIVL Swedish Environmental Research Institute Ltd., P.O. Box 53021, SE-400 14
Göteborg, Sweden

^bInternational Centre for Indoor Environment and Energy, Department of Civil
Engineering, Technical University of Denmark, DK-2800 Lyngby, Denmark

^cEnvironmental and Occupational Health Sciences Institute, Rutgers University,
Piscataway, New Jersey, 08854, United States

^dOccupational and Environmental Medicine, Sahlgrenska Academy at University of
Gothenburg, Box 414, SE-405 30, Göteborg, Sweden

^eIVL Swedish Environmental Research Institute Ltd., P.O. Box 210 60, SE-100 31
Stockholm, Sweden

*Corresponding author:

sarka.langer@ivl.se; IVL Swedish Environmental Research Institute Ltd., P.O. Box 53021,
SE-400 14 Göteborg, Sweden; phone: +46 (0)10 788 6897

Keywords: exposure, flame retardants, indoor environment, plasticizers, SVOC

Highlights

- Organophosphates were determined in dust from 497 homes and 151 daycare centers
- Mass fractions in dust from daycare centers were larger than in dust from homes
- Estimated exposures to organophosphates via dust ingestion were relatively low
- The country-to-country distribution of organophosphates is more variable than that of phthalate esters

Abstract

Organophosphates are used in a wide range of materials and consumer products and are ubiquitous in indoor environments. Certain organophosphates have been associated with various adverse health effects. The present paper reports mass fractions of organophosphates in dust samples collected from 500 bedrooms and 151 daycare centers of children living in Odense, Denmark. The identified compounds include: tris(isobutyl) phosphate (TIBP), tri-n-butyl phosphate (TNBP), tris(2-chloroethyl) phosphate (TCEP), tris(2-chloroisopropyl) phosphate (TCIPP), tris(1,3-dichloroisopropyl) phosphate (TDCIPP), tris(2-butoxyethyl) phosphate (TBOEP), triphenylphosphate (TPHP), 2-ethylhexyl-diphenyl phosphate (EHDPP), tris(2-ethylhexyl) phosphate (TEHP) and tris(methylphenyl) phosphate (TMPP). Both the number of organophosphates with median values above the limit of detection and the median values were higher for samples from daycare centers than for samples from homes. Organophosphates with median mass fractions above the limit of detection were: TCEP from homes ($6.9 \mu\text{g g}^{-1}$), and TCEP ($16 \mu\text{g g}^{-1}$), TCIPP ($5.6 \mu\text{g g}^{-1}$), TDCIPP ($7.1 \mu\text{g g}^{-1}$), TBOEP ($26 \mu\text{g g}^{-1}$), TPHP ($2.0 \mu\text{g g}^{-1}$) and EHDPP ($2.1 \mu\text{g g}^{-1}$) from daycare centers. When present, TBOEP was typically the most abundant of the identified OPs. The sum of the organophosphate dust mass fractions measured in this study was roughly in the mid-range of summed mass fractions reported for dust samples collected in other countries. On a global scale, the geographical distribution of organophosphates in indoor dust is quite variable, with higher concentrations in industrialized countries. This trend differs from that for phthalate esters, whose geographic distribution is more homogeneous. Exposure to organophosphates via dust ingestion is relatively low, although there is considerable uncertainty in this assessment.

1. Introduction

Semivolatile organic compounds (SVOC) are ubiquitous in indoor environments, and, depending on their chemical properties, can persist for a very long time (Weschler and Nazaroff, 2008; Shin et al., 2013). Indoor SVOCs commonly include organophosphate triesters. The use of the organophosphates is increasing because of their favorable properties as both flame retardants and plasticizers (Marklund et al., 2003). The halogenated organophosphates are mostly used as flame retardants while the non-halogenated compounds are generally used as plasticizers (Andresen et al., 2004). Organophosphorus flame retardants are found in a wide range of commercial products including textiles, rubber, polyurethane foam (PUF), antistatic formulations, cellulose, cotton, cutting oils, electronic equipment, glues, engineering thermoplastics, epoxy resins and phenolic resins (Marklund et al., 2005). Polyvinyl chloride (PVC) is an example of a product where phosphorus flame retardants also function as plasticizers (Marklund et al., 2003).

Organophosphates have been implicated in various adverse health effects, including skin irritation and contact dermatitis in humans, and neurological and carcinogenic effects in rats (WHO 1991 a,b, 1998, 2000). Araki et al. (2014) have recently found associations between organophosphates flame retardants and atopic dermatitis, asthma and allergic rhinitis. Organophosphates have also been associated with altered hormone levels and decreased semen quality in men (Meeker and Stapleton, 2010).

The widespread use of organophosphates in everyday products results in redistribution from their original source into other indoor compartments, including indoor air, airborne particles, settled dust and all exposed indoor surfaces (*e.g.* Marklund et al., 2005; Wensing et al., 2005). They were first reported in indoor airborne particles in 1980 (Weschler, 1980). Their concentrations have been found to be much higher indoors than outdoors, reflecting the much greater presence of their sources indoors compared to outdoors (Weschler et al., 1984; Staaf and Östman, 2005; Wensing et al., 2005). The mass fractions of various

organophosphates in settled dust have been determined over the past 14 years in samples collected from homes, daycare centers and workplaces in various countries throughout the world (Brommer and Harrad, 2015; He et al., 2015; Hoffman et al., 2015; Luongo and Östman, 2015; Abdallah and Covaci, 2014; Araki et al., 2014; Cequier et al., 2014; Fan et al., 2014; Fromme et al., 2014; Shin et al., 2014; Tajima et al., 2014; Ali et al., 2013; Brommer et al., 2012; Kim et al., 2013; Dodson et al., 2012; Ali et al., 2012a; Ali et al., 2012b, Bergh et al., 2011, Van den Eede et al., 2011; Dirtu et al., 2010; Kazanawa et al., 2010; Meeker and Stapleton, 2010; Stapleton et al., 2009; García et al., 2007; Wensing et al., 2005; Marklund et al., 2003; Nagorka and Ullrich, 2003, Kersten and Reich, 2003; Becker et al., 2002; Ingerowski et al., 2001). Human exposure to organophosphates is apparent from biomonitoring studies, which have detected various organophosphate metabolites in urine samples (e.g., Reemtsma et al., 2011; Fromme et al., 2014; Hoffman et al., 2014). A recent study found that levels of such metabolites in urine correlated with levels of organophosphates in dust, and that children's exposure to organophosphates can be related to levels found in their indoor environments (Cequier et al., 2015).

The aim of the present paper is threefold: to report the mass fractions of selected organophosphate esters (OPEs) measured in the dust samples collected from children's bedrooms ($n = 500$) and daycare centers ($n = 151$) as part of the IECH investigation; to examine potential correlations between levels of the individual organophosphates, both in the children's bedrooms and daycare facilities; and to compare the results with those reported in other studies to derive a sense of variations over geographic location and time. The information presented in this paper can be used to improve understanding of Danish children's exposure to organophosphate esters.

2. Material and Methods

2.1 Study design

The Danish study “Indoor Environment and Children’s Health (IECH)” has investigated the relationship between children’s health and their indoor environments. The design of the IECH study and the methods used in the investigations have been described by Clausen et al. (2012). Questionnaires were sent to 17,486 families that had children between the ages of one and five. All children in the study lived on the Danish island of Fyn (482,310 inhabitants). The final database contained questionnaires from 11,082 children, *i.e.* the response rate was 63%. Using the information from the survey, 500 children between the ages of three and five and living in Odense (166,000 inhabitants) were selected for the case-base study: 200 “cases” with asthma/allergies and 300 randomly selected “bases”, *i.e.* both healthy and sick children. All of the daycare facilities attended by these children were also included in the study (n = 151). Settled dust was collected from the children’s bedrooms from non-floor surfaces such as shelves, ledges, and window sills as well as from horizontal non-floor surfaces in the daycare centers. Dust from floors was avoided, since it may be contaminated by direct contact with organophosphorus flame retardants or plasticizers present at the surfaces of flooring materials that contain these additives. The dust samples were subsequently analyzed for five phthalate esters and three polycyclic aromatic hydrocarbons (Langer et al., 2010) as well as for squalene and cholesterol (Weschler et al., 2011).

2.2 Dust collection

Dust was collected from non-floor surfaces in the children’s bedrooms and daycares using ALK dust filters (ALK-Abelló A/S, Hørsholm, Denmark) mounted in a holder connected to a vacuum cleaner. Collection of dust from plastic surfaces and textiles was

avoided. In 5% of the homes (randomly chosen) a field blank was obtained (filter was mounted in the holder but no sampling was performed).

2.3 Chemical analysis

Prior to sampling, the virgin filters were pre-conditioned at a temperature of 23 ± 1 °C and relative humidity of $50 \pm 5\%$, weighed and assigned an ID number. The loaded filters or blanks were returned to the laboratory (shipped at ambient temperature) where they were conditioned, weighed, wrapped in aluminum foil and re-packed in their original bag. The net dust mass on the loaded filters ranged from 43 to 1396 mg (average 198 mg; median 160 mg). Of the total number of more than 700 filters that were analyzed, 33 (~ 5%) were field blanks, and 42 were laboratory blanks.

The organic compounds in the dust were extracted using the procedure described by Rudel et al. (2003). Logistics and details of the chemical analysis are presented in the Supplementary Material.

The organophosphate esters evaluated in this study, in order of retention times, were: tris(isobutyl) phosphate (TIBP), tri-n-butyl phosphate (TNBP), tris(2-chloroethyl) phosphate (TCEP), tris(2-chloroisopropyl) phosphate (TCIPP), tris(1,3-dichloroisopropyl) (TDCIPP), tris(2-butoxyethyl) phosphate (TBOEP), tri-phenylphosphate (TPHP), 2-ethylhexyl-diphenyl phosphate (EHDPP), tris(2-ethylhexyl) phosphate (TEHP) and tris(methylphenyl) phosphate (TMPP). The compounds were named and abbreviated according to the nomenclature proposed by Bergman et al. (2012).

The original target compounds in these dust samples were phthalate esters and polycyclic aromatic hydrocarbons (Langer et al., 2010). After the samples were extracted and analyzed, we recognized peaks in the Total Ion Chromatograms (TIC) that corresponded to organophosphate esters. As the organophosphates were not targeted from the beginning, their

recoveries could not be determined using appropriate isotopically labelled standards. Given that phthalate esters and organophosphate esters span similar ranges of physical and chemical properties (Wensing et al., 2005), we assume that the efficiency with which they are extracted from the dust, using 6% diethyl ether in hexane for 16 h, is similar (Toda et al., 2004). To support this, we have determined the recoveries of a phthalate (d_4 -di-isononyl phthalate (DnOP- d_4); c.f. Langer et al., 2010) and a phosphate (triarylphosphate (TAP); c.f. Brommer et al., 2012) from dust samples collected onboard the Swedish icebreaker Oden. Seven dust samples were extracted and analyzed by the same procedure as the samples from this study. The recoveries were similar: 0.96 ± 0.09 and 0.99 ± 0.09 for TAP and DnOP- d_4 , respectively. Furthermore, using methods similar to those used in the present study, good procedural recoveries of 90-110 % for the organophosphates have been reported in the literature (e.g., Van den Eede et al., 2011).

The organophosphate esters were first identified by their mass spectra in the TICs and matched against their mass spectra in the NIST (National Institute of Standards and Technology) Mass Spectral Library. They were later also matched against the mass spectra of the authentic compounds used to prepare the calibration standards. The 10 target compounds were then quantified with the “Extraction Ion” procedure from the full scan chromatograms using the m/z -ratio (mass-to-charge) for the most abundant ion in each compound (Table 1); $m/z = 266$ was selected for 1,2,3,4-tetrachloronaphthalene. For the purpose of quantification, a five-point standard curve for each of the ten organophosphates was prepared, in the range of $0.1 - 5.0 \text{ ng } \mu\text{L}^{-1}$ with $2.0 \text{ ng } \mu\text{L}^{-1}$ of 1,2,3,4-tetrachloronaphthalene as the injection standard, also using the m/z ratio for the most abundant ions (Table 1). The limits of detection (LOD) were calculated from the standard deviations of the repeated injections of the analytical standards plotted against their concentrations (Taylor, 1987). The value of the standard deviation as the concentration approaches zero (S_0) was determined as the intercept of the

best-fit straight line of this plot. Limits of detection were then calculated as $3 \times S_0$. These limits of detection are on the high end of those reported in the literature (see studies summarized in Table 4). The magnitude of the detection limits using GC/MS is partially a consequence of the operating mode of the mass spectrometer. Selected Ion Monitoring (SIM) is more sensitive (lower amounts detected) than using extracted ions from the TICs. Nonetheless, some studies report comparable limits of detection (e.g. Tajima et al. (2014).

The organophosphates in the blanks were all below the limits of detection, with the exception of TIBP ($0.06 \pm 0.05 \mu\text{g}/\text{filter}$). After correcting for the amount of TIBP found in the blank samples, the values of all the other organophosphates below the LOD were replaced with a value one-half the LOD. The amounts of the organophosphates in the dust samples were expressed as their mass fractions in units of μg of the compound per gram of dust ($\mu\text{g g}^{-1}$).

3. Results and Discussion

Reliable measurements were obtained for dust samples from 497 homes (1 sample lost, 2 with analytical issues) and from all 151 daycare centers. Not all the targeted compounds were found in all samples. Only TCEP, TCIPP and TDCIPP were identified in more than a third of the home samples, while only TCEP, TCIPP, TDCIPP, TBOEP, TPHP and EHDPP were identified in more than a third of the daycare samples. Descriptive statistics of the results are presented in Table 2 for both the homes and daycare centers. For each organophosphate the table lists detection frequency (DF; i.e., the percentage of samples with values above the LOD), LOD, and the median, 90th percentile, and maximum mass fractions.

3.1 The organophosphates in dust from homes and daycare centers

Figure 1 (a - c) presents plots of the natural logarithm of selected organophosphate mass fractions in homes and daycare centers against the cumulative distribution function, with the x-axis scaled according to the normal error function. The figures show only the organophosphates evaluated in this study that had median values above $\frac{1}{2}$ the LOD. For each compound, in both homes and daycare centers, the plots are roughly linear, indicating that the distribution of the concentrations is better described as “log-normal” than “normal”. This is consistent with expectations (Ott, 1995) and earlier studies of indoor volatile organic compounds (*e.g.* Shields et al., 1996).

Table 3 presents Spearman correlation coefficients for the individual organophosphates. While there are many statistically significant correlations, there are only a few compounds that are strongly correlated. These are TCEP/TIBP in homes and TDCIPP/TCIPP and EHDPP/TPHP in daycare centers. The mass fractions of all organophosphates except TIBP, TEHP and TMPP were significantly different between homes and daycare centers ($p < 0.05$; non-parametric two-sample Wilcoxon rank-sum (Mann Whitney) test, as the data were not normally distributed; STATA software, release 11.2 for Windows, StataCorp LP, College Station, Texas, USA).

TCEP was found in $\geq 50\%$ of the dust samples from both the children’s homes and daycare centers. The median mass fraction of TCEP from the daycare centers ($16 \mu\text{g g}^{-1}$) was more than twice as high as the concentration from homes ($6.9 \mu\text{g g}^{-1}$). The median mass fractions of TCIPP and TDCIPP were noticeably different for the daycare center dust ($5.6 \mu\text{g g}^{-1}$ and $7.1 \mu\text{g g}^{-1}$, respectively) and the home dust ($0.5 \mu\text{g g}^{-1}$ and $0.6 \mu\text{g g}^{-1}$, respectively; both $= \frac{1}{2}$ LOD).

These findings indicate the presence of more flame retarded products at the daycare centers than in the homes. TCEP and TCIPP, which are chlorinated organophosphorus flame

retardants, are mostly used in flexible foams such as polyurethane foam (PUF) that can be found in products such as cushions, upholstered furniture, pillows and mattresses (WHO, 1998). Bergh et al. (2011) also found TCEP and TCIPP to have higher concentrations in daycare centers compared to homes, with TCEP being the second most common organophosphate in the daycare centers. In the study by Marklund et al. (2003), TCEP and TCIPP had relatively low concentrations in the homes as well as the one daycare center they sampled, but the concentration was still higher in the daycare center compared with the homes. Concentrations of chlorinated OPEs in dust from homes in general seem to be lower compared to other public places such as work environments (Bergh et al., 2011). Marklund et al. (2003) found higher levels of TDCIPP ($1.8 \mu\text{g g}^{-1}$) in a daycare center compared to the average value for two homes ($0.75 \mu\text{g g}^{-1}$); the number of samples in that study is, however, very limited. The findings of Bergh et al. (2011) differ from the results of this study: they detected similar levels in the homes and daycare centers ($10 \mu\text{g g}^{-1}$ vs. $9.1 \mu\text{g g}^{-1}$). Fromme et al. (2014) report TCEP and TCIPP from German daycare centers with TCIPP being the second most abundant organophosphate in their study. In the present study it should be noted that even though TDCIPP was detected in a smaller percentage of samples from homes (DF = 41%) than in samples from daycare centers (DF = 67%), when it was detected in homes the mass fractions tended to be high ($P90 = 54 \mu\text{g g}^{-1}$, $\text{max} = 860 \mu\text{g g}^{-1}$).

It is apparent from Figure 1c that the range of mass fractions is largest for TBOEP in dust samples from daycare centers, with several extreme values exceeding $1\,000 \mu\text{g g}^{-1}$. Although the detection frequency was much smaller in homes (28% in homes vs. 68% in daycares), when it was detected, its mass fraction was often high ($P90 = 44 \mu\text{g g}^{-1}$, $\text{max} = 1\,300 \mu\text{g g}^{-1}$). TBOEP is often used in floor polishes, acting as a leveling agent as well as a plasticizer. In homes or daycare centers where the mass fraction is high, TBOEP may be a

constituent of the floor care product (Weschler, 1980; Marklund et al., 2003; Araki et al., 2014; Cequier et al., 2014).

For small children, dust ingestion is common and, for certain SVOCs, can be a substantial fraction of a child's total intake. Assuming a 15 kg child ingesting 60 mg of dust a day (US EPA, 2009) in either indoor environment (home or daycare), daily intakes of the median mass fractions of organophosphates would be several orders of magnitude below the reference doses (RfD) presented in Van den Eede et al. (2011, and references therein). The mass fractions in dust required to exceed the RfD under such a hypothetical exposure are presented for each organophosphate in Table S1 of the Supplementary Material. The fractions of dust samples from homes with organophosphate mass fractions exceeding these limits were 0.6% for TDCIPP and 0.8% for TBOEP; in the case of daycare centers they were 4.0% for TCEP and 13% for TBOEP.

The organophosphates identified in the dust from bedrooms and daycare facilities are also anticipated to be present in the gas phase (Weschler and Nazaroff, 2010), associated with airborne particles (Weschler et al., 2008), and present on indoor surfaces. Enriched surface concentrations are specifically anticipated in the case of materials that contain organophosphates as flame retardants or plasticizers or are treated with polishes that contain such compounds. Hence, in addition to dust ingestion, intakes also occur via inhalation and dermal absorption. The latter can occur following contact with organophosphates on surfaces or, for certain compounds, directly from the gas phase (Weschler and Nazaroff, 2012; 2014). Based on estimated transdermal permeability coefficients (final column of Table 1), dermal absorption from the gas phase may only be meaningful ($k_{p-g} > 0.2 \text{ m h}^{-1}$) for TBOEP, TDCIPP, TPHP and TMPP. However, the gas phase concentrations of these low-vapor pressure compounds tend to be small, limiting the magnitude of their direct uptake from air. The overarching point is that, when inhalation intake and dermal absorption are also

considered, the percentage of daily intakes exceeding the RfD is anticipated to be larger than those presented in the previous paragraph.

3.2 Comparison with other studies

To our knowledge, in terms of the number of samples (497 homes and 151 daycare centers) in combination with the number of the analyzed organophosphates (10 compounds), this is the largest study of its kind to date. A larger number of samples were collected in a few studies, but they targeted fewer organophosphates (Becker et al., 2002; Ingerowski et al., 2001).

A comparison of median mass fractions of organophosphates from this study with mass fractions previously reported in the literature is presented in Table 4. Compound by compound comparisons of our results with those reported in the literature are not appropriate, since different studies target different compounds. Hence, the following discussion is based on “when available” values from the referenced literature. Additionally, when comparing the present results with a number of earlier studies from around the world, it should be borne in mind that those studies may have relied on different sampling and analytical methods. For the purpose of direct comparison, values reported in the literature as below the limit of detection or quantification were replaced with LOD/LOQ values.

In the case of the tributyl phosphate isomers, TNBP was identified in most studies, while TIBP was identified in a few studies. Median TNBP or TIBP values exceeded $1 \mu\text{g g}^{-1}$ in samples from Germany (Nagorka and Ulrich, 2003), Belgium (van den Eede et al., 2011), Sweden (Bergh et al., 2011; Luongo and Östman, 2015) and Japan (Kanazawa et al., 2010; Araki et al., 2014), while the median values for the homes and daycare centers in the present study were low (for both homes and daycares 0.01 and $0.05 \mu\text{g g}^{-1}$ for TIBP and TNBP,

respectively). TPHP was also found in most studies. Values for both the Danish homes and daycare centers were comparable with the Swedish (Bergh et al., 2011) and Norwegian (Cequier et al., 2014) homes and daycare centers. TPHP values $> 4 \mu\text{g g}^{-1}$ were found in dust samples from German, U.S., Japanese, English and Swedish homes (Nagorka and Ulrich, 2003; Stapleton et al., 2009; Araki et al., 2014; Kanazawa et al., 2010; Brommer and Harrad, 2015; Luongo and Östman, 2015) and Swedish workplaces (Bergh et al., 2011). EHDPP, TEHP and TMPP were reported in approximately half of all the studies. The median values were low, and some of the median values that are seemingly elevated reflect high LODs.

The chlorinated organophosphates were found in more than 2/3 of the studies summarized in Table 4. In the case of TCEP, the highest mass fractions were found in the Swedish (Bergh et al., 2011) and Danish (this work) daycare centers. Comparable median concentrations were measured in homes in Japan (Araki et al., 2014; Kanazawa et al., 2010), USA (Dodson et al., 2012) and Denmark (this work), as well as in the Swedish workplaces (Berg et al., 2011). The chlorinated organophosphates are mainly used as flame retardants and commonly added to polyurethane foam (WHO, 1998). The levels of the chlorinated organophosphates in the dust samples are expected to mirror the level of fire protection mandated by law for the furniture and furnishings in the respective countries.

In Table 4 the organophosphate whose levels in indoor dust stand out is TBOEP (tris-butoxyethyl phosphate). It was found in the majority of the tabulated studies. This compound was already detected in size-selected indoor aerosols in the very early 1980s (Weschler, 1980), and is common in floor polishes (see above). Median values as high as $\sim 1600 \mu\text{g g}^{-1}$ were found in samples from Swedish daycare centers (Bergh et al., 2011) and in Japanese homes (Kanazawa et al., 2010). Araki et al. (2014) report median mass fractions of TBOEP of $\sim 500 \mu\text{g g}^{-1}$ in floor dust; houses with wooden floors had significantly higher levels of TBOEP in indoor dust, probably due to more frequent floor care.

Figure 2 shows the sum of organophosphate mass fractions in dust from various countries based on data from Table 4. Only entries with three or more individual organophosphates are included. The highest totals are for Swedish and Japanese homes and are due to high mass fractions of TBOEP. The sum of organophosphate levels from the Danish daycare centers was almost five times higher than the corresponding value from the Danish homes, which was comparable to most sums for other countries. The sums for daycare centers were also higher in Germany, Sweden, Norway and the UK compared to their counterparts for the homes. Subtracting TBOEP from the sums leads to a more even geographic distribution. Very low levels were found in samples from Egypt, Kuwait, New Zealand, Pakistan, the Philippines and Romania. The median mass fractions of the individual organophosphates from these countries were generally an order of magnitude lower than from the other countries. In general, the mass fraction of organophosphate esters in dust displayed greater country-to-country variation than observed for the mass fraction of phthalate esters in dust (Langer et al., 2010), reflecting less extensive global use of the organophosphates.

4. Summary

The mass fractions of organophosphates in dust were higher in samples from daycare centres than in samples from homes. The targeted compounds were present at levels below the limit of detection in the majority of the dust samples. Median mass fractions above the $\frac{1}{2}$ the limit of detection were measured for TCEP from the home samples and TCEP, TCIPP, TDCIPP, TBOEP, TPHP and EHDPP from the daycare center samples. Although TBOEP was not the most frequently detected of the ten targeted organophosphates, its median mass fraction was the highest of all the organophosphates. The values measured in this study were in line with values reported for other industrialized countries.

The organophosphates show larger geographic variation than phthalates, which are more or less evenly distributed over the globe (Langer et al., 2010). For the children targeted in this study, based on mass fractions found in dust from their homes and daycare centers, intake of organophosphates through dust ingestion was in most instances below current guidelines. However, inhalation and dermal absorption are also anticipated to be substantial exposure pathways for organophosphates in indoor environments.

Acknowledgement

We would like to thank the VILLUM FOUNDATION for generous support of this study and the participating families for their cooperation. We would also like to acknowledge Andreas Fischer for the analytical work with the samples.

References

- Abdallah, M.A.-E., Covaci, A., 2014. Organophosphate flame retardants in indoor dust from Egypt: Implications for human exposure. *Environ. Sci. Technol.* 48, 4782-4789.
- Ali, N., Dirtu, A.C., Van den Eede, N., Goosey, E., Harrad, S., Neels, H., Mannetje, A., Coakley, J., Douwes, J., Covaci A., 2012a. Occurrence of alternative flame retardants in indoor dust from New Zealand: Indoor sources and human exposure assessment. *Chemosphere* 88, 1276-1282.
- Ali, N., Van den Eede N., Dirtu A.C., Neels H., Covaci A., 2012b. Assessment of human exposure to indoor organic contaminants via dust ingestion in Pakistan. *Indoor Air* 22, 200-211.
- Ali, N., Ali, L., Mehdi, T., Dirtu, A.C., Al-Shammari, F., Neels, H., Covaci ,A., 2013. Levels and profiles of organochlorines and flame retardants in car and house dust from Kuwait and Pakistan: Implication for human exposure via dust ingestion. *Environ.Int.* 55, 62-70.
- Andresen, J.A., Grundmann, A., Bester, K., 2004. Organophosphorus flame retardants and plasticisers in surface waters. *Sci. Total Environ.* 332, 155-166.

375 Araki, A., Saito, I., Kazanawa, A., Morimoto, K., Nakayama, K., Shibata, E., Tanaka, M.,
376 Takigawa, T., Yoshimura, T., Chikara, H., Saiko, Y., Kishi R., 2014. Phosphorus flame
377 retardants in indoor dust and their relation to asthma and allergies of inhabitants. *Indoor Air*
378 24, 3-15.

379 Becker, K., Seiwert, M., Kaus, S., Krause, C., Schults, C., Seifert B., 2002. German
380 Environmental Survey (GerES III): Pesticides and other pollutants in indoor dust. *Proceedings*
381 *Indoor Air* 2002, 883-886.

382 Bergh, C., Torgrip, R., Emenius, G., Östman, C., 2011. Organophosphate and phthalate esters
383 in air and settled dust – a multi-location indoor study. *Indoor Air* 21, 67-76.

384 Bergman, Å., Rydén, A., Law, R.J., de Boer, J., Covaci, A., Alae, M., Birnbaum, L., Petreas,
385 M., Rose, M., Sakai, S., Van den Eede, N., van der Veen, I., 2012. A novel abbreviation
386 standard for organobromine, organochlorine and organophosphorus flame retardants and
387 some characteristics of the chemicals. *Environ. Int.* 49, 57-82.

388 Brommer, S., Harrad, S., Van den Eede, N., Covaci, A., 2012. Concentrations of
389 organophosphate esters and brominated flame retardants in German indoor dust samples. *J.*
390 *Environ. Monit.* 14, 2482-2487.

391 Brommer, S., Harrad, S., 2015. Sources and human exposure implications of concentrations
392 of organophosphate flame retardants in dust from UK cars, classrooms, living rooms, and
393 offices. *Environ.Int.* 83, 202-207.

394 Cequier, E., Ionas, A.C., Covaci, A., Marcé, R.M., Becher, G., Thomsen, C., 2014. Occurrence
395 of a broad range of legacy and emerging flame retardants in indoor environments in Norway.
396 *Environ.Sci. Technol.* 48, 6827-6835.

397 Cequier, E., Sakhi A.K., Marcé, R.M., Becher, G., Thomsen, K., 2015. Human exposure
398 pathways to organophosphate triesters – A biomonitoring study of mother-child pairs.
399 *Environ. Int.* 75, 159-165. Clausen, G., Høst, A., Toftum, J., Bekö, G., Weschler, C., Callesen,
400 M., Buhl, S., Ladegaard, M., Langer, S., Andersen, B., Sundell, J., Bornehag, C.G.,
401 Sigsgaard, T., 2012. Children's health and its association with indoor environments in Danish
402 homes and daycare centres – methods. *Indoor Air* 22, 467-475.

403 Dirtu, A.C., Ali, N., van den Eede, N., Neels, H., Covaci, A., 2012. Country specific
404 comparison for profile of chlorinated, brominated and phosphate organic contaminants in
405 indoor dust. Case study for eastern Romania. *Environ. Int.* 49, 1-8.

406 Dodson, R.E., Perowich, L.J., Covaci, A., van den Eede, N., Ionas, A.C., Dirtu, A.C., Brody,
407 J.G., Rudel, R.A., 2012. After the PBDE phase-out: a broad suite of flame retardants in repeat
408 house dust samples from California. *Environ. Sci. Technol.* 46, 13056-13066.

409 Fan, X., Kubwabo C., Rasmussen, P.E., Wu F., 2014. Simultaneous determination of thirteen
410 organophosphate esters in settled indoor house dust and comparison between two sampling
411 techniques. *Sci. Total Environ.* 491-492, 80-86.

412 Fromme, H., Lahrz, T., Kraft, M., Fembacher, L., Mach, C., Dietrich, S., Burkardt, R.,
 413 Völkel, W., Göen, T., 2014. Organophosphate flame retardants and plasticizers in the air and
 414 dust in German daycare centers and human biomonitoring in visiting children (LUOE 3).
 415 *Environ.Int.* 71, 158-163.

416 García, M., I. Rodríguez, Cela R., 2007. Microwave-assisted extraction of organophosphate
 417 flame retardants and plasticizers from indoor dust samples. *J. Chromatography A*, 1152, 280-
 418 286.

419 Hansen, D., Volland, G., Krause, G., Zöltze, D., 2001. Determination and occurrence of
 420 phosphororganic compounds (POV) in dust and indoor air. *Gefahrstoffe – Reinhaltung der*
 421 *luft* 61, 13-17 (in German).

422 He, C.-T., Zheng, J., Qiao, L., Chen, S.-J., Yang, J.-Z., Yuan, J.-C., Yang, Z.-Y., Mai, B.-X.,
 423 2015. Occurrence of organophosphorus flame retardants in indoor dust in multiple
 424 environments of southern China and implications for human exposure. *Chemosphere* 133, 47-
 425 52.

426 Hoffman, K., Daniels, J.L., Stapleton, H.M., 2014. Urinary metabolites of organophosphate
 427 flame retardants and their variability in pregnant women. *Environ. Int.* 63, 169-172.

428 Hoffman, K., Garantziotis, S., Birnbaum, L.S., Stapleton, H., 2015. Monitoring indoor
 429 exposure to organophosphate flame retardants: Hand wipes and house dust. *Environ. Health*
 430 *Perspect.* 123, 160-165.

431 Ingerowski, G., Friedle, A., Thummulla, J., 2001. Chlorinated ethyl and isopropyl phosphoric
 432 acid triesters in the indoor environment – an interlaboratory exposure study. *Indoor Air* 11,
 433 145-149.

434 Kazanawa, A., Saito, I., Araki, A., Takeda, M., Ma, M., Saijo, Y., Kishi R., 2010. Association
 435 between indoor exposure to semi-volatile organic compounds and building-related symptoms
 436 among the occupants of residential dwellings. *Indoor Air* 20, 72-84.

437 Kersten, W., Reich, T., 2003. Non-volatile organic substances in Hamburg indoor dust.
 438 *Gefahrstoffe Reinhaltung der Luft* 63, 85-91 (in German).

439 Kim, J.-W., Isobe, T., Sudaryanto, A., Malarvannan, G., Chang, K.-H., Muto, M., Prudente,
 440 M., Tanabe, S., 2013. Organophosphorus flame retardants in house dust from the Philippines:
 441 occurrence and assessment of human exposure. *Environ. Sci. Poll. Res.* 20, 812-822.

442 Langer, S., Weschler, C.J., Fischer, A., Bekö, G., Toftum, J., Clausen, G., 2010. Phthalate
 443 and PAH concentrations in dust collected from Danish homes and daycare centers. *Atmos.*
 444 *Environ.* 44, 2294-2301.

445 Luongo, G., Östman, C., 2015. Organophosphate and phthalate esters in settled dust from
 446 apartment building in Stockholm. *Indoor Air*, DOI: 10.1111/ina.12217.

447 Marklund, A., Andersson, B., Haglund, P., 2003. Screening of organophosphorus compounds
448 and their distribution in various indoor environments. *Chemosphere* 53, 1137-1146.

449 Marklund, A., Andersson, B., Haglund, P., 2005. Organophosphorus flame retardants and
450 plasticizers in air from various indoor environments. *J. Environ. Monit.* 7, 814-819.

451 Meeker, J.D., Stapleton, H.M., 2010. House dust concentration of organophosphate flame
452 retardants in relation to hormone levels and semen quality parameters. *Environ. Health*
453 *Perspect.* 118, 318-323.

454 Nagorka, R., Ullrich, D., 2003. Determination of organophosphorus flame retardants in indoor
455 dust: Screening method with GC/NPD. *Gefahrstoffe Reinhaltung der Luft* 63, 79-84. (in
456 German).

457 Ott, W. R., 1995. *Environmental Statistics and Data Analysis*. Ann Arbor, Lewis Publishers.

458 Reemtsma, T., Lingott, J., Roegler, S., 2011. Determination of 14 monoalkyl phosphates,
459 dialkyl phosphates and dialkyl thiophosphates by LC-MS/MS in human urinary samples. *Sci.*
460 *Total Environ.* 409, 1990-1993.

461 Rudel, R.A., Camann, D.E., Spengler, J.D., Korn, L.R., Brody, J.G., 2003. Phthalates,
462 alkylphenols, pesticides, polybrominated diphenyl ethers, and other endocrine-disrupting
463 compounds in indoor air and dust. *Environ. Sci. Technol.* 37, 4543-4553.

464 Sagunski, H., Ingerowski, G., Mattulat, A., Scheutvinkel, M., 1997. Exposure and
465 environmental health evaluation of tris(2-chloroethyl)phosphate. *Umweltmedizin in*
466 *Forschung und Praxis* 2, 185-192 (in German).

467 Shields, H.C., Fleischer, D.M., Weschler, C.J., 1996. Comparisons among VOCs measured in
468 three types of US commercial buildings with different occupant densities. *Indoor Air* 6, 2-17.

469 Shin, H.M., McKone, T.E., Tulse, N.S., Clifton, M.S., Bennett, D.H., 2013. Indoor residence
470 times of semivolatile organic compounds: Model estimation and field evaluation. *Environ. Sci.*
471 *Technol.* 47, 859-867.

472 Shin, H.M., McKone, T.E., Nishioka, M.G., Fallin, M.D., Croen, L.A., Hertz-Picciotto, I.,
473 Newschaffer, C.J., Bennett, D.H., 2014. Determining source strength of semivolatile organic
474 compounds using measured concentrations in indoor dust. *Indoor Air* 24, 206-271.

475 Staaf, T., Östman, C., 2005. Organophosphate triesters in indoor environments. *J. Environ.*
476 *Monit.* 7, 883-887.

477 Stapleton, H.M., Klosterhaus, S., Eagle, S., Fuh, J., Meeker, J.D., Blum, A., Webster, T.F.,
478 2009. Detection of Organophosphate Flame Retardants in Furniture Foam and U.S. House
479 Dust. *Environ. Sci. Technol.* 43, 7490-7495.

480 Tajima, S., Araki, A., Kawai, T., Tsuboi, T., Bamai, Y.A., Yoshioka, E., Kanazawa, A.,
481 Cong, S., Kiski, R., 2014. Detection and intake assessment of organophosphate flame
482 retardants in house dust in Japanese dwellings. *Sci. Total Environ.* 478, 190-199.

483 Taylor, J., 1987. Quality assurance of chemical measurements. Lewis Publishers. p. 79-81.
 484 ISBN 0-87371-097-5.

485 Toda, H., Sako, K., Yagome, Y., Nakamura, T., 2004. Simultaneous determination of
 486 phosphate esters and phthalate esters in clean room air and indoor air by gas chromatography-
 487 mass spectrometry. *Anal. Chim. Acta* 519, 213–218.

488 US EPA, 2009. Highlights of the Child-specific exposure factors handbook (Final Report).
 489 US Environmental Protection Agency, Washington, DC, EPA/600/R-08/135, Available:
 490 [http://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=200445&CFID=55467099&CFTOKEN](http://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=200445&CFID=55467099&CFTOKEN=81128915)
 491 [=81128915](http://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=200445&CFID=55467099&CFTOKEN=81128915). Accessed 2016 January 13.

492 Van den Eede N., Dirtu A.C., Neels H., Covaci A., 2011. Analytical developments and
 493 preliminary assessment of human exposure to organophosphate flame retardants from indoor
 494 dust. *Environ. Int.* 37, 454-461.

495 Wensing, M., Uhde, E., Salthammer, T., 2005. Plastics additives in the indoor environment –
 496 flame retardants and plasticizers. *Sci. Total Environ.* 339, 19-40.

497 Weschler, C.J., 1980. Characterization of selected organics in size-fractionated indoor
 498 aerosols. *Environ. Sci. Technol.* 14, 428-431.

499 Weschler, C.J., 1984. Indoor-outdoor relationships for nonpolar organic constituents of
 500 aerosol particles. *Environ. Sci. Technol.* 18, 648-652.

501 Weschler, C.J., Nazaroff, W.W., 2008. Semivolatile organic compounds in indoor
 502 environments. *Atmos. Environ.* 42, 9018-9040.

503 Weschler, C.J., Salthammer, T., Fromme, H., 2008. Partitioning of phthalates among the gas
 504 phase, airborne particles and settled dust in indoor environments. *Atmos. Environ.* 42, 1449-
 505 1460.

506 Weschler, C.J., Nazaroff, W.W., 2010. SVOC partitioning between the gas phase and settled
 507 dust indoors. *Atmos. Environ.* 44, 3609-3620.

508 Weschler, C.J., Langer, S., Fischer, A., Bekö, G., Toftum, J., Clausen, G., 2011. Squalene and
 509 cholesterol in dust from Danish homes and daycare centers. *Environ. Sci. Technol.* 45, 3872-
 510 3879.

511 Weschler, C.J., Nazaroff, W.W. 2012. SVOC exposure indoors: fresh look at dermal
 512 pathways. *Indoor Air* 22, 356-377.

513 Weschler, C.J., Nazaroff, W.W., 2014. Dermal uptake of organic vapors commonly found in
 514 indoor air. *Environ. Sci. Technol.* 48, 1230-1237.

515 WHO (1991a). Tri-n-butyl Phosphate, Environmental Health Criteria 112, World Health
 516 Organization, Geneva.

517 WHO (1991b). Triphenyl Phosphate, Environmental Health Criteria 111, World Health
 518 Organization, Geneva.

519 WHO (1998). Flame retardants: Tris(chloropropyl) Phosphate and Tris(2-chloroethyl)
520 Phosphate, Environmental Health Criteria 209, World Health Organization, Geneva.

521 WHO (2000). Flame retardants: Tris(2-butoxyethyl) Phosphate and Tris(2-ethylhexyl)
522 Phosphate and Tetrakis(hydroxymethyl) Phosphonium salts, Environmental Health Criteria
523 218, World Health Organization, Geneva.

Table 1. Names, abbreviations, CAS numbers, molecular weight (MW), the m/z of the most abundant ion and transdermal permeability coefficients k_{p-g} ($m\ h^{-1}$) of the organophosphates investigated in this work.

Chemical name	Abbreviation*	CAS #	MW	m/z	k_{p-g}^{**} ($m\ h^{-1}$)
Tris(isobutyl) phosphate	TIBP	126-71-6	266	99	0.0002
Tri-n-butyl phosphate	TNBP	126-73-8	266	99	0.002
Tris(2-chloroethyl) phosphate	TCEP	115-96-8	285	63	0.039
Tris(2-chloroisopropyl) phosphate	TCIPP	13674-84-5	327	125	0.011
Tris(1,3-dichloroisopropyl) phosphate	TDCIPP	13674-87-8	431	75	1.3
Tris(2-butoxyethyl) phosphate	TBOEP	78-51-3	398	57	5.3
Tri-phenyl phosphate	TPHP	115-86-6	326	326	0.68
2-Ethylhexyl-diphenyl phosphate	EHDPP	1241-94-7	362	251	0.046
Tris(2-ethylhexyl) phosphate	TEHP	78-42-2	434	99	0.002
Tris(methylphenyl) phosphate	TMPP	78-32-0	368	368	0.69

*The compounds are abbreviated according to the nomenclature presented by Bergman et al. (2012).

**Calculated using the procedure outlined in Weschler and Nazaroff (2012).

Table 2. Mass fractions ($\mu g\ g^{-1}$) of the organophosphates found in settled dust from homes (n = 497) and daycare centers (n = 151). Values in each cell are listed as “homes/daycare centers”. Detection frequency (DF) is percentage of samples found in concentrations above the detection limit (LOD). LOD values are the same for both homes and daycare centers. Median mass fractions and P90, respectively, above the LOD are reported in the table.

Compound	DF (%)	LOD	Median	P90	Max.
TIBP	31/30	0.027	*/*	0.95/0.37	37/2.5
TNBP	22/14	0.11	*/*	0.26/0.13	4.0/4.3
TCEP	69/78	0.60	6.9/16	42/230	230/1 800
TCIPP	37/72	1.1	*/5.6	6.1/41	100/350
TDCIPP	41/67	1.2	*/7.1	54/69	860/320
TBOEP	28/68	0.40	*/26	44/420	1 300/11 000
TPHP	21/65	0.94	*/2.0	1.8/16	91/210
EHDPP	29/77	0.14	*/2.1	1.2/28	11/540
TEHP	0.80/1.3	2.5	*/*	*/*	11/3.8
TMPP	1.6/1.6	0.89	*/*	*/*	18/36

*below LOD

Table 3. Spearman correlation coefficients for the individual organophosphates.

Homes	TIBP	TNBP	TCEP	TCIPP	TDCIPP	TBOEP	TPHP	EHDPP	TEHP	TMPP
TIBP	1									
TNBP	0.400*	1								
TCEP	0.572*	0.408*	1							
TCIPP	0.333*	0.375*	0.395*	1						
TDCIPP	0.192*	0.175*	0.376*	0.119*	1					
TBOEP	0.283*	0.145*	0.316*	0.248*	0.154*	1				
TPHP	0.314*	0.214*	0.366*	0.251*	0.098*	0.217*	1			
EHDPP	0.260*	0.242*	0.315*	0.351*	0.084	0.312*	0.422*	1		
TEHP	-0.002	0.105*	0.089*	0.020	0.049	0.046	0.018	0.073	1	
TMPP	0.048	-0.010	0.026	0.032	0.014	0.094*	0.104*	0.106*	-0.018	1
Daycare centers	TIBP	TNBP	TCEP	TCIPP	TDCIPP	TBOEP	TPHP	EHDPP	TEHP	TMPP
TIBP	1									
TNBP	0.491*	1								
TCEP	0.378*	0.253*	1							
TCIPP	0.436*	0.302*	0.478*	1						
TDCIPP	0.406*	0.204*	0.367*	0.593*	1					
TBOEP	0.281*	0.104	0.261*	0.282*	0.233*	1				
TPHP	0.219*	0.190*	0.313*	0.492*	0.407*	0.209*	1			
EHDPP	0.184*	0.193*	0.265*	0.438*	0.264*	0.178*	0.597*	1		
TEHP	0.041	0.079	-0.009	0.020	-0.066	0.027	0.085	0.083	1	
TMPP	0.222*	0.129	0.117	0.175*	0.071	0.039	0.231*	0.108	-0.019	1

*p < 0.05

Table 4. Organophosphate mass fractions ($\mu\text{g g}^{-1}$) in dust samples collected from residences and other indoor spaces as reported in relevant studies. Present results listed for comparison; the values in bold are organophosphates in this study with the median above LOD. All values are medians unless stated otherwise.

Reference	Country	Location	n	TIBP	TNBP	TCEP	TCIPP	TDCIPP	TBOEP	TPHP	EHDPP	TEHP	TMPP
Sagunski et al. (1997)	Germany	Homes	59	---	---	0.9	---	---	---	---	---	---	---
Ingerowski et al. (2001)	Germany	Homes	983	---	---	0.66	---	---	---	---	---	---	---
Ingerowski et al. (2001)	Germany	Homes	436	---	---	---	0.57	---	---	---	---	---	---
Becker et al. (2002)	Germany	Homes	199	---	---	< 0.1	---	---	5.8	0.3	---	< 0.1	< 0.1
Kersten and Reich (2003)	Germany	Homes	65	---	0.4	1.6	1.4	1.2	5	2.9	---	0.2	2.2
Marklund et al. (2003)	Sweden	Various indoor spaces	15	---	0.35	1.4	2.4	1.1	31	3.1	---	0.16	---
Nagorka and Ulrich (2003)	Germany	Homes	28	---	2.5	2.5	---	1.7	16.1	6.5	0.8	0.8	---
García et al. (2007)	Spain	Homes	8	0.22	0.23	0.51	3.8	---	9.4	1.9	---	---	---
Stapleton et al. (2009)	USA	Homes	50	---	---	---	0.57	1.9	---	7.4	---	---	---
Kanazawa et al. (2010)	Japan	Homes	41	---	1.4	7.5	18.7	4	1 570	5.4	---	4.3	---
Meeker and Stapleton (2010)	USA	Homes	50	---	---	---	---	1.8	---	5.5	---	---	---
Van den Eede et al. (2011)	Belgium	Homes	33	3.0	0.13	0.23	1.4	0.36	2.0	0.50	---	---	0.20
Van den Eede et al. (2011)	Belgium	Shops	15	1.0	0.21	0.59	2.9	0.76	3.6	2.0	---	---	0.02
Bergh et al. (2011)	Sweden	Home	10	1.1	0.3	2.1	1.6	10	4	1.2	0.5	n.d.	---
Bergh et al. (2011)	Sweden	Day-care centers	10	0.7	1.2	30	3.1	9.1	1 600	1.9	0.8	0.1	---
Bergh et al. (2011)	Sweden	Work	10	1.3	0.2	6.7	19	17	87	5.3	1.0	n.d.	---
Ali et al. (2012a)	New Zealand	Homes (Floor dust)	34	---	0.08	0.11	0.35	---	4.02	0.6	---	---	0.1
Ali et al. (2012a)	New Zealand	Homes (Mattresses)	16	---	0.07	0.04	0.25	---	1.55	0.24	---	---	0.2
Ali et al. (2012b)	Pakistan	Homes	31	---	0.014	---	---	---	0.027	0.094	---	---	---
Brommer et al. (2012)*	Germany	Homes	6	---	0.13	0.20	0.74	< 0.080	0.73	0.38	---	---	0.1
Brommer et al. (2012)*	Germany	Offices	10	---	0.22	0.12	3.0	0.15	7.0	2.5	---	---	0.4
Dirtu et al. (2012)	Romania	Homes	47	0.39	0.045	0.10	0.86	0.06	1.5	0.50	---	---	0.50
Dodson et al. (2012)	USA 2006	Homes	16	0.084	0.032	5.1	2.1	2.8	12	---	0.61	<0.2	1
Dodson et al. (2012)	USA 2011	Homes	16	< 0.080	< 0.080	2.7	2.2	2.1	11	---	0.56	<0.2	0.68

Ali et al. (2013)	Kuwait	Homes	15	0.054	0.058	0.71	1.46	0.36	0.86	0.43	0.19	0.065	0.16
Ali et al. (2013)	Pakistan	Homes	15	0.025	<0.020	0.015	<0.020	<0.005	0.017	0.175	0.067	0.02	0
Kim et al. (2013)	Philippines	Homes (Malate)	37	---	0.019	0.034	---	---	---	0.089	0.11	0.14	0
Kim et al. (2013)	Philippines	Homes (Payatas)	20	---	0.020	0.016	---	---	---	0.074	0.034	0.041	0
Araki et al. (2014)	Japan	Homes (Floor)	148	---	1.0	5.8	8.7	2.8	508	4.5	---	2.1	<4.0
Araki et al. (2014)	Japan	Homes (Multi-surface)	120	---	1.2	8.3	25.8	10.8	111	11.5	---	1.5	<4.0
Tajima et al. (2014)	Japan	Homes (Floor)	48	---	< 0.36	< 0.65	0.74	< 0.59	31	0.9	---	< 0.67	<4.0
Tajima et al. (2014)	Japan	Homes (Upper surfaces)	128	---	0.74	1.17	2.23	< 0.59	26.55	3.13	---	< 0.67	<4.0
Abdallah and Covaci (2014)	Egypt	Homes	20	0.023	0.017	0.022	0.028	0.072	0.018	0.067	0.042	---	---
Abdallah and Covaci (2014)	Egypt	Offices	20	0.028	0.023	0.031	0.080	0.049	0.143	0.073	0.048	---	---
Cequier et al. (2014)	Norway	Homes	48	---	0.06	0.41	2.68	0.50	13.4	0.98	0.62	---	0.31
Cequier et al. (2014)	Norway	Classrooms	6	---	0.04	1.21	2.04	1.49	87.2	1.54	2.34	---	0.06
Fan et al., 2014	Canada	Homes (fresh dust)	134		0.25	0.80	1.4	2.7	31.9	1.70	0.54	---	2.60
Fromme et al., 2014	Germany	Day-care centers	63	<0.30	<0.30	0.40	2.68	---	225	0.50	1.10	0.50	---
Shin et al., 2014	USA	Homes	30	---	---	0.50	---	3.6	---	2.0	---	---	---
Brommer & Harrad, 2015	UK	Homes (living rooms)	32	---	<0.03	0.81	21	0.71	---	3.3	1.6	---	0
Brommer & Harrad, 2015	UK	Offices	61	---	<0.03	0.87	33	0.48	---	4.3	5.3	---	<0.01
Brommer & Harrad, 2015	UK	Schools, day-care centers	28	---	0.12	0.86	16	0.51	---	4.1	29	---	<0.01
He et al., 2015	China	Rural homes	25	---	0.14	1.93	1.22	0.15	0.20	1.09	0.31	0.19	---
He et al., 2015	China	Urban homes	11	---	0.08	3.78	0.75	0.13	0.32	0.15	0.36	0.14	---
Hoffman et al., 2015	USA	Homes	49	---	---	---	---	1.39	---	1.02	---	---	---
Luongo and Östman, 2015	Sweden	Homes	62	5.3	5.6	4.0	11	2.0	11	4.3	2.7	---	2.7
This work	Denmark	Homes	497	0.03	0.1	6.9	1.1	1.2	0.4	0.9	0.1	2.5	0.9
This work	Denmark	Day-care centers	151	0.03	0.1	16.2	5.6	7.1	26.2	2.0	2.1	2.5	0.9

n.d. not detected; --- not available/investigated; * mean values.

Figure 1.

Figure 2.

Figure Captions

Figure 1. Cumulative frequency distributions of measured mass fractions of a) TCEP in dust from homes (filled circle) and daycare centers (open circles); b) TCIPP (filled triangles) and TDCIPP (open circles) in dust from daycare centers; c) TBOEP (filled triangles), TPHP (open circles) and EHDPP (open triangles) in dust from daycare centers.

Figure 2. Geographic distribution of the sum of organophosphates reported in indoor dust from homes, daycare centers (DCC), classrooms (school) and various work environments (offices, shops, work). The data are selected from Table S1; entries with two or more individual organophosphates are included. The references can be identified by mapping country and number of samples with corresponding information in Table 4. Results from this work are marked with an “x”.