



Marine fish traits follow fast-slow continuum along coastal-offshore gradient

Beukhof, Esther; Frelat, Romain ; Pécuchet, Lauréne; Fock, Heino; Punzón, Antonio; Sólmundsson, Jón; Moellmann, Christian; Lindegren, Martin

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Beukhof, E., Frelat, R., Pécuchet, L., Fock, H., Punzón, A., Sólmundsson, J., Moellmann, C., & Lindegren, M. (2017). *Marine fish traits follow fast-slow continuum along coastal-offshore gradient*. Poster session presented at 3rd International Workshop on Trait-based Approaches to Ocean Life, Bergen, Norway.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Marine fish traits follow environmental gradients across European seas

Esther Beukhof¹, Romain Frelat², Laurène Pécuchet¹, Heino Fock³, Antonio Punzon⁴, Jón Sólmundsson⁵, Christian Möllmann² and Martin Lindegren¹

¹Centre for Ocean Life, DTU Aqua, Lyngby, Denmark; ²Institute of Hydrobiology and Fisheries, University of Hamburg, Hamburg, Germany; ³Institute of Sea Fisheries, Thünen Institute, Hamburg, Germany; ⁴Instituto Español de Oceanografía, Madrid, Spain; ⁵Marine Research Institute, Reykjavik, Iceland

Introduction

One of the major goals in biogeography is describing and understanding species distributions. However, when focusing on species-environment relationships, one may miss the mechanistic understanding of what underlies these distributions. Therefore, trait-environment relationships are useful in explaining where species occur, since traits determine which environment a species can inhabit¹. In this study, we apply this trait-based approach to Europe's marine fish communities.

Aims of this study

- 1) To identify key traits for marine fish that explain fish species distributions;
- 2) To identify the most important relationships between marine fish traits and the environment.


Data

SITES & SPECIES:


~2200 sampling sites and ~250 species from scientific surveys done in the continental shelf seas of Europe (combination of publicly available and institutional data).

ENVIRONMENT: depth, temperature, salinity, Chlorophyll (Chl) concentration, seasonal variability in temperature and monthly variability in Chl-concentration (NOAA; GlobColour).

TRAITS: 7 quantitative traits



Methods



• RLQ analysis²

Multivariate ordination approach combining all three datasets (R, L and Q). Gives a score to all data onto the same ordination axes.


• Fourth-corner analysis²

Makes use of species distributions when testing for correlations between traits and environmental variables.

Results


TRAITS

The first axis of the **RLQ analysis** (RLQ 1) explained 95% of the variation. In terms of traits, RLQ 1 represents a **fast-slow continuum**, mainly characterized by age at maturity, lifespan and the growth coefficient K.




ENVIRONMENT

RLQ 1 represents a **depth gradient**, along which also vary: temperature (Temp.), seasonal variability in temperature (Temp.ssn), chlorophyll concentration (Chl) and seasonal variability in Chl (Chl.sd).



SPECIES

RLQ 1 identified **specialist species** having typical fast or slow life histories, e.g. anchovy and common skate. **Generalist species**, e.g. Atlantic cod and Atlantic herring, were not strongly characterized by RLQ 1 in terms of their traits and distribution. This indicates that the trait values they have allow them to inhabit a wide range of environments. Indeed, the most common species (i.e. that were present at most sampling sites) were generalist species, whereas typical specialist species were less abundant.



SAMPLING SITES

The RLQ 1 scores of the sampling sites follow a **coastal-to-offshore gradient**, thereby corresponding to the depth gradient that was found to be an important determinant for fish species distributions.

The **fourth-corner analysis** confirmed the importance of the three traits that characterize the fast-slow continuum as well as depth as an important environmental variable. The analysis also revealed significant relationships of the fast-slow traits with temperature and with the variability in temperature and Chl-concentration.

	Temp.ssn	Chl.sd	Chl	Depth	Temp.	Salinity
Age at maturity	**	**	*	**	**	
Lifespan	**	**	*	**	**	
K	**	**	*	**	**	
Max. length	*					
Trophic level						
Offspring size						
Fecundity						

Significance: + (red), + (blue)

Conclusions

We demonstrated that marine fish species can be characterized according to their traits along a fast-slow continuum. Traits in particular related to growth and maturity are key for explaining fish species distributions. The trait continuum is strongly determined by a depth gradient. Along this gradient, other factors vary, such as temperature, productivity and seasonality, which help in explaining species distributions and the structure of marine fish communities.

References

1. Verberk, W. C. E. P. *et al.* Delivering on a promise: integrating species traits to transform descriptive community ecology into a predictive science. *Freshw. Sci.* 32, 531-547 (2007).
2. Dray, S. *et al.* Combining the fourth-corner and the RLQ methods for assessing trait responses to environmental variation. *Ecology* 95, 14-21 (2014).



Esther Beukhof
PhD student
estb@aqu.aqua.dtu.dk
@EstherBeukhof



The results of this poster reflect only the author's view and the Commission is not responsible for any use that may be made of the information it contains.