Adaptive feeding behavior and functional responses in pelagic copepods

Kjørboe, Thomas; Saiz, Enrico; Tiselius, Peter; Andersen, Ken Haste

Published in:
Limnology and Oceanography

Link to article, DOI:
[10.1002/lno.10632](https://doi.org/10.1002/lno.10632)

Publication date:
2018

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Kjørboe, T., Saiz, E., Tiselius, P., & Andersen, K. H. (2018). Adaptive feeding behavior and functional responses in pelagic copepods. *Limnology and Oceanography*, 63(1), 308-321. <https://doi.org/10.1002/lno.10632>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ADAPTIVE FEEDING BEHAVIOR AND FUNCTIONAL RESPONSES IN ZOOPLANKTON

Thomas Kiørboe¹, Enric Saiz², Peter Tiselius³, Ken H. Andersen¹

¹Centre for Ocean Life, DTU Aqua, Technical University of Denmark

⁵University of Gothenburg, Department of Biological and Environmental Sciences – Kristineberg,
Fiskebäckskil, Sweden

⁴Institut de Ciències del Mar – CSIC, Barcelona, Catalunya, Spain

For Review Only

Abstract

Zooplankton may modify their feeding behavior in response to prey availability and presence of predators with implications to populations of both predators and prey. Optimal foraging theory predicts that such responses result in a type II functional response for passive foragers and a type III response for active foragers, with the latter response having a stabilizing effect on prey populations. Here, we test the theoretical predictions and the underlying mechanisms in pelagic copepods that are actively feeding (feeding-current feeders), passively feeding (ambushers), or that can switch between the two feeding modes. In all cases individual behaviors are consistent with the resulting functional response. Passive ambushing copepods have invariant foraging behavior and a type II functional response, as predicted. When foraging actively, the species with switching capability change its functional response from type II to III and modify its foraging effort in response to prey density and predation risk, also as predicted by theory. The obligate active feeders, however, follow a type II response inconsistent with the theoretical prediction. A survey of the literature similarly finds consistent type II response in ambush feeding copepods, but variable (II or III) responses in active feeders. We examine reasons for why observed behaviors at times deviate from predictions, and discuss the population dynamics and food web implications of the two types of functional responses and their underlying mechanisms.

28 ***Introduction***

29 The functional response in feeding rate to prey concentration is the primary quantification of
30 predator-prey interactions and it has direct implications for population dynamics and stability of
31 both prey and predators (Holling 1965; Murdoch 1977). A type II functional response is typically
32 described by some saturating function of prey concentration in which the parameters (maximum
33 ingestion and prey search rates) are assumed constant. However, the assumption of constant search
34 rate may not be true, and at both low and high prey densities it may be advantageous for the
35 predator or grazer to reduce its food searching effort to minimize its exposure to predators and
36 energetic costs of food acquisition (Abrams 1982, 1990; Werner and Anholt 1993; Visser 2007). At
37 low prey densities, the costs are simply not warranted by the gains and the grazer may reduce or
38 stop searching for prey, which may lead to a sigmoid type III functional response. At high prey
39 densities, feeding rate is limited by digestion rather than by encounter rate and the grazer may
40 therefore reduce its foraging effort. The presence of predators may induce a further reduction in
41 foraging effort, particularly at high prey densities. Reduced feeding efforts at high prey densities,
42 whether due to predation risk or energy saving, will not necessarily change the type of the
43 functional response (i.e., a type II may remain a type II, Abrams 1990), but predator-induced
44 changes in behaviors may cause behavioral cascading effects that are very important in shaping
45 food webs and organism abundances, as has been demonstrated in both terrestrial (Suraci et al.
46 2016) and freshwater environments (Peacor and Werner 2001; Romare and Hansson 2003; Biro et
47 al. 2005).

48 Zooplankton, the main consumers of oceanic primary production, may also modify their foraging
49 effort in response to prey concentration and presence of predators with important implications to
50 their function in pelagic food webs. The significance of behavioral adaptations to fundamental

properties of pelagic food webs, such as primary production and material fluxes (Anderson et al. 2010), diversity of the phytoplankton prey (Prowse et al. 2012), and the seasonal successions in plankton communities (Visser 2007; Mariani et al. 2013; Visser and Fiksen 2013) has been demonstrated in multiple modelling studies. Yet, actual demonstrations of the behavioral adaptations to prey and predators assumed in these models are rare, particularly in marine zooplankton (Price and Paffenhöfer 1986; Saiz et al. 1993; Saiz 1994; Duren and Videler 1995; Van Duren and Videler 1996). Early zooplankton-specific models of optimal foraging (Lam and Frost 1976; Lehman 1976) were consistent with more general models (Abrams 1982, 1990; Werner and Anholt 1993) in typically predicting reduced feeding effort at low prey concentration, but were based on fundamentally wrong assumptions about the feeding behavior and considered, similar to the more recent model of Pahlow and Prowse (2010), only the energetic costs of feeding, not predation. Finally, experimental studies of functional responses in zooplankton are abundant but are inconsistent in their findings, reporting both type II and type III responses (Online Appendix 1), and without the mechanistic underpinning that is important in assessing the food web effects predicted by models.

Here, we test the predictions of a simple fitness optimization model through direct observations of the behavioral basis of the functional response in pelagic copepods, the dominating group of metazoan zooplankton in the oceans. We consider two contrasting foraging modes: ambush feeding, in which the copepod waits for prey to arrive, and active feeding, in which the copepod generates a feeding current or swims to encounter prey. The active foraging modes are more efficient in terms of prey encounter than the passive mode (Kjørboe 2011). We demonstrate that ambush feeders consistently have invariant foraging behavior and type II responses, while active feeders may modify their foraging effort in response to prey concentration and predation risk and have a type III response, but that the predation response is “hard-wired” and not plastic.

75

76 ***Material and methods***77 **Fitness optimization model**

78 We take a Holling type II functional response as a starting point:

79
$$F = F_{max} \frac{\beta R}{\beta R + F_{max}} \quad (1)$$

80 where F is the ingestion rate and F_{max} the maximum ingestion rate (biomass per time), R is the prey
 81 concentration (biomass per volume), and β is the ‘instantaneous rate of prey discovery’ or
 82 maximum clearance rate (volume per time). Assume that the zooplankter feeds only a fraction of
 83 the time, p (non-dimensional foraging effort), and that p vary in response to food availability and
 84 predation risk in order to optimize the zooplankter’s fitness. The foraging effort, p , modulates the
 85 effective clearance rate ($p\beta$) and the resulting functional response becomes (Werner and Anholt
 86 1993):

87
$$F(p) = F_{max} \frac{p\beta R}{p\beta R + F_{max}} \quad (2)$$

88 Note that if feeding rate is ultimately limited by handling of the prey during the capture process
 89 (with $1/F_{max}$ being the “handling time”), as is typically assumed (Abrams 1982; Abrams 1990;
 90 Pahlow and Prowe 2010) the resulting functional response then becomes proportional to foraging
 91 effort:

92
$$F(p) = F_{max} p \frac{\beta R}{\beta R + F_{max}} \quad (3)$$

93 However, in suspension feeding zooplankton and many other organisms, the handling of prey is
 94 rarely, if ever, the limiting factor (Tiselius et al. 2013). Rather, ingestion is limited by the capacity
 95 of the gut to process food, and the appropriate equation in our case is therefore (2) and not (3).

We now examine what foraging effort (p) optimizes the fitness of the zooplankter when considering the energetic cost as well as the predation risk of feeding. To estimate energetic (metabolic) expenses and mortality risk, we assume

$$\text{Metabolism } M(p) = m_0 + pm_f \quad (4)$$

$$\text{Mortality } \mu(p) = \mu_0 + p\mu_f \quad (5)$$

where m_0 and μ_0 are background metabolism (mass per time) and mortality (per time), and m_f and μ_f are metabolic costs and mortality risk of feeding, respectively. There is both theoretical and experimental evidence that mortality risk increases with foraging activity in zooplankton (Tiselius et al. 1997, Kiørboe et al. 2014; Almeda et al. 2016). The interpretation of p as the fraction of time spent feeding makes it natural to assume a linear dependence of foraging metabolism and predation mortality risk on p .

The optimal behavior is the one that optimizes life-time reproductive output. To avoid a full life-time calculation, two different fitness proxies are frequently used: either to optimize the difference between birth ($\propto$ net energy gain, $F(p) - M(p)$) and mortality rates (e.g. Abrams 1982, 1990), or the ratio between the two (behavioral optimization; e.g. Werner and Anholt 1993; Visser 2007; Gillam's rule: Gilliam and Fraser 1987). If the environment is constant, it has been demonstrated that Gillam's rule optimizes life-time reproductive output (Sainmont et al. 2015). We therefore use Gillam's rule as a fitness proxy. However, a special case occurs when the resource levels are insufficient to ensure a positive net energy gain, where $F - M < 0$. In that case survival is limited and life-time reproductive output would be less than zero. We argue that under such adverse prey conditions the organism will adjust its behavior to maximize the time it can survive by minimizing energy losses. Thus, under low resource concentration the fitness proxy is to minimize the net energy loss. The two fitness proxies are now:

$$119 \quad \frac{F(p)-M(p)}{\mu(p)} \quad \text{for } F - M \geq 0$$

$$120 \quad (6)$$

$$121 \quad F(p) - M(p) \text{ for } F - M < 0 \quad (7)$$

122 The optimal value of p is the one that maximizes the relevant fitness criterion. For the specific
 123 choices of functional response (2) and linear costs of metabolism (4) and mortality (5), the optimal
 124 foraging effort is:

$$p^* = \frac{1}{\rho} \frac{f_c(m - \mu) - \sqrt{\rho\mu - f_c(\rho - \mu)(\mu - m)}}{f_c(\mu - m) - \mu} \quad \text{for } F - M \geq 0, \quad (8)$$

125 where $\rho = R\beta/F_{\max}$ is the scaled resource concentration, $f_c = m_0/F_{\max}$ is scaled standard
 126 metabolism and $m = m_f/m_0$ and $\mu = \mu_f/\mu_0$ are scaled costs of feeding. This expression shows
 127 that foraging effort generally declines with resource concentration (the $1/\rho$ term). Feeding is,
 128 however, limited by the condition that p^* should be ≤ 1 (Fig 1A).

129 The critical resource concentration R_c , where gains equal losses even while feeding all the time
 130 ($p = 1$), is found by equating gains from eq. (2) with metabolic losses from eq. (4), $F(1) = M(1)$:

$$131 \quad R_c = \frac{F_{\max}(m_0 + m_f)}{F_{\max} - m_0 - m_f} \frac{1}{\beta}. \quad (9)$$

132 Below this concentration the functional response is approximately linear, $F \approx p\beta R$ and, hence, the
 133 optimization problem (7) is linear. The organism will then feed continuously ($p = 1$) as long as the
 134 net energy loss is less the loss while not feeding, i.e., $F(1) - M(1) > -M(0) = -m_0$:

$$135 \quad R_0 \approx \frac{m_f}{\beta}. \quad (10)$$

136 Note that this result is general, i.e., it does not rely on the cost of feeding effort being linear in p
 137 (eq. 4) since m_f is the feeding cost of feeding at the maximal rate. Using these fitness measures and

depending on the magnitude of the tradeoffs, the model predicts that feeding effort (p) is highest at intermediate prey concentrations, and declines at both lower and higher concentrations (Fig. 1A), which results in a type III functional response (Fig 1C). It also predicts that the feeding effort is further reduced in the presence of predators (compare black and grey lines). Specifically, for an ambush feeder, since $\mu_f = m_f = 0$ we would not expect a feeding threshold but predict that $p = 1$ and independent prey concentration and, hence, a type II functional response (dashed lines in Fig. 1).

Note the difference between the realized clearance rate ($\neq \beta$), which is $F(p)/R$, and the foraging effort, p (Fig. 1B). Both may vary with resource concentration: At low resource concentrations, they show similar dependencies on the resource (the clearance rate $= F(p)/R \approx p$); at high resource concentration, the clearance rate varies with $1/R$, independent of the variation in p . This follows directly from the optimization and equation (2).

Parameter estimates: Based on observations in Kiørboe et al. (1985) we provide rough estimates of F_{max} ($0.65 \mu\text{g C } (\mu\text{g dry body weight})^{-1}\text{d}^{-1}$), β ($1.65 \text{ mL } (\mu\text{g dry body weight})^{-1}\text{d}^{-1}$), m_0 ($0.015 \mu\text{g C } (\mu\text{g dry body weight})^{-1}\text{d}^{-1}$), and m_f ($0.1 \mu\text{g C } (\mu\text{g dry body weight})^{-1}\text{d}^{-1}$) for one of the study species, *Acartia tonsa*, feeding on one of the prey, *Rhodomonas baltica*. The two metabolic rate estimates were taken as starvation metabolism (m_0) and maximum metabolism (m_f) of feeding individuals, respectively. The latter includes also the cost of processing and metabolizing the food and we added also losses to defecation, which are not strictly ‘costs of feeding’. We have no estimates of mortality rates for the study species but assume $\mu_0 = \mu_f = 0.05 \text{ d}^{-1}$. This implies a mortality rate of $\sim 0.1 \text{ d}^{-1}$ for a small, feeding copepod in the ocean, a magnitude typical for mm-sized feeding-current feeding copepods (Hirst and Kiørboe 2002).

The predictions of this model of optimal foraging, as illustrated in Fig. 1, are the hypotheses that we test experimentally here.

162

163 **Experimental organisms**

164 We quantified feeding behavior and feeding rate as a function of prey concentration in the adult
165 females of 4 species of copepods: an ambush feeder, *Oithona davisae* (cephalothorax length 0.3
166 mm), a copepod that can switch between ambush and active feeding, *Acartia tonsa* (0.8 mm), and
167 two species that are obligate active feeders, *Temora longicornis* (0.8 mm) and *Centropages*
168 *hamatus* (1.0 mm). *A. tonsa* generates a feeding current when offered small prey, and acts as an
169 ambush feeder when offered large, motile prey (Jonsson and Tiselius 1990). All copepods were
170 taken from our continuous cultures; the two first species were from our long-term cultures (> 10
171 years), the latter two had been in culture for < ½ year. We used three different types of prey, the
172 flagellate *Rhodomonas baltica* (6.5 µm equivalent spherical diameter) and the dinoflagellates
173 *Oxyrrhis marina* (16.5 µm) and *Akashiwo sanguinea* (42 µm), all in exponential growth. We did not
174 do all possible predator-prey combinations, and some data were taken from our earlier work (Table
175 1).

176

177 **Measures of foraging effort**

178 The active feeders beat their cephalic appendages to generate a feeding current and/or to propel
179 themselves through the water during shorter or longer ‘feeding bouts’ (Tiselius and Jonsson 1990)
180 and we used the fraction of time that the organism beats the appendages as the main measure of
181 foraging effort, but note that some minimum beat-activity may be necessary to keep the copepod
182 suspended, irrespective of feeding. Additional, but secondary measures of foraging effort are
183 appendage beat frequency and swimming speed. The interpretation of swimming speed is not
184 straightforward, because with a given beat frequency (~ force), a hovering copepod feeds more
185 efficiently than one that cruises through the water, but it also produces a stronger fluid signal and

becomes more detectable to rheotactic predators (Kjørboe and Jiang 2013). The ambush feeding *O. davisae* only moves the cephalic appendages in connection with prey capture, but relocates occasionally in short, rapid jumps using the swimming legs (Kjørboe et al. 2009); in this species there is no foraging effort and, hence, no measure of foraging effort, but we recorded jump frequency as a measure of activity.

Feeding behavior

Adult females were isolated from the cultures and starved overnight. 50 individuals were then added to each observation aquaria (250 mL NUNC bottles) containing fresh food suspension and acclimated for 2 h before filming commenced. We used seven different food concentrations for each prey, including no food (Table 1), selected to yield similar prey biovolume ranges for each prey ($\sim 0\text{-}5\text{ mm}^3\text{ L}^{-1}$). Prey concentration was adjusted just prior to filming. We used a high-speed (200 fps), high resolution (1280 x 800 pixels) Phantom v210 camera equipped with optics to yield fields of view ranging between ca. 20x32 and 52x78 mm depending on the size of the copepods. Collimated infrared light was shined through the aquarium toward the camera. Three 27 s sequences were filmed during a 15 min period for each treatment. The water in the aquaria was then replaced with water containing fish smell (see below) and the appropriate prey concentration, and after 5 min the animals were filmed again during the subsequent ca. 15 min.

Swimming trajectories were extracted from the movies using the tracking software LabTrack (DiMedia). The movies were analyzed both at 20 Hz to construct time budgets (fraction of time feeding) and estimate event durations (feeding bouts) and speeds (swimming speed), and at 33 or 67 Hz to estimate jump frequencies. Between 20 and 150 tracks were analyzed per treatment; in a few cases we retrieved only 10 tracks. The output from the tracking analyses were run through an R-script that distinguished between jump, sinking, and swimming events based on species-specific

thresholds that were selected to match what a manual frame-by-frame analysis would yield, but the patterns observed were very robust to choice of thresholds. We finally also estimated appendage beat-frequencies in 1-s long sequences by frame-by-frame analyses of 20 random, active animals per treatment.

Exposure to predation risk

Predation risk was mimicked by the addition of fish smell to the observational aquaria. Fish smell was produced by a mixture of small, coastal planktivorous fish: juveniles of corkwing (*Symphosus melops*), sea stickleback (*Spinacia spinachia*), and black goby (*Gobius niger*) that had all been feeding on copepods. 13 g wet weight of this coastal fish assemblage were incubated in 2-L of filtered seawater for > 2 h. This water was then filtered through a GF/C filter and diluted to 50 % by adding an algal suspension of twice the nominal concentration, thus having a fish smell solution with the nominal phytoplankton concentration. Effect of fish smell was only tested with *R. baltica* as prey.

Functional response

All functional responses were measured in our laboratories, either for the purpose of this study or earlier (Table 1), and followed the same protocol: Adult females were incubated in 325-650 mL bottles at 5-6 different prey concentrations with 3 bottles with copepods and 3 controls at each concentration. The bottles were mounted on a slowly rotating wheel for ca. 24 h at ~17°C. *O. davisae* were incubated at 21°C and we had only two 72-mL bottles with copepods and two control at each concentration. We added enough copepods to get a reduction in phytoplankton concentration of ~25 % during the incubation period. Cell concentrations were measured both at start and termination of the experiments, and clearance and ingestion rates were computed as in Kiørboe et al. (1982). Cell concentrations were either measured electronically on a Coulter Counter

and in most cases also in the microscope. At low prey concentrations electronic counts were always verified by microscopic counts.

We fitted functional response functions to the observed ingestion rates, either the disc equation (eq. 1) or a descriptive sigmoid equation (Kjørboe et al 1982):

$$F = \alpha \beta e^{1-\alpha/R} \quad , \quad (11)$$

where α is the prey concentration where the clearance rate is the highest and equals β , and the maximum ingestion rate $F_{\max} = \alpha \beta e^1$. We also fitted the same functional response functions expressed as clearance rates (F/R) to the estimated clearance rates. We decided on the most appropriate description of the functional response (type II or III) based on (i) whether or not observed clearance rates decline at low prey concentrations, and (ii) which formulation yielded the best description (highest R^2).

Literature survey

Expanding on the data base of Kjørboe and Hirst (2014) we compiled from the literature experimental observations of functional responses in marine suspension feeding copepods offered a variety of prey sizes (Online Appendix 1). We used the measured clearance rates to decide on the shape of the functional response type, cf. above. We also computed maximum clearance rates (β) and maximum ingestion rate ($F_{\max}$) for each data set, either by using the functional response fits in the original paper or by our own fits to the data after correcting for differences in temperature, all as described in Kjørboe and (Hirst 2014). In some cases only maximum clearance rates were reported. We finally computed the feeding and maintenance thresholds, R_0 (only species with active foraging behavior) and R_c for each dataset using equation 9 and 10 and the computed maximum clearance and ingestion rates and assuming the above default values for the mass-specific metabolism (m_0) and mass-specific metabolic cost of feeding (m_f). We find the latter assumptions justified by the fact

that both metabolism and clearance scale approximately in proportion with body mass in zooplankton (Kjørboe and Hirst 2014), and that the metabolic cost of feeding is likely to be proportional to the clearance rate.

Results

Feeding behavior and functional response

The behavioral responses to prey concentration varied significantly between copepod species and was also dependent on the prey type (Figs. 2-4, Table 2). The most diverse behavioral repertoire is shown by *A. tonsa*. This copepod beats its feeding appendages to generate a feeding current during feeding bouts each lasting between ca. 0.2-0.8 s on average (Fig. 2D); between feeding bouts it is inactive and sinks slowly. When offered the small flagellate (*R. baltica*) its foraging effort varied with prey concentration pretty much as predicted by the optimization model (with predation): the highest feeding effort (~80 %) is at an intermediate prey concentration and it declines at both higher and lower concentrations (Fig. 2A, 2B). There is no distinct feeding threshold and even in filtered seawater the animals are active for ~25 % of the time, probably simply to remain suspended. This is accomplished by frequent, but short-lasting feeding bouts, whereas the longer feeding bouts at intermediate concentration mainly serve the purpose of feeding (Fig. 2D). Swimming speed shows an inverse pattern with the slowest speeds at intermediate concentrations (Fig. 2E), while appendage beat frequency was independent of prey concentration and only reduced in the absence of food (Fig. 3A). The patterns in these two secondary foraging effort measures thus also suggest the most efficient and potentially most risky feeding behavior at intermediate concentrations and declining at both higher and lower concentrations. There is no significant effect on the behavior of adding fish smell (Fig. 2A, D, E).

281 The measured functional response in clearance rate of *A. tonsa* to the concentration of *R. baltica*
282 showed peak clearance at an intermediate prey concentration and lower clearance at both lower and
283 higher resource concentrations (Fig. 2C). This translates directly to a type III sigmoid functional
284 response in ingestion rate to prey concentration (Fig. 2F), and the observed functional response is
285 thus consistent with the observed variation in foraging effort.

286 When *A. tonsa* is offered large motile prey (*Ox. marina* and *Ak. sanguinea*) it changes its behavior
287 and functional response compared to when offered the small *R. baltica* prey in a way that is
288 consistent with a change in foraging mode from feeding-current feeding to ambush feeding (Fig. 4).
289 This is most evident with the largest prey (*Ak. sanguinea*): The feeding bouts are short and the
290 foraging effort remains low, independent of prey concentration, and not different from a situation
291 with no prey (Fig. 4B); the ‘feeding bouts’ thus mainly serve to keep the animal suspended while
292 waiting for prey to pass within its sensory reach. With the intermediately sized prey (*Ox. marina*),
293 the behavioral changes with prey concentration are intermediate between that observed with the
294 smaller and the larger prey (Fig. 4D,E), suggesting partial active and partial passive feeding. The
295 functional response in clearance and ingestion rates when offered the largest prey (*Ak. sanguinea*)
296 is of type II (Fig. 4 C, F), and thus consistent with the observed behavior.

297 None of the 3 other copepod species showed consistent behavioral changes with prey concentration,
298 prey type, or presence of fish cues except that appendage beat frequencies were slightly reduced at
299 the lowest prey concentrations (Fig. 3) and that *C. hamatus* jumps more frequently when offered the
300 large prey as compared to the small prey (Table 2). In a few cases small differences in the other
301 parameters are statistically significant, but the patterns are inconsistent (Table 2; all data are plotted
302 in Online Appendix 2). Consistent with the concentration-independent behavior, all predator-prey
303 combinations for the 3 species showed a type II functional response (Online Appendix 2). This was
304 unexpected, since type II responses were *a priori* only predicted for the ambush feeding *O. davisae*.

The behavior and foraging effort of the 4 studied species are, however, very different (Table 2). *T. longicornis* is actively beating its feeding appendages almost constantly while cruising or hovering, whereas *C. hamatus* has a feeding behavior that resembles that of *A. tonsa*, i.e., alternating between short upwards-directed feeding bouts (~ about 50 % of the time) and sinking with occasionally longer swimming events, although in this species the pattern is invariant with prey concentration. Finally, the ambush feeding *O. davisae* never beats the feeding appendages but just performs relocation jumps every 2-3 s (Table 2).

Comparison between modelled and observed responses

For actively foraging *A. tonsa* the modelled pattern in foraging effort describes both qualitatively and quantitatively very well the observations, but only when predator presence is assumed (Fig. 2C). This resemblance to observations is robust to at least a ± 50 % variation in the magnitude of the default parameters. However, in the absence of a foraging induced predation risk the predicted foraging effort remains 100% at concentrations exceeding the feeding threshold. A decline in foraging effort with increasing prey concentration and in the absence of predators is only predicted to occur at very high prey concentrations, $> 6 \text{ mm}^3 \text{L}^{-1}$, far beyond what is examined here and typically found in the ocean, and a pattern in foraging effort that resembles the observed pattern is never achieved with any combination of parameters (Fig. 2C).

For passively feeding *A. tonsa* (i.e. when fed *Ak. sanguinea*) and *O. davisae* the prey concentration-invariant foraging behavior and type II responses are consistent with model predictions (Fig. 4 and Table 2), while the invariant foraging behavior of actively feeding *T. longicornis* and *C. hamatus* (Table 2) deviated from the expected (i.e. adaptive foraging effort and type III response).

Feeding and maintenance thresholds

The predicted concentration threshold, R_c , where the net energy gain is zero, is a minimum measure of the lowest prey concentration at which the copepod is able to survive in the long run, whether or not it stops feeding at the lower concentration R_0 . For ambush feeders there is no feeding threshold ($R_0 = 0$), and for active feeders the two thresholds are related and rather similar (using default parameters, $R_c = 1.4 \times R_0$). Both thresholds were computed to be able to utilize a larger number of observations. The feeding and maintenance thresholds vary over several orders of magnitude and scatter around a phytoplankton concentration of $\sim 1 \text{ mm}^3 \text{ L}^{-1}$, corresponding to $\sim 100 \mu\text{g C L}^{-1}$, and not very different between active and passive feeders (Fig. 5A, B). The thresholds are dependent on the size of the prey relative to the size of the copepod: with small relative prey sizes the feeding and maintenance threshold are high, and *vice versa* (Fig. 5 C, D)

Discussion

Mechanistic underpinning of the functional response

Overall, there is consistency between the observed feeding behaviors and the measured functional responses for all predator-prey combinations examined here, and the former thus provides a mechanistic underpinning of the latter. We found both type II and type III functional responses among the copepods studied, partly in agreement with the predictions from the optimization model. Type III responses may arise in several ways, including through prey switching in mixed diet environments (Murdoch and Oaten 1975; Elliott 2004; Leeuwen et al. 2007), but here we show that the type III is due to a change in foraging effort with prey concentration. We are unaware of previous reports that provide a direct behavioral underpinning of the observed functional response in copepod or other zooplankton although there are a few studies that have examined how relevant components of feeding behavior vary with prey density. For example, the copepod *Eucalanus*

elongates spends the highest fraction of time feeding at intermediate prey concentrations, (Price and Paffenhöfer 1986) and a previous study of *A. tonsa* showed that, similarly to clearance rates, the time spent in long feeding bouts peaked at intermediate concentrations of a small diatom (Saiz 1994), consistently with our finding. There are also reports that the copepod *Temora longicornis* modifies its swimming speed (Duren and Videler 1995; Van Duren and Videler 1996; Moison et al. 2013) or appendage beat frequency (Gill and Poulet 1988) in response to concentration of food or presence of dissolved amino acids or predators. We are unaware of studies in other zooplankton groups. The scarcity of such evidence hampers a complete understanding of the role of zooplankton in pelagic food webs.

Functional response

Both type II and III functional responses have previously been reported for pelagic copepods, and a survey of ~ 120 functional response experiments reported in the literature (Online Appendix 1) reveals that ambush feeding copepods consistently show type II responses (43 experiments) as predicted, while active cruising or feeding-current feeding species either showed a type II (45 cases) or a type III response (30 cases). The latter result is consistent with the finding here of variable responses in the active feeders. For at least 21 out of the 45 reported type II responses in active feeders, prey concentrations lower than the predicted feeding threshold were not tested and consequently those experiments are inconclusive with respect to the actual type of functional response (see Online Appendix 1). Moreover, clearance measurements at low prey concentrations, where a type III response is best distinguished from a type response II, can be challenging, and some of the reports may not provide very strong evidence for the type of response. For this reason, in our experiments we intentionally examined behaviors also at very low prey concentrations, including the absence of prey, and still found high foraging effort at low and no prey in two species. We must therefore conclude that the observed differences among species are real and that some

374 active feeders, contrary to our expectations, have a high resource-independent level of foraging
375 activity and a type II functional response.

376 What causes the deviation from prediction in some species? Swimming in active feeders is
377 accomplished by the beating of the feeding appendages and the need to swim to areas with more
378 food or fewer predators may override other effects, a tradeoff that has not been considered here and
379 that may vary between species. Kinetic motility responses to prey concentration has been reported
380 in copepods (Tiselius 1992) and other zooplankters (Buskey and Stoecker 1988; Fenchel and
381 Jonsson 1988; Menden-Deuer and Grünbaum 2006), thus potentially explaining why several active
382 feeders appear to have no lower feeding threshold and type II functional responses.

383 **Induced responses and phenotypic plasticity**

384 None of the three examined species showed a response to the presence of a predator cue, and the
385 reduced foraging effort at high prey concentrations demonstrated in one species (*A. tonsa*) occurred
386 both in the absence and presence of predator cues. This suggests limited behavioral plasticity and
387 that any adaptation to predation risk is wired into the genes of *A. tonsa* rather than being triggered
388 in response to the actual presence of predators. One may argue that a copepods susceptibility to
389 visual predators (fish) is less dependent on the feeding activity than its susceptibility to rheotactic
390 predators, but it is well documented that feeding copepods are much more prone to visual predators
391 than non-feeding individuals due to the elevated visual contrast that a full gut implies in an
392 otherwise near transparent copepod (Tsuda et al. 1998; Torgersen 2003). Thus, a fish cue seems to
393 be relevant.

394 The lack of a response to predator cues is surprising in light of the commonly documented effect of
395 predator kairomones in freshwater zooplankton, including induction of reduced feeding efforts in
396 freshwater copepods (see reviews by Lass and Spaak 2003; Heuschele and Selander 2014). Our

397 protocol to produce chemical fish cues appear not to deviate from what is typically used in
398 freshwater studies. One may argue that the overnight starvation of the copepods and the only 2 h
399 acclimation to prey concentration prior to behavioral observations may lead to a more bold behavior
400 of the copepod, where the need for food dominates over predator avoidance behavior. Indeed,
401 starvation may result in significantly increased feeding in *Acartia* spp, but the effect is reduced or
402 has disappeared within 100 min of feeding (Tiselius 1998), consistent with gut turnover times at
403 the experimental temperature of just 20-25 min (Kiørboe and Tiselius 1987; Dam and Peterson
404 1988) allowing the animals to fill their guts 5-6 times during the acclimation period. Also, predator
405 avoidance behavior is in fact observed in *A. tonsa* at the high prey concentrations. The lack of
406 response to cues in our experiments is, however, consistent with the almost entirely lack of reports
407 on behavioral effects of kairomones in marine zooplankton and copepods. Thus, Buskey et al.
408 (2012) in a review failed to find evidence of predator-induced responses for marine zooplankton,
409 and only three studies were identified in the review by Heuschele and Selander (2014) in addition to
410 Bjærke et al. (2014), of which only two report effects on feeding-related behavioral changes
411 (reduced swimming speed or reduced gut fullness with predator cues (Van Duren and Videler 1996;
412 Cieri and Stearns 1999). There is also one report that diurnal vertical behavior can be induced by
413 the presence of fish, but the cue that elicited the response was not identified, except that it was not
414 of chemical nature (Bollens and Frost 1989). The literature may be biased towards negative results
415 not being reported, and therefore the scarcity of evidence may thus reflect rarity of responses at
416 ecological time scales. Freshwater systems, in particular smaller lakes and ponds, may vary with
417 respect to the presence of fish predators, while marine systems are all large and interconnected and,
418 hence, always contain planktivorous fish. Thus, adaptations to predator avoidance is commonly
419 found among marine zooplankton, including vertical migration (Ringelberg 2010; Ohman and
420 Romagnan 2016) and reduced feeding during daytime to reduce susceptibility to visual predators

(review by Torgersen 2003), but they are typically elicited by light intensity (Stearns 1986; Buskey et al. 1989) or some other proximate cue, not by the actual presence of predators. The apparent lack of phenotypic plasticity may simply reflect the constant need for a behavior that reduces predation risk.

Ecological implications

The functional response in feeding rate to prey concentration provides the fundamental description of predator-prey interactions and thus is fundamental to the understanding of population dynamics and food web structure in several ways. First, a type III functional response may stabilize prey populations due to the density-dependent prey mortality that it implies, while a type II response may drive the prey population to (local) extinction (Holling 1965). The often significant impact of the choice of functional response type in models of both simple pelagic food chains and more complex food webs has been recognized by many authors (Anderson et al. 2010) and “inappropriate choices may incorrectly quantify biologically mediated fluxes and predict spurious dynamics” (Gentleman et al. 2003). This realization warrants the search for a fundamental understanding of the mechanisms that are generating one functional response or another to allow the ‘correct’ choice. The attempt in this study to find such ‘rules’ for a very important group of phytoplankton consumers has been partly successful and has demonstrated the utility of optimal foraging theory in this endeavor. However, it has also pointed to gaps in our understanding and identified possible additional mechanisms, particularly kinetic motility responses, which may lead to more robust predictions of the functional response in copepods and other zooplankton.

Secondly, the behavior that generates the functional response may have implications beyond prey mortality and grazer growth rates. Specifically, behavioral adaptations to the presence of predators (or their cues) may lead to a behavioral cascade and a ‘landscape of fear’ (Suraci et al. 2016), where grazing and growth rates are determined as much by the presence of predators as by the availability

of food, as demonstrated in freshwater zooplankton (Gliwicz and Maszczyk 2007) and many other organisms. The scarcity of predator-induced behavioral responses in copepods and other marine zooplankton suggests that ‘behavioral cascades’ are less important in controlling marine planktonic systems than what has been demonstrated for higher trophic levels in freshwater systems (Romare and Hansson 2003; Biro et al. 2005) and assumed in models (e.g., Visser 2007).

Finally, the dependency of the maintenance resource concentration on the prey:predator size ratio, which is a function of the copepod prey size spectra (Kjørboe 2016), suggests that environmental food conditions may put predictable constraints on the size distribution and biogeography of copepods. Thus, the smaller the relative size of the prey, the higher the required prey concentration, and therefore large copepods are constrained to regions with high concentration of large (phytoplankton) prey. This prediction accords well with observed body-size biogeographies of copepods, where the larger species occur in polar and temperate regions characterized by seasonal high concentrations of large diatoms; and smaller species dominate in tropical and subtropical regions, characterized by lower biomasses of small phytoplankton (Brun et al. 2016). Because maximum clearance rates and metabolic rates scale approximately in proportion to body mass when considered over the entire range of pelagic organisms in the ocean (Makarieva et al. 2008; Kjørboe and Hirst 2014), and assuming that the cost of clearing water for food is proportional to the volume cleared, this prediction may be generalized to zooplankton in general, not just copepods.

464 **Acknowledgements**

465 The Centre for Ocean Life is supported by the Villum Foundation. ES was funded by grant
466 CGL2014-59227-R (MINECO/FEDER, UE), and P.T. by a sabbatical grant from University of
467 Gothenburg.

468

For Review Only

References

- Abrams, P. A. 1982. Functional response of optimal foragers. *Am. Nat.* **120**: 382–390.
doi:10.2307/2678832
- Abrams, P. A. 1990. The effects of adaptive behavior on the type-2 functional response. *Ecology* **71**: 877–885.
- Almeda, R., H. van Someren Gréve, and T. Kiørboe. 2016. Behavior is a major determinant of predation risks in zooplankton. *Ecosphere* **8**. doi:10.1002/ecs2.1668
- Anderson, T. R., W. C. Gentleman, and B. Sinha. 2010. Influence of grazing formulations on the emergent properties of a complex ecosystem model in a global ocean general circulation model. *Prog. Oceanogr.* **87**: 201–213. doi:10.1016/j.pocean.2010.06.003
- Biro, P. a, J. R. Post, and M. V Abrahams. 2005. Ontogeny of energy allocation reveals selective pressure promoting risk-taking behaviour in young fish cohorts. *Proc. Biol. Sci.* **272**: 1443–1448. doi:10.1098/rspb.2005.3096
- Bjærke, O., T. Andersen, and J. Titelman. 2014. Predator chemical cues increase growth and alter development in nauplii of a marine copepod. *Mar. Ecol. Prog. Ser.* **510**: 15–24.
doi:10.3354/meps10918
- Bollens, S. M., and B. W. Frost. 1989. Predator-induced diel vertical migration in a planktonic copepod. *J. Plankton Res.* **11**: 1047–1065. doi:10.1093/plankt/11.5.1047
- Brun, P., M. R. Payne, and T. Kiørboe. Trait biogeography of marine copepods – an analysis across scales. *Ecol. Lett.* in review.
- Buskey, E., K. Baker, R. Smith, and E. Swift. 1989. Photosensitivity of the oceanic copepods

- 490 *Pleuromamma gracilis* and *Pleuromamma xiphias* and its relationship to light penetration and
491 daytime depth distribution . Mar. Ecol. Prog. Ser. **55**: 207–216. doi:10.3354/meps055207
- 492 Buskey, E. J., P. H. Lenz, and D. K. Hartline. 2012. Sensory perception, neurobiology, and
493 behavioral adaptations for predator avoidance in planktonic copepods. Adapt. Behav. **20**: 57–
494 66. doi:10.1177/1059712311426801
- 495 Buskey, E. J., and D. K. Stoecker. 1988. Locomotory patterns of the planktonic ciliate *Favella* sp.:
496 Adaptations for remaining within food patches. Bull. Mar. Sci. **43**: 783–796.
- 497 Cieri, M. D., and D. E. Stearns. 1999. Reduction of grazing activity of two estuarine copepods in
498 response to the exudate of a visual predator. Mar. Ecol. Prog. Ser. **177**: 157–163.
499 doi:10.3354/meps177157
- 500 Dam, H. G., and W. T. Peterson. 1988. The effect of temperature on the gut clearance rate constant
501 of planktonic copepods. J. Exp. Mar. Bio. Ecol. **123**: 1–14. doi:10.1016/0022-0981(88)90105-
502 0
- 503 Duren, L. A. van, and J. J. Videler. 1995. Swimming behavior of developmental stages of the
504 calanoid copepod *Temora longicornis* at different food concentrations. Mar. Ecol. Prog. Ser.
505 **126**: 153–161.
- 506 Van Duren, L. a, and J. J. Videler. 1996. The trade-off between feeding, mate seeking and predator
507 avoidance in copepods: behaviour responses to chemical cues. J. Plankton Res. **18**: 805–818.
508 doi:10.1093/plankt/18.5.805
- 509 Elliott, J. M. 2004. Prey switching in four species of carnivorous stoneflies. Freshw. Biol. **49**: 709–
510 720. doi:10.1111/j.1365-2427.2004.01222.x

- 511 Fenchel, T., and P. Jonsson. 1988. The functional biology of *Strombidium sulcatum*, a marine
512 oligotrich ciliate (Ciliophora, Oligotrichina). Mar. Ecol. Prog. Ser. **48**: 1–15.
513 doi:10.3354/meps048001
- 514 Gentleman, W., A. Leising, B. Frost, S. Strom, and J. Murray. 2003. Functional responses for
515 zooplankton feeding on multiple resources: A review of assumptions and biological dynamics.
516 Deep. Res. Part II Top. Stud. Oceanogr. **50**: 2847–2875. doi:10.1016/j.dsr2.2003.07.001
- 517 Gill, C., and S. Poulet. 1988. Responses of copepods to dissolved free amino acids. Mar. Ecol.
518 Prog. Ser. **43**: 269–276. doi:10.3354/meps043269
- 519 Gilliam, J. F., and D. F. Fraser. 1987. Habitat selection under predation hazard: test of a model with
520 foraging minnows. Ecology **68**: 1856–1862.
- 521 Gliwicz, Z. M., and P. Maszczyk. 2007. *Daphnia* growth is hindered by chemical information on
522 predation risk at high but not at low food levels. Oecologia **150**: 706–715.
523 doi:10.1007/s00442-006-0528-7
- 524 Gonçalves, R., H. van Someren Gréve, D. Couespel, and T. Kiørboe. 2014. Mechanisms of prey
525 size selection in a suspension-feeding copepod, *Temora longicornis*. Mar. Ecol. Prog. Ser. **517**:
526 61–74. doi:10.3354/meps11039
- 527 Heuschele, J., and E. Selander. 2014. The chemical ecology of copepods. J. Plankton Res. **36**: 895–
528 913. doi:10.1093/plankt/fbu025
- 529 Hirst, a. G., and T. Kiørboe. 2002. Mortality of marine planktonic copepods: Global rates and
530 patterns. Mar. Ecol. Prog. Ser. **230**: 195–209. doi:10.3354/meps230195
- 531 Holling, C. S. 1965. The Functional response of predators to prey density and its role in mimicry

- 532 and population regulation. Mem. Entomol. Soc. Canada **97**: 1–60. doi:10.4039/entm9745fv
- 533 Jonsson, P., and P. Tiselius. 1990. Feeding behaviour, prey detection and capture efficiency of the
534 copepod *Acartia tonsa* feeding on planktonic ciliates. Mar. Ecol. Prog. Ser. **60**: 35–44.
535 doi:10.3354/meps060035
- 536 Kiørboe, T. 2011. How zooplankton feed: Mechanisms, traits and trade-offs. Biol. Rev. **86**: 311–
537 339. doi:10.1111/j.1469-185X.2010.00148.x
- 538 Kiørboe, T. 2016. Foraging mode and prey size spectra in suspension feeding copepods and other
539 zooplankton. Mar. Ecol. Prog. Ser. **558**: in press. doi:10.3354/meps11877
- 540 Kiørboe, T., a. Andersen, V. Langlois, H. H. Jakobsen, and T. Bohr. 2009. Mechanisms and
541 feasibility of prey capture in ambush feeding zooplankton. Proc. Natl. Acad. Sci. **106**: 12394–
542 12399.
- 543 Kiørboe, T., and A. G. Hirst. 2014. Shifts in mass scaling of respiration, feeding, and growth rates
544 across life-form transitions in marine pelagic organisms. Am. Nat. **183**: E118-30.
545 doi:10.1086/675241
- 546 Kiørboe, T., and H. Jiang. 2013. To eat and not be eaten: optimal foraging behaviour in suspension
547 feeding copepods. J. R. Soc. Interface **10**: 20120693. doi:10.1098/rsif.2012.0693
- 548 Kiørboe, T., H. Jiang, R. J. Gonçalves, L. T. Nielsen, and N. Wadhwa. 2014. Flow disturbances
549 generated by feeding and swimming zooplankton. Proc. Natl. Acad. Sci. **111**: 11738–11743.
550 doi:10.1073/pnas.1405260111
- 551 Kiørboe, T., F. Møhlenberg, and K. Hamburger. 1985. Bioenergetics of the planktonic copepod
552 *Acartia tonsa*: relation between feeding, egg production and respiration, and composition of

- 553 specific dynamic action . Mar. Ecol. Prog. Ser. **26**: 85–97. doi:10.3354/meps026085
- 554 Kiørboe, T., and P. Tiselius. 1987. Gut clearance and pigment destruction in a herbivorous copepod,
555 *Acartia tonsa*, and the determination of in situ grazing rates. J. Plankton Res. **9**: 525–534.
- 556 Lam, R. K., and B. W. Frost. 1976. Model of copepod filtering response to changes in size and
557 concentration of food. Limnol. Oceanogr. **21**: 490–500. doi:10.4319/lo.1976.21.4.0490
- 558 Lass, S., and P. Spaak. 2003. Chemically induced anti-predator defences in plankton: A review.
559 Hydrobiologia **491**: 221–239. doi:10.1023/A:1024487804497
- 560 Leeuwen, E. van, V. Jansen, and P. Bright. 2007. How population dynamics shape the functional
561 response. Ecology **88**: 1571–1581.
- 562 Lehman, J. T. 1976. The filter-feeder as an optimal forager, and the predicted shapes of feeding
563 curves. Limnol. Oceanogr. **21**: 501–516. doi:10.4319/lo.1976.21.4.0501
- 564 Makarieva, A. M., V. G. Gorshkov, B.-L. Li, S. L. Chown, P. B. Reich, and V. M. Gavrillov. 2008.
565 Mean mass-specific metabolic rates are strikingly similar across life's major domains:
566 Evidence for life's metabolic optimum. Proc. Natl. Acad. Sci. U. S. A. **105**: 16994–16999.
567 doi:10.1073/pnas.0802148105
- 568 Mariani, P., K. H. Andersen, A. W. Visser, A. D. Barton, and T. Kiørboe. 2013. Control of plankton
569 seasonal succession by adaptive grazing. Limnol. Oceanogr. **58**: 173–184.
570 doi:10.4319/lo.2013.58.1.0173
- 571 Menden-Deuer, S., and D. Grünbaum. 2006. Individual foraging behaviors and population
572 distributions of a planktonic predator aggregating to phytoplankton thin layers. Limnol.
573 Oceanogr. **51**: 109–116. doi:10.4319/lo.2006.51.1.0109

- 574 Moison, M., F. G. Schmitt, and S. Souissi. 2013. Differences in feeding activity between females
575 and males of *Temora longicornis*. Ecol. Res. **28**: 459–467. doi:10.1007/s11284-013-1034-0
- 576 Murdoch, W. W. 1977. Stabilizing effects of spatial heterogeneity in predator-prey systems. Theor.
577 Popul. Biol. **11**: 252–273.
- 578 Murdoch, W. W., and A. Oaten. 1975. Predation and Population Stability. Adv. Ecol. Res. **9**: 1–
579 131.
- 580 Ohman, M. D., and J.-B. Romagnan. 2016. Nonlinear effects of body size and optical attenuation on
581 diel vertical migration by zooplankton. Limnol. Oceanogr. **61**: 765–770.
582 doi:10.1002/lno.10251
- 583 Pahlow, M., and A. E. F. Prowe. 2010. Model of optimal current feeding in zooplankton. Mar. Ecol.
584 Prog. Ser. **403**: 129–144. doi:10.3354/meps08466
- 585 Peacor, S. D., and E. E. Werner. 2001. The contribution of trait-mediated indirect effects to the net
586 effects of a predator. Proc. Natl. Acad. Sci. U. S. A. **98**: 3904–3908.
587 doi:10.1073/pnas.071061998
- 588 Price, H. J., and G.-A. Paffenhöfer. 1986. Effects of concentration on the feeding of a marine
589 copepod in algal monocultures and mixtures. J. Plankton Res. **8**: 119–128.
- 590 Prowe, A. E. F., M. Pahlow, S. Dutkiewicz, M. Follows, and A. Oschlies. 2012. Top-down control
591 of marine phytoplankton diversity in a global ecosystem model. Prog. Oceanogr. **101**: 1–13.
592 doi:10.1016/j.pocean.2011.11.016
- 593 Ringelberg, J. 2010. Diel vertical migration of zooplankton in lakes and oceans: Causal
594 explanations and adaptive significances, Springer, Netherlands.

- 595 Romare, P., and L.-A. Hansson. 2003. A behavioral cascade: Top-predator induced behavioral
596 shifts in planktivorous fish and zooplankton. *Limnol. Oceanogr.* **48**: 1956–1964.
597 doi:10.4319/lo.2003.48.5.1956
- 598 Sainmont, J., K. H. Andersen, U. H. Thygesen, Ø. Fiksen, and A. W. Visser. 2015. An effective
599 algorithm for approximating adaptive behavior in seasonal environments. *Ecol. Modell.* **311**:
600 20–30. doi:10.1016/j.ecolmodel.2015.04.016
- 601 Saiz, E. 1994. Observations of the free-swimming behavior of *Acartia tonsa*: Effects of food
602 concentration and turbulent water motion. *Limnol. Oceanogr.* **39**: 1566–1578.
603 doi:10.4319/lo.1994.39.7.1566
- 604 Saiz, E., A. Calbet, and E. Broglio. 2003. Effects of small-scale turbulence on copepods: The case
605 of *Oithona davisae*. *Limnol. Oceanogr.* **48**: 1304–1311. doi:10.4319/lo.2003.48.3.1304
- 606 Saiz E., Tiselius P., Jonsson P.R., Verity P., and G.-A. Paffenhöfer. 1993. Experimental records of
607 the effects of food patchiness and predation on egg production of *Acartia tonsa*. *Limnol.*
608 *Oceanogr.* **38**: 280–289.
- 609 Stearns, D. E. 1986. Copepod grazing behavior in simulated natural light and its relation to
610 nocturnal feeding. *Mar. Ecol. Prog. Ser.* **30**: 65–76.
- 611 Suraci, J. P., M. Clinchy, L. M. Dill, D. Roberts, and L. Y. Zannette. 2016. Fear of large carnivores
612 causes a trophic cascade. *Nat. Commun.* **7**: 10698. doi:10.1038/ncomms10698
- 613 Tiselius, P. 1992. Behavior of *Acartia tonsa* in patchy food environments. *Limnol. Oceanogr.* **37**:
614 1640–1651.
- 615 Tiselius, P. (1998) Short term feeding responses to starvation in three species of small calanoid

- 616 copepods. Mar. Ecol. Prog. Ser. 168: 119-126
- 617 Tiselius, P., and P. Jonsson. 1990. Foraging behaviour of six calanoid copepods: observations and
618 hydrodynamic analysis. Mar. Ecol. Prog. Ser. **66**: 23–33. doi:10.3354/meps066023
- 619 Tiselius, P., Jonsson, P.R., Kaartvedt, S., Olsen, E.M., Jørstad, T. (1997) Effects of copepod
620 foraging behavior on predation risk: An experimental study of the predatory copepod
621 *Pareuchaeta norvegica* feeding on *Acartia clausi* and *A. tonsa* (Copepoda). Limnol. Oceanogr.
622 42: 164-170
- 623 Tiselius, P., E. Saiz, and T. Kiørboe. 2013. Sensory capabilities and food capture of two small
624 copepods, *Paracalanus parvus* and *Pseudocalanus* sp. Limnol. Oceanogr. **58**: 1657–1666.
625 doi:10.4319/lo.2013.58.5.1657
- 626 Torgersen, T. 2003. Proximate causes for anti-predatory feeding suppression by zooplankton during
627 the day: Reduction of contrast or motion - Ingestion or clearance? J. Plankton Res. **25**: 565–
628 571. doi:10.1093/plankt/25.5.565
- 629 Tsuda, A., H. Saito, and T. Hirose. 1998. Effect of gut content on the vulnerability of copepods to
630 visual predation. Limnol. Oceanogr. **43**: 1944–1947.
- 631 Visser, A., and Ø. Fiksen. 2013. Optimal foraging in marine ecosystem models: selectivity,
632 profitability and switching. Mar. Ecol. Prog. Ser. **473**: 91–101. doi:10.3354/meps10079
- 633 Visser, A. W. 2007. Motility of zooplankton: fitness, foraging and predation. J. Plankton Res. **29**:
634 447–461. doi:10.1093/plankt/fbm029
- 635 Werner, E. E., and B. R. Anholt. 1993. Ecological consequences of the trade-off between growth
636 and mortality rates mediated by foraging activity. Am. Nat. **142**: 242–272.

637

638

For Review Only

FIGURE LEGENDS

Fig 1. Foraging patterns as a function of scaled resource concentration. Three cases are shown: foraging constant with $p = 1$ (dashed); optimal foraging under no predation risk ($\mu_f = 0$, $\mu_0 = 0.05 \text{ d}^{-1}$, grey); optimal foraging under a predation risk ($\mu_f = \mu_0 = 0.05 \text{ d}^{-1}$, black full line). A) Foraging effort, p . B) Realized clearance rate normalized by maximum clearance rate. C) The functional response in ingestion rate normalized by the maximum ingestion rate ($F_{\max}$).

Fig. 2. *Acartia tonsa* feeding on *Rhodomonas baltica* as a function of cell density in the presence and absence of fish cues. A. Foraging effort (% time swimming); D. Feeding bout duration; C. Modelled foraging effort at various values of predator induced mortality risk and metabolic costs of feeding. The black line is for the default parameters estimated from (Kjørboe et al. 1985); E. Swimming speed; C and F. Observed clearance and ingestion rates, from Kjørboe et al. (1985); The curves are fits of a type III functional response model to the data with $\alpha = 0.58 \pm 0.04$, $\beta = 12.4 \pm 0.6$, and $R^2 = 0.83$ (panel B), and $\alpha = 0.561 \pm 0.04$, $\beta = 12.1 \pm 0.6$, and $R^2 = 0.98$ (Panel F) (estimates of coefficients with standard error). Experimental values are given as averages $\pm 95\%$ Confidence limits.

Fig. 3. Beating frequencies (average $\pm 95\%$ confidence limits) of feeding appendages as a function of prey concentration in 3 species of copepods. A. *A. tonsa* feeding on *R. baltica*. B. *C. hamatus* feeding on *R. baltica*; C. *C. hamatus* feeding on *Ak. sanguinea*. D. *T. longicornis* feeding on *R. baltica*. Averages $\pm 95\%$ confidence limits.

Fig. 4. *Acartia tonsa* feeding on and *Akashiwo sanguinea* (A-C, F) and *Oxyrrhis marina* (B, D) as a function of prey density. A and D: Foraging effort (% time swimming); B and E: Duration of individual feeding bouts; E and F: Observed clearance and ingestion rates and (curves) Hollings disk equation fitted to the data, with $F_{\max} = 22.3 \pm 12.8$, $\beta = 23.5 \pm 4.6$, and $R^2 = 0.30$ (panel C), and

662 $F_{\max} = 11.6 \pm 2.4$, $\beta = 46.6 \pm 32.7$, and $R^2 = 0.28$ (panel F). Values of behavioral parameters are given
663 as averages $\pm 95\%$ Confidence limits; clearance and ingestion rates are individual values (grey) and
664 averages (black).

665 **Fig. 5.** Feeding thresholds (R_0) and maintenance threshold (R_c) estimated for pelagic copepods with
666 a ‘active’ (black symbols) and ‘passive’ (grey symbols) feeding strategies. Maximum clearance (β)
667 and ingestion ($F_{\max}$) rates were estimated from literature data on functional responses (see Online
668 Appendix 1), and we assumed body mass specific metabolic cost of feeding of ($m_f = 0.1 \mu\text{g C (mg}$
669 $\text{dry body weight)}^{-1}\text{d}^{-1}$) $\sim 0.01 \mu\text{g C (mg body C)}^{-1}\text{h}^{-1}$) identical to that estimated for *A. tonsa*. A:
670 Frequency distribution of feeding thresholds threshold for active feeders; B: Frequency distribution
671 of maintenance thresholds for active and passive feeders; C: Feeding threshold as a function of the
672 prey:predator carbon-mass ratio for active feeders; the regression line is $\log(R_0) = -1.1 - 0.22 \text{ Log}$
673 $(\text{prey:predator mass ratio})$; $R^2 = 0.35$, $n = 182$; D: Maintenance thresholds as a function of the
674 prey:predator carbon-mass ratio for active and passive feeders; the regression line is $\log(R_0) = -1.0$
675 $- 0.18 \text{ Log}(\text{prey:predator mass ratio})$; $R^2 = 0.19$, $n = 209$.

676
677

678

Species of grazer\prey	<i>Rhodomonas baltica</i>	<i>Oxyrrhis marina</i>	<i>Akashiwo sanguinea</i>
Prey size, Equivalent spherical diameter, μm	6.5	16.5	42
<i>Acartia tonsa</i>	Behavior: this study Functional response: Kjørboe et al. (1985)	Behavior: this study	Behavior: this study Functional response: This study
<i>Temora longicornis</i>	Behavior: this study Functional response: Gonçalves et al. (2014)		
<i>Centropages hamatus</i>	Behavior: this study Functional response: Sommeren-Greve et al. unpublished		Behavior: this study Functional response: Sommeren-Greve et al. unpublished
<i>Oithona davisae</i>		Behavior: this study Functional response: Saiz et al. (2003)	

679

Table 1: Predator-prey combinations examined for feeding behavior and functional responses.

680

681

682

Predator-Prey	# tracks	Total time, s	% Active	Swim speed, mm s ⁻¹	Bout duration, s	Jump frequency, s ⁻¹
<i>Acartia-Oxyrrhis</i>	292	3052	Fig. 3	1.4 ± 0.1	Fig. 3	0.68 ± 0.09
<i>Acartia-Akashiwo</i>			Fig. 3	1.7 ± 0.1	Fig. 3	0.75 ± 0.07
<i>Temora-Rhodomonas</i>	163	1235	96 ± 1.7	1.7 ± 0.1	3.3± 0.5	0.12 ± 0.05
<i>Temora-Rhodomonas</i> +Fish	206	1116	94 ± 1.2	1.9 ± 0.2	2.6 ± 0.3	0.16 ± 0.06
<i>Centropages-Rhodomonas</i>	442	2263	50 ± 2.5	3.0 ± 0.1	0.6 ± 0.1	0.01 ± 0.01
<i>Centropages-Rhodomonas</i> + Fish	482	2235	48 ± 2.5	3.1 ± 0.1	0.6± 0.1	0.00 ± 0.00
<i>Centropages-Akashiwo</i>	809	4654	43 ± 1.3	3.8 ± 0.1	0.3 ± 0.02	0.11± 0.03
<i>Oithona davisae</i>	163	2101	0	-	-	0.39 ± 0.07

683

684 Table 2. Summary statistics of behaviors for copepod-prey (± Fish) combinations where behavior is
685 (near) independent on prey concentration and prey type. Values given are averages ± 95 %
686 Confidence limits. All the data have been plotted in Online Appendix 2.

687

Fig. 1

Fig 2

Fig. 3

Fig. 4

Fig. 5

Online Appendix I

Species	Prey type	Feeding mode	FR	Temp	Body mass	$F_{\max}$	β	prey mass/cop mass	R_0	R_c	Source
				C	μgC	$\mu\text{g C h}^{-1}$	ml h^{-1}	$\mu\text{gC}(\mu\text{gC})^{-1}$	$\text{mm}^3 \text{L}^{-1}$	$\text{mm}^3 \text{L}^{-1}$	
<i>Acartia grani</i> naupl	<i>Heterocapsa</i> sp	Ambush	II	20	0.040	0.00137	0.0094	0.0047		0.143	(Henriksen et al. 2007)
<i>Acartia grani</i> naupl	<i>Thalassiosira weissflogii</i>	Ambush	II	20	0.040	0.00108	0.0072	0.0055		0.185	(Henriksen et al. 2007)
<i>Acartia tonsa</i>	<i>Strombidium sulcatum</i>	Ambush	II	18	3.000	0.10638	5.6141	0.00028		0.018	(Saiz and Kiørboe 1995)
<i>Corycaeus angelicus</i>	<i>Acartia clausi</i> male	Ambush	II	11	2.400		0.2412	0.9375			(Landry et al. 1985)
<i>Oithona davisae</i>	<i>Isochrysis galbana</i>	Ambush	II	18	0.390	0.00106	0.0150	0.00002		0.871	(Saiz et al. 2014)
<i>Oithona davisae</i>	<i>Tetraselmis chuii</i>	Ambush	II	18	0.384	0.00557	0.0130	0.00012		0.987	(Saiz et al. 2014)
<i>Oithona davisae</i>	<i>Heterocapsa</i> sp.	Ambush	II	18	0.368	0.01813	0.0303	0.00047		0.405	(Saiz et al. 2014)
<i>Oithona davisae</i>	<i>Prorocentrum minimum</i>	Ambush	II	18	0.375	0.01417	0.0218	0.00046		0.575	(Saiz et al. 2014)
<i>Oithona davisae</i>	<i>Oxyrrhis marina</i>	Ambush	II	18	0.336	0.01928	0.1416	0.00138		0.079	(Saiz et al. 2014)
<i>Oithona davisae</i>	<i>Scrippsiella trochoidea</i>	Ambush	II	18	0.389		0.0452	0.00192			(Saiz et al. 2014)
<i>Oithona davisae</i>	<i>Prorocentrum micans</i>	Ambush	II	18	0.383		0.0653	0.00338			(Saiz et al. 2014)

<i>Oithona davisae</i>	<i>Strombidium sulcatum</i>	Ambush	II	18	0.351	0.02597	0.3593	0.00522		0.033	(Saiz et al. 2014)
<i>Oithona davisae</i>	<i>Strombidium sulcatum</i>	Ambush	II	18	0.390		0.2940	0.0052			(Saiz et al. 2014)
<i>Oithona davisae</i>	<i>Akashiwo sanguinea</i>	Ambush	II	18	0.393	0.01268	0.0112	0.01097		1.166	(Saiz et al. 2014)
<i>Oithona davisae</i>	<i>Paracartia grani nauplii</i>	Ambush	II	18	0.369	0.00241	0.0310	0.07429		0.397	(Saiz et al. 2014)
<i>Oithona davisae</i>	<i>Thalassiosira weissflogii</i>	Ambush	II	18	0.348		0.0278	0.00029			(Saiz et al. 2014)
<i>Oithona davisae</i>	<i>Oxhyrris marina</i>	Ambush	II	21	0.250	0.00494	0.1078	0.00106		0.077	(Saiz et al. 2003)
<i>Oithona davisae</i> male	<i>Oxhyrris marina</i>	Ambush	II	22	0.188	0.00253	0.0477	0.00142		0.131	(Kiørboe 2008)
<i>Oithona davisae</i> naupl	<i>Heterocapsa sp</i>	Ambush	II	20	0.032	0.00060	0.0047	0.00585		0.229	(Henriksen et al. 2007)
<i>Oithona davisae</i> naupl	<i>Isochrysis galbana</i>	Ambush	II	18	0.051	0.00020	0.0016	0.00017		1.042	(Saiz et al. 2014)
<i>Oithona davisae</i> naupl	<i>Tetraselmis chuii</i>	Ambush	II	18	0.056	0.00090	0.0053	0.00097		0.351	(Saiz et al. 2014)
<i>Oithona davisae</i> naupl	<i>Heterocapsa sp.</i>	Ambush	II	18	0.055	0.00137	0.0071	0.00349		0.256	(Saiz et al. 2014)
<i>Oithona davisae</i> naupl	<i>Oxyrrhis marina</i>	Ambush	II	18	0.059	0.00221	0.0477	0.00615		0.041	(Saiz et al. 2014)
<i>Oithona davisae</i> naupl	<i>Thalassiosira weissflogii</i>	Ambush	II	18	0.058		0.0045	0.00173			(Saiz et al. 2014)
<i>Oithona davisae</i> , CII-III	<i>Oxyrrhis marina</i>	Ambush	II	20.5	0.233	0.00325	0.0263	0.00114		0.296	(Almeda et al. 2010)
<i>Oithona davisae</i>	<i>Oxyrrhis marina</i>	Ambush	II	18	0.250	0.00600	0.2233	0.00106		0.037	(Zamora-Terol and Saiz 2013)

<i>Oithona davisae</i> , NI-II	<i>Oxyrrhis marina</i>	Ambush	II	20.5	0.089	0.00074	0.0090	0.00298		0.332	(Almeda et al. 2010)
<i>Oithona davisae</i> , NI-II	<i>Oxyrrhis marina</i>	Ambush	II	20.5	0.092	0.00088	0.0092	0.00288		0.334	(Almeda et al. 2010)
<i>Oithona davisae</i> , NII-III	<i>Oxyrrhis marina</i>	Ambush	II	20.5	0.101	0.00099	0.0109	0.00263		0.310	(Almeda et al. 2010)
<i>Oithona davisae</i> , NV-VI	<i>Oxyrrhis marina</i>	Ambush	II	20.5	0.147	0.00186	0.0135	0.00181		0.364	(Almeda et al. 2010)
<i>Oithona nana</i>	<i>Acartia nauplii</i>	Ambush	II	10	0.230	0.00192	0.6226	0.12609		0.012	(Lampitt 1978)
<i>Oithona nana</i>	<i>Isochysis galbana</i>	Ambush	II	10	0.230	0.00064	0.0119	0.00001		0.650	(Lampitt and Gamble 1982)
<i>Oithona nana</i> male	<i>Acartia nauplii</i>	Ambush	II	10	0.210	0.00673	0.7197	0.13810		0.010	(Lampitt 1978)
<i>Oithona nana</i> male	<i>Dunaliella euchlora</i>	Ambush	II	10	0.230	0.00293	0.0112	0.00012		0.688	(Lampitt and Gamble 1982)
<i>Oithona nana</i> male	<i>Chricosphaera elongata</i>	Ambush	II	10	0.230	0.00359	0.0467	0.00048		0.164	(Lampitt and Gamble 1982)
<i>Oithona nana</i> male	<i>Thalassiosira weissflogii</i>	Ambush	II	10	0.230	0.00146	0.0077	0.00018		1.002	(Lampitt and Gamble 1982)
<i>Oithona nana</i> male	<i>Prorocentrum micans</i>	Ambush	II	10	0.230	0.00128	0.0202	0.00282		0.380	(Lampitt and Gamble 1982)
<i>Oithona nana</i> male	<i>Acartia clausi</i> NI	Ambush	II	10	0.230	0.00603	0.7195	0.12609		0.011	(Lampitt and

											Gamble 1982)
<i>Oithona nana</i> male	<i>Calanus finmarchicus</i> NI	Ambush	II	10	0.230	0.00530	0.2106	0.86957		0.036	(Lampitt and Gamble 1982)
<i>Oithona nana</i> male	<i>Calanus finmarchicus</i> NII	Ambush	II	10	0.230	0.00006	0.0906	1.73913		0.090	(Lampitt and Gamble 1982)
<i>Oithona similis</i>	<i>Prorocentrum micans</i>	Ambush	II	8.5	0.360	0.00931	0.1369	0.00321		0.088	Drits & Semenova 1984
<i>Oithona similis</i>	<i>Peridinium trochoideum</i>	Ambush	II	8.5	0.360	0.01051	0.1665	0.00193		0.072	Drits & Semenova 1984
<i>Oithona similis</i>	<i>Platymonas viridis</i>	Ambush	II	8.5	0.360	0.01250	0.1543	0.00009		0.078	Drits & Semenova 1984
<i>Acartia clausi</i>	<i>Rhodomonas baltica</i>	Active	(II)	15	5.000	0.23000	0.3335	0.00001	1.499251	1.728	(Dutz 1998)
<i>Acartia clausi</i>	<i>Alexandrium lusitanicum</i>	Active	(II)	15	5.000	0.27000	0.6244	0.00020	0.800818	0.923	(Dutz 1998)
<i>Acartia Erythraea</i>	<i>Chattonella antiqua</i>	Active	II	20	4.580	0.14462	0.7889	0.00033	0.580591	0.670	Uye 1986
<i>Acartia hudsonica</i>	<i>Thalassiosira constricta</i>	Active	III	4.5	6.790	0.38937	2.7416	0.00003	0.247669	0.285	(Durbin and Durbin 1992)
<i>Acartia hudsonica</i>	<i>Thalassiosira constricta</i>	Active	III	8	5.940	0.18246	1.7887	0.00003	0.332091	0.383	(Durbin and Durbin 1992)
<i>Acartia hudsonica</i>	<i>Thalassiosira constricta</i>	Active	III	12	4.420	0.18159	1.3619	0.00003	0.324545	0.374	(Durbin and Durbin 1992)

<i>Acartia hudsonica</i>	<i>Thalassiosira constricta</i>	Active	III	16	3.880	0.15750	0.8751	0.00004	0.44338	0.511	(Durbin and Durbin 1992)
<i>Acartia tonsa</i>	<i>Thalassiosira weisflogii</i>	Active	III	20	3.710		3.3586	0.00003	0.110463		(Durbinl and Durbinl 1990)
<i>Acartia tonsa</i>	<i>Thalassiosira weisflogii</i>	Active	(II)	18	3.000	0.09545	5.8007	0.00009	0.051718	0.060	(Saiz and Kjørboe 1995)
<i>Acartia tonsa</i>	<i>Isochrysis galbana</i>	Active	III	18	2.484	0.40385	0.1247	0.00001	1.99254	2.293	(Støttrup and Jensen 1990)
<i>Acartia tonsa</i>	<i>Dunaliella tertiolecta</i>	Active	III	18	2.484	0.14685	0.4004	0.00001	0.620434	0.715	(Støttrup and Jensen 1990)
<i>Acartia tonsa</i>	<i>Rhodomonas baltica</i>	Active	III	18	2.484	0.19825	0.3750	0.00001	0.662394	0.763	(Støttrup and Jensen 1990)
<i>Acartia tonsa</i>	<i>Thalassiosira weifsflogii</i>	Active	III	18	2.484	0.17622	1.0415	0.00007	0.238507	0.275	(Støttrup and Jensen 1990)
<i>Acartia tonsa</i>	<i>Ditylum brightwellii</i>	Active	III	18	2.484	0.14685	0.7210	0.00033	0.344499	0.397	(Støttrup and Jensen 1990)
<i>Acartia tonsa</i> , copepodites	<i>Pavlova lutheri</i>	Active		17	0.898		0.0619	0.00001	1.451902		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Pavlova lutheri</i>	Active		17	1.239		0.1341	0.00001	0.923956		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Pavlova lutheri</i>	Active		17	0.555		0.1827	0.00002	0.303728		(Berggreen et al. 1988)

<i>Acartia tonsa</i> , copepodites	<i>Pavlova lutheri</i>	Active		17	0.448		0.2132	0.00002	0.210038		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Pavlova lutheri</i>	Active		17	0.898		0.3391	0.00001	0.26493		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Pavlova lutheri</i>	Active		17	1.174		0.3221	0.00001	0.364585		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Pavlova lutheri</i>	Active		17	3.248		0.5679	0.00000	0.571992		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Isochrysis galbana</i>	Active		17	0.555		0.0378	0.00005	1.466256		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Isochrysis galbana</i>	Active		17	0.448		0.0620	0.00006	0.722605		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Isochrysis galbana</i>	Active		17	1.535		0.0731	0.00002	2.100661		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Isochrysis galbana</i>	Active		17	1.174		0.1015	0.00002	1.156698		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Isochrysis galbana</i>	Active		17	1.239		0.1335	0.00002	0.927744		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Isochrysis galbana</i>	Active		17	0.852		0.1756	0.00003	0.48483		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Isochrysis galbana</i>	Active		17	1.901		0.1335	0.00001	1.423884		(Berggreen et al. 1988)

<i>Acartia tonsa</i> , copepodites	<i>Isochrysis galbana</i>	Active		17	3.615		0.3210	0.00001	1.126139		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Dunaliella tertiolecta</i>	Active		17	0.852		0.0748	0.00004	1.137956		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Rhodomonas baltica</i>	Active		17	0.420		0.0316	0.00011	1.331749		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Rhodomonas baltica</i>	Active		17	0.551		0.0339	0.00009	1.624873		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Rhodomonas baltica</i>	Active		17	0.850		0.0696	0.00006	1.220547		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Rhodomonas baltica</i>	Active		17	1.056		0.0864	0.00004	1.221581		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Rhodomonas baltica</i>	Active		17	1.114		0.1331	0.00004	0.837436		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Rhodomonas baltica</i>	Active		17	1.628		0.4208	0.00003	0.386952		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Rhodomonas baltica</i>	Active		17	3.477		0.1907	0.00001	1.823137		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Rhodomonas baltica</i>	Active		17	2.022		0.0561	0.00002	3.60392		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Amphidinium carterae</i>	Active		17	0.425		0.0561	0.00025	0.757912		(Berggreen et al. 1988)

<i>Acartia tonsa</i> , copepodites	<i>Amphidinium carterae</i>	Active		17	0.554		0.0477	0.00019	1.161111		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Amphidinium carterae</i>	Active		17	1.163		0.1016	0.00009	1.144427		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Amphidinium carterae</i>	Active		17	1.515		0.1132	0.00007	1.338614		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Amphidinium carterae</i>	Active		17	0.803		0.1482	0.00013	0.541509		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Amphidinium carterae</i>	Active		17	1.103		0.1565	0.00010	0.704854		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Amphidinium carterae</i>	Active		17	1.776		0.2990	0.00006	0.593945		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Amphidinium carterae</i>	Active		17	3.352		0.7090	0.00003	0.472724		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Thalassiosira weissflogii</i>	Active		17	0.439		0.1289	0.00061	0.340893		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Thalassiosira weissflogii</i>	Active		17	0.611		0.3083	0.00044	0.198051		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Thalassiosira weissflogii</i>	Active		17	0.947		0.5316	0.00028	0.17806		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Thalassiosira weissflogii</i>	Active		17	1.245		0.7786	0.00021	0.159932		(Berggreen et al. 1988)

<i>Acartia tonsa</i> , copepodites	<i>Thalassiosira weissflogii</i>	Active		17	1.245		1.0225	0.00021	0.121784		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Thalassiosira weissflogii</i>	Active		17	1.638		0.9169	0.00016	0.178639		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Thalassiosira weissflogii</i>	Active		17	2.040		0.8222	0.00013	0.248057		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Thalassiosira weissflogii</i>	Active		17	3.529		2.7270	0.00008	0.129401		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Scripsiella faroense</i>	Active		17	0.555		0.0399	0.00144	1.390723		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Scripsiella faroense</i>	Active		17	0.425		0.0949	0.00189	0.447189		(Berggreen et al. 1988)
<i>Acartia tonsa</i> , copepodites	<i>Scripsiella faroense</i>	Active		17	0.852		0.2385	0.00094	0.357122		(Berggreen et al. 1988)
<i>Aetideus divergens</i>	<i>Thalassiosira fluviatilis</i>	Active	II	12	21.2		0.4903	0.00001	4.332161		(Robertson and Frost 1977)
<i>Aetideus divergens</i>	<i>Coscinodiscus angatii</i>	Active	II	12	21.2		4.6032	0.00012	0.461414		(Robertson and Frost 1977)
<i>Aetideus divergens</i>	<i>Coscinodiscus angatii</i>	Active	(II)	12	21.2		9.4789	0.00055	0.224077		(Robertson and Frost 1977)
<i>Aetideus divergens</i>	<i>Artemia nauplii</i>	Active	(II)	12	21.2		16.4053	0.03578	0.12947		(Robertson and Frost 1977)

<i>Calanus finmarchicus</i> CV + female	<i>Emiliana huxley</i>	Active		13	104.3		0.2156	0.00000	48.39151		(Nejstgaard et al. 1995)
<i>Calanus finmarchicus</i> CV + female	<i>Emiliana huxley</i>	Active		13	104.3		0.5658	0.00000	18.43486		(Nejstgaard et al. 1995)
<i>Calanus finmarchicus</i> CV + female	<i>Emiliana huxley</i>	Active		13	104.3		1.4280	0.00000	7.304379		(Nejstgaard et al. 1995)
<i>Calanus finmarchicus</i> CV + female	<i>Prymnesium patelliferum</i>	Active		13	104.3		0.6467	0.00000	16.1305		(Nejstgaard et al. 1995)
<i>Calanus finmarchicus</i> CV + female	<i>Thallasiosira nordenskioldii</i>	Active		13	104.3		6.4127	0.00000	1.626605		(Nejstgaard et al. 1995)
<i>Calanus finmarchicus</i> CV + female	<i>Thallasiosira nordenskioldii</i>	Active		13	104.3		9.0533	0.00000	1.152179		(Nejstgaard et al. 1995)
<i>Calanus finmarchicus</i> CV + female	<i>Chaetoceros calcitrans</i>	Active		13	104.3		0.8622	0.00000	12.09788		(Nejstgaard et al. 1995)
<i>Calanus finmarchicus</i> CV + female	<i>Pavlova lutheri</i>	Active		13	104.3		0.4311	0.00000	24.19576		(Nejstgaard et al. 1995)
<i>Calanus finmarchicus</i> CV + female	<i>Rhodomonas baltica</i>	Active		13	104.3		3.1525	0.00000	3.308821		(Nejstgaard et al. 1995)
<i>Calanus finmarchicus</i> CV + female	<i>Rhodomonas baltica</i>	Active	II	13	104.3	1.27473	3.4418	0.00000	3.030697	3.518	(Nejstgaard et al. 1995)
<i>Calanus finmarchicus</i> CV + female	<i>Emiliana huxley</i>	Active	III	13	104.3	0.92661	1.4826	0.00000	7.035695	8.197	(Nejstgaard et al. 1995)

<i>Calanus pacificus</i> C I	<i>Chlamydomonas</i> sp	Active	III	15	1.506	0.04001	0.1480	0.00012	1.01725	1.175	(Fernandez 1979)
<i>Calanus pacificus</i> C I	<i>Thallasiosira weisflogii</i>	Active	III	15	1.506	0.08810	0.3612	0.00006	0.416952	0.480	(Fernandez 1979)
<i>Calanus pacificus</i> C I	<i>Lauderia borealis</i>	Active	(II)	15	1.506	0.06943	1.2289	0.00032	0.122549	0.141	(Fernandez 1979)
<i>Calanus pacificus</i> C I	<i>Gymnodinium splendens</i>	Active	III	15	1.506	0.07824	0.5320	0.00150	0.28306	0.326	(Fernandez 1979)
<i>Calanus pacificus</i>	<i>Coscinodiscus angstii</i>	Active	III	12.5	76.5	1.59109	11.7792	0.00001	0.649451	0.751	(Frost 1972)
<i>Calanus pacificus</i>	<i>Coscinodiscus eccentricus</i>	Active	III	12.5	76.5	1.31944	17.0348	0.00002	0.44908	0.520	(Frost 1972)
<i>Calanus pacificus</i>	<i>Centric diatom</i>	Active	II/III	12.5	76.5	1.50054	22.4979	0.00004	0.340031	0.393	(Frost 1972)
<i>Calanus pacificus</i> N V	<i>Isochrysis galbana</i>	Active	III	15	0.823	0.00460	0.0322	0.00002	2.558218	3.004	(Fernandez 1979)
<i>Calanus pacificus</i> N V	<i>Chlamydomonas</i> sp	Active	III	15	0.823		0.0645	0.00023	1.27531		(Fernandez 1979)
<i>Calanus pacificus</i> N V	<i>Thallasiosira weisflogii</i>	Active	III	15	0.823	0.03795	0.3985	0.00011	0.206537	0.238	(Fernandez 1979)
<i>Calanus pacificus</i> N V	<i>Peridinium trochoideum</i>	Active	(II)	15	0.823	0.04748	0.1329	0.00061	0.619123	0.713	(Fernandez 1979)
<i>Calanus pacificus</i> N V	<i>Lauderia borealis</i>	Active	III	15	0.823	0.04690	0.7083	0.00059	0.116201	0.134	(Fernandez 1979)
<i>Calanus pacificus</i> N V	<i>Gymnodinium splendens</i>	Active	III	15	0.823	0.04364	0.4582	0.00274	0.179601	0.207	(Fernandez 1979)
<i>Calanus pacificus</i> N V	<i>Gonyaulax polyedra</i>	Active	(II)	15	0.823	0.03708	0.3670	0.00373	0.224225	0.259	(Fernandez 1979)
<i>Calanus pacificus</i> N VI	<i>Isochrysis galbana</i>	Active	II	15	1.168	0.01823	0.0091	0.00001	12.80731	14.838	(Fernandez 1979)
<i>Calanus pacificus</i> N VI	<i>Chlamydomonas</i> sp	Active	(II)	15	1.168	0.04971	0.0795	0.00016	1.468064	1.693	(Fernandez 1979)

<i>Calanus pacificus</i> N VI	<i>Thallasiosira weisflogii</i>	Active	III	15	1.168	0.05757	0.3685	0.00008	0.316892	0.365	(Fernandez 1979)
<i>Calanus pacificus</i> N VI	<i>Lauderia borealis</i>	Active	III	15	1.168	0.07000	0.7490	0.00041	0.155879	0.180	(Fernandez 1979)
<i>Calanus pacificus</i> N VI	<i>Gymnodinium splendens</i>	Active	III	15	1.168	0.06206	0.6641	0.00193	0.175825	0.203	(Fernandez 1979)
<i>Calanus pacificus</i> NIII	<i>Thallasiosira weisflogii</i>	Active	III	15	0.384	0.02096	0.0629	0.00025	0.610736	0.704	(Fernandez 1979)
<i>Calanus pacificus</i> NIII	<i>Gymnodinium splendens</i>	Active	(II)	15	0.384	0.02478	0.0644	0.00587	0.596034	0.687	(Fernandez 1979)
<i>Calanus pacificus</i> NIII	<i>Gonyaulax polyedra</i>	Active	(II)	15	0.384	0.02096	0.0629	0.00799	0.610736	0.704	(Fernandez 1979)
<i>Calanus pacificus</i> NIV	<i>Chlamydomonas</i> sp	Active	III	15	0.479	0.01494	0.0777	0.00039	0.616672	0.712	(Fernandez 1979)
<i>Calanus pacificus</i> NIV	<i>Thallasiosira weisflogii</i>	Active	III	15	0.479	0.02374	0.1377	0.00020	0.347915	0.401	(Fernandez 1979)
<i>Calanus pacificus</i> NIV	<i>Peridinium trochoideum</i>	Active	(II)	15	0.479	0.01998	0.2218	0.00105	0.215946	0.249	(Fernandez 1979)
<i>Calanus pacificus</i> NIV	<i>Lauderia borealis</i>	Active	III	15	0.479	0.05403	0.2539	0.00101	0.188644	0.217	(Fernandez 1979)
<i>Calanus pacificus</i> NIV	<i>Gymnodinium splendens</i>	Active	III	15	0.479	0.03342	0.1704	0.00471	0.281062	0.324	(Fernandez 1979)
<i>Calanus sinicus</i>	<i>Alexandrium tamarense</i> ARC101	Active	II	18	30.2	0.68654	2.5333	0.00002	1.190136	1.376	(Liu and Wang 2002)
<i>Calanus sinicus</i>	<i>Alexandrium tamarense</i> CCMP1771	Active	II	18	30.2	0.50077	1.2966	0.00004	2.325241	2.693	(Liu and Wang 2002)
<i>Calanus sinicus</i>	<i>Thallasiosira weissflogii</i>	Active		18	30.2		1.2850	0.00000	2.346371		(Liu and Wang 2002)
<i>Calanus sinicus</i>	<i>Chattonella antiqua</i>	Active	II	20	51.8	0.75228	3.4303	0.00003	1.510069	1.750	(Uye 1986)

<i>Centropages yamadaii</i>	<i>Chattonella antiqua</i>	Active	II	20	9.6	0.22960	1.6255	0.00016	0.590584	0.682	(Uye 1986)
<i>Euchaete elongata</i>	<i>Pseudocalanus</i> sp	Active	II	8	637	8.53215	117.8054	0.00890	0.540722	0.627	(Yen 1985)
<i>Euchaete elongata</i>	<i>Acartia clausii</i>	Active	II	8	637	6.80516	72.9859	0.00396	0.872771	1.015	(Yen 1985)
<i>Euchaete norvegica</i>	Larval cod	Active	II	7.5	1350	13.00897	224.0314	0.01667	0.602594	0.701	(Yen 1985)
<i>Paracalanus crassirostris</i>	<i>Alexandrium tamarense</i> ARC101	Active	II	18	4.19	0.08371	0.4338	0.00014	0.964658	1.116	(Liu and Wang 2002)
<i>Paracalanus crassirostris</i>	<i>Alexandrium tamarense</i> CCMP1771	Active	II	18	4.19	0.08077	0.4124	0.00027	1.014834	1.174	(Liu and Wang 2002)
<i>Paracalanus crassirostris</i>	<i>Thallasiosira weissflogii</i>	Active		18	4.19		0.4773	0.00001	0.876858		(Liu and Wang 2002)
<i>Paracalanus parvus</i>	<i>Chattonella antiqua</i>	Active	II	20	2.70	0.07225	0.5001	0.00056	0.53991	0.624	(Uye 1986)
<i>Pseudocalanus marinus</i>	<i>Chattonella antiqua</i>	Active	II	20	4.62	0.10231	0.6747	0.00032	0.684742	0.792	(Uye 1986)
<i>Tortanus dextrilobatus</i>	<i>Oithona davisae</i>	Active	II	14	24.4	0.15518	6.9186	0.00983	0.352963	0.413	(Hooff and Bollens 2004)
<i>Tortanus dextrilobatus</i>	<i>Oithona davisae</i>	Active	II	19	18.7	0.11979	2.9947	0.01070	0.6241	0.731	(Hooff and Bollens 2004)
<i>Tortanus dextrilobatus</i>	<i>Acartia</i> sp	Active	II	14	24.4	1.94713	22.2983	0.12244	0.109515	0.126	(Hooff and Bollens 2004)
<i>Tortanus dextrilobatus</i>	<i>Acartia</i> sp	Active	II	19	18.7	0.97222	16.7483	0.12734	0.111593	0.129	(Hooff and Bollens 2004)

<i>Tortanus discaudatus</i>	<i>Calanus pacificus</i> NIII	Active	II	12.5	18.1	0.66554	56.5936	0.01626	0.031949	0.037	(Ambler and Frost 1974)
<i>Tortanus discaudatus</i>	<i>Calanus pacificus</i> NV	Active	II	12.5	18.1	0.92772	86.5515	0.03794	0.02089	0.024	(Ambler and Frost 1974)
<i>Tortanus forcipatus</i>	<i>Pseudodiaptomus nauplii</i>	Active	II	21	6.28	0.04241	3.1810	0.01273	0.197555	0.231	(Uye and Kayano 1994a)
<i>Tortanus forcipatus</i>	<i>Oithona davisae</i>	Active	II	21	6.28	0.08626	4.3132	0.03819	0.145697	0.169	(Uye and Kayano 1994a)
<i>Tortanus forcipatus</i>	<i>Artemia nauplii</i>	Active	II	21	6.16	0.34308	4.5468	0.13473	0.135488	0.156	(Uye and Kayano 1994a)
<i>Tortanus</i> spp CI-III	<i>Oithona davisae</i> CV-VI	Active	(II)	25	1.10	0.01637	0.9673	0.20000	0.113723	0.132	(Uye and Kayano 1994b)
<i>Tortanus</i> spp CI-III	<i>Oithona davisae</i> CV-VI	Active	(II)	20	1.10	0.02739	0.8715	0.20000	0.126216	0.146	(Uye and Kayano 1994b)
<i>Tortanus</i> spp CVIF	<i>Oithona davisae</i> CV-VI	Active	(II)	25	5.62	0.06515	3.2574	0.03915	0.172528	0.200	(Uye and Kayano 1994b)
<i>Tortanus</i> spp CVIF	<i>Oithona davisae</i> CV-VI	Active	(II)	20	5.62	0.07669	3.4861	0.03915	0.161212	0.187	(Uye and Kayano 1994b)
<i>Tortanus</i> spp IV-V	<i>Oithona davisae</i> CV-VI	Active	(II)	25	2.23	0.02848	1.5536	0.09865	0.14354	0.167	(Uye and Kayano 1994b)
<i>Tortanus</i> spp IV-V	<i>Oithona davisae</i> CV-VI	Active	(II)	20	2.23	0.03944	1.9721	0.09865	0.113076	0.131	(Uye and Kayano 1994b)

<i>Tortanus</i> spp male	<i>Oithona davisae</i> CV-VI	Active	(II)	25	3.40	0.04616	1.4688	0.06471	0.231489	0.268	(Uye and Kayano 1994b)
<i>Tortanus</i> spp male	<i>Oithona davisae</i> CV-VI	Active	(II)	20	3.40	0.05807	1.8476	0.06471	0.18402	0.213	(Uye and Kayano 1994b)

Appendix Table 1. Summary of functional response experiments reported in the literature, mainly compiled by Kiørboe and Hirst (2014). The copepods are adult females, unless otherwise noted (N = nauplii, C = copepodites). Feeding mode differentiates between ‘passive’ ambush feeders and ‘active’ cruise and feeding-current feeders. Maximum ingestion rates (F_{max}) and maximum clearance rates (β) were estimated from fitted functional response curves to observational data and temperature corrected to 15 °C as described in Kiørboe and Hirst (2014). FR refers to functional response type II or III evaluated from the observational data, preferentially plots of clearance rate versus prey concentration. Functional response (II) reported in parentheses are experiments where the lowest concentration used was larger than the threshold concentration and, thus, inconclusive with respect to functional response type. R_0 and R_c are the threshold concentrations for feeding and maintenance, respectively, computed from the data using equations 9 and 10, as explained in the main text.

References

Almeda, R., C. B. Augustin, M. Alcaraz, A. Calbet, and E. Saiz. 2010. Feeding rates and gross growth efficiencies of larval developmental stages of *Oithona davisae* (Copepoda, Cyclopoida). J. Exp. Mar. Bio. Ecol. **387**: 24–35. doi:10.1016/j.jembe.2010.03.002

Ambler, J. W., and B. W. Frost. 1974. The feeding behavior of a predatory planktonic copepod, *Tortanus discaudatus*. Limnol. Oceanogr. **19**: 446–451.

Berggreen, U., B. Hansen, and T. Kiørboe. 1988. Food size spectra, ingestion and growth of the copepod *Acartia tonsa* during development: Implications for determination of copepod production. Mar. Biol. **99**: 341–352.

- Durbin, E. G., and A. G. Durbin. 1992. Effects of temperature and food abundance on grazing and short-term weight change in the marine copepod *Acartia hudsonica*. *Limnol. Oceanogr.* **37**: 361–378. doi:10.4319/lo.1992.37.2.0361
- Durbinl, A. G., and E. G. Durbinl. 1990. Diel feeding behavior in the marine copepod *Acartia tonsa* in relation to food availability. *Mar. Ecol. Prog. Ser.* **68**: 23–45. doi:10.3354/meps068023
- Dutz, J. 1998. Repression of fecundity in the neritic copepod *Acartia clausi* exposed to the toxic dinoflagellate *Alexandrium lusitanicum*: Relationship between feeding and egg production. *Mar. Ecol. Prog. Ser.* **175**: 97–107. doi:10.3354/meps175097
- Fernandez, F. 1979. Nutrition Studies in the Nauplius Larva of *Calanus pacificus* (Copepoda : Calanoida). *Mar. Biol.* **53**: 131–147. doi:10.1007/BF00389185
- Frost, B. W. 1972. Effects of size and concentration of food particles on the feeding behavior of the marine planktonic copepod *Calanus pacificus*. *Limnol. Oceanogr.* **17**: 805–815.
- Henriksen, C., E. Saiz, A. Calbet, and B. Hansen. 2007. Feeding activity and swimming patterns of *Acartia grani* and *Oithona davisae* nauplii in the presence of motile and non-motile prey. *Mar. Ecol. Prog. Ser.* **331**: 119–129. doi:10.3354/meps331119
- Hooff, R. C., and S. M. Bollens. 2004. Functional response and potential predatory impact of *Tortanus dextrilobatus*, a carnivorous copepod recently introduced to the San Francisco Estuary. *Mar. Ecol. Prog. Ser.* **277**: 167–179. doi:10.3354/meps277167
- Kjørboe, T. 2008. A mechanistic approach to plankton ecology, Princeton University Press.
- Kjørboe, T., and A. G. Hirst. 2014. Shifts in mass scaling of respiration, feeding, and growth rates across life-form transitions in marine pelagic organisms. *Am. Nat.* **183**: E118–30. doi:10.1086/675241
- Lampitt, R. S. 1978. Carnivorous feeding by a small marine copepod. *Limnol. Oceanogr.* **23**: 1228–1231. doi:10.4319/lo.1978.23.6.1228
- Lampitt, R. S., and J. C. Gamble. 1982. Diet and respiration of the small planktonic marine copepod *Oithona nana*. *Mar. Biol.* **66**: 185–190. doi:10.1007/BF00397192
- Landry, M. R., J. M. Lehner-Fournier, and V. L. Fagerness. 1985. Predatory feeding behavior of the marine cyclopoid copepod *Corycaeus anglicus*. *Mar. Biol.* **85**: 163–169. doi:10.1007/BF00397435

Liu, S., and W. X. Wang. 2002. Feeding and reproductive responses of marine copepods in South China Sea to toxic and nontoxic phytoplankton. *Mar. Biol.* **140**: 595–603. doi:10.1007/s00227-001-0714-4

Nejstgaard, J., U. Båmstedt, E. Bagøien, and T. Solberg. 1995. Algal constraints on copepod grazing. Growth state, toxicity, cell size, and season as regulating factors. *ICES J. Mar. Sci.* **52**: 347–357.

Robertson, S. B., and B. W. Frost. 1977. Feeding by an omnivorous planktonic copepod *Aetideus divergens* Bradford. *J. Exp. Mar. Bio. Ecol.* **29**: 231–244.

Saiz, E., A. Calbet, and E. Broglio. 2003. Effects of small-scale turbulence on copepods: The case of *Oithona davisae*. *Limnol. Oceanogr.* **48**: 1304–1311. doi:10.4319/lo.2003.48.3.1304

Saiz, E., K. Griffell, A. Calbet, and S. Isari. 2014. Feeding rates and prey : predator size ratios of the nauplii and adult females of the marine cyclopoid copepod *Oithona davisae*. *Limnol. Oceanogr.* **59**: 2077–2088. doi:10.4319/lo.2014.59.6.2077

Saiz, E., and T. Kiørboe. 1995. Predatory and suspension feeding of the copepod *Acartia tonsa* in turbulent environments. *Mar. Ecol. Prog. Ser.* **122**: 147–158. doi:10.3354/meps122147

Støttrup, J. G., and J. Jensen. 1990. Influence of algal diet on feeding and egg-production of the calanoid copepod *Acartia tonsa* Dana. *J. Exp. Mar. Bio. Ecol.* **141**: 87–105. doi:10.1016/0022-0981(90)90216-Y

Uye, S. 1986. Impact of copepod grazing on the red-tide flagellate *Chattonella antiqua*. *Mar. Biol. Int. J. Life Ocean. Coast. Waters* **92**: 35–43. doi:10.1007/BF00392743

Uye, S., and Y. Kayano. 1994a. Predatory feeding behavior of *Tortanus forcipatus* on three different prey. *Bull. Plankt. Soc. Japan* **40**: 173–176.

Uye, S., and Y. Kayano. 1994b. Predatory feeding behavior of *Tortanus* (Copepoda: Calanoida): Life-stage differences and the predation impact on small planktonic crustaceans. *J. Crustac. Biol.* **14**: 473–483. doi:10.1017/CBO9781107415324.004

Yen, J. 1985. Selective predation by the carnivorous marine copepod *Euchaeta elongata* : Laboratory measurements of predation rates verified by field observations of temporal and spatial feeding patterns. *Limnol. Oceanogr.* **30**: 577–597. doi:10.4319/lo.1985.30.3.0577

Zamora-Terol, S., and E. Saiz. 2013. Effects of food concentration on egg production and feeding rates of the cyclopoid copepod *Oithona*

davisae. Limnol. Oceanogr. **58**: 376–387. doi:10.4319/lo.2013.58.1.0376

For Review Only

Online Appendix II

Here, all data reported as averages in Table 2 of the main paper are plotted

Appendix Fig. 1. *Centropages hamatus* feeding on *Akashiwo sanguinea*. All values are averages $\pm$ 95 % confidence limits.

Appendix Fig. 2. *Centropages hamatus* feeding on *Rhodomonas baltica*. All values are averages $\pm$ 95 % confidence limits.

Appendix Fig. 3. *Temora longicornis* feeding on *Rhodomonas baltica*. All values are averages $\pm$ 95 % confidence limits.

Appendix Fig. 4. *Acartia tonsa* feeding on *Oxyrrhis marina* (A,B) and *Akashiwo sanguinea* (C, D). All values are averages $\pm$ 95 % confidence limits.

Appendix Figure 5. Functional response in clearance rate and ingestion rate for the copepod-prey combinations examined here. All observations are for adult females. Plots of clearance rates offer the best way to distinguish between a functional response type II and III. Most data are taken from our previous work: *C. hamatus* feeding on *R. baltica* (A,B) and *A. sanguinea* (C,D) (Sommeren Grève, Almeda, Kiørboe *unpublished*); *A. tonsa* feeding on *R. baltica* (E, F) (Kiørboe et al. 1985). *A. tonsa* feeding on *A. sanguinea* is from the present work (G, H); *T. longicornis* feeding on *R. baltica* (I, J) (Gonçalves et al. 2014), *O. davisae* feeding on *O. marina* (K, L) (Saiz et al. 2003). The curves represent fits of type II or type III functional response models to the data, see appendix Table 1 for parameters.

Appendix Table 1. Parameter estimates for curve fits in Appendix figure 5

Copepod	Prey	Clearance	Ingestion
<i>Centropages hamatus</i>	<i>Rhodomonas baltica</i>	$F_{\max} = 8.3 \pm 4.9$ $\beta = 2.5 \pm 0.4$ $R^2 = 0.68$	$F_{\max} = 4.9 \pm 0.65$ $\beta = 3.7 \pm 0.8$ $R^2 = 0.97$
<i>Centropages hamatus</i>	<i>Akashiwo sanguinea</i>	$F_{\max} = 22.2 \pm 3.1$ $\beta = 162 \pm 13$ $R^2 = 0.96$	$F_{\max} = 13.7 \pm 1.0$ $\beta = 290 \pm 79$ $R^2 = 0.67$
<i>Acartia tonsa</i>	<i>Rhodomonas baltica</i>	$\alpha = 0.58 \pm 0.04$ $\beta = 12.4 \pm 0.62$ $R^2 = 0.83$	$\alpha = 0.61 \pm 0.04$ $\beta = 12.1 \pm 0.60$ $R^2 = 0.98$
<i>Acartia tonsa</i>	<i>Akashiwo sanguinea</i>	$F_{\max} = 10.7 \pm 2.7$ $\beta = 9.7 \pm 1.1$ $R^2 = 0.90$	$F_{\max} = 10.1 \pm 1.1$ $\beta = 10.3 \pm 1.9$ $R^2 = 0.97$
<i>Temora longicornis</i>	<i>Rhodomonas baltica</i>	$F_{\max} = 2.6 \pm 0.6$ $\beta = 5.7 \pm 0.4$ $R^2 = 0.91$	$F_{\max} = 2.7 \pm 0.3$ $\beta = 5.5 \pm 0.7$ $R^2 = 0.99$
<i>Oithona davisae</i>	<i>Oxyrrhis marina</i>	$F_{\max} = 19.6 \pm 10.0$ $\beta = 23.4 \pm 3.9$ $R^2 = 0.81$	$F_{\max} = 19.6 \pm 10.0$ $\beta = 23.4 \pm 3.9$ $R^2 = 0.81$