



## Offshore Extreme Wind Atlas Using Wind-Wave Coupled Modeling

Larsén, Xiaoli Guo; Du, Jianting; Bolanos, Rodolfo; Imberger, Marc; Badger, Merete

*Publication date:*  
2018

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*

Larsén, X. G., Du, J., Bolanos, R., Imberger, M., & Badger, M. (2018). *Offshore Extreme Wind Atlas Using Wind-Wave Coupled Modeling*. Abstract from EGU General Assembly: 2018, Vienna, Austria.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Offshore Extreme Wind Atlas Using Wind-Wave Coupled Modeling

Authors: Xiaoli Guo Larsén<sup>1</sup>, Jianting Du<sup>1</sup>, Rodolfo Bolanos<sup>2</sup>, Marc Imberger<sup>1</sup> and Merete Badger<sup>1</sup>

1: Wind Energy Department, Technical University of Denmark; 2: DHI Denmark

Extreme wind is a required design parameter to be estimated for offshore and coastal structures such as wind turbines and platforms and floating objects, to avoid structures obtaining damages from severe wind conditions.

Mid-latitude storms contribute to offshore extreme wind over the North Sea. It is often understood that offshore winds are relatively homogenous over space and the water surface can be described through the roughness length through the Charnock formulation. Though knowledge was gained that the Charnock formulation is not valid during storm conditions or in coastal regions where surface waves are breaking. In the last decades, researchers have been trying to introduce the wave impact to the atmospheric modeling through parameterization of derived wave parameters such as significant wave height, wave length and steepness, however all for open sea conditions.

It is shown in this study that during extreme wind storms over the North Sea, the wind field can be far from spatially homogeneous. About half of the time, open cellular structures are present, corresponding to strong and highly fluctuating (spatially and temporally) wind. It is also shown that for light to moderate winds, the many parameterization schemes in the literature give similar wave effect on the offshore wind, however, the differences between these schemes increase with increasing wind speed and become significant for winds greater than about 20 m/s. A wave boundary layer model (WBLM) was developed here to act as the interface between the atmospheric and wave modeling. WBLM uses one consistent set of equations, namely the momentum and kinetic energy conservations, to link the calculation in the wave model (here Spectral Wave model Nearshore SWAN) and that in the atmospheric model (here the Weather Research and Forecasting model WRF), thus avoiding the inconsistency that often occurs in parameterization schemes of the roughness length.

The modeling system WRF-WBLM-SWAN has been used to model 429 storms from 1994 to 2016 that contributed to the calculation of the extreme wind over Denmark and surrounding waters. The WRF model was setup with consideration of climatological storm path and wind variability, which helped the decision of the domain size and position, simulation length and resolution. The extreme wind was obtained using the selective dynamical downscaling method in Larsén et al. (2013).

Case studies show that WBLM outperforms the parameterization schemes for coastal areas and storm winds. Validation with measurements from several offshore sites in the North Sea suggests that WRF-WBLM-SWAN model system reproduces better offshore extreme winds over Denmark than using WRF alone.

Reference:

Larsén X., Badger J., Hahmann A. and Mortensen N. (2013): The selective dynamical downscaling method for extreme-wind atlases, *Wind Energy*, 16:1167-1182.