The influence of generalist predators in spatially extended predator-prey systems

Chakraborty, Subhendu

Published in:

Ecological Complexity: An International Journal on Biocomplexity in the Environment and Theoretical Ecology

Link to article, DOI:

[10.1016/j.ecocom.2015.06.003](https://doi.org/10.1016/j.ecocom.2015.06.003)

Publication date:

2015

Document Version

Early version, also known as pre-print

[Link back to DTU Orbit](#)

Citation (APA):

Chakraborty, S. (2015). The influence of generalist predators in spatially extended predator-prey systems. *Ecological Complexity: An International Journal on Biocomplexity in the Environment and Theoretical Ecology*, 23, 50-60. <https://doi.org/10.1016/j.ecocom.2015.06.003>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The influence of generalist predators in spatially extended predator-prey systems

Subhendu Chakraborty^{a,b,*}

^a*ICBM, Carl von Ossietzky University Oldenburg, Carl von Ossietzky Str. 9-11, 26111 Oldenburg, Germany*

^b*VKR Centre for Ocean Life, National Institute of Aquatic Resources, Technical University of Denmark, Charlottenlund Slot, Jægersborg Allé, DK-2920 Charlottenlund, Denmark*

Abstract

The presence of generalist predators is known to have important ecological impacts in several fields. They have wide applicability in the field of biological control. However, their role in the spatial distribution of predator and prey populations is still not clear. In this paper, the spatial dynamics of a predator-prey system is investigated by considering two different types of generalist predators. In one case, it is considered that the predator population has an additional food source and can survive in the absence of the prey population. In the other case, the predator population is involved in intraguild predation, i.e., the source of the additional food of the predator coincides with the food source of the prey population and thus both prey and predator populations compete for the same resource. The conditions for linear stability and Turing instability are analyzed for both the cases. In the presence of generalist predators, the system shows different pattern formations and spatiotemporal chaos which has important implications for ecosystem functioning not only in terms of their predictability, but also in influencing species persistence and ecosystem stability in response to abrupt environmental changes. This study establishes the importance of the consideration of spatial dynamics while determining optimal strategies for biological control through generalist predators.

Keywords: Generalist predator, additional food, intraguild predation, Turing instability, pattern formation, biological control

1. Introduction

Predator-prey interactions are determinants of the composition and distribution of species in a community. These interactions mainly depend on the type of predators and their activities. Generalist

*Corresponding author

Email addresses: subc@aquas.dtu.dk (Subhendu Chakraborty), Tel.: +45 3588 3347, Fax: +45 3396 3434 (Subhendu Chakraborty)

4 predators, that utilize a possibly wide variety of food sources, play a crucial role in determining the
5 dynamics of such communities. For example, raccoons (a medium-sized mammal native to North
6 America) are an important part of our ecosystem as they feed on insects, small mammals and birds,
7 eggs, and plant foods. For the last couple of decades, generalist predators have received considerable
8 attention in the context of invasion ecology and pest control, which are important for sustainable and
9 integrated pest-management strategies (Rosenheim et al., 1995; Symondson et al., 2002; Magal et al.,
10 2008; Crowder and Snyder, 2010). Generalist predators affect pest populations in various ways. The
11 ability of generalist predators to ingest new invasive pests can have drastic effects on the local pest
12 populations. For example, the control of the local tomato pest *Bemisia tabaci* populations enhances
13 by the generalist predator *Macrolophus pygmaeus* in the presence of invasive alien pest *Tuta absoluta*
14 (Jaworski et al., 2013). However, predator-prey interactions generally occur over a wide range of spatial
15 and temporal scales and the spatial components of ecological interactions play an important role in
16 shaping ecological communities. In this respect, spatial patterns are ubiquitous in nature and often
17 change the temporal dynamics of the system (Malchow et al., 2008; Seurout, 2009; Chakraborty et al.,
18 2015). But, till now, very less attention has been paid to investigate the role of generalist predators
19 under the influence of heterogeneous environments.

20 In the past, several researchers used mathematical models to investigate the role of generalist preda-
21 tors on ecological dynamics. Most of them modeled generalist predators simply by using a sigmoidal
22 Holling type III response (which reflects prey switching at low prey concentrations) without considering
23 another food source (Rosenzweig, 1971; Steele and Henderson, 1992; Hesaaraki and Moghadas, 2001;
24 Xu et al., 2004; Kar and Matsuda, 2007; Morozov and Petrovskii, 2009; Chakraborty and Feudel,
25 2014). However, this is inconsistent with the fact that generalist predators can survive in the ab-
26 sence of focal prey. Only a few studies investigated the role of generalist predators in the presence
27 of additional food source in predator-prey systems. Spencer and Collie (1995) and Chakraborty and
28 Chattopadhyay (2008) considered a linear growth term to represent the growth of a predator due to
29 the additional food source apart from the growth due to focal prey species. van Baalen et al. (2001)
30 examined the switching between a focal prey and alternative food source by considering the alternative
31 food density as constant. van Leeuwen et al. (2007) discussed the validation of different functional

32 responses for generalist predators and found that generalist predators can have both stabilizing and
 33 destabilizing effects on the system dynamics. Similar to Spencer and Collie (1995), Magal et al. (2008)
 34 also considered additional food for a generalist predator, but Holling type II functional response for
 35 the uptake of focal prey rather than a sigmoidal functional response. Recently, Erbach et al. (2013)
 36 modeled a generalist predator by density-dependent birth rate of the predator and a linear death rate.
 37 Moreover, there are also few studies where generalist predators are modeled in the presence of spatial
 38 heterogeneity. Some of them did not consider an extra food source for the generalist predator (Rosen-
 39 zwig, 1973; Segel and Levin, 1976) whereas others did not investigate different pattern formations due
 40 to the presence of generalist predators (Magal et al., 2008; Kumari, 2013). In the present paper, I
 41 investigate how a generalist predator affects the spatial distribution of the populations and results in
 42 different pattern formations.

43 Here, a two-dimensional reaction-diffusion predator-prey system is considered where the predator
 44 is a generalist predator and has additional food source apart from the focal prey population. The main
 45 focus of the paper is to investigate how the presence of a generalist predator affects the spatial distri-
 46 bution of the predator and prey populations. The dynamics with linear as well as density-dependent
 47 birth rate of the predator as considered in Spencer and Collie (1995) and Erbach et al. (2013), respec-
 48 tively, is investigated. Furthermore, the situation when the additional food source coincides with the
 49 food source of the focal prey is also examined. This kind of predation is known as intraguild predation
 50 (also mixotrophy), a special case of generalist predation (Gagnon et al., 2011; Kang and Wedekin,
 51 2013). In this case, the predator is involved in competition for the common resources with the prey in
 52 addition to predate on them. For example, the scorpion *Paruroctonus mesaenis* eats smaller arachnid
 53 and insect predators together with the prey of these predators (Polis and McCormick, 1987). Several
 54 other examples of intraguild predation from natural communities can be found in Polis et al. (1989).

55 The rest of the article is organized as follows: Section 2 deals with the model considering linear
 56 and density dependent birth rate of the predator due to the additional food source. Specifically, the
 57 model with linear birth rate of the predator due to the additional food and diffusion is presented in
 58 Section 2.1. Section 2.2 and 2.3 consist of the linear stability analysis of the model without diffusion
 59 and Turing instability conditions of the model with diffusion, respectively. In Section 2.4, different

dynamics of the system are examined numerically and different types of pattern formation are shown in subsection 2.5 and 2.6. In subsection 2.7, a system with density-dependent birth rate of the predator is stated and the results are compared with the results from the previous section. A model with an intraguild predator is presented and analyzed in Section 3. Finally, the paper ends with a discussion.

2. A predator-prey model with a generalist predator

2.1. Basic model structure

Here, a reaction-diffusion system with a prey and a generalist predator in the presence of additional food for the predator is considered in the following form:

$$\begin{aligned}\frac{\partial n}{\partial t} &= r_1 n \left(1 - \frac{n}{K}\right) - \frac{gnp}{h+n} + D_1 \left(\frac{\partial^2 n}{\partial x^2} + \frac{\partial^2 n}{\partial y^2}\right), \\ \frac{\partial p}{\partial t} &= r_2 p + \frac{egnp}{h+n} - mH(p)p + D_2 \left(\frac{\partial^2 p}{\partial x^2} + \frac{\partial^2 p}{\partial y^2}\right),\end{aligned}\tag{1}$$

where $n(x, y, t)$ and $p(x, y, t)$ denote the densities of the prey and the predator, respectively, at location $(x, y) \in \mathbb{R}^2$ and time $t \geq 0$, r_1 and K are the intrinsic growth rate and carrying capacity of the prey population, respectively, g is the prey capturing rate by the predator, h is the corresponding handling time, e is the efficiency of converting prey into predator biomass ($e < 1$), r_2 is the growth rate of the predator due to the additional food source, D_1 and D_2 are diffusion coefficients of prey and predator, respectively, $mH(p)$ is the death rate of the predator. Concerning the form of $H(p)$, several functions are used in literature with various ecological interpretations (Steele and Henderson, 1992). However, in the present work, to take into account the predation of higher-order predators on the generalist predator that is not explicitly included in the model, a quadratic closure term is chosen, i.e., $H(p) = p$. This form of $H(p)$ assumes that the higher predator population changes in proportion with the generalist predator (Steele and Henderson, 1981).

Let, Ω be the two-dimensional bounded connected square domain with $\partial\Omega$ as boundary, and $\frac{\partial}{\partial\eta}$ be the outward drawn normal derivative on the boundary. In Ω , the following initial conditions are taken for system (1)

$$n(0, x, y) = n_0(x, y) > 0, \quad p(0, x, y) = p_0(x, y) > 0, \quad \forall (x, y) \in \Omega$$

and the zero-flux boundary conditions are chosen as

$$\frac{\partial n}{\partial \eta}|_{(x,y)} = \frac{\partial p}{\partial \eta}|_{(x,y)} = 0, \text{ where } (x, y) \in \partial\Omega.$$

It is to be noted here that the general model structure of system (1) is similar with the model of Magal et al. (2008) where a host-parasitoid model was considered to search for the conditions to restrict the growth of the host population. However, the motivation of the present work is completely different; here different pattern formations in a predator-prey system are investigated depending on the additional food source. In the following, the conditions for local asymptotic stability and Turing instability will be derived.

2.2. Linear stability analysis

To study Turing instability, first we need to analyze the stability criteria of the non-diffusive version of system (1). The corresponding non-diffusive model is

$$\begin{aligned} \frac{dn}{dt} &= r_1 n \left(1 - \frac{n}{K}\right) - \frac{gnp}{h+n}, \\ \frac{dp}{dt} &= r_2 p + \frac{egnp}{h+n} - mp^2. \end{aligned} \quad (2)$$

System (2) possesses four different equilibrium points: (i) the population free equilibrium $E_0 = (0, 0)$, (ii) the predator free equilibrium $E_1 = (K, 0)$, (iii) the prey free equilibrium $E_2 = (0, \frac{r_2}{m})$, and (iv) the interior equilibrium $E_*(n_*, p_*)$ with $p_* = \frac{r_1}{g} \left(1 - \frac{n_*}{K}\right)(h + n_*)$, and n_* is a positive root of the equation

$$n^3 + an^2 + bn + c = 0,$$

where

$$a = 2h - K, \quad b = \frac{gK}{r_1 m} + h^2 - 2hK, \quad c = hK \left(\frac{r_2 g}{r_1 m} - h\right).$$

It is clear that the equilibrium points E_0 , E_1 and E_2 always exist. Let us denote

$$\alpha = a^2 - b \text{ and } \beta = 2a^2 - 3ab + c.$$

Then the existence conditions of the interior equilibrium are obtained by using the criteria given by Murray (1989) as:

(i) If $\alpha > 0$ and either $\beta = 0$ or $|\beta| \leq 2\alpha^{\frac{2}{3}}$, there is a possibility of the existence of zero, one, two or three non-trivial equilibria. It is to be mentioned here that this is a necessary but not sufficient condition to obtain three non-trivial equilibria.

107 (ii) If $\alpha > 0$ and $|\beta| > 2\alpha^{\frac{2}{3}}$ or $\alpha \leq 0$, we have at most one non-trivial equilibrium.

108 From the biological point of view (regarding pattern formation), the most interesting thing would be
 109 to study the stability of the interior equilibrium point E_* . The Jacobian matrix corresponding to E_*
 110 can be written as:

$$111 \quad J = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix},$$

112 where $a_{11} = -\frac{r_1 n_*}{K} + \frac{g n_* p_*}{(h+n_*)^2}$, $a_{12} = -\frac{g n_*}{h+n_*}$, $a_{21} = \frac{e g h p_*}{(h+n_*)^2}$, $a_{22} = -m p_*$.

113 The corresponding characteristic equation of J is

$$114 \quad \lambda^2 + A\lambda + B = 0,$$

115 where

$$116 \quad A = -(a_{11} + a_{22}) = \frac{r_1 n_*}{K} + m p_* - \frac{g n_* p_*}{(h+n_*)^2},$$

$$117 \quad B = a_{11} a_{22} - a_{12} a_{21} = m p_* \left(\frac{r_1 n_*}{K} - \frac{g n_* p_*}{(h+n_*)^2} \right) + \frac{e g^2 h n_* p_*}{(h+n_*)^3}.$$

118 Here A and B are the trace and determinant of J , respectively. Our main interest is to investigate
 119 the Turing instability of the system where the uniform steady state of the system without diffusion is
 120 stable, but it is unstable in the partial differential equations with diffusion terms. Now, the condition
 121 for the uniform steady state to be stable for the corresponding ordinary differential equation (2) is
 122 given by

$$123 \quad A > 0 \text{ and } B > 0.$$

124 2.3. Turing instability

125 Here, the condition for Turing instability of the spatially positive steady state E_* of system (1)
 126 will be investigated. Although, the Turing instability criterion is obtained following the standard
 127 analysis (Murray, 2003; Edelstein-Keshet, 1988; Okubo and Levin, 2001; Segel and Jackson, 1972), it
 128 is included here for the completeness of the text. To study this, let us consider the linearized form of
 129 system (1) about $E_*(n_*, p_*)$ as follows:

$$\begin{aligned} \frac{\partial n_1}{\partial t} &= a_{11} n_1 + a_{12} p_1 + D_1 \left(\frac{\partial^2 n_1}{\partial x^2} + \frac{\partial^2 n_1}{\partial y^2} \right), \\ \frac{\partial p_1}{\partial t} &= a_{21} n_1 + a_{22} p_1 + D_2 \left(\frac{\partial^2 p_1}{\partial x^2} + \frac{\partial^2 p_1}{\partial y^2} \right), \end{aligned} \quad (3)$$

130 where, $n = n_* + n_1, p = p_* + p_1$. Here, (n_1, p_1) are small perturbations of (n, p) about the interior
 131 equilibrium point $E_*(n_*, p_*)$. Now consider the solution of system (3) in the form

$$132 \quad \begin{pmatrix} n_1 \\ p_1 \end{pmatrix} = \begin{pmatrix} N_k \\ P_k \end{pmatrix} e^{\lambda_1 t + i(\kappa_x x + \kappa_y y)}$$

133 where λ_1 is the growth rate of perturbation in time t , κ_x and κ_y represent the wave numbers of the
 134 solution. The Jacobian matrix of the linearized system can be written as:

$$135 \quad \tilde{J} = \begin{pmatrix} a_{11} - D_1(\kappa_x^2 + \kappa_y^2) & a_{12} \\ a_{21} & a_{22} - D_2(\kappa_x^2 + \kappa_y^2) \end{pmatrix}.$$

136 In the spatial model, the value of λ_1 depends on the sum of the square of wave numbers $\kappa_x^2 + \kappa_y^2$
 137 (Baurmann et al., 2004). As a result, both wave numbers affect the eigenvalues. This makes clear
 138 that some Fourier modes will vanish in the long-term limit whereas others will amplify. For the sake
 139 of simplicity, we can make use of λ_1 being rotational symmetric function on the (κ_x, κ_y) -plane and
 140 substitute $\kappa^2 = \kappa_x^2 + \kappa_y^2$ and obtain the results for the two-dimensional case from the one-dimensional
 141 formulation. Thus, the corresponding characteristic equation of system (1) is given by

$$\lambda_1^2 + \tilde{A}\lambda_1 + \tilde{B} = 0, \quad (4)$$

142 where

$$143 \quad \tilde{A} = A + \kappa^2(D_1 + D_2),$$

$$144 \quad \tilde{B} = B - (a_{11}D_2 + a_{22}D_1)\kappa^2 + D_1D_2\kappa^4.$$

145 Using the Routh-Hurwitz criterion, it appears that the equilibrium point E_* is locally asymptotically
 146 stable in the presence of diffusion iff $\tilde{A} > 0$ and $\tilde{B} > 0$. Clearly, $A > 0$ implies $\tilde{A} > 0$. Therefore,
 147 diffusive instability occurs only in the case when $B > 0$, but $\tilde{B} < 0$. Hence, the condition for diffusive
 148 instability is given by

$$H(\kappa^2) = D_1D_2\kappa^4 - (a_{11}D_2 + a_{22}D_1)\kappa^2 + B < 0. \quad (5)$$

149 This shows that diffusion can induce the loss of stability with respect to perturbations of certain wave
 150 numbers. Here, H is a quadratic function of κ^2 and the graph of $H(\kappa^2) = 0$ is a parabola. Let, the
 151 minimum of $H(\kappa^2) = 0$ is reached at $\kappa^2 = \kappa_c^2$, where κ_c^2 is given by

152

$$\kappa_c^2 = (a_{11}D_2 + a_{22}D_1)/2D_1D_2.$$

153

Therefore, with the above value of κ_c^2 , the condition for diffusive instability given in Eq. (5) can be

154

written as

155

$$(a_{11}D_2 + a_{22}D_1)^2 > 4D_1D_2B.$$

156

In explicit form, the condition becomes

$$\left\{ mp_*D_1 + \left(\frac{r_1n_*}{K} - \frac{gn_*p_*}{(h+n_*)^2} \right) D_2 \right\}^2 > 4D_1D_2 \left\{ mp_* \left(\frac{r_1n_*}{K} - \frac{gn_*p_*}{(h+n_*)^2} \right) + \frac{eg^2hn_*p_*}{(h+n_*)^3} \right\}. \quad (6)$$

157

Since it is not prominent from analytic conditions how the local asymptotic stability and the Turing

158

instability depend on r_2 , further investigation in the form of numerical simulation is carried out in the

159

following.

160

2.4. Numerical simulation

161

In this section, numerically it is examined how a generalist predator influences the system dynamics

162

depending on the availability of the additional food source. Specifically, the growth rate of the predator

163

due to the additional food, r_2 , is varied and observe the changes in the dynamics of the system where

164

the other parameter values are fixed at $r_1 = 2$, $K = 10$, $g = 2$, $h = 5$, $e = 0.25$, $m = 0.016$. The

165

bifurcation results are obtained by using the software XPPAUT and plotted in MATLAB, whereas the

166

other figures are drawn by writing code in MATLAB.

167

First, the existence of equilibria (marked with filled black circles) and their stability are observed

168

in the phase plane starting at $(n, p) = (2, 4)$ (marked with open black circles) for different r_2 . In

169

Figure 1(a), n and p-nullclines, marked by the dashed blue and green lines, respectively, are plotted at

170

$r_2 = 0$. There exist three different equilibria: (i) $E_0 = (0, 0)$, $E_1 = (10, 0)$ (not shown in the figure) and

171

$E^* = (1.05, 5.41)$. Here, the eigenvalues of E_0 are 0 and 2, and therefore it is unstable. The eigenvalues

172

of E_1 are -2 and 0.33 , and therefore it is an unstable saddle. The eigenvalues of E^* are $0.007 \pm 0.3456i$,

173

and therefore it is an unstable focus surrounded by a limit cycle. The trajectory approaching the limit

174

cycle is shown by the red line. Figure 1(b) is drawn at $r_2 = 0.05$ having four different equilibria: (i)

175

$E_0 = (0, 0)$, $E_1 = (10, 0)$ (not shown in the figure), $E_2 = (0, 3.12)$ and $E^* = (0.35, 5.16)$. Here, the

176

eigenvalues of E_0 are 2 and 0.05, and therefore it is an unstable node. The eigenvalues of E_1 are -2

177

and 0.3833 , and therefore it is an unstable saddle. The eigenvalues of E_2 are 0.75 and -0.05 , and

Figure 1: Phase plane of the model system (2) at different values of r_2 : (a) $r_2 = 0$, (b) $r_2 = 0.05$ and (c) $r_2 = 0.1$. Blue and green dashed lines are the n and p -nullclines, respectively. Different equilibria are marked by the filled black circles. Red lines are the corresponding trajectories starting at $(n, p) = (2, 4)$, marked with open black circles. Parameter values used: $r_1 = 2$, $K = 10$, $g = 2$, $h = 5$, $e = 0.25$, $m = 0.016$. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Figure 2: Two-dimensional bifurcation plot of model system (1) in $r_2 - m$ plane which divides the $r_2 - m$ parametric space into different regions; R_E (blue): stable E_2 ; R_S (white): stable E^* ; R_O (red): oscillating E^* ; and R_T (green): Turing instability. Here TH and TB are Turing-Hopf and Takens-Bogdanov bifurcations, respectively. The upper part of R_O starting from the dashed-dot line is corresponding to the Turing-Hopf domain. Along the lines L_1 , L_2 and L_3 , Figs. 3 (a), -(c) are drawn, respectively. Parameter values used $D_1 = 0.01$ and $D_2 = 0.6$ and the other parameter values are same as in Fig. 1. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

therefore it is also an unstable saddle. The eigenvalues of E^* are $-0.0132 \pm 0.2324i$, and therefore it is a stable focus and the corresponding trajectory reaching towards E^* is shown by the red line. Next, Figure 1(c) is drawn at $r_2 = 0.1$. In this case, E^* does not exist. The other equilibria are: (i) $E_0 = (0, 0)$, $E_1 = (10, 0)$ (not shown in the figure), and $E_2 = (0, 6.25)$. The eigenvalues of E_0 are 2 and 0.1, and therefore it is an unstable node. The eigenvalues of E_1 are -2 and 0.43 , and therefore it is an unstable saddle. The eigenvalues of E_2 are -0.5 and -0.1 , and therefore it is a stable node and the corresponding trajectory reaching E_2 is shown by the red line.

To get a clearer view on how the presence of additional food source influences different dynamical behavior of the system, a two-parameter bifurcation diagram is drawn by varying the growth rate of the predator (r_2) due to the additional food and the mortality of the predator (m) (Figure 2). There are four different dynamical behaviors of the system marked by different regions R_E , R_S , R_O , and R_T . In region R_E (marked by the blue color), E_2 is locally asymptotically stable (LAS), i.e., in this parametric region prey population becomes extinct due to high predation pressure and the predator

191 population survives solely on the additional food source. Region R_S (marked by the white color) is
 192 corresponding to the stable E^* , i.e., both the populations stably coexist in this parametric region. In
 193 region R_O (marked by the red color), E^* becomes unstable, and both the populations coexist with
 194 fluctuating densities. Region R_T (marked by the green color) is the Turing space, i.e., in this region,
 195 E^* remains stable for the system without diffusion, but becomes unstable in the presence of diffusion.
 196 As a result, different stationary spatially inhomogeneous patterns of predator and prey populations
 197 emerge within this region. The existence of two codimension-2 bifurcations are also observed, where
 198 the bifurcation curves interact. The first one is the Takens-Bogdanov bifurcation (TB) where the
 199 Hopf bifurcation and transcritical bifurcation meet. The other one is Turing-Hopf bifurcation (TH)
 200 where the Turing bifurcation and Hopf bifurcation meet. The backward extended lower boundary of
 201 the Turing space, marked by the dash-dot line, divides the region R_O into two parts. The upper part
 202 of this region is the Turing-Hopf domain where the inhomogeneous stationary patterns caused by the
 203 Turing instability interacts with the oscillations due to the Hopf bifurcation. Clearly, at lower rates
 204 of predator mortality, the presence of additional food to the predator helps in the stabilization of the
 205 system, whereas very high growth due to additional food results in prey extinction. On the other
 206 hand, when the mortality rate is comparatively high, the presence of additional food can make the
 207 distribution of the prey and predator inhomogeneous in space.

208 To get an overview of how prey abundance changes with r_2 , three one-dimensional bifurcation
 209 diagrams are plotted (Figure 3) by varying r_2 continuously at (a) $m = 0.01$, (b) $m = 0.017$, and
 210 (c) $m = 0.025$, which are drawn along the lines L_1 , L_2 and L_3 , respectively, as indicated in Figure
 211 2. Specifically, the steady-state values of the abundances of the prey population are plotted with
 212 r_2 . The black and red (dashed) lines indicate that the interior steady state is stable and unstable,
 213 respectively. The magenta (dashed) lines indicate that the steady state corresponding to the extinction
 214 of prey is stable. Additionally, the green lines represent the maximum and minimum abundances of
 215 the populations for the stable limit cycle. Color coding of the ranges of r_2 is same as in Figure 2. From
 216 Figure 3(a) it is clear that the prey population shows high fluctuation at low values of r_2 . However, an
 217 increase in r_2 stabilizes system dynamics and finally prey population goes extinct from the system. In
 218 this case Turing instability does not occur. Figure 3(b) shows a similar kind of behavior except for the

Figure 3: One-dimensional bifurcation diagrams to show how prey abundances change with r_2 at (a) $m = 0.01$, (b) $m = 0.017$ and (c) $m = 0.025$. They are drawn along the lines L_1 , L_2 and L_3 , respectively, of Fig. 2. Color coding represents similar regions as that of Fig. 2. The black and red (dashed) lines indicate that E^* is stable and unstable, respectively. The magenta (dashed) line is corresponding to stable E_2 . The green lines represent the maximum and minimum abundances of the populations for the stable limit cycle. The other parameter values are same as in Fig. 2. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Figure 4: (a) The graph of the function $H(\kappa^2)$ at $r_2 = 0$ (blue), 0.02 (magenta), 0.03 (cyan), 0.04 (green) and 0.05 (red). The other parameter values are same as in Fig. 3(c). An increase in the value of r_2 increases the possibility of diffusive instability by increasing the interval of negativity of $H(\kappa^2)$. (b) Dispersion relation plotting the largest real part of the eigenvalues at different r_2 ; $r_2 = 0$ (blue) and $r_2 = 0.05$ (red). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

range of r_2 just after the Hopf bifurcation where Turing instability occurs. For comparatively higher values of m , Figure 3(c) shows the non-existence of oscillating and prey-extinction regions. However, the range of r_2 for Turing instability is much larger compared to the previous case. In this case, the conditions of Turing instability obtained analytically (Eq. (5)) are also checked by plotting $H(\kappa^2)$ for different values of r_2 . Turing instability condition $\min(H(\kappa^2)) < 0$ is satisfied within the range $r_2 \in (0.28, 0.9)$. In Figure 4(a), the curve $H(\kappa^2) = 0$ is plotted for $r_2 = 0$ (blue), 0.02 (magenta), 0.03 (cyan), 0.04 (green) and 0.05 (red). The largest real parts of the eigenvalues of the characteristic equation (4) of system (1) are also drawn (Figure 4(b)) for $r_2 = 0$ (blue) and $r_2 = 0.05$ (red). The length of the interval of κ^2 within which the largest real part of the eigenvalues are positive provides the existence of diffusive instability.

In the following, different pattern formations are investigated at different values of r_2 .

2.5. Pattern formation

Here, extensive numerical simulations of the spatial model system (1) are performed in two dimensional space using the forward finite difference method, and the results of different pattern formations due to the variation of r_2 are shown.

234 To analyze the dynamic behavior of system (1), the stationary distributions of the prey population
 235 are plotted in two-dimensional spaces. Here, the system is studied on a squared spatial grid of 50×50
 236 points with the Neumann boundary conditions and run the simulation up to the time $t = 5000$ for
 237 different values of r_2 . The space step is taken as 0.2, and the time step as 0.005. It is assumed that the
 238 prey and predator populations are spread over the whole domain at the beginning of the simulation. We
 239 know that the choice of the initial distribution of the populations greatly affects the spatial dynamics
 240 of a system. If the initial spatial distributions of the prey and predator are homogeneous, then the
 241 species distribution remains homogeneous forever, which is not so interesting (Petrovskii and Malchow,
 242 1999). Apart from that, from a biological point of view, it is reasonable to consider a scattered non-
 243 uniform initial distribution of populations over the space under consideration. Here, such scattered
 244 initial distribution has been employed by considering a random sampling of the prey and predator
 245 populations around the equilibrium values of the corresponding non-spatial model. It is assured that
 246 the time at which simulations are stopped is sufficient for the patterns to attain the stationary state
 247 and they do not change further with time.

248 Figure 5 plots the stationary distribution of prey over the spatial domain for four different values
 249 of r_2 ($r_2 = 0.032, 0.037, 0.045, 0.08$) keeping m fixed at 0.025. Specifically, r_2 is varied along the line L_3
 250 in Figure 2 in such a way that r_2 lies within the Turing domain. It is to be mentioned here that, the
 251 distribution of the prey and predator remains homogeneous in space in the absence of additional food
 252 ($r_2 = 0$) (the figure is not shown). Clearly, as r_2 increases, different types of dynamics emerge and
 253 it is observed that the distributions of prey and predator are always of the same type. Consequently,
 254 it is enough to show only the distributions of the prey for different r_2 . At $r_2 = 0.032$, a cold spot
 255 pattern is observed. As we increase r_2 , at $r_2 = 0.037$, the stripe pattern dominates the space. Again,
 256 at $r_2 = 0.045$, a mixture of hot spot and stripe patterns can be found, although hot spots dominate in
 257 this case. Finally, at $r_2 = 0.08$, we see stable hot spots with high prey densities in isolated zones.

258 *2.6. Spatiotemporal chaos*

259 Next, the spatial pattern formations of system (1) are examined by considering the parameters lying
 260 outside the Turing domain and inside the Hopf domain. Following the insightful work of Medvinsky
 261 et al. (2002), Wang et al. (2010), and Upadhyay et al. (2010), three different initial distributions are

Figure 5: Stationary pattern formations of prey population over space at different values of r_2 ; $r_2 = 0.032$: cold spots; $r_2 = 0.037$: stripes; $r_2 = 0.045$: mixture of stripes and hot spots; $r_2 = 0.08$: hot spots. Parameter values used $m = 0.025$, $D_1 = 0.01$, $D_2 = 0.6$ and the other parameter values are same as in Fig. 2.

Figure 6: Formation of spiral pattern and its destruction for prey population at $t = 0, 400, 800$, and 3000 . The parameter values used $m = 0.01$, $r_2 = 0.01$, $D_1 = 0.1$, $D_2 = 0.2$ and the other parameters are same as in Fig. 2 and the initial distribution is given in Eq. (7).

Figure 7: (a) Plot of spatial average values of prey population against time with parameter values same as in Fig. 6 showing chaotic oscillation. (b) Time evolution of prey population at the spatial location $(25, 25)$.

chosen to investigate the evolutionary process of the prey population in pattern formation. In this case, the system is studied on a squared spatial grid of 200×200 points and the parameters used are $r_2 = 0.01$, $m = 0.01$, $D_1 = 0.1$, and $D_2 = 0.2$, whereas the other parameters are same as in Fig. 5.

In the first case, the initial distribution of the populations is chosen as

$$\begin{aligned} n(x, y, 0) &= n^* - \varepsilon_1(x - 100), \\ p(x, y, 0) &= p^* - \varepsilon_2(y - 100), \end{aligned} \tag{7}$$

with $\varepsilon_1 = 2 \times 10^{-3}$ and $\varepsilon_2 = 3 \times 10^{-3}$. Snapshots of the spatial distributions are shown in Figure 6 for $t = 0, 400, 800$, and 3000 . Clearly, the formation of the irregular patchy structure can be preceded by the evolution of a regular spiral pattern. Here, the occurrence of the spiral is not due to the initial conditions. The center of the spiral is situated at the critical point $(x^*, y^*) = (100, 100)$ with $n(x^*, y^*) = n^*$, $v(x^*, y^*) = p^*$. After the formation of the spiral, it grows upto a certain time, following the destruction of the spiral by making an irregular patchy pattern all over the domain.

Here, the distribution of the prey population does not converge to any stationary state. The spatial average of the prey population with time is plotted in Figure 7(a) which shows chaotic oscillation. The prey abundance at the spatial position $(25, 25)$ is also plotted with respect to time in Figure 7(b) which also shows an irregular oscillation with time.

In the second case, a different set of initial distribution of the populations is chosen as

$$\begin{aligned} n(x, y, 0) &= n^* - \varepsilon_1(x - 40)(x - 160) - \varepsilon_2(y - 60)(y - 140), \\ p(x, y, 0) &= p^* - \varepsilon_3(x - 90) - \varepsilon_4(y - 100), \end{aligned} \tag{8}$$

with $\varepsilon_1 = 3 \times 10^{-6}$, $\varepsilon_2 = 8 \times 10^{-6}$, $\varepsilon_3 = 3 \times 10^{-4}$, and $\varepsilon_4 = 6 \times 10^{-4}$. Snapshots of the spatial distribution are shown in Figure 8 for $t = 0, 600, 900$, and 3000 . Here, the initial distribution contains two critical points, which are $(40, 140)$ and $(160, 60)$. As a result, two spirals emerge with centers situated at the above mentioned points. In this case also the spiral pattern is destroyed and an irregular patchy pattern is formed all over the domain.

Finally, another set of initial distribution of the populations is considered as mentioned in the following

$$n(x, y, 0) = n^* - \varepsilon_1(x - 40)(x - 160),$$

Figure 8: Formation of spiral pattern and its destruction for prey population at $t = 0, 600, 900$, and 3000 with parameter values same as in Fig. 6. The initial distribution is given in Eq. (8).

Figure 9: Formation of spiral pattern and its destruction for prey population at $t = 0, 400, 800$, and 3000 with parameter values same as in Fig. 6. The initial distribution is given in Eq. (9).

$$p(x, y, 0) = p^* - \varepsilon_2(y - 40)(y - 160), \quad (9)$$

with $\varepsilon_1 = 2 \times 10^{-5}$ and $\varepsilon_2 = 3 \times 10^{-5}$. Snapshots of the spatial distribution are shown in Figure 9 for $t = 0, 400, 800$, and 3000 . Here, the occurrence of four spirals is observed, which are finally destroyed and makes the spatial domain patchy.

2.7. Density dependent birth rate for the generalist predator

The behavior of system (1) is also checked by considering a density dependent birth rate of the generalist predator due to the additional food source (Erbach et al., 2013) in the form $\frac{r_2 p}{h_1 + p}$ where h_1 represents the half saturating constant for the growth of the predator due to the additional food source. In the absence of focal prey, the reproduction term of the predator population looks like Beverton-Holt function.

The behavior of the new system is checked at $h_1 = 1$. It is observed that the new system shows qualitatively similar spatial behavior as system (1). Only the difference is that the region of oscillation, R_O (comparing with Figure 2) is relatively bigger and the prey extinction occurs at larger values of r_2 .

3. Model with intraguild predation

In this section, a particular type of generalist predator is considered, called intraguild predator. In the case of intraguild predation, the additional food source of the predator coincides with the food source of the prey (Gagnon et al., 2011; Kang and Wedekin, 2013). System (1) can be modified in the presence of intraguild predation as:

$$\begin{aligned} \frac{\partial n}{\partial t} &= r_1 n \left(1 - \frac{n + \varepsilon p}{K} \right) - \frac{g n p}{h + n} + D_1 \left(\frac{\partial^2 n}{\partial x^2} + \frac{\partial^2 n}{\partial y^2} \right), \\ \frac{\partial p}{\partial t} &= r_2 \varepsilon p \left(1 - \frac{n + \varepsilon p}{K} \right) + \frac{e g n p}{h + n} - m p^2 + D_2 \left(\frac{\partial^2 p}{\partial x^2} + \frac{\partial^2 p}{\partial y^2} \right), \end{aligned} \quad (10)$$

where ε is the fraction of the predator population involved in intraguild predation. Clearly, $\varepsilon = 0$ represents the situation where p is not an intraguild (generalist) predator.

It is to be noted here that the intraguild predators share the same food as that of the prey population and as a result, they are involved in competition with the prey population for the common food source in addition to predate on them. A special kind of intraguild predation is known as mixotrophy where

mixotrophs use a mix of different sources of energy and carbon, and because of that they compete with their prey organisms. Our mathematical form of intraguild predation is similar with the form used by Hammer and Pitchford (2005) where mixotrophy was explained in a phytoplankton-zooplankton system.

In the absence of diffusion, system (10) possesses four different equilibrium points: (i) the population free equilibrium $\bar{E}_0 = (0, 0)$, (ii) the predator free equilibrium $\bar{E}_1 = (K, 0)$, (iii) the prey free equilibrium $\bar{E}_2 = (0, \frac{r_2 \varepsilon}{m})$, and (iv) the interior equilibrium $\bar{E}_*(\bar{n}_*, \bar{p}_*)$ which can be obtained by solving the equations

$$\begin{aligned} r_1 \left(1 - \frac{n + \varepsilon p}{K} \right) - \frac{gp}{h + n} &= 0, \\ r_2 \varepsilon \left(1 - \frac{n + \varepsilon p}{K} \right) + \frac{egn}{h + n} - mp &= 0. \end{aligned}$$

Here, the condition for LAS of the non-diffusive version of system (10) is

$$A_1 > 0 \text{ and } B_1 > 0,$$

where

$$\begin{aligned} A_1 &= -(a_{11} + a_{22}) = \frac{r_1 \bar{n}_*}{K} + m\bar{p}_* - \frac{g\bar{n}_*\bar{p}_*}{(h + \bar{n}_*)^2} + \frac{r_2 \varepsilon^2 \bar{p}_*}{K}, \text{ and} \\ B_1 &= a_{11}a_{22} - a_{12}a_{21} = (m\bar{p}_* + \frac{r_2 \varepsilon^2 \bar{p}_*}{K}) \left(\frac{r_1 \bar{n}_*}{K} - \frac{g\bar{n}_*\bar{p}_*}{(h + \bar{n}_*)^2} \right) + \left(\frac{r_1 \varepsilon \bar{n}_*}{K} + \frac{g\bar{n}_*}{(h + \bar{n}_*)} \right) \left(-\frac{r_2 \varepsilon \bar{p}_*}{K} + \frac{egh\bar{p}_*}{(h + \bar{n}_*)^2} \right). \end{aligned}$$

Here, A_1 and B_1 are the trace and determinant of the corresponding Jacobian, respectively. The condition for diffusive instability is given by

$$H_1(\kappa^2) = D_1 D_2 \kappa^4 - \left((m\bar{p}_* + \frac{r_2 \varepsilon^2 \bar{p}_*}{K}) D_1 + \left(\frac{r_1 \bar{n}_*}{K} - \frac{g\bar{n}_*\bar{p}_*}{(h + \bar{n}_*)^2} \right) D_2 \right) \kappa^2 + B_1 < 0. \quad (11)$$

Following the same method as previous, it is possible to write down the explicit form of the condition for diffusive instability as

$$\begin{aligned} \left\{ (m\bar{p}_* + \frac{r_2 \varepsilon^2 \bar{p}_*}{K}) D_1 + \left(\frac{r_1 \bar{n}_*}{K} - \frac{g\bar{n}_*\bar{p}_*}{(h + \bar{n}_*)^2} \right) D_2 \right\}^2 &> \\ 4D_1 D_2 \left\{ (m\bar{p}_* + \frac{r_2 \varepsilon^2 \bar{p}_*}{K}) \left(\frac{r_1 \bar{n}_*}{K} - \frac{g\bar{n}_*\bar{p}_*}{(h + \bar{n}_*)^2} \right) + \left(\frac{r_1 \varepsilon \bar{n}_*}{K} + \frac{g\bar{n}_*}{(h + \bar{n}_*)} \right) \left(-\frac{r_2 \varepsilon \bar{p}_*}{K} + \frac{egh\bar{p}_*}{(h + \bar{n}_*)^2} \right) \right\}. \end{aligned} \quad (12)$$

First, the condition of Turing instability obtained analytically in Eq. (11) is checked by plotting $H_1(\kappa^2)$ for different values of ε . In Figure 10 (top), the curve $H_1(k^2) = 0$ is plotted for $\varepsilon = 0$ (blue), 0.01 (magenta), 0.02 (cyan), 0.03 (green) and 0.04 (red). Clearly, the Turing instability condition

Figure 10: (Top) The graph of the function $H_1(\kappa^2)$ for system (10) at $\varepsilon = 0$ (blue), 0.01 (magenta), 0.02 (cyan), 0.03 (green) and 0.04 (red). (Bottom) Stationary pattern formations of prey population over space at $\varepsilon = 0.07$. The parameter values used $r_1 = 0.8$, $D_1 = 0.001$, $D_2 = 0.1$ and the other parameter values are same as in Fig. 4. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

327 $\min(H_1(\kappa^2)) < 0$ is satisfied for higher values of ε which results in Turing pattern formation. Next,
 328 a numerical example of Turing pattern formation is shown. Figure 10 (bottom) is drawn at $\varepsilon = 0.07$
 329 which clearly shows stationary pattern formation by the prey population in the presence of intraguild
 330 predator.

331 4. Discussion

332 Predator-prey interactions affect species composition and community dynamics. The complexity
 333 in a community depends on the type of predation, which differs for different predators. Generalist
 334 predators increase such complexity by feeding on a variety of prey items. In the present work, the

335 influences of two different types of generalist predators are investigated: (i) the predator is having
 336 an additional food source apart from the focal prey, and (ii) the predator is an intraguild predator
 337 where the additional food source coincides with the food of the prey, which results in a competition
 338 between the prey and the predator for the common food. Here, a separate growth term for the
 339 generalist predator is considered to represent its growth due to the additional food sources. The
 340 non-spatial version of the model shows stabilizing effect of generalist predators on system dynamics.
 341 However, the most interesting result occurs after considering diffusion in the model system in order to
 342 investigate the role of the generalist predator in the presence of spatial movements of both predator
 343 and prey populations. Although, the presence of the generalist predator assures temporal stability, the
 344 distribution of both prey and predator populations can become inhomogeneous in space and results
 345 in different patterns, like stripes, spots, and the mixture of them depending on the availability of
 346 the additional food to the generalist predator. Moreover, spatiotemporal chaotic patterns have also
 347 been observed for a certain range of the availability of additional food and mortality of the generalist
 348 predator.

349 Most of the previous modeling studies revealed the stabilizing role of generalist predators (Ander-
 350 sson and Erlinge, 1977; Turchin and Hanski, 1997; van Baalen et al., 2001; Smout et al., 2010). The
 351 presence of generalist predators results in the dampening or elimination of the cyclical interactions
 352 between predators and their prey (Hanski et al. 1991). Several empirical evidences also support this
 353 claim (Erlinge et al., 1983; Hanski et al., 1991). However, under certain conditions, it can also have
 354 destabilizing effects (Chakraborty and Chattopadhyay, 2008). Matthiopoulos et al. (2007) studied the
 355 interaction between a generalist predator Hen Harrier (*Circus cyaneus*) and three of its prey species
 356 in the United Kingdom, the Meadow Pipit (*Anthus pratensis*), the field vole (*Microtus agrestis*), and
 357 the Red Grouse (*Lagopus lagopus scoticus*). They found that the generalist predator can damp or
 358 suppress the cyclic oscillation in grouse population when the alternative prey density remains low.
 359 But, the presence of high alternative prey results in an increase in the oscillation. The present spatial
 360 system can also show a similar destabilizing effect on system dynamics in the presence of additional
 361 food. However, in this case, the destabilization occurs in space, whereas the temporal dynamics still
 362 remain stable. Under different conditions, additional food can also stabilize the system in both time

363 and space.

364 The presence of generalist predators can make the system dynamics very complex. Previously,
365 bistability between two alternative stable coexistence states, and bistability between a coexistence
366 state and a stable limit cycle have been observed in a single prey-generalist predator system (Spencer
367 and Collie, 1995). Magal et al. (2008) found the existence of homoclinic loops in the presence of
368 generalist predator. Moreover, Erbach et al. (2013) found bistability, limit cycles and several global
369 bifurcations in a simple predator-prey system with generalist predator. The present paper shows
370 that, in spite of having less complex dynamics in the temporal model, the consideration of spatial
371 inhomogeneity can result in different complex behaviors due to the presence of generalist predators.
372 Addition of diffusion results in Turing instability, where the prey and predator populations oscillate in
373 space although remain stationary in time. Similar kind of Turing instability was previously observed
374 in a host-parasitoid model with generalist predation on host population by Wilson et al. (1999).
375 They extended the Nicholson and Bailey model (1935) by incorporating the dispersal of both host
376 and parasitoid offspring and found either stable pattern or rapid host extinction depending on the
377 initial conditions. Generalist predation has also been observed to produce spatially varying stable
378 patterns in the context of the McArthur-Resenzweig predator-prey model (Rosenzweig, 1973; Segel
379 and Levin, 1976). However, the consideration of additional food source for the generalist predator
380 which helps generalist predator to survive in the absence of focal prey makes the present approach
381 more realistic and unique. Moreover, the existence of Turing-Hopf bifurcation and Takens-Bogdanov
382 bifurcation is also observed, which are codimension-2 bifurcations resulting due to the interaction of
383 Hopf and Turing bifurcations, and Hopf and transcritical bifurcations, respectively. The existence of
384 spatiotemporal chaos in the presence of generalist predator is another interesting finding of the present
385 work. Previously, Kumari (2013) observed the existence of chaos in a spatial prey-predator-top predator
386 system where the top predator was considered as the generalist predator. In the present case, chaos
387 occurs in a parametric range that falls outside the Turing domain. Such generation of chaotic patterns
388 outside the Turing domain was found in some of the previous studies without generalist predators
389 (Baurmann et al., 2007; Banerjee and Petrovskii, 2011; Banerjee and Abbas, 2014).

390 Spatial variations in population densities due to the variation of extrinsic factors such as nutrient

391 concentration, moisture and temperature, are normal phenomena in ecological systems. In comparison,
 392 empirical evidences of intrinsically generated fixed spatial patterns are difficult to identify as it is hard
 393 to neglect the extrinsic factors as well as the difficulty in accurately estimating the key interactions
 394 and dispersal parameters. In spite of such difficulty, researchers found several evidences of spatial
 395 pattern formations due to biological factors. For example, the clustered spatial pattern of ant nests
 396 emerges from the natural history of the ant/scale/beetle interaction (Liere et al. 2012). With the help
 397 of experimental and modeling studies, Shiyomi (1980) showed that the spatial pattern of a population
 398 of *Galleria mellonella* is affected by the frequency of attack by the predator *Podisus maculiventris*
 399 (attack ability), the homogeneity of the attack ability within a predator population and the mobility
 400 of the predator. There are also evidences of spatial pattern formation due to the predation by generalist
 401 predators. In a field study, Winder et al. (2005) found a deep impact of spatial distribution of cereal
 402 aphids in the presence of two generalist predators, *Pterostichus melanarius* and *P. madidus*. These
 403 observations support the findings of the present study regarding the possibility of pattern formation
 404 in the presence of generalist predators.

405 Generalist predators have important ecological impacts and wide applicability in the field of bi-
 406 ological control. In practice, generalist predators are used to control the populations of ecologically
 407 damaging species, particularly of agricultural weed and insect pests (DeBach, 1974; Holt and Hochberg,
 408 1997). Such biological controls are environment friendly alternatives for the use of insecticides. How-
 409 ever, the success in controlling damaging species depends on the preferences of the generalist predator
 410 for the focal prey and alternative food (Koss and Snyder, 2005) as well as on the spatial and tem-
 411 poral scales at which the process is studied (Walde, 1994). In this respect, theoretical studies can
 412 provide significant insights in finding optimal strategies for control mechanisms. Previously, Magal et
 413 al. (2008) examined conditions under which the invasion of leafminers can be stopped and reversed
 414 by generalist parasitoid in spatial scale. The present study reveals that the theoretical prediction of
 415 a temporal model can go horribly wrong in real systems where populations are involved in spatial
 416 movements. In the presence of a generalist predator, the system can show different pattern formations
 417 and spatiotemporal chaos which has important implications for ecosystem functioning not only in
 418 terms of their predictability, but also in influencing species persistence (Huisman and Weissing, 1999)

and ecosystem's stability in response to abrupt environmental changes (Petrovskii et al., 2004). The relevance of investigating the role of generalist predators in spatially extended domain was recently mentioned by Erbach et al. (2013).

To the best of our knowledge, the present paper is the first possible theoretical work showing different pattern formations due to the presence of generalist predators. In nature, predator-prey systems are more complex than what a simple two dimensional model can capture. Further investigation and empirical support are needed to confirm the importance of generalist predators in spatial scale. Our next step would be to investigate the effects of generalist predators in the presence of a specialist predator. In that case, the generalist predator would be either sharing food with the focal prey or simply depend on the additional food different from the food source of the focal prey in addition to compete with the specialist prey for the focal prey. To determine proper optimal strategy for biological control we need to examine different mechanisms of pattern formation as they mimic the processes of ecological patterning in real world ecosystems.

Acknowledgment: The author wishes to thank the Alexander von Humboldt Foundation for financial support in the form of a postdoctoral fellowship at the ICBM, Carl von Ossietzky University, Oldenburg, Germany. The author also received support from HCØ postdoctoral fellowship program for a part of this work. The author is grateful to the reviewers for their insightful comments and suggestions on the previous versions of the manuscript.

- [1] Andersson, M., S. Erlinge., 1977. Influence of predation on rodent populations. *Oikos* 29, 591-597.
- [2] Banerjee, M., Abbas, S., 2014. Existence and non-existence of spatial patterns in a ratio-dependent predator-prey model. *Ecological Complexity*. doi:10.1016/j.ecocom.2014.05.005
- [3] Banerjee, M., Petrovskii, S., 2011. Self-organised spatial patterns and chaos in a ratio-dependent predator-prey system. *Theor. Ecol.* 4, 37-53.
- [4] Baurmann, M., Ebenhoh, W., Feudel, U., 2004. Turing instabilities and pattern formation in a benthic nutrient-microorganism system. *Math. Biosci. Eng.* 1, 111-130.
- [5] Baurmann, M., Gross, T., Feudel, U., 2007. Instabilities in spatially extended predator-prey

- 446 systems: Spatio-temporal patterns in the neighborhood of Turing-Hopf bifurcations. J. Theor.
447 Biol. 245, 220-229.
- 448 [6] Chakraborty, S., Chattopadhyay, J., 2008. Nutrient-phytoplankton-zooplankton dynamics in the
449 presence of additional food sourceA mathematical study. Journal of Biological Systems 16 (04),
450 547-564.
- 451 [7] Chakraborty, S., Feudel, U., 2014. Harmful algal blooms: combining excitability and competition.
452 Theoretical Ecology 7 (3), 221-237.
- 453 [8] Chakraborty, S., Tiwary, P., Misra, A., Chattopadhyay, J., 2015. Spatial dynamics of a nutrient-
454 phytoplankton system with toxic effect on phytoplankton. Mathematical Biosciences 264, 94-100.
- 455 [9] Crowder, D.W., Snyder, W.E., 2010. Eating their way to the top? Mechanisms underlying the
456 success of invasive generalist predators. Biol. Invas. 12, 2857-2876.
- 457 [10] DeBach, P., 1974. Biological Control by Natural Enemies, Cambridge University Press, UK.
- 458 [11] Edelstein-Keshet, L., 1988. Mathematical Models in Biology, Birkhauser Mathematical Series.
- 459 [12] Erbach, A., Lutscher, F., Seo. G., 2013. Bistability and limit cycles in generalist predator-prey
460 dynamics. Ecological Complexity 14, 48-55.
- 461 [13] Erlinge, S., Gransson, G., Hansson, L., 1983. Predation as regulating factor on small rodent
462 populations in southernmost Sweden. Oikos 40, 36-52.
- 463 [14] Gagnon, AÉ., Heimpel, GE., Brodeur, J., 2011. The Ubiquity of Intraguild Predation among
464 Predatory Arthropods. PLoS ONE 6(11), e28061.
- 465 [15] Hammer, A.C., Pitchford, J.W., 2005. The role of mixotrophy in plankton bloom dynamics, and
466 the consequences for productivity. ICES Journal of Marine Science 62, 833-840.
- 467 [16] Hanski, I., Hansson, L., Henttonen, H., 1991. Specialist predators, generalist predators, and the
468 microtine rodent cycle. Journal of Animal Ecology 60, 353-367.
- 469 [17] Hesaaraki, M., Moghadas, S.M., 2001. Existence of limit cycles for predator-prey systems with
470 a class of functional responses. Ecological Modelling 142, 1-9.

- [18] Holt, R.D., Hochberg, M., 1997. When is biological control evolutionarily stable (or is it)? *Ecology* 78, 1673-1683.
- [19] Huisman, J., Weissing, F.J., 1999. Biodiversity of plankton by oscillations and chaos. *Nature* 402, 407-410
- [20] Jaworski, CC., Bompard, A., Genies, L., Amiens-Desneux, E., Desneux, N., 2013. Preference and Prey Switching in a Generalist Predator Attacking Local and Invasive Alien Pests. *PLoS ONE* 8(12), e82231.
- [21] Kang Y., Wedekin L., 2013. Dynamics of a intraguild predation model with generalist or specialist predator. *J. Math. Biol.* 67, 1227-1259.
- [22] Kar, T.K., Matsuda, H., 2007. Global dynamics and controllability of a harvested prey-predator system with Holling type III functional response. *Nonlinear Analysis: Hybrid Systems* 1, 59-62.
- [23] Koss, A.M., Synder, W.E., 2005. Alternative prey disrupt biocontrol by a guild of generalist predators. *Biol. Control.* 32, 243-251.
- [24] Kumari, N., 2013. Pattern Formation in Spatially Extended Tritrophic Food Chain Model Systems: Generalist versus Specialist Top Predator. *ISRN Biomathematics*, Article ID 198185, 12 pages
- [25] Liere, H., Jackson, D., Vandermeer, J., 2012. Ecological Complexity in a Coffee Agroecosystem: Spatial Heterogeneity, Population Persistence and Biological Control. *PLoS ONE* 7(9), e45508.
- [26] Magal, C., Cosner, C., Ruan, S., Casas, J., 2008. Control of invasive hosts by generalist parasitoids. *Mathematical Medicine and Biology* 25, 1-20.
- [27] Malchow, H., Petrovskii, S., Venturino, E., 2008. Spatiotemporal patterns in Ecology and Epidemiology: Theory, Models, Simulations. Chapman & Hall / CRC Press.
- [28] Matthiopoulos, J., Graham, K., Smout, S., Asseburg, C., Redpath, S., Thirgood, S., Hudson, P. and Harwood, J., 2007. Sensitivity to assumptions in models of generalist predation on a cyclic prey. *Ecology* 88, 2576-2586.

- 496 [29] Medvinsky, A.B., Petrovskii, S.V., Tikhonova, I.A., Malchow, H., Li, B.L., 2002. Spatiotemporal
497 complexity of plankton and fish dynamics. *SIAM Rev.* 44, 311-370.
- 498 [30] Morozov, A., Petrovskii, S., 2009. Excitable population dynamics, biological control failure, and
499 spatiotemporal pattern formation in a model ecosystem. *Bulletin of Mathematical Biology* 71,
500 863-887.
- 501 [31] Murray, J.D., 1989. *Mathematical Biology*. New York: Springer.
- 502 [32] Murray, J.D., 2003. *Mathematical Biology*, Springer-Verlag, 3rd edition.
- 503 [33] Nicholson, A.J., and Bailey, V.A., 1935. The balance of animal population. *Proceedings of the*
504 *zoological society of London* 1, 551-598.
- 505 [34] Okubo, A., and Levin, S.A., 2001. *Diffusion and Ecological Problems: modern Perspectives* (2nd
506 edition). New York, Springer-Verlag.
- 507 [35] Petrovskii, S.V., Li, B.L., Malchow, H., 2004. Transition to spatiotemporal chaos can resolve the
508 paradox of enrichment. *Ecol. Complexity.* 1, 37-47.
- 509 [36] Petrovskii, S.V., Malchow, H., 1999. A minimal model for pattern formation in a prey-predator
510 system. *Math. Comput. Model.* 29, 49-63.
- 511 [37] Polis, G., McCormick, S. 1987. Intraguild predation and competition among desert scorpions.
512 *Ecology* 68, 332-343.
- 513 [38] Polis, G.A., Meyers, C.A., Holt, R.D., 1989. The ecology and evolution of intraguild predation:
514 potential competitors that eat each other. *Annual Review of Ecology and Systematics* 20, 297-
515 330.
- 516 [39] Rosenheim, J.A., Kaya, H.K., Ehler, L.E., Marois, J.J., Jaffee, B.A., 1995. Intraguild predation
517 among biological control agents: theory and evidence. *Biol. Control* 5, 303-335.
- 518 [40] Rosenzweig, M.L., 1971. Paradox of enrichment: destabilization of exploitation ecosystems in
519 ecological time. *Science* 171, 385-387.
- 520 [41] Rosenzweig, M.L., 1973. Evolution of the predator isocline. *Evolution* 27, 84-94.

- [42] Segel, L.A., and Jackson, J.L., 1972. Dissipative structure: An explanation and an ecological example. *J. Theor. Biol.* 37, 545-559.
- [43] Segel, L.A., Levin, S.A., 1976. Application of nonlinear stability theory to the study of the effects of diffusion on predator-prey interactions. *Selected topics in statistical mechanics and biophysics* (ed. R. Piccirelli) Conference Proceedings No. 27. AIP, New York.
- [44] Seuront, L., 2009. *Fractals and multifractals in ecology and aquatic science*. Chapman & Hall, London.
- [45] Shiyom, M., 1980. Predation-affected spatial pattern changes in a prey population. *Ecol. Modelling.* 11, 1-14.
- [46] Smout, S., Asseburg, C., Matthiopoulos, J., et al., 2010. The Functional Response of a Generalist Predator. *PLoS ONE* 5(5), e10761. doi:10.1371/journal.pone.0010761
- [47] Spencer, P.D., Collie, J.S., 1995. A simple predator-prey model of exploited marine fish populations incorporating alternative prey. *ICES Journal of Marine Science* 53, 615-628.
- [48] Steele, J.H., and Henderson, E.W., 1981. A simple plankton model. *Am. Nat.* 117, 676-691.
- [49] Steele, J.H., Henderson, E.W., 1992. The role of predation in plankton models. *Journal of Plankton Research* 14, 157-172.
- [50] Symondson, W.O.C., Sunderland, K.D., Greenstone, M.H., 2002. Can generalist predators be effective biocontrol agents? *Annu. Rev. Entomol.* 47, 561-594.
- [51] Turchin, P., Hanski, I., 1997. An empirically based model for latitudinal gradient in vole population dynamics. *American Naturalist* 149, 842-874.
- [52] Upadhyay, R.K., Wang, W., Thakur, N.K., 2010. Spatiotemporal dynamics in a spatial Plankton system. *Math. Modelling Nature Phenom* 5, 102-122.
- [53] van Baalen, M., Křivan, V., van Rijn, P.C.J., Sabelis, M.W., 2001. Alternative Food, Switching Predators, and the Persistence of Predator-Prey Systems. *American Naturalist* 157, 512-524.

- 545 [54] van Leeuwen, E., Jansen, V.A.A., Bright, P.W., 2007. How population dynamics shape the
546 functional response in a one-predator-two-prey system. *Ecology* 88, 1571-1581.
- 547 [55] Walde, S., 1994. Immigration and the dynamics of a predator-prey interaction in biological
548 control. *J. Anim. Ecol.* 63, 337-346.
- 549 [56] Wang, W., Zhang, L., Wang, H., Li, Z., 2010. Pattern formation of a predator-prey system with
550 Ivlev-type functional response. *Ecol. Modelling* 221, 131-140.
- 551 [57] Wilson, W.G., Harrison, S.P., Hastings, A., McCann, K., 1999. Exploring stable pattern formation
552 in models of tussock moth populations. *J. Anim. Ecol.* 68, 94-107.
- 553 [58] Winder, L., Alexander, C.J., Holland, J.M. Symondson, W.O.C., Perry, J.N., Woolley, C., 2005.
554 Predatory activity and spatial pattern: the response of generalist carabids to their aphid prey.
555 *Journal of Animal Ecology* 74, 443-454.
- 556 [59] Xu, R., Chen, L., Chaplain, M., 2004. Persistence and global stability in a delayed predator-prey
557 system with Holling-type functional response. *ANZIAM Journal* 46, 121-141.