Bioaugmentation of activated sludge with *Achromobacter denitrificans* PR1 for enhancing the biotransformation of sulfamethoxazole and its human conjugates in real wastewater: Kinetic tests and modelling

Nguyen, P.Y.; Carvalho, G.; Polesel, Fabio; Torresi, Elena; Rodrigues, Alexandre M.; Rodrigues, João E.; Cardoso, Vitor Vale; Benoliel, Maria João; Plósz, Benedek G.; Reis, M.A.M.

Total number of authors:
11

Published in:
Chemical Engineering Journal

Link to article, DOI:
[10.1016/j.cej.2018.07.011](https://doi.org/10.1016/j.cej.2018.07.011)

Publication date:
2018

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Nguyen, P. Y., Carvalho, G., Polesel, F., Torresi, E., Rodrigues, A. M., Rodrigues, J. E., Cardoso, V. V., Benoliel, M. J., Plósz, B. G., Reis, M. A. M., & Oehmen, A. (2018). Bioaugmentation of activated sludge with *Achromobacter denitrificans* PR1 for enhancing the biotransformation of sulfamethoxazole and its human conjugates in real wastewater: Kinetic tests and modelling. *Chemical Engineering Journal*, 352, 79-89. <https://doi.org/10.1016/j.cej.2018.07.011>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Bioaugmentation of activated sludge with *Achromobacter denitrificans* PR1 for enhancing the biotransformation of sulfamethoxazole and its human conjugates in real wastewater: Kinetic tests and modelling

P.Y. Nguyen, G. Carvalho, Fabio Polesel, Elena Torresi, Alexandre M. Rodrigues, João E. Rodrigues, Vitor Vale Cardoso, Maria João Benoliel, Benedek Gy. Plósz, M.A.M. Reis, A. Oehmen

PII: S1385-8947(18)31244-0
DOI: <https://doi.org/10.1016/j.cej.2018.07.011>
Reference: CEJ 19409

To appear in: *Chemical Engineering Journal*

Received Date: 22 April 2018
Revised Date: 28 June 2018
Accepted Date: 1 July 2018

Please cite this article as: P.Y. Nguyen, G. Carvalho, F. Polesel, E. Torresi, A.M. Rodrigues, J.E. Rodrigues, V.V. Cardoso, M.J. Benoliel, B.G. Plósz, M.A.M. Reis, A. Oehmen, Bioaugmentation of activated sludge with *Achromobacter denitrificans* PR1 for enhancing the biotransformation of sulfamethoxazole and its human conjugates in real wastewater: Kinetic tests and modelling, *Chemical Engineering Journal* (2018), doi: <https://doi.org/10.1016/j.cej.2018.07.011>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Bioaugmentation of activated sludge with *Achromobacter denitrificans* PR1 for enhancing the biotransformation of sulfamethoxazole and its human conjugates in real wastewater: Kinetic tests and modelling

P.Y. Nguyen¹, G. Carvalho^{1,Δ}, Fabio Polesel², Elena Torresi^{2,4}, Alexandre M. Rodrigues³, João E. Rodrigues³, Vitor Vale Cardoso³, Maria João Benoliel³, Benedek Gy. Plósz^{2,5}, M.A.M. Reis¹, A. Oehmen^{1,*,∞}

¹ UCIBIO, REQUIMTE, Departamento de Química, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal

² DTU Environment, Technical University of Denmark, Bygningstorvet 115, Kongens Lyngby, Denmark

³ EPAL – Empresa Portuguesa das Águas Livres, S.A., Laboratório de Análises de Água, Avenida de Berlim, 15, 1800-031 Lisboa, Portugal

⁴ Veolia Water Technologies AB, AnoxKaldnes, Klosterängsvägen 11A, SE-226 47 Lund, Sweden

⁵ Department of Chemical Engineering, University of Bath, Claverton Down, Bath BA2 7AY, UK

* Correspondence to:

Email: a.ohmen@uq.edu.au

Phone: +61 7 3365 3729

^Δ Present address: Advanced Water Management Centre, The University of Queensland, Brisbane, QLD, 4072, AUSTRALIA

[∞] Present address: School of Chemical Engineering, The University of Queensland, Brisbane, QLD, 4072, AUSTRALIA

Abstract

Achromobacter denitrificans PR1 has previously shown potential to degrade the antibiotic sulfamethoxazole, whereby sulfamethoxazole biotransformation was stimulated in the presence of biogenic substrates. This study examined the biotransformation kinetics of sulfamethoxazole and its two main conjugates, N₄-acetyl-SMX and SMX-N₁-Glucuronide, by activated sludge and activated sludge bioaugmented with *A. denitrificans* PR1. SMX biotransformation under both anoxic and aerobic conditions was tested, with and without the addition of acetate as growth substrate, to understand the range of applicable conditions for bioaugmentation purposes. Biological process models, such as the pseudo-first order kinetic and cometabolic models, were also applied and, following the estimation of kinetic parameters, could well describe data measured in bioaugmented and non-bioaugmented AS batch experiments under various test conditions. Experimental and modelling results suggest that (i) retransformation of the two conjugates to SMX in AS occurred under both aerobic and anoxic conditions, and (ii) biotransformation kinetics of SMX can vary significantly depending on redox conditions, e.g., SMX was biotransformed by AS only under aerobic conditions. Notably, SMX biotransformation was significantly enhanced when PR1 was bioaugmented in AS. Addition of acetate as biogenic substrate is not necessary, as PR1 was capable of enhancing the SMX biotransformation by using the carbon sources present in wastewater. Overall, bioaugmentation by means of *A. denitrificans* PR1 could be a viable strategy for enhancing SMX removal in AS wastewater treatment plants (WWTPs).

Keywords: antibiotics, cometabolism, N₄-acetyl-SMX, SMX-N₁-Glucuronide, modelling, retransformation

1. Introduction

The intensive use of antibiotics for human and veterinary therapy has led to their continuous discharge, also in the form of conjugates, in the environment. WWTPs are not designed to remove these and other xenobiotic chemicals, and discharge of treated effluents into the environment has been a major concern due to the risk of a worldwide dispersal of antibiotic resistance genes [1].

Amongst antibiotics, sulfamethoxazole (SMX) is one of the most widely used synthetic sulfonamides worldwide. SMX enters WWTPs via human excretion in the forms of unchanged SMX (15-25% of the excreted dose) as well as the conjugated forms N₄-acetyl-SMX (Ac-SMX) (> 40%) and SMX-N₁-Glucuronide (SMX-Glu) (9-15%) [2]. The two human conjugates have been detected in wastewater influent and effluent, and were observed to rapidly deconjugate during wastewater treatment [3,4] which was considered to likely explain the reported 'negative removal' of SMX in wastewater treatment [3,5]. This suggests the importance of investigating the retransformation of the two major human conjugates to parent SMX, in order to explain the reported differences in removal efficiencies in WWTPs [6,7].

SMX removal has been shown to vary greatly, i.e. from negative (-138%) to very high (>90%) [1] in full-scale WWTPs, and with variability in SMX biotransformation kinetics. SMX was also shown to be not readily biodegradable during the 28-day test period in a closed bottle test [8].

Biotransformation has been recognized as the major elimination mechanism of SMX and its conjugates during biological treatment of domestic wastewater, with minor contribution of sorption onto sludge (due to the polar nature of these compounds). Overall, literature reports of inconsistent and incomplete SMX elimination suggest that novel technologies/strategies would be required if more stringent discharge limits for SMX and other antibiotics are enforced. Bioaugmentation can be an alternative WWTP operational strategy to enable or enhance xenobiotics removal by inoculating specialized degrading bacteria [9]. Despite the fact that bioaugmentation has been studied for years in wastewater treatment to reinforce biological processes, few studies have tested the use of bioaugmentation for enhancing the removal of xenobiotics, e.g. 17 β -estradiol [10], estradiol [11], fungicides [12]. With respect to antibiotics, bioaugmentation resulted in

limited SMX removal when applying *Microbacterium* sp. strain BR1 in full-scale membrane bioreactors [13], except for SMX concentrations far higher than the ones normally found in municipal wastewater.

Previously, we showed that a pure culture of *Achromobacter denitrificans* PR1 could exhibit faster biotransformation kinetics (up to two to three orders of magnitude higher) of SMX compared to AS alone [14], even at the low SMX concentrations typical of wastewater effluents. Given its ability to degrade SMX in the presence and/or absence of other additional carbon sources (acetate and succinate) at environmentally relevant concentrations (typical of e.g., wastewater effluents), the strain likely has potential for treating SMX in wastewater upon bioaugmentation. Therefore, the overall objective of this work was to investigate whether PR1 can enhance SMX biotransformation kinetics when bioaugmented to AS with real wastewater feed. Specifically, we (i) investigated the effect of redox conditions, i.e. aerobic and anoxic conditions, on the transformation rates of targeted compounds; (ii) assessed the potential influence of retransformation processes of the two main human conjugates, i.e. Ac-SMX and SMX-Glu, on the fate of sulfamethoxazole under the testing conditions; and (iii) evaluated the need for supplementation with a biogenic substrate (e.g. acetate) or whether the availability of carbon sources in wastewater could serve as biogenic substrates to achieve a sufficiently interesting kinetic for SMX removal upon bioaugmentation of AS with PR1. Modelling the fate of xenobiotics in WWTPs can be a useful tool to understand their removal mechanisms, predict and reduce their emissions with treated effluent through process optimization. Specifically, the Activated Sludge Modelling framework for Xenobiotics (ASM-X), has been previously used to predict the fate of SMX in biological treatment systems [7] and to identify factors (influent concentration of conjugates, solid residence time) possibly explaining the variability in SMX removal efficiencies [15]. In this context, suitable mathematical models were developed to examine the metabolic mechanism and predict kinetics of SMX and human conjugates biotransformation upon bioaugmentation of *A. denitrificans* PR1 into AS.

2. Materials and methods

2.1. Chemicals and reagents

Reagent grade (purity $\geq 99\%$) SMX was purchased from Sigma-Aldrich. Ac-SMX, SMX-Glu and isotopically labelled Ac-SMX-d4, SMX-d4-Glu, SMX-d4 were obtained from Toronto Research Chemicals,

Inc. (TRC, Canada). Individual stock standard solutions were prepared on a weight basis in methanol and stored at -20°C. HPLC-grade methanol was supplied by Merck (Darmstadt, Germany).

2.2. *Laboratory-scale experiments - Bioaugmentation of AS with A. denitrificans PR1*

2.2.1. *Culture media*

Bacterial inoculum was grown in mineral medium B, supplemented with ammonium phosphate at concentration of 400 mM (designated as MMBN), as previously described by [16]. The cells were harvested by centrifugation (7000 x g for 10 min at 20°C using a Sigma® 4-16KS centrifuge), and rinsed three times with fresh MMBN medium to remove the trace amount of SMX remaining from the culture medium before augmenting to the reactors to get an initial cell suspension concentration of approximately 0.05-0.06 mg_{biomass} L⁻¹.

2.2.2. *Batch tests*

Biotransformation of SMX and the two main human conjugates by AS and bioaugmented AS was assessed in a series of batch experiments in 1 L jacketed glass reactors. Dried compressed atmospheric air or pure nitrogen were continuously sparged by a diffuser placed at the bottom of each reactor to create aerobic or anoxic conditions, respectively. Temperature was controlled at 20°C using an external recirculating bath and pH was monitored and maintained between 7.0-7.4 by the addition of HCl (0.2 M) or NaOH (0.2 M), using pH controllers (HI8711, Hanna Instruments, US) with dual set point.

For all experiments, primary effluent wastewater and AS (from a Modified Ludzack Ettinger system) collected from the Chelas WWTP (Lisbon, Portugal) were used. More information about Chelas WWTP is provided in Section S2 and Table S3, Supplementary Information (SI). AS and primary effluent were seeded to the 1 L glass reactors at an initial biomass concentration of approximately 3 gTSS L⁻¹ for all the experiments. Overall, four types of batch tests were performed: (i) abiotic control tests; (ii) sorption tests; (iii) bioaugmentation tests; (iv) nitrification inhibition tests. The testing conditions are presented in Table 1. All the tests were performed in duplicate, except for the anoxic bioaugmented AS test (An2, Table 1), the control 1 and the allylthiourea (ATU) nitrification inhibition tests.

Abiotic control test (control 1)

The goal of this experiment was to determine the contribution of abiotic removal mechanisms (stripping, sorption onto reactor walls and equipment, and abiotic chemical reactions). In this test, the 1 L-glass-reactor was filled with Milli-Q water that was spiked with the three compounds, e.g. SMX, Ac-SMX and SMX-Glu at the concentrations of $10 \mu\text{g L}^{-1}$, $15 \mu\text{g L}^{-1}$ and $15 \mu\text{g L}^{-1}$, respectively. The experiment lasted 6 hours.

Sorption tests (control 2)

Sodium azide (NaN_3) is a well know respiration inhibitor used for negative control tests in AS studies. A wide range of concentrations from 0.5 to $720 \text{ mg}_{\text{azide}} \text{g}_{\text{TSS}}^{-1}$ were used in previous studies for this purpose [17].

In this test, a concentration of $\sim 650 \text{ mg}_{\text{azide}} \text{g}_{\text{TSS}}^{-1}$ was used to inhibit AS activity. SMX, Ac-SMX and SMX-Glu were spiked into the reactors at the initial concentrations of $5 \mu\text{g.L}^{-1}$, $10 \mu\text{g.L}^{-1}$ and $10 \mu\text{g.L}^{-1}$, respectively. The tests were performed in duplicate.

Bioaugmentation tests

The goal of these tests was to assess biotransformation of the targeted compounds with non-bioaugmented and bioaugmented AS with *A. denitrificans* PR1. Batch experiments were performed during 12 to 14 hours under aerobic and anoxic conditions. In aerobic tests, the influence of a biogenic substrate on SMX biotransformation was assessed by adding acetate at an initial concentration of ~ 137 to $152 \text{ mg}_{\text{COD}} \text{L}^{-1}$ that is similar to the level of soluble COD typically found in many activated sludge WWTPs.

In anoxic batch tests, reactors were supplemented with an initial nitrate concentration of $80 \text{ mg NO}_3\text{-N L}^{-1}$ in the form of KNO_3 . Aqueous stock solutions of SMX and the two target conjugates were spiked to obtain an initial concentration of approximately $5 \mu\text{g L}^{-1}$ and $10 \mu\text{g L}^{-1}$, respectively.

Table 1. Overview of the different tested conditions in the bioaugmented and non-bioaugmented AS experiments

		Batch	Feed	AS (gTSS L ⁻¹)	<i>A.denitrificans</i> PR1 (g _{biomass} L ⁻¹)	Acetate (mg _{COD} L ⁻¹)	NaN ₃ (mg _{nitrite} L ⁻¹)	ATU (mg L ⁻¹)
Control	Control 1	C1	MilliQ-water	-	-	-		
	Control 2	C2	WW*	~ 3	-	-	~ 650	
Nitrification-inhibition		ATU	WW*	~ 3	-	-		30
		Without ATU	WW*	~ 3	-	-	-	-
Bioaugmentation tests	Aerobic	A1	WW*	~ 3	0	0		
		A2	WW*	~ 3	0	137-152		
		A3	WW*	~ 3	~ 0.05-0.06	0		
		A4	WW*	~ 3	~ 0.05-0.06	137-152		
	Anoxic	An1	WW*	~ 3	0	0		
		An2	WW*	~ 3	~ 0.05-0.06	0		

WW*: wastewater from the effluent of a primary sedimentation tank was centrifuged at 10000 x g for 15 min at 4°C, and then filtered through Whatman® Glass microfiber filters, pore size 1.2 µm binder free, Grade GF/C before feeding to the reactors.

Nitrification inhibition tests

To determine the contribution of ammonia oxidizing bacteria and heterotrophs to the SMX biotransformation in AS communities, biomass was inactivated using ATU at 30 mg L⁻¹ [18], a copper chelator that depletes copper ions from the active center of ammonia monooxygenases (AMO), therefore inhibiting ammonia oxidizing activity.

2.3. *Sample preparation and analytical procedures*

Samples collected along the tests were centrifuged for 5 min at 8000 xg, followed by syringe filtration through 0.2 µm cellulose Whatman filters and stored at -20°C prior to analysis of soluble chemicals.

The acetate concentrations were determined by high-performance liquid chromatography (HPLC) using an IR detector and a BioRad Aminex HPX-87H column. 0.01 N sulfuric acid was used as eluent, with an elution rate of 0.6 mL/min and a 50°C operating temperature.

Total and volatile suspended solids (TSS, VSS) were determined according to Standard Methods (APHA, 1995). Ammonium, nitrate and nitrite concentrations were measured using a segmented flow analyzer through the Skalar San++ system. Samples were also analyzed for soluble COD (sCOD) using HACH-lange test kits and a DR2800 spectrophotometer (HACH, Germany).

Analysis of SMX and the two human conjugates was performed on a high performance liquid chromatography coupled to tandem mass spectrometry (HPLC-MS/MS) using a Dionex Ultimate 3000 system from Thermo Scientific. Detailed descriptions of the sample preparation and analytical methods used are provided as Section S1 (SI).

2.4. *Determination of kinetic parameters-modelling approach*

2.4.1. *Modelling assumptions*

In this study, we hypothesized that (i) only retransformation of the two conjugates, e.g. Ac-SMXI and SMX-Glu, will occur through deconjugation to form the parent compound SMX and that (ii) the dissolved compounds are the only biodegradable fractions. Thus, the biotransformation of SMX includes two processes: (i) formation of SMX due to the retransformation (deconjugation) of Ac-SMX and SMX-Glu; (ii) simultaneous elimination of SMX.

2.4.2. Model implementation and estimation of parameters

In this study, the biotransformation rate of the three target compounds was calibrated using the ASM-X modelling framework [7,15,19].

Deconjugation of the two human conjugates (Ac-SMX and SMX-Glu) to form the parent compound SMX is described by a pseudo-first order kinetic model (Table 2, process (1) for aerobic and process (9) for anoxic removal), thus allowing the estimation of the biotransformation rate coefficients, e.g. $k_{Dec,Ox}$ or $k_{Dec,Ax}$ ($L\ gTSS^{-1}\ d^{-1}$).

For the biotransformation of SMX under aerobic conditions, both pseudo-first order and cometabolic models were implemented to test which one could appropriately predict SMX biotransformation (Table 2). The cometabolic biotransformation model [19] consisted of two biotransformation rates: the enhanced rate in the presence (q_{bio} , $L\ d^{-1}\ g^{-1}$) and the pseudo-first order rate in the absence (k_{bio} , $L\ d^{-1}\ g^{-1}$) of growth substrates. Accordingly, biotransformation kinetics of the cometabolic substrate (e.g. micropollutants) depend on the readily biodegradable growth substrates, S_s ($mgCOD\ L^{-1}$). S_s was determined as the difference between soluble COD (sCOD, measured during the experiments) and soluble inert COD (S_I – calculated according to [20]). The initial S_s concentration of the pre-clarified municipal wastewater used in this study ranged between 41 and 128 $mgCOD\ L^{-1}$. Parameters that could not be identified through model calibration to experimental results (i.e. heterotrophic yields Y_H , substrate affinity constant K_S) were adopted from literature [21]. Concentration profiles of acetate, expressed as sCOD, were used to calibrate the maximum specific growth rate of heterotrophs μ_H (Table S5 and Fig. S1, SI). The estimated parameters included: (i) biotransformation rate constants of the AS ($k_{bio,AS}$) and the bioaugmented strain ($k_{bio,PR1}$) in the absence of primary substrate; and (ii) the cometabolic biotransformation rate constants of the AS ($q_{bio,AS}$) and the bioaugmented strain ($q_{bio,PR1}$) in the presence of the primary substrates. Each batch test was designed to determine a specific kinetic constant and is described in Table 3. The model was implemented in Aquasim 2.1d [22] and the embedded secant method was used for parameter estimation.

In our study, experimental data from A1, A4, An1 and An2 tests were used for the model calibration and estimation of the biotransformation rate constants of SMX and the two human conjugates by AS and *A. denitrificans* PR1 under aerobic and anoxic conditions (Table 3). More details on the model calibration

procedure are presented in Supplementary Information (section S3).

2.4.3. *Model validation*

Two different sets of experimental results (A2 and A3) were used to validate the cometabolic kinetic models calibrated with the data sets of A1 and A4.

ACCEPTED MANUSCRIPT

Table 2. Stoichiometric (Gujer) matrix of the ASM-X for processes of parent compound retransformation, biotransformation and the alternative cometabolic biotransformation model. Parameters and state variables for determination of micropollutants kinetics are described in the main text.

Processes i → j process ↓	C_{LI}	C_{CJ}	S_s	X_{AS}	X_{PRI}^*	Process rate
Pseudo-first order kinetics – Aerobic processes						
(1) Parent compound formation due to retransformation of human	F	-1		*		$\frac{k_{Dec} C_{CJ} X_{AS}}{1 + K_D X_{AS}}$
(2) Pseudo-first order kinetics – biotransformation transformation of parent compound C_{LI} by AS	-1			*		$\frac{k_{bio_AS} C_{LI} X_{AS}}{1 + K_D X_{AS}}$
(3) Pseudo-first order kinetics – biotransformation of parent compound C_{LI} by the bioaugmentation strain, i.e. A. denitrificans PRI	-1				*	$\frac{k_{bio_PRI} C_{LI} X_{PRI}}{1 + K_D X_{PRI}}$
Cometabolic model – Aerobic processes						
4) Cometabolic biotransformation of C_{LI} by AS	-1			*		$\frac{(q_{bio_AS} \frac{S_s}{S_s + K_S} + k_{bio_AS}) C_{LI} X_{AS}}{1 + K_D X_{AS}}$
(5) Cometabolic enhancement biotransformation of C_{LI} by the bioaugmentation strain, i.e. A. denitrificans strain PRI					*	$\frac{(q_{bio_PRI} \frac{S_s}{S_s + K_S} + k_{bio_PRI}) C_{LI} X_{PRI}}{1 + K_D X_{AS}}$
(6) Aerobic growth				*	*	$\mu_H \frac{S_s}{S_s + K_S} X_H$

Pseudo first order kinetics – Anoxic processes

(7) Parent compound formation due to retransformation of human conjugates C_{CJ}	F	-1	*	$\frac{k_{Dec} C_{CJ} X_{AS}}{1 + K_D X_{AS}}$
(8) Biotransformation of parent compound C_{LI} by AS	-1		*	$\frac{k_{bio_AS} C_{LI} X_{AS}}{1 + K_D X_{AS}}$
(9) Biotransformation of parent compound C_{LI} by the bio-augmentation strain, i.e. <i>A. denitrificans</i> PR1	-1		*	$\frac{k_{bio_PR1} C_{LI} X_{PR1}}{1 + K_D X_{PR1}}$

*Due to short duration of the batch experiment and low S/X ratio, negligible biomass growth was assumed.

F = ratio between molecular mass of parent compound and metabolite undergoing deconjugation.

S_S : primary substrate concentration (e.g., organic matter or acetate in some of these experiments, expressed as readily soluble biodegradable COD) considering a co-substrate (gCOD L^{-1}).

C_{LI} and C_{CJ} : the aqueous concentrations of the parent compound and the human conjugates undergoing deconjugation to the parent compound, respectively ($\mu\text{g L}^{-1}$).

k_{Dec} : retransformation rate constant of deconjugation of the human conjugates to parent compound ($\text{L gTSS}^{-1} \text{d}^{-1}$).

k_{bio_AS} : is the reaction rate coefficient of biotransformation of parent compound ($\text{L gTSS}^{-1} \text{d}^{-1}$) by AS.

k_{bio_PR1} : is the reaction rate coefficient of biotransformation of parent compound ($\text{L gTSS}^{-1} \text{d}^{-1}$) by the bioaugmented *A. denitrificans* strain PR1.

K_S : half-saturation coefficient for S_S

K_D : sorption coefficient ($0.256 \text{ L gbiomass}^{-1}$ for SMX - [23]). The values are not available for N_4 -acetyl-SMX and SMX- N_1 -Glucuronide, and were therefore assumed to be equal to 0.

X_{PR1} or X_{AS} : biomass concentration of bio-augmented strain *A. denitrificans* or AS, expressed in gTSS L⁻¹;

X_H is expressed in gCOD L⁻¹ by assuming biomass-to-COD ratio of 0.75

Table 3. Model calibration and parameter estimation procedures for the batch tests performed under aerobic and anoxic conditions

Batch	Goal	Process used	Input parameters	Estimated parameters
A1 (non-bioaugmented)	Model calibration	Process (1) Process (2) Process (1), (4) and (6)	$k_{dec_N4_Ox}$, $k_{dec_Glu_Ox}$, $k_{bio_AS_Ox}$	$k_{dec_N4_Ox}$, $k_{dec_Glu_Ox}$ $k_{bio_AS_Ox}$ $Q_{bio_AS_Ox}$
A4 (bioaugmented)	Model calibration	Process (3) Process (1), (4), (5) and (6)	$k_{dec_N4_Ox}$, $k_{dec_Glu_Ox}$, $k_{bio_AS_Ox}$, $Q_{bio_AS_Ox}$, $k_{bio_PR1_Ox}$	$k_{bio_PR1_Ox}^{**}$ $Q_{bio_PR1_Ox}$
A2 (non-bioaugmented)	Model validation	Process (1), (4) and (6)	$k_{dec_N4_Ox}$, $k_{dec_Glu_Ox}$, $k_{bio_AS_Ox}$, $Q_{bio_AS_Ox}$	None
A3 (bioaugmented)	Model validation	Process (1), (4), (5) and (6)	$k_{dec_N4_Ox}$, $k_{dec_Glu_Ox}$, $k_{bio_AS_Ox}$, $Q_{bio_AS_Ox}$, $k_{bio_PR1_Ox}$, $Q_{bio_PR1_Ox}$	None
An1 (non-bioaugmented)	Model calibration	Process (7) Process (8)	$k_{dec_N4_Ax}$, $k_{dec_Glu_Ax}$	$k_{dec_N4_Ax}$, $k_{dec_Glu_Ax}$ $k_{bio_AS_Ax}$
An2 (bioaugmented)	Model calibration	Process (7), (8) and (9)	$k_{dec_N4_Ax}$, $k_{dec_Glu_Ax}$, $k_{bio_AS_Ax}$	$k_{bio_PR1_Ax}$

** $k_{\text{bio_PR1_Ox}}$ was determined in our previous study [14], from the test with the pure culture (i.e. *A.denitrificans* PR1 biodegradation test) conducted in mineral medium supplemented with SMX as the only substrate.

3. Results and discussions

3.1. Abiotic and sorption processes

Fig. 1 shows the evolution of SMX, Ac-SMX and SMX-Glu concentrations over the test period of 6 h in control test 1, revealing 1.8%, 11.4% and 11.8% removal for Ac-SMX, SMX-Glu and SMX, respectively. This suggests that abiotic processes had minor contribution to the removal of the tested compounds, in agreement with previous studies [24].

To investigate sorption to AS, sodium azide (NaN_3) was used to inhibit the aerobic respiration and suppress the microbial activity of the AS (control 2). The results showed that Ac-SMX and SMX-Glu were transformed concomitantly with an increase in SMX concentration (Fig. 1), indicating that the retransformation of the parent SMX from the two human conjugates occurred even with inactivated biomass, likely via extracellular enzymes. This is in agreement with previous studies for other conjugates [25]. In terms of mass balance, supposing that all the human conjugates were converted back to SMX, there was approx. $0.02 \mu\text{mol}$ SMX formed after 4.5 hours, while there was a removal of approx. $0.02 \mu\text{mol}$ of the two human conjugates. Therefore, no SMX removal was observed in the presence of NaN_3 (control 2). Since sodium azide was present at concentrations previously observed to be sufficient to inhibit the fraction of aerobic biomass [26], biotransformation of the two human conjugates could be due to the activity of facultative anaerobic bacteria, which was not sufficiently inhibited by the addition of NaN_3 .

Due to the impossibility of determining the partitioning coefficient for SMX, the sorption fraction was assessed considering the sorption coefficient K_D obtained from previous literature. A K_D value of $0.256 \text{ L gTSS}^{-1}$ [23] was chosen as the tests in this study were performed with fresh AS and real wastewater, which was representative of the real WWTPs where the K_D was obtained. At circumneutral pH typical of activated sludge systems, SMX is predominantly speciated as an anion ($\text{pK}_a = 5.7$). Possibly due to repulsion with negatively charged sludge particles, sorption of SMX has been generally found to be limited ($K_d < 0.4 \text{ L g}^{-1}$) but not negligible. Notably, the K_d value used is in agreement with other determinations in activated sludge (see, e.g. [7], [27], [23]). Sorption of the two

human conjugates onto AS was not considered in these experiments as no reference values of sorption coefficient were reported. Indeed, the pH of the mixed liquor in the tests was between 7.0-7.4, which is well above the pK_a of SMX-Glu and Ac-SMX ($pK_{a2} = 2.7$ and 5.6, respectively). Under these experimental conditions, Ac-SMX and SMX-Glu exist predominantly or completely as negatively charged species in the aqueous phase. Hence, negligible sorption of Ac-SMX and SMX-Glu was assumed due to their high solubility and polar nature.

3.2. Biotransformation of SMX, SMX-Glu and Ac-SMX in bioaugmented and non-bioaugmented AS tests

Aerobic batch experiments (A1-A4)

In general, biotransformation of the two human conjugates was almost complete for all the batch tests performed under aerobic conditions. This is in agreement with previous studies, showing >85% removal of the two conjugates in laboratory-scale [3] and full-scale [4] AS processes, or even fully eliminated in a pilot membrane bioreactor [28]. In the aerobic non-bioaugmented reactors (A1, A2), an increase in SMX concentration was observed in the first 4–6 hours followed by a slow decrease during the rest of the tests when biotransformation of the two human conjugates was complete (Fig. 2a and 2b). Negative SMX removal was observed, i.e. -43.1% and -63.8%, for the AS in tests A1 and A2, respectively. The decrease in Ac-SMX and SMX-Glu concentrations corresponded to increases in SMX concentrations (Fig. 2a, 2b), strongly suggesting that the two human conjugates deconjugated rapidly to form the parent compound SMX under aerobic conditions. There is relatively limited knowledge on the environmental fate and behavior of the conjugated pharmaceuticals, but these conjugates can undergo deconjugation reactions where deconjugation enzymes are present, with cleavage of the conjugated moiety, resulting in the formation of the parent pharmaceuticals [15,29].

For bioaugmentation of AS with PR1, experimental results obtained in the two batches A3 and A4, show a comparably high rate of SMX biotransformation in the first 4 hours, followed by a relatively lower SMX removal rate after the growth substrates were completed (Fig. 2c and 2d). Similar SMX

removal was observed in A3 and A4 after 12 hours, i.e. $92.5 \pm 1.0\%$ and $89.4 \pm 1.5\%$, respectively. Concomitantly, complete removal of the SMX-Glu and Ac-SMX in the first 4 hours of the test was observed, which supposedly was converted back to SMX (Fig. 2c and 2d) as suggested by other studies [3,15].

Relative contribution of heterotrophs and ammonia oxidizing bacteria

Batch tests performed in the presence of ATU showed no removal of ammonium and no formation of nitrate (Fig. 3b), suggesting nitrification was completely suppressed. There was 42 % removal of SMX after 12 hours (Fig. 3a). No appreciable differences were observed in the removal efficiency of SMX with and without ATU (Fig. 3a), suggesting negligible contribution of AOB to the biotransformation of SMX. Also, for all the remaining aerobic batch tests (A1, A2, A3, and A4) performed, no ammonium removal and no nitrate formation could be observed (Fig. S2) during the testing periods, confirming that no nitrifying activity occurred in the tested AS. Even though biotransformation of SMX in AS was previously shown to correlate with both nitrifying activity [30,31] and heterotrophic bacteria [32,33], no appreciable differences were observed in the removal efficiency of SMX with and without ATU (Fig. 3a) in our study, which could be due to the fact that (i) the SMX biotransformation rate by heterotrophic aerobic degradation was reported to be much faster compared to autotrophic nitrification (e.g. $k_{\text{bio, h}} = 0.09 \text{ L/g}_{\text{VSS}} \text{ d}$ vs. $k_{\text{bio, a}} = 0.01 \text{ L/g}_{\text{VSS}} \text{ d}$) [32]; and (ii) the possible higher abundance of heterotrophs compared to nitrifiers. Thus, heterotrophs seem to be the dominant organisms responsible for the biotransformation of SMX in the current study and the cometabolic model applied for all the batch tests could be based on only organic carbon (i.e. readily biodegradable substrates and supplemental acetate) as the primary substrates.

Effect of acetate addition on the biotransformation of SMX

We previously showed that the presence of a biogenic substrate, e.g. acetate or succinate, provided for an 8-fold increase of SMX biotransformation kinetics by PR1 [14]. In the current study, the initial concentration of acetate was approximately 137 to 152 mg COD L⁻¹, supplemented with preclarified wastewater to (i) the bioaugmented reactor (A4) (Fig. S8b) as a biogenic substrate to enhance the

kinetics of SMX biotransformation by PR1, and (ii) the non-bioaugmented reactor (A2) (**Error! Reference source not found.**S8a) as a control for comparison purposes. Upon bioaugmentation of AS with PR1 (reactor A3, Fig. 2c), with only wastewater (no acetate addition), SMX was biotransformed steadily and almost completely without any lag phase. In addition, the profiles of SMX in the tests fed with acetate were comparable to the one in the tests without acetate (**Error! Reference source not found.**S8a (A1 versus A2) and S8b (A3 versus A4)). From these observations, we hypothesized that (i) PR1 could use other available carbon sources present in wastewater as biogenic substrates to enhance the SMX biotransformation kinetics and the addition of acetate is unnecessary for the bioaugmentation with PR1; (ii) there is no enhancement effect due to acetate addition on the biotransformation of SMX by AS when fed with real wastewater. These hypotheses were also justified with the modelling results in section 3.3 and 3.4.

Anoxic experiments

Pure culture biotransformation tests had previously shown that the *A.denitrificans* PR1 is capable of biotransformation of SMX under both aerobic [14] and anoxic conditions (data not shown). Hence the extent of SMX removal under anoxic conditions was also assessed and compared to those obtained in aerobic conditions.

In the two anoxic batch tests with non-bioaugmented (An1) and bioaugmented AS (An2), most of the two human conjugates were removed in the first 6 hours (Fig. 4a and 4b), which is in agreement with Stadler et al. 2015 [3] that observed >90% of Ac-SMX and SMX-Glu removals under anoxic condition.

For the non-bioaugmented reactor An1, the consumption of nitrate (Fig. 4c) revealed denitrifying activity, while no net SMX removal could be observed (Fig. 4a) overall. SMX concentration increased in the first 6 hours simultaneously with the deconjugation of the two human conjugates, and remained constant for the rest of the experiment. This is opposed to what was observed in denitrifying AS [7] and denitrifying MBBR sludge [34,35]. In contrast, in the bioaugmented reactor (An2), a slight decrease in SMX concentration was observed after the retransformations of the two human conjugates

had completed (after 4 hours) (Fig. 4b). This suggests a biotransformation of SMX associated with the activity of the bioaugmented strain PR1, but at a rather slow rate.

3.3. Model-based assessment of biotransformation kinetics

Experimental data obtained in the batches A1, A4 and An1, An2 was used for the estimation of the biotransformation rate constants for AS and the *A. denitrificans* PR1 under aerobic and anoxic conditions. Predicted dissolved concentration profiles of SMX and its two human conjugates during batch experiments are compared with measured data and shown in Fig. 2a, 2c, Fig. 4a and 4b. The estimated parameters are summarized in Table 4. The model predictions were evaluated using the R-squared (R^2) coefficient, shown in Fig. S3-S5, and summarized in Table S4 (SI). Confidence intervals were obtained by the estimation of standard deviations for a level of confidence of 95% (Fig. S3-S5, SI).

Kinetics of deconjugation

Deconjugation kinetics of the two conjugates could be described with pseudo-first order kinetics, in processes for aerobic (1) and anoxic (9) conditions. For the conjugated Ac-SMX, fitting of measured data resulted in k_{Dec} values of $8.9 \pm 0.53 \text{ L gTSS}^{-1} \text{ d}^{-1}$ for aerobic conditions, which was almost 2-fold higher than anoxic conditions ($5.30 \pm 0.21 \text{ L gTSS}^{-1} \text{ d}^{-1}$) (Table 4). These data agree well with values reported in literature for k_{Dec} of $5.9\text{-}7.6 \text{ L gTSS}^{-1} \text{ d}^{-1}$ [7,36] under aerobic conditions or $7.9 \text{ L gTSS}^{-1} \text{ d}^{-1}$ [7] under anoxic conditions. No difference in the rate constants of SMX-Glu under aerobic and anoxic conditions was obtained (4.76 ± 0.4 and $4.74 \pm 0.31 \text{ L gTSS}^{-1} \text{ d}^{-1}$, respectively). No data for the biotransformation rate coefficients of SMX-Glu were available in literature for comparison. Good agreement between experimental data and model simulations for Ac-SMX and SMX-Glu was shown, as confirmed by high R^2 coefficients (≥ 0.98), indicating that pseudo-first order equations describe well the biotransformation kinetics of the two human conjugates. These results suggest that (i) deconjugation rate constants are well above $1 \text{ L gTSS}^{-1} \text{ d}^{-1}$, thus indicating high degradability for conjugates; and (ii) deconjugation kinetics depend on redox conditions for Ac-SMX only, being faster under aerobic conditions.

Kinetics of SMX biotransformation under aerobic conditions

The removal of SMX under aerobic conditions was predicted using different mathematical models for comparison (i) a pseudo-first order kinetic model; and (ii) cometabolic models. Fig. 2 and Fig. S6 (SI) compare predicted and measured concentrations of SMX in the batches A1-A4, using a cometabolic model and a pseudo first order model, respectively. According to the data plotted in these figures, the prediction of SMX biotransformation was significantly improved by adopting the cometabolic model (R^2 ranged from 0.79 to 0.99, Table S4, SI) compared to pseudo-first order biotransformation model (R^2 ranged from 0.044 to 0.94). These results: (i) show that the cometabolic model was able to consistently describe the experimental data, with measured concentrations that fall well within the 95% confidence interval (Fig. S3, S4, SI), making the cometabolic model the relevant choice for description of the removal of SMX; (ii) support our hypothesis of the deconjugation of Ac-SMX and SMX-Glu results in the formation of parent compound SMX, which likely explains the previously observed variability/negative SMX removal efficiencies in biological treatment. In this study, we also simulated two other scenarios for the fate of SMX, i.e.: (i) biotransformation of the two human conjugates leading to the formation of a compound different from SMX (model simulations presented as blue dashed lines in Fig. 2 and 4); and (ii) all of the Ac-SMX and SMX-Glu are converted back to parent SMX, but no SMX is biodegraded, and the model simulation is presented as black dashed lines in Fig. 2 and Fig. 4. However, the model simulations in these scenarios were far different from the respective observed concentrations of SMX in all the batch experiments (Fig. 2 and 4), indicating that neither situation is applicable in the current study.

According to our simulation results, a SMX biotransformation rate constant $k_{\text{bio,AS}}$ of $0.47 \pm 0.03 \text{ L gTSS}^{-1} \text{ d}^{-1}$ and a cometabolic biotransformation rate constant $q_{\text{bio_AS}}$ of $7.97 \pm 0.51 \text{ L gTSS}^{-1} \text{ d}^{-1}$ (Table 4) were obtained by AS under aerobic conditions. Previous studies reported $k_{\text{bio,SMX}}$ of $0.14 - 0.41 \text{ L gTSS}^{-1} \text{ d}^{-1}$ [7,27,32,37,38], in agreement with our $k_{\text{bio,AS}}$ result.

For the bioaugmented AS, test A4 – with acetate supplementation, values of measured SMX concentration data plotted as a function of time elapsed show comparably high biotransformation rate in

the first 4-6 hours (when primary substrate was available), followed by lower removal rate during the remaining time (following primary substrate depletion) (Fig. 2d). By using the obtained parameters $k_{\text{bio,AS}}$, $q_{\text{bio,AS}}$ from the A1 test, $k_{\text{bio,PR1}}$ ($56.20 \pm 3.70 \text{ L gTSS}^{-1} \text{ d}^{-1}$) from our previous study [14], fitting of measured data in the substrate depletion phase using cometabolic model as described in Table 2 resulted in estimation of $q_{\text{bio,PR1}} = 528.39 \pm 6.78 \text{ L gTSS}^{-1} \text{ d}^{-1}$. This $q_{\text{bio,PR1}}$ value is consistent with rate constants ($k_{\text{bio,PR1}} = 445.6 - 570.1 \text{ L gTSS}^{-1} \text{ d}^{-1}$) previously obtained with PR1 when acetate was supplemented as a biogenic substrate to enhance the biotransformation rate of SMX in pure culture biodegradation tests [14]. Higher biotransformation kinetics of SMX by PR1 ($q_{\text{bio,PR1}}$ and $k_{\text{bio,PR1}}$), compared to the retransformation kinetics of the two human conjugates (k_{Dec}), likely lead to the observation of no increase in SMX concentration in bioaugmented AS tests (A3 and A4), differently than what was observed in non-bioaugmented AS tests (A1 and A2).

In general, two different kinetic rates of the removal of SMX are obtained for AS as well as for *A. denitrificans* PR1: a fast rate q_{bio} when primary substrate was available and a slower rate k_{bio} when primary substrate was depleted. These results can likely explain the two patterns of SMX biotransformation observed in bioaugmented batch tests A3 and A4 (Fig. 2c and 2d). As a result of cometabolism, SMX removal was enhanced in the presence of primary substrates (as characterized by q_{bio}), with a subsequent decrease of biotransformation kinetics upon primary substrate limitation (characterized by the k_{bio}) at the end of the A3 and A4 tests (Fig. 2c and 2d).

Also, the significant differences between k_{bio} and q_{bio} imply that growth substrates (readily biodegradable substrates) availability can substantially impact the removal of SMX as a result of cometabolism. In fact, typically present in wastewater at very low concentrations (ng L^{-1} to $\mu\text{g L}^{-1}$), micropollutants are unable to support cell replication and primary substrates (e.g. readily biodegradable carbon sources or ammonium) are essential for biomass growth and to induce enzymes for assimilation or co-factors for biotransformation [39]. As wastewater is a complex medium where not only micropollutants but also organic matter and nutrients are present, which could be degraded simultaneously by AS, cometabolism kinetics could be suitable to predict the behavior of micropollutants in real WWTPs.

In addition, the SMX biotransformation rate constants by *A. denitrificans* PR1, e.g. $q_{\text{bio,PR1}}$ in the presence and $k_{\text{bio,PR1}}$ in the absence of growth substrates, are three and two orders of magnitude higher, respectively, than that estimated for AS, confirming a specialized biotransformation capability by PR1 in comparison to the mixed AS community. Thus, bioaugmentation of AS with PR1 substantially enhanced the biotransformation rate of SMX.

Kinetics of SMX biotransformation under anoxic conditions

SMX retransformation and removal under anoxic conditions can be predicted using pseudo-first order kinetics (processes (7) – (9)), thereby allowing for the estimation of $k_{\text{bio,Ax}}$ ($\text{L gTSS}^{-1} \text{d}^{-1}$). In Fig. 4a-b, simulated and corresponding measured concentrations of the three compounds in An1 and An2 batch experiments are plotted. High R^2 values (≥ 0.98) were obtained, and measured concentrations always fell within the 95% confidence interval (Fig. S5, SI), indicating that the pseudo-first order model was able to predict the fates of SMX and the two human conjugates obtained in the anoxic experiments. The estimated $k_{\text{bio,Ax}}$ for SMX biotransformation are 13.57 ± 2.10 and $0 \text{ L gTSS}^{-1} \text{d}^{-1}$ for *A. denitrificans* PR1 and AS, respectively (Table 4). The latter value is in contrast with other studies. Plósz et al. [7] obtained a SMX biotransformation rate constant of $0.41 \text{ L gTSS}^{-1} \text{d}^{-1}$ under anoxic conditions with AS. In other studies, values of 0.1 and $0.05 \text{ L gTSS}^{-1} \text{d}^{-1}$ were reported for SMX biotransformation rate constants of heterotrophic denitrification and autotrophic denitrification, respectively [32]. Torresi et al. [34] reported a rate constant k_{bio} of $0.1 \pm 0.1 \text{ L gTSS}^{-1} \text{d}^{-1}$ and q_{bio} of 1.7 and 3.2 for SMX biotransformation in a post-denitrification MBBR system dosed with methanol and ethanol, respectively. The obtained results suggest that, upon bioaugmentation to AS, PR1 could also be able to degrade SMX under anoxic conditions but at a significantly lower rate as compared to aerobic conditions – decreasing by 4-fold in terms of the rate constant ($k_{\text{bio,PR1}}$) under anoxic conditions as compared to aerobic conditions (Table 4).

In the current study, we also provided a detailed description of SMX removal in AS processes when assessing the biotransformation of the parent compound and the deconjugation of the two major human conjugates (Ac-SMX and SMX-Glu) back to SMX. SMX formation from the deconjugation of the two

human conjugates was experimentally observed and confirmed by model-based predictions. Ac-SMX and SMX-Glu were detected at levels that are comparable to the SMX concentrations in WWTPs [6,40] (see also Table S3, SI). Significant retransformation of SMX can take place in WWTPs at a higher rate compared to its removal rate (Table 4), resulting in the sometimes negative or varied SMX removal that have been observed in many studies. It implies that deconjugation of human conjugates should be taken into account to thoroughly understand the fate and removal of SMX during wastewater treatment.

In general, the results of these tests highlight the potential application of *A. denitrificans* PR1 for bioaugmentation for SMX removal in WWTPs. One criterion for a successful bioaugmentation is the metabolically active inoculum of a microorganism or consortium. Inability of the inoculated strains to degrade the xenobiotic chemicals once augmented into AS has been reported [41]. One explanation given for such failure in bioaugmentation was the presence of alternative readily biodegradable substrates [41]. In our experiments, enhancement of SMX biotransformation upon bioaugmentation of AS with PR1 was observed. Upon bioaugmentation of AS with *A. denitrificans*, without lag phase, a fast biotransformation of SMX was observed at rates similar to those obtained in pure culture biodegradation tests when acetate was supplemented as biogenic substrate. In addition, the SMX reaction rate constant and cometabolic biotransformation rate of PR1 were about two orders of magnitude higher than the kinetics of AS, regardless of the presence of additional acetate. The fact that the strain was able to use the complex substrates present in real wastewater to stimulate the activity and provide energy for growth and maintenance, suggests that PR1 has a great potential to survive in AS communities upon bioaugmentation. Overall, bioaugmentation with PR1 appears to be a feasible solution for enhancing SMX removal in wastewater, while further studies should focus on long-term biotransformation activity and stability of the bioaugmentation strain in wastewater systems in order to make bioaugmentation applicable.

3.4. Model validation

The models for retransformation of SMX from the two human conjugates and cometabolic biotransformation of SMX were validated using the two sets of experimental results, A2 and A3, for the

non-bioaugmented and bioaugmented cases, respectively (Fig. 2b and 2c). The set of estimated parameter values ($k_{Dec,N4}$, $k_{Dec,Glu}$, $k_{bio,AS}$, $q_{bio,AS}$, $k_{bio,PR1}$, $q_{bio,PR1}$) was used to test the capability of the proposed models to predict the behaviour of the three compounds (SMX, Ac-SMX and SMX-Glu) in the reactors A2 and A3, providing measured data independent from those used for model identification. Measured and predicted concentrations were compared and R^2 was calculated to determine the extent of correlation. Good agreement between the experimental data and model simulations could be observed with high R^2 (≥ 0.95 , Table S4). This indicates the applicability of the model towards the prediction of the fate of SMX and human metabolite biotransformation by both AS and PR1 (Fig. 2b and 2c), even in the presence of an externally dosed carbon source (in this case acetate).

As cometabolic biotransformation depend on the readily biodegradable growth substrates, S_s (mgCOD L^{-1}), the addition of acetate to AS would affect the biotransformation of SMX. From our previous study [14] as well as the modelling result of the bioaugmentation test A4 (section 3.3.2), there is no doubt that acetate is a biogenic substrate to enhance the SMX biotransformation by *A. denitrificans* PR1, i.e. primary substrate for the cometabolism of SMX. For the non-bioaugmented activated sludge (A2), we hypothesized above (section 3.2.1.2) that there is no enhanced effects of acetate on the biotransformation of SMX by AS. To test this hypothesis, for modelling of SMX biotransformation of non-bioaugmented AS (batch A2 – with the supplementation of acetate), we tested, (i) both acetate and other readily biodegradable substrates that are present in wastewater (expressed as sCOD); (ii) only readily biodegradable substrates that are present in wastewater (expressed as sCOD) were considered as the primary substrates (S_s) in the cometabolic model to enhance the SMX biotransformation by AS. However, only the latter option gave good fitting between measured and model-based prediction (Fig. S7, SI vs. Fig. 2b), suggesting that the readily biodegradable substrates that are present in wastewater (expressed as sCOD) acted and were sufficient as primary substrates for the cometabolism of SMX by AS. Acetate was measured in the wastewater and it was typically below 7 mg/L, therefore the microorganisms were probably not particularly adapted to it. Müller et al. [33] observed that SMX cometabolism with acetate by AS occurred only after a sufficient adaptation time, meaning that the supplementation of additional acetate might have still enhanced the SMX further if

sufficient adaptation time was allowed, although little is known about which easily biodegradable compounds are used as primary substrates by the AS community. Although the R^2 calculated for the SMX in the A2 test (Fig. 2b) was equal to 0.79, the difference between measured and predicted SMX concentrations were still within the standard deviations of the measured concentrations and falls in between confidence interval boundaries (Fig. S3), making the cometabolic model term still relevant.

Model calibration and validation results revealed that the applied models could predict accurately the fate of SMX, Ac-SMX and SMX-Glu. Kinetic parameter values describing the biotransformation of SMX, Ac-SMX and SMX-Glu under aerobic and anoxic conditions could therefore be implemented in AS models linking organic carbon removal (heterotrophic activity) and xenobiotic biotransformation to predict the fate of SMX in WWTPs. Overall, modelling is fundamental to understand the kinetics and the contribution of different members in bioaugmented AS communities with respect to xenobiotic biodegradation. Thus, combining modeling and experimental data offers the opportunity for a thorough understanding of elimination mechanisms of micropollutants in WWTPs to facilitate optimization of wastewater treatment processes and reduce emissions of xenobiotics.

4. Conclusions

In this study, six different batch tests with non-bioaugmented and bio-augmented (with *A. denitrificans* PR1) AS, operated under different redox conditions and primary substrate addition levels, were used for in-depth assessment of SMX removal in combination with its two main human conjugates (Ac-SMX and SMX-Glu). The extent of SMX removal varied depending on the experimental conditions, and process models were used for interpretation of experimental results and determination of biotransformation kinetics. The results from the experimental and model-based predictions show that the conversion of Ac-SMX and SMX-Glu back to parent SMX was confirmed in activated sludge, whereby deconjugation in non-bioaugmented AS was significantly faster than SMX biotransformation. This likely explains the previously observed variability of SMX removal efficiencies, including net formation of SMX, in full-scale biological WWTPs. The biotransformation of SMX and the deconjugation of Ac-SMX clearly depend on the redox conditions, with the highest removal occurring

under aerobic conditions. Deconjugation of SMX-Glu was however independent of redox conditions. Also, cometabolic models were successfully used to predict the biological transformation kinetics of SMX in both bioaugmented and non-bioaugmented reactors under different test conditions. Estimation of kinetic parameters allowed for the assessment of the success of the bioaugmentation strategy and for the identification of best conditions for its applicability. Furthermore, estimated kinetic parameters obtained from this study could be integrated in AS models to predict the fate of SMX during the biological treatment in WWTPs. Overall, bioaugmentation of AS with *A. denitrificans* strain PR1 led to enhanced (100-fold) biotransformation kinetics of SMX compared to the non-bioaugmented AS, within a complex carbon environment found at a WWTP without an addition of another C-source (acetate) as specific substrate for the biotransformation of SMX. These results prospect the use of *A. denitrificans* PR1 for bioaugmentation as a feasible and efficient option to improve SMX elimination in WWTPs.

5. Acknowledgement

This research was supported by MERMAID, an Initial Training Network funded by the People Programme (Marie Skłodowska-Curie Actions) of the European Union's Seventh Framework Programme FP7/2007-2013/ under REA grant agreement n°607492, and by UCIBIO (FCT/MEC UID/Multi/04378/2013, POCI-01-0145-FEDER-007728). P.Y. Nguyen wishes to acknowledge and thank EPAL Water Laboratory for allowing her to perform all the analyses with LC-MS/MS.

References

- [1] I. Michael, L. Rizzo, C.S. McArdell, C.M. Manaia, C. Merlin, T. Schwartz, C. Dagot, D. Fatta-Kassinos, Urban wastewater treatment plants as hotspots for the release of antibiotics in the environment: A review, *Water Res.* 47 (2013) 957–995. doi:10.1016/j.watres.2012.11.027.
- [2] A. Van der Ven, M. Mantel, T. Vree, P. Koopmans, J. Van der Meer, Formation and elimination of sulphamethoxazole hydroxylamine after oral administration of sulphamethoxazole., *Br. J. Clin. Pharmacol.* 38 (1994) 147–150. doi:10.1111/j.1365-2125.1994.tb04339.x.

- [3] L.B. Stadler, L. Su, C.J. Moline, A.S. Ernstoff, D.S. Aga, N.G. Love, Effect of redox conditions on pharmaceutical loss during biological wastewater treatment using sequencing batch reactors, *J. Hazard. Mater.* 282 (2015) 106–115. doi:10.1016/j.jhazmat.2014.08.002.
- [4] A. Göbel, C.S. McArdell, A. Joss, H. Siegrist, W. Giger, Fate of sulfonamides, macrolides, and trimethoprim in different wastewater treatment technologies, *Sci. Total Environ.* 372 (2007) 361–371. doi:http://dx.doi.org/10.1016/j.scitotenv.2006.07.039.
- [5] B.G. Plósz, H. Leknes, H. Lilved, K. V Thomas, Diurnal variations in the occurrence and the fate of hormones and antibiotics in activated sludge wastewater treatment in Oslo, Norway, *Sci. Tot. Environ.* 408 (2010) 1915. doi:10.1016/j.scitotenv.2010.01.042.
- [6] A. Joss, E. Keller, A.C. Alder, A. Göbel, C.S. McArdell, T. Ternes, H. Siegrist, Removal of pharmaceuticals and fragrances in biological wastewater treatment, *Water Res.* 39 (2005) 3139–3152. doi:10.1016/j.watres.2005.05.031.
- [7] B.G. Plósz, H. Leknes, K. V. Thomas, Impacts of competitive inhibition, parent compound formation and partitioning behavior on the removal of antibiotics in municipal wastewater treatment, *Environ. Sci. Technol.* 44 (2010) 734–742. doi:10.1021/es902264w.
- [8] R. Alexy, T. Kümpel, K. Kümmerer, Assessment of degradation of 18 antibiotics in the Closed Bottle Test, *Chemosphere.* 57 (2004) 505–512. doi:https://doi.org/10.1016/j.chemosphere.2004.06.024.
- [9] H. Van Limbergen, E.M. Top, W. Verstraete, Bioaugmentation in activated sludge: current features and future perspectives, *Appl. Microbiol. Biotechnol.* 50 (1998) 16–23. doi:https://doi.org/10.1007/s002530051250.
- [10] H. Roh, K.H. Chu, Effects of solids retention time on the performance of bioreactors bioaugmented with a 17 β -estradiol-utilizing bacterium, *Sphingomonas* strain KC8, *Chemosphere.* 84 (2011) 227–233. doi:10.1016/j.chemosphere.2011.04.029.
- [11] L. Iasur-Kruh, Y. Hadar, D. Minz, Isolation and Bioaugmentation of an Estradiol-Degrading

- Bacterium and Its Integration into a Mature Biofilm, *Appl. Environ. Microbiol.* 77 (2011) 3734–3740. doi:10.1128/aem.00691-11.
- [12] H. Wu, J. Shen, X. Jiang, X. Liu, X. Sun, J. Li, W. Han, L. Wang, Bioaugmentation strategy for the treatment of fungicide wastewater by two triazole-degrading strains, *Chem. Eng. J.* 349 (2018) 17–24. doi:10.1016/J.CEJ.2018.05.066.
- [13] A. Fenu, B.M.R. Donckels, T. Beffa, C. Bemfohr, M. Weemaes, Evaluating the application of *Microbacterium* sp. strain BR1 for the removal of sulfamethoxazole in full-scale membrane bioreactors, *Water Sci. Technol.* 72 (2015) 1754–1761. doi:10.2166/wst.2015.397.
- [14] P.Y. Nguyen, G. Carvalho, A.C. Reis, O.C. Nunes, M.A.M. Reis, A. Oehmen, Impact of biogenic substrates on sulfamethoxazole biodegradation kinetics by *Achromobacter denitrificans* strain PR1, *Biodegradation*. (2017) 1–13. doi:10.1007/s10532-017-9789-6.
- [15] F. Polesel, H.R. Andersen, S. Trapp, B.G. Plósz, Removal of Antibiotics in Biological Wastewater Treatment Systems—A Critical Assessment Using the Activated Sludge Modeling Framework for Xenobiotics (ASM-X), *Environ. Sci. Technol.* 50 (2016) 10316–10334. doi:10.1021/acs.est.6b01899.
- [16] P.J.M. Reis, A.C. Reis, B. Ricken, B.A. Kolvenbach, C.M. Manaia, P.F.X. Corvini, O.C. Nunes, Biodegradation of sulfamethoxazole and other sulfonamides by *Achromobacter denitrificans* PR1, *J. Hazard. Mater.* 280 (2014) 741–749. doi:http://dx.doi.org/10.1016/j.jhazmat.2014.08.039.
- [17] P. Hamon, M. Villain, B. Marrot, Determination of sorption properties of micropollutants: What is the most suitable activated sludge inhibition technique to preserve the biomass structure?, *Chem. Eng. J.* 242 (2014) 260–268. doi:10.1016/J.CEJ.2013.07.117.
- [18] J. Park, N. Yamashita, G. Wu, H. Tanaka, Removal of pharmaceuticals and personal care products by ammonia oxidizing bacteria acclimated in a membrane bioreactor: Contributions of cometabolism and endogenous respiration, *Sci. Total Environ.* 605–606 (2017) 18–25.

doi:<https://doi.org/10.1016/j.scitotenv.2017.06.155>.

- [19] B.G. Plósz, K.H. Langford, K. V. Thomas, An activated sludge modeling framework for xenobiotic trace chemicals (ASM-X): Assessment of diclofenac and carbamazepine, *Biotechnol. Bioeng.* 109 (2012) 2757–2769. doi:10.1002/bit.24553.
- [20] P.J. Roeleveld, M.C.M. Van Loosdrecht, Experience with guidelines for wastewater characterisation in The Netherlands, *Water Sci. Technol.* 45 (2002) 77–87.
- [21] M. Henze, W. Gujer, T. Mino, M.C.M. van Loosdrecht, Activated Sludge Models ASM1, ASM2, ASM2d and ASM3, IWA Publ. (2000) 121. doi:10.1007/s13398-014-0173-7.2.
- [22] P. Reichert, Aquasim – A tool for simulation and data analysis of aquatic systems, *Water Sci. Technol.* 30 (1994) 21–30. <http://wst.iwaponline.com/content/ppiawast/30/2/21.full.pdf>.
- [23] A. Göbel, A. Thomsen, C.S. McArdell, A. Joss, W. Giger, Occurrence and sorption behavior of sulfonamides, macrolides, and trimethoprim in activated sludge treatment, *Environ. Sci. Technol.* 39 (2005) 3981–3989. doi:10.1021/es048550a.
- [24] B. Li, T. Zhang, Biodegradation and Adsorption of Antibiotics in the Activated Sludge Process, *Environ. Sci. Technol.* 44 (2010) 3468–3473. doi:10.1021/es903490h.
- [25] P. Ramin, A. Libonati Brock, F. Polesel, A. Causanilles, E. Emke, P. de Voogt, B.G. Plósz, Transformation and Sorption of Illicit Drug Biomarkers in Sewer Systems: Understanding the Role of Suspended Solids in Raw Wastewater, *Environ. Sci. Technol.* (2016) acs.est.6b03049. doi:10.1021/acs.est.6b03049.
- [26] É. Barbot, I. Seyssiecq, N. Roche, B. Marrot, Inhibition of activated sludge respiration by sodium azide addition: Effect on rheology and oxygen transfer, *Chem. Eng. J.* 163 (2010) 230–235. doi:10.1016/j.cej.2010.07.050.
- [27] C. Abegglen, A. Joss, C.S. McArdell, G. Fink, M.P. Schlüsener, T.A. Ternes, H. Siegrist, The fate of selected micropollutants in a single-house MBR, *Water Res.* 43 (2009) 2036–2046. doi:10.1016/j.watres.2009.02.005.

- [28] J. Mamo, M.J. García-Galán, M. Stefani, S. Rodríguez-Mozaz, D. Barceló, H. Monclús, I. Rodríguez-Roda, J. Comas, Fate of pharmaceuticals and their transformation products in integrated membrane systems for wastewater reclamation, *Chem. Eng. J.* 331 (2018) 450–461. doi:10.1016/J.CEJ.2017.08.050.
- [29] V. Kumar, A.C. Johnson, N. Nakada, N. Yamashita, H. Tanaka, De-conjugation behavior of conjugated estrogens in the raw sewage, activated sludge and river water, *J. Hazard. Mater.* 227–228 (2012) 49–54. doi:10.1016/J.JHAZMAT.2012.04.078.
- [30] E. Kassotaki, G. Buttiglieri, L. Ferrando-Climent, I. Rodríguez-Roda, M. Pijuan, Enhanced sulfamethoxazole degradation through ammonia oxidizing bacteria co-metabolism and fate of transformation products, *Water Res.* 94 (2016) 111–119. doi:http://dx.doi.org/10.1016/j.watres.2016.02.022.
- [31] E. Torresi, S.J. Fowler, F. Polesel, K. Bester, H.R. Andersen, B.F. Smets, B.G. Plósz, M. Christensson, Biofilm thickness influences biodiversity in nitrifying MBBRs - Implications on micropollutant removal, *Environ. Sci. Technol.* 50 (2016) 9279–9288. doi:10.1021/acs.est.6b02007.
- [32] T. Alvarino, P. Nastold, S. Suarez, F. Omil, P.F.X. Corvini, H. Bouju, Role of biotransformation, sorption and mineralization of ¹⁴C-labelled sulfamethoxazole under different redox conditions, *Sci. Total Environ.* 542, Part (2016) 706–715. doi:https://doi.org/10.1016/j.scitotenv.2015.10.140.
- [33] E. Müller, W. Schüssler, H. Horn, H. Lemmer, Aerobic biodegradation of the sulfonamide antibiotic sulfamethoxazole by activated sludge applied as co-substrate and sole carbon and nitrogen source, *Chemosphere.* 92 (2013) 969–978. doi:10.1016/j.chemosphere.2013.02.070.
- [34] E. Torresi, M. Escolà Casas, F. Polesel, B.G. Plósz, M. Christensson, K. Bester, Impact of external carbon dose on the removal of micropollutants using methanol and ethanol in post-denitrifying Moving Bed Biofilm Reactors, *Water Res.* 108 (2017) 95–105.

doi:<https://doi.org/10.1016/j.watres.2016.10.068>.

- [35] F. Polesel, E. Torresi, L. Loreggian, M.E. Casas, M. Christensson, K. Bester, B.G. Plósz, Removal of pharmaceuticals in pre-denitrifying MBBR – Influence of organic substrate availability in single- and three-stage configurations, *Water Res.* 123 (2017) 408–419. doi:<https://doi.org/10.1016/j.watres.2017.06.068>.
- [36] A. Joss, S. Zabczynski, A. Göbel, B. Hoffmann, D. Löffler, C.S. McArdell, T.A. Ternes, A. Thomsen, H. Siegrist, Biological degradation of pharmaceuticals in municipal wastewater treatment: Proposing a classification scheme, *Water Res.* 40 (2006) 1686–1696. doi:[10.1016/j.watres.2006.02.014](https://doi.org/10.1016/j.watres.2006.02.014).
- [37] S. Suarez, J.M. Lema, F. Omil, Removal of Pharmaceutical and Personal Care Products (PPCPs) under nitrifying and denitrifying conditions, *Water Res.* 44 (2010) 3214–3224. doi:<http://dx.doi.org/10.1016/j.watres.2010.02.040>.
- [38] X. Min, W. Li, Z. Wei, R. Spinney, D.D. Dionysiou, Y. Seo, C.-J. Tang, Q. Li, R. Xiao, Sorption and biodegradation of pharmaceuticals in aerobic activated sludge system: A combined experimental and theoretical mechanistic study, *Chem. Eng. J.* 342 (2018) 211–219. doi:[10.1016/J.CEJ.2018.01.012](https://doi.org/10.1016/J.CEJ.2018.01.012).
- [39] D.J. Arp, C.M. Yeager, M.R. Hyman, Molecular and cellular fundamentals of aerobic cometabolism of trichloroethylene, *Biodegradation.* 12 (2001) 81–103. doi:[10.1023/A:1012089908518](https://doi.org/10.1023/A:1012089908518).
- [40] J. Wang, P.R. Gardinali, Identification of phase II pharmaceutical metabolites in reclaimed water using high resolution benchtop Orbitrap mass spectrometry, *Chemosphere.* 107 (2014) 65–73. doi:[10.1016/j.chemosphere.2014.03.021](https://doi.org/10.1016/j.chemosphere.2014.03.021).
- [41] N.C. McClure, J.C. Fry, A.J. Weightman, Survival and catabolic activity of natural and genetically engineered bacteria in a laboratory-scale activated sludge unit, *Appl. Environ. Microbiol.* 57 (1991) 366–373.

ACCEPTED MANUSCRIPT

Fig. 1. Measured concentrations of SMX, Ac-SMX, and SMX-Glu as a function of time for the control batch tests, i.e. control 1-with Milli-Q water (continuous lines), control 2-with NaN_3 (dashed lines) as an inhibitor. Error bars indicate the standard deviations for duplicates

Fig. 2. Illustration of measured concentrations of SMX, Ac-SMX, and SMX-Glu (markers) and simulated (lines) as a function of time for aerobic batch tests (A1): non-bioaugmented AS test (2a); (A2): non-bioaugmented AS test with supplementation of acetate as additional C-source (2b); (A3): bioaugmented AS with *A. denitrificans* PR1 test (2c); and (A4): bioaugmented AS with *A. denitrificans* PR1 supplemented with acetate test (2d). Orange dashed lines represent the SMX simulation if no SMX formation from the retransformation of Ac-SMX, and SMX-Glu. Black dashed lines represent the SMX simulation when all of Ac-SMX and SMX-Glu are converted back to parent SMX, but no SMX biodegraded. Error bars indicate the standard deviations for duplicates.

Fig. 3. Effect of ATU inhibition on removal of SMX (3a) and nitrogen (3b)

Fig. 4. Illustration of measured concentrations of SMX, N4-acetyl-SMX, and SMX-N1-Glucuronide (markers) and simulated (lines) as a function of time under anoxic conditions (An1): non-bioaugmented AS (4a); (An2): bioaugmented AS with *A. denitrificans* strain PR1 (4b). Orange dashed lines represent the SMX simulation if no SMX is formed from the retransformation of Ac-SMX, and SMX-Glu. Black dashed lines represent the SMX simulation when all of the Ac-SMX and SMX-Glu are converted back to SMX, but no SMX is biodegraded. Nitrate consumption for the batch tests is also shown (4c). Error bars indicate the standard deviations for duplicates.

Table 4. Model parameters and information of estimated kinetics for the biotransformations of SMX and the two human conjugates by activated sludge and *A. denitrificans* PR1 (PR1). Values in brackets indicate literature references.

Symbol	Definition	Unit	Compound					
			SMX-Glu		N4--SMX		SMX	
			AS	PR1	AS	PR1	AS	PR1
Aerobic								
$k_{Dec,Ox}$	Aerobic biotransformation rate coefficient for the human conjugates, C_{CJ}	L g TSS^{-1} d ⁻¹	4.76 ± 0.38 (n.a)	-	8.9 ± 0.53 (5.9-7.6 [36]) (6.8 ^[7])	-		
$k_{bio,Ox}$	Aerobic biotransformation rate coefficient for the parent compound, C_{LI}	L g TSS^{-1} d ⁻¹	-	-	-	-	0.47 ± 0.03 (0.19-0.2[27]) (0.14±0.07[38]) (0.3[37]) (0.41 ^[7])	56.20 ± 3.70

Parameter		Unit	Value	Uncertainty	Reference	Notes
$q_{bio, Ox}$	Aerobic cometabolic-biotransformation rate constant for parent compound, C_{LI}	$L \cdot g \cdot TSS^{-1} \cdot d^{-1}$	-	-	-	$(0.1 \pm 0.1)^{[6]}$ $(0.1 \pm 0.1[34])^j$ 7.97 ± 0.51 $(1.7 \pm 0.2[34])$ 528.39 ± 6.78
	Anoxic	-	-	-	-	-
$k_{Dec, Ax}$	Anoxic biotransformation rate coefficient for the human conjugates, C_{CJ}	$L \cdot g \cdot TSS^{-1} \cdot d^{-1}$	4.74 ± 0.31	-	5.30 ± 0.21	-
			(n.a)		$(7.9^{[13]})$	-
$k_{bio, Ax}$	Anoxic biotransformation rate coefficient for the parent compound, C_{LI}	$L \cdot g \cdot TSS^{-1} \cdot d^{-1}$	-	-	-	0 13.57 ± 2.10
			-	-	-	-

n.a.: not available

Graphical abstract

Highlights

- Activated sludge bioaugmented with *A. denitrificans* PR1 enhanced SMX transformation
- Retransformation of human metabolites occurred under aerobic and anoxic conditions
- *A. denitrificans* PR1 metabolised wastewater carbon sources as co-substrates
- Co-metabolic models described well the biotransformation kinetics