Activity of type i methanotrophs dominates under high methane concentration: Methanotrophic activity in slurry surface crusts as influenced by methane, oxygen, and inorganic nitrogen

Duan, Yun Feng; Reinsch, Sabine; Ambus, Per; Elsgaard, Lars; Petersen, Søren O.

Published in:
Journal of Environmental Quality

Link to article, DOI:
[10.2134/jeq2017.02.0047](https://doi.org/10.2134/jeq2017.02.0047)

Publication date:
2017

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Duan, Y. F., Reinsch, S., Ambus, P., Elsgaard, L., & Petersen, S. O. (2017). Activity of type i methanotrophs dominates under high methane concentration: Methanotrophic activity in slurry surface crusts as influenced by methane, oxygen, and inorganic nitrogen. *Journal of Environmental Quality*, 46(4), 767-775.
<https://doi.org/10.2134/jeq2017.02.0047>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Article (refereed) - postprint

Duan, Yun-Feng; Reinsch, Sabine; Ambus, Per; Elsgaard, Lars; Petersen, Søren O. 2017. **Activity of type I methanotrophs dominates under high methane concentration: methanotrophic activity in slurry surface crusts as influenced by methane, oxygen, and inorganic nitrogen.** *Journal of Environmental Quality*, 46 (4). 767-775. [10.2134/jeq2017.02.0047](https://doi.org/10.2134/jeq2017.02.0047)

© 2017

This version available <http://nora.nerc.ac.uk/517125/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. There may be differences between this and the publisher's version. You are advised to consult the publisher's version if you wish to cite from this article.

The definitive version is available at <https://dl.sciencesocieties.org/>

Contact CEH NORA team at
noraceh@ceh.ac.uk

Methanotrophic activity in slurry surface crusts as influenced by CH₄, O₂, and inorganic N

Yun-Feng Duan ^{1*}, Sabine Reinsch ^{2#}, Per Ambus ^{2§}, Lars Elsgaard ¹, and Søren O. Petersen ¹

¹ Department of Agroecology, Aarhus University, Blichers Allé 20, DK-8830 Tjele, Denmark

² Department of Chemical and Biochemical Engineering, Technical University of Denmark, Frederiksborgvej 399, DK-4000 Roskilde, Denmark

* Corresponding Author: Yun-Feng Duan, Department of Agroecology, Aarhus University, Blichers Allé 20, DK-8830 Tjele, Denmark; E-mail: Kevin.YF.Duan@gmail.com.

Present address: Centre for Ecology and Hydrology, Deiniol Rd, Bangor Gwynedd, LL57 2 UW, UK

§ Present address: Department of Geosciences and Natural Resource Management, University of Copenhagen, Øster Voldgade 10, DK-1350 Copenhagen, Denmark

Abbreviations

MOB: Methane oxidizing bacteria; PLFA: Phospholipid fatty acid; FAME: Fatty acid methyl esters

Keywords

Methane oxidation; Inorganic nitrogen; Microsensor; PLFA stable-isotope probing; Methane oxidizing bacteria.

Core Ideas

- Oxygen penetration into surface crusts is shallow.
- Nitrous oxide accumulates at oxic-anoxic interfaces in surface crusts.
- Oxygen availability is important to high-concentration methane oxidation.
- Microbial methane oxidation is affected by interactions of inorganic N and oxygen.
- Activity of Type I methanotrophs dominates under high methane concentration.

Abstract

Livestock slurry is a major source of atmospheric CH₄, but surface crusts harboring methane oxidizing bacteria (MOB) could mediate against CH₄ emissions. This study examined conditions for methane oxidation by *in-situ* measurements of O₂ and N₂O, as a proxy for inorganic N transformations, in intact crusts using microsensors. This was combined with laboratory incubations of crust material to investigate effects of O₂, CH₄, and inorganic N on methane oxidation, using ¹³CH₄ to trace C incorporation into lipids of MOB. Oxygen penetration into the crust was 2–14 mm, confining the potential for aerobic methane oxidation to a shallow layer. Nitrous oxide accumulated within or below the zone of O₂ depletion. With 10² ppmv CH₄ there was no O₂ limitation on methane oxidation at O₂ concentrations as low as 2%, whereas methane oxidation at 10⁴ ppmv CH₄ was reduced at ≤ 5% O₂. As hypothesized, methane oxidation was in general inhibited by inorganic N, especially NO₂⁻, and there was an interaction between N inhibition and O₂ limitation at 10² ppmv CH₄, as indicated by consistently stronger inhibition of methane oxidation by NH₄⁺ and NO₃⁻ at 3% compared to 20% O₂. Recovery of ¹³C in phospholipid fatty acids suggested that both Type I and Type II MOB were active, with Type I dominating high-concentration methane oxidation. Given the structural heterogeneity of crusts, methane oxidation activity likely varies spatially as constrained by the combined effects of CH₄, O₂, and inorganic N availability in microsites.

Introduction

In regions with intensive livestock production such as Western Europe and North America, up to 40% of livestock CH₄ emissions may be related to manure management (Francesco et al., 2013). In most cases, CH₄ capture and/or biofiltration is neither technically feasible nor economical (Melse and van der Werf, 2005), and more cost-effective alternatives must be considered. When manure is stored as liquid slurry, a dense floating crust is often formed either naturally from dry matter in the slurry, or by facilitation of admixing with chopped straw (Hansen et al., 2009). Studies have demonstrated a potential for aerobic methane oxidation in such surface crusts (Petersen and Ambus, 2006; Petersen et al., 2005), where diverse communities of methane-oxidizing bacteria (MOB) were also documented (Duan et al., 2014). These findings

26 suggest that surface crusts could act as a low-cost filter for manure-derived CH₄, but the physical, chemical,
27 and biological regulation of methane oxidation inside crusts is largely unknown.

28 Methane oxidation potential depends on CH₄ and O₂ availability, which are highly variable due to the
29 heterogeneous structure of the crust. Since surface crusts overlie liquid manure with a high methanogenic
30 potential, they are typically high-CH₄ environments with concentrations far above the atmospheric level. We
31 have observed up to 200 ppmv CH₄ in the stagnant atmosphere immediately above the surface crust (Y.F.
32 Duan, unpublished data), and headspace CH₄ concentrations of 10² and 10⁴ ppmv were used previously to
33 simulate this range of CH₄ availability for MOB in laboratory incubations (Duan et al., 2013). Due to the often
34 loose structure of the crust, sectioning for extraction and determination of *in situ* CH₄ availability by
35 sampling are impractical. While a CH₄ biosensor has been described (Damgaard and Revsbech, 1997), it is
36 not commercially available and has not been tested in heterogeneous environments such as surface crusts.
37 In contrast to CH₄, the distribution of O₂ in surface crusts can be readily determined using a microsensor
38 (Revsbech, 2005), as demonstrated by Hansen et al. (2009) and Nielsen et al. (2010) who investigated O₂
39 penetration into various slurry crusts.

40 Inhibition of methane oxidation by inorganic N is known from many environments (Bosse et al., 1993;
41 Dunfield and Knowles, 1995; Wang and Ineson, 2003), as well as in surface crusts (Duan et al., 2013), but
42 stimulation or no effect can also occur (Liu and Greaver, 2009). Livestock slurry and surface crusts are highly
43 enriched in inorganic N (Table 1): the slurry phase may contain up to 200 mM ammoniacal N (NH₃/NH₄⁺)
44 (Nielsen et al., 2010; Sommer et al., 2007), while up to 35 mM NO₃⁻ (Hansen et al., 2009) and 98 mM NO₂⁻
45 (Nielsen et al., 2010) have been found at the oxic-anoxic interface in surface crusts. These concentration
46 ranges (Table 1) represent a high variability of inorganic N at microsites within surface crusts due to
47 fluctuations in water content as a result of precipitation and insolation (Nielsen et al., 2010). Due to this
48 heterogeneity, bulk N concentrations are insufficient to characterize N distribution and thus potential
49 interference with methane oxidation in microsites. While specialized sensors are available to determine
50 micro-scale concentrations of NH₄⁺, NO₂⁻, and NO₃⁻ in aqueous environments (De Beer et al., 1997; De Beer

et al., 1991; Larsen et al., 1996), they are difficult to apply in unsaturated matrices such as surface crusts, where contact to liquid phase may be intermittent. Instead, the accumulation of N₂O in the crust was used in this study as a proxy for the distribution of N transformations and thus presence of inorganic N (Larsen et al., 1996), as both nitrification and denitrification can produce N₂O under sub-oxic conditions (Bollmann and Conrad, 1998).

The effects of environmental factors on methane oxidation in surface crusts will ultimately depend on the MOB present (Bodelier, 2011; Hu and Lu, 2015). Aerobic MOB have been conventionally categorized into Type I and Type II based on phylogenetic and functional traits, including the presence of signature 16-carbon (C₁₆) or 18-carbon (C₁₈) phospholipid fatty acids (PLFAs) (Hanson and Hanson, 1996). Type I MOB have been reported to thrive in N-sufficient, high-O₂ and low-CH₄ environments, whereas Type II MOB seem to favor the opposite (Amaral et al., 1995; Amaral and Knowles, 1995; Graham et al., 1993). It is still unclear to what extent CH₄, O₂, and inorganic N conditions will select one type of MOB over another in surface crusts, but previous studies found that Type I MOB are more abundant and diverse than Type II MOB in this environment (Duan et al., 2014; Hansen et al., 2009; Nielsen et al., 2013). In recent years, novel methane oxidation pathways, such as NO₂⁻ dependent anaerobic methane oxidation (Ettwig et al., 2010; Welte et al., 2016), as well as aerobic methane oxidation coupled with partial denitrification (Kits et al., 2015a; Kits et al., 2015b), have also been described, but the importance of these processes and the presence of relevant microorganisms in surface crusts remain unclear.

A main objective of this study was to examine the potential for microbial methane oxidation under realistic storage conditions by characterizing *in situ* distributions of O₂ and inorganic N transformations in a cattle slurry surface crust using microsensors. Effects and interactions of O₂ and inorganic N species with respect to methane oxidation could not be quantified *in situ* where microbial activities occur in micro-sites, and instead this was investigated under controlled laboratory conditions. Here, ¹³CH₄ was used as substrate, allowing ¹³C stable isotope probing of PLFAs to study the involvement of Type I and Type II MOB in methane oxidation in surface crusts. Based on previous results (Duan et al., 2014; Duan et al., 2013) we hypothesized that

methane oxidation activity in surface crusts would be determined by both CH₄ and O₂ availability, and inhibited by inorganic N, and that Type I MOB would be primarily responsible for methane oxidation.

Materials and Methods

Microsensor Measurement of O₂ and N₂O Distribution

Dairy cattle slurry was collected from a full-scale manure storage facility in May, 2012, and transferred to two tanks at a pilot-scale storage system (Petersen et al., 2009) at Aarhus University (Foulum, Denmark). The slurry was stored for six weeks prior to the measurement in June, by which time a 5–6 cm thick surface crust with a stable structure had developed on top of the slurry. The development of surface crust reflected typical storage conditions, where a new crust is formed following the mixing of slurry in spring for field application.

Oxygen and N₂O concentration profiles in the surface crust were determined using, respectively, an O₂ and a N₂O microsensor with a tip diameter of 0.5 mm (both produced by Unisense, Aarhus, Denmark). Both microsensors were calibrated according to manufacturer's instructions. Detection limits for O₂ and N₂O were 0.3 μmol L⁻¹ and 0.1 μmol L⁻¹, respectively.

A custom-made mounting system was used to place the microsensors over the surface crust (Supplemental Fig. S1). Oxygen profiles were recorded at 20-cm intervals from 20 to 180 cm along the 200-cm diameter of the storage tank. During measurement, the microsensor was introduced stepwise into the crust at 0.5 mm increments to a maximum depth of 30 mm using a motorized micromanipulator (Unisense). At each depth, the microsensor was stationary for 3 s to allow gas equilibration, and then the O₂ concentration was determined as the average of a 3 s reading. The signal was amplified by a multimeter (Unisense), and registered by the SensorTrace PRO v3.0 software (Unisense). Nitrous oxide profiles were determined at the same locations as O₂ profiles, but with a +2 mm offset to avoid any disturbance to the crust caused by the O₂ microsensor. The N₂O profiles were recorded with the same procedure as O₂ profiles, and initially over the

99 same depth. However, due to incidences of significant N₂O accumulation at 30 mm depth, the maximum
100 depth was extended to 60 mm starting from the 5th profile (at 100 cm distance from the edge).

101 Due to the uneven surface of the crust, a fixed depth cannot accurately define the crust-air interface. For O₂,
102 an abrupt decrease from atmospheric concentration defined the crust-air interface. For N₂O, which was
103 produced inside the crust and diffused towards the atmosphere, the crust-air interface was defined as the
104 depth where N₂O concentration dropped below the detection limit. Using these two criteria, the O₂ and N₂O
105 concentration profiles were aligned.

106 ***Methane Oxidation in Response to O₂ and N Amendments***

107 Surface crust was collected from a full-scale storage tank at the biogas plant of Aarhus University in March,
108 2012. The 10-cm surface crust had developed on slurry co-digested with maize silage. Homogenized crust
109 samples were stored in closed plastic containers at 2°C until used for experiments within four weeks. Duan
110 et al. (2013) showed that MOB can survive and recover activity under these storage conditions for at least
111 three months.

112 Methane oxidation rates were determined by incubating 3-g crust samples in liquid media under a controlled
113 atmosphere. To reduce background N, crusts were washed three times by vortexing with 20 mL deionized
114 water followed by centrifugation at 10,000 × *g* for 10 min. The washing did not remove all NH₄⁺, but the
115 residual NH₄⁺ was negligible compared to the received NH₄⁺ amendment (Duan et al., 2013). Washed crust
116 materials were transferred to 125 mL serum bottles, and resuspended in 20 mL basal salt (BS) medium
117 prepared according to Whittenbury et al. (1970), but excluding N salts. The BS medium was dispensed using
118 a customized system designed to remove dissolved O₂ and maintain anoxia during sample preparation
119 (Supplemental Fig. S2).

120 Various treatments with combinations of different CH₄, O₂, and inorganic N were prepared (Table 2). For N
121 amendments, pre-made solutions of (NH₄)₂SO₄, KNO₃, or KNO₂ were injected into the serum bottles to

122 achieve the desired N concentrations. The bottles were purged using a vacuum pump and refilled with
123 helium, and this step was repeated three times to exhaust residual O₂. Then, air and ¹³C-labeled CH₄ (99
124 atom% ¹³C, ISOTEC, Miamisburg, OH, USA) were injected to achieve the desired headspace gas
125 concentrations. To meet the requirement for CO₂ by some MOB, 4 mL pure CO₂ was also added (Acha et al.,
126 2002). After gas injection, the bottles were mounted on a rotary shaker at 150 rpm for 30 min to allow for
127 liquid-gas equilibration. Headspace O₂ concentrations were verified using an Agilent 3000A MicroGC
128 (Hørsholm, Denmark) as described by (Petersen et al., 2009). For treatments with 0% O₂, residual headspace
129 O₂ concentration was undetectable (< 10–20 ppmv). The bottles were then incubated on a rotary shaker at
130 200 rpm at ca. 21°C; Duan et al. (2013) had shown that there is no gas diffusion limitation under these
131 incubation conditions. Headspace CH₄ concentrations were measured after 0, 2, 4, 6, 8, 24, 48, and 72 h
132 using a Shimadzu 14B GC as described by Duan et al. (2013).

133 In each batch of assays, a control with crust material in N-free BS medium at atmospheric O₂ concentration
134 was included to check for batch-to-batch variations in crust MOB activity, and a blank control without crust
135 material to correct for loss of pressure during repeated gas samplings.

136 First-order rate constants for the first 8 h of incubation and relative activities were calculated according to
137 Duan et al. (2013). First-order rate constants were compared between treatments using R v3.2.2 (R Core
138 Team, 2015). For each CH₄ concentration, effects of O₂ and inorganic N, and their interaction, were analyzed
139 by a two-way ANOVA. Differences between treatments were determined by Duncan's post-hoc multiple
140 comparison test.

141 ***PLFA Extraction and GC-c-IRMS Analysis***

142 After incubation with ¹³C-labeled CH₄, selected crust samples were processed for PLFA analysis (Table 2).
143 Prior to lipid extraction it was necessary to reduce the organic load since otherwise the humic material
144 would bind the chloroform phase and prevent isolation of lipid-soluble compounds. Crust samples were
145 vortexed and centrifuged for 10 min at 3000 × g to extract microbial cells. The supernatant was filtered

through 0.2 µm chloroform-soluble polycarbonate filters, and the material retained on the filter was used for lipid extraction. Hence, lipid results refer to the fraction of extractable low particle size material only. Polar lipid fatty acid methyl esters (FAMES) from each filter were prepared as previously described by Petersen et al. (2002).

FAMES were analyzed using a HP6890 GC (Agilent, Santa Clara, CA, USA) coupled via a GC combustion interface (Thermo Scientific, Bremen, Germany) in continuous flow mode to a Finnigan Delta^{PLUS} isotope ratio mass spectrometer (Thermo Scientific, Bremen, Germany). The oxidation reactor on the interface was maintained at 940 °C, the reduction reactor at 650 °C. Samples (1 µL) were injected at 240 °C in splitless mode. The column temperature was held at 50 °C for 2 min, then increased at 15 °C min⁻¹ to 100 °C, subsequently at 2 °C min⁻¹ to 220 °C, and finally at 15 °C min⁻¹ to 240 °C, where the final temperature was held for 5 min. Separated compounds were measured against a CO₂ reference gas calibrated with reference to Vienna PeeDee belemnite. PLFAs were identified by relative retention time comparing samples against a FAME standard mix (Supelco 37 component FAME mix, 47885-U, Sigma Aldrich). All δ¹³C values were corrected for the methanol C added during methanolysis:

$$\delta^{13}\text{C}_{\text{FAME}} = \frac{(N_{\text{FA}} + 1)\delta^{13}\text{C}_{\text{FAME}} - \delta^{13}\text{C}_{\text{MeOH}}}{N_{\text{FA}}}$$

where N_{FA} refers to the number of carbon atoms of the fatty acid component, $\delta^{13}\text{C}_{\text{FAME}}$ is the observed δ¹³C value of the FAME, and $\delta^{13}\text{C}_{\text{MeOH}}$ is the δ¹³C value of the methanol used for methanolysis (−37.7 ‰ ± 3.2 ‰). The δ¹³C isotope ratios were converted to atom%, and atom% excess was then calculated by subtraction of an unlabeled control. The incorporation of ¹³CH₄ into membrane PLFAs ($n_{13\text{C}}$, nmol) was calculated as $n_{13\text{C}} = (\text{PA}_{\text{FAME}} / \text{PA}_{\text{ISME}}) \times n_{\text{ISME}} \times (\text{atom\% } ^{13}\text{C excess})$, where PA_{FAME} and PA_{ISME} are peak areas of the FAME and internal standard *Me19:0*, respectively, and n_{ISME} (nmol C) is the concentration of the internal standard fatty acid. The lower limit of identified peaks corresponded to 0.1 ng ¹³C g⁻¹ crust material.

168 In view of the uncertain recovery of microbial cells following centrifugation and filtration, statistical testing
169 of treatment effects was not performed, and results will only be presented as means $\pm$ standard errors.

170 **Results**

171 ***Oxygen and N₂O Distribution in Natural Surface Crusts***

172 Surface crusts from the two storage tanks showed a high spatial variability in shape and penetration depth of
173 individual O₂ and N₂O profiles, yet the distribution patterns were qualitatively similar between the two
174 tanks. Thus, results presented here are from one of the storage tanks only (Fig. 1).

175 Oxygen penetration depth varied from 2 to 14 mm, with either a steep or more gradual decline in
176 concentration. Irregularities such as a secondary increase following the initial decline were also observed
177 (e.g., at 60 cm).

178 Nitrous oxide profiles showed peak concentrations at 5–25 mm depth below the crust-air interface. Some
179 N₂O profiles consisted of more than one zone of N₂O accumulation (e.g., at 60, 100, and 160 cm). Maximum
180 N₂O accumulation often coincided with sub-oxic or anoxic zones indicated by O₂ profiles. In a few cases (e.g.,
181 at 60 and 180 cm), N₂O production took place where O₂ availability was relatively high.

182 ***Response of CH₄ Oxidation to O₂ Concentrations***

183 Figure 2 shows the changes in headspace CH₄ concentrations during the 72-h incubation study with different
184 initial O₂ levels and two initial CH₄ levels. For both 10² and 10⁴ ppmv CH₄, consistent methane consumption
185 throughout the incubation was observed only at 20% initial O₂, with a > 90% decline in headspace CH₄
186 concentrations after 72 h. In treatments with $\leq$ 5% initial O₂, CH₄ consumption was generally observed within
187 the first 24 h, followed by net CH₄ accumulation. At 0% initial O₂, net CH₄ production was observed
188 throughout the incubation.

189 The strength of methane oxidation activity was expressed as first-order rate constants during the first 8 h of
190 incubation (Fig. 3). At 10^2 ppmv initial CH_4 there were no significant differences in first-order rate constants
191 at O_2 levels from 20% down to 2%, but at 1% O_2 the rate was significantly impaired. At 10^4 ppmv initial CH_4 ,
192 methane oxidation rates were significantly reduced at lower O_2 levels, though not significantly different
193 between 2% and 1% O_2 .

194 ***Response of CH_4 Oxidation to Interactions between O_2 and Inorganic N***

195 When samples with manipulated O_2 and CH_4 concentrations were amended with inorganic N, complex
196 patterns in microbial methane oxidation were observed. For each of the two initial CH_4 concentrations, two-
197 way ANOVA showed significant effects of O_2 concentrations and N amendments, as well as their interaction,
198 on methane oxidation rates.

199 At 10^2 ppmv CH_4 , first-order rate constants in all N-amended samples were lower than in the N-free control
200 (Table 3). The inhibition was strengthened at increasing concentrations of both NH_4^+ and NO_3^- . Samples
201 treated with NO_3^- consistently showed less inhibition than treatments receiving other N salts, whereas NO_2^-
202 was a potent inhibitor as indicated by similar inhibitions with 1 mM NO_2^- and 50 mM NH_4^+ . At 3% O_2 ,
203 inhibition by individual N species and concentrations was slightly stronger than at 20% O_2 , but the difference
204 was not always statistically significant.

205 At 10^4 ppmv CH_4 the order of inhibition by different N species was similar to that at 10^2 ppmv CH_4 , with NO_2^-
206 as the strongest inhibitor (Table 3). Yet, several NO_3^- treatments caused a weak stimulation rather than
207 inhibition as compared to the N-free control. Generally, the N amendments inhibited methane oxidation at
208 20% O_2 , but not at 3% O_2 , where low rates were already observed in the N-free control. One exception,
209 though, was the inhibition caused by the treatment with 50 mM NH_4^+ .

210 ***¹³C Incorporation into C₁₆ and C₁₈ PLFAs***

211 The yields of PLFA varied considerably between samples, and ¹³C incorporation was below the detection limit
212 for several PLFAs, precluding a detailed quantitative analysis. Instead of absolute PLFA concentrations, an
213 index based on peak area was calculated (Fig. 4, a1–a4). Recovery of ¹³C PLFAs was consistently low in
214 incubations with 10² ppmv CH₄ (Fig. 4, a1). At 10⁴ ppmv CH₄ there was an 8–9 times higher ¹³C recovery in
215 PLFAs at 20% and 3% O₂ than at 1% O₂ (Fig. 4, a2). While NH₄⁺ and NO₃[−] considerably reduced ¹³C recovery,
216 NO₂[−] caused no or only moderate inhibition of ¹³C recovery (Fig. 4, a3 and a4). The total recovery of ¹³C was
217 5–15 times higher with NO₂[−] amendment than with the other two N species.

218 Two ¹³C-labeled PLFA clusters, C₁₆ and C₁₈, were defined in accordance with the predominance of these
219 PLFAs in Type I and Type II MOB, respectively (Bodelier et al., 2009). The C₁₆ cluster included peaks identified
220 as 16:0, 16:1ω6, 16:1ω7, and 16:1ω8, whereas the C₁₈ cluster included 18:0, 18:1ω7, 18:1ω9, and probably
221 also small peaks of 18:1ω8. Also, 16:1ω6 probably co-eluted with 10Me16:0, and 16:1ω8 with i17:0, but this
222 did not influence the calculated ¹³C incorporation for the cluster.

223 Due to the low recovery of total ¹³C PLFA (Fig. 4, a1) and low percentage (3–9%) of ¹³C in C₁₆ and C₁₈ clusters
224 at 10² ppmv CH₄ (Fig. 4, b1), no detailed interpretation of these results was possible. At 10⁴ ppmv CH₄ the
225 recovery of ¹³C in the C₁₆ and C₁₈ clusters together accounted for an average of 52% of ¹³C recovered in
226 PLFAs. A higher percentage of ¹³C was always recovered in the C₁₆ than in the C₁₈ cluster, and the percentage
227 of the C₁₆ cluster was particularly high in treatments with NO₂[−] amendment (Fig. 4, b3 and b4).

228 **Discussion**

229 Due to the heterogenous nature of surface crusts, microbial activities within this environment are controlled
230 by physical and chemical properties of individual microsites rather than overall bulk properties. However,
231 detailed analysis of surface crusts is challenged by the often loose and fibrous structure of the material, and
232 this was also the case with the straw-containing cattle slurry crust used in this study. Therefore, we chose to

233 characterize O₂ and inorganic N distributions *in situ* by microsensors, while regulation of microbial activities
234 were investigated by controlled laboratory incubations. For logistic reasons, the *in situ* gas measurements
235 and laboratory incubations were performed using different surface crusts. However, previous studies have
236 shown that crusts of different origin are qualitatively similar with respect to, e.g., depth of O₂ penetration
237 and the presence of MOB (Duan et al., 2014; Hansen et al., 2009; Nielsen et al., 2010; Nielsen et al., 2013),
238 and results from the two parts are therefore analyzed and discussed together.

239 ***Effects of O₂ Limitation and N Amendments***

240 Methane oxidation kinetics are complex as the reaction involves two substrates, CH₄ and O₂ (Cai and Yan,
241 1999). We were not able to monitor O₂ concentrations during incubation, but instead calculated O₂
242 consumption by MOB based on the amounts of CH₄ consumed, the stoichiometry of methane oxidation
243 (Urmann et al., 2007), and the diffusion coefficients of CH₄ and O₂ in water (Broecker and Peng, 1974)
244 (Supplemental Table S1). These calculations suggested that there was no diffusional limitation of O₂ for
245 methane oxidation, and that the amounts of O₂ used for methane oxidation were < 10% of the available O₂
246 even at 1% initial O₂. Thus, depletion of O₂ during incubation must have been mainly due to aerobic
247 processes other than methane oxidation, i.e., any O₂ limitation for MOB activity reflected competition for O₂
248 against other aerobes. Headspace CH₄ concentrations decreased exponentially during the first 8 h, indicating
249 that the rates of CH₄ uptake depended mainly on CH₄ availability. Therefore, the reaction was approximated
250 by first-order reaction kinetics, which have also been used previously in studies of microbial methane
251 oxidation (De Visscher et al., 1999; King and Schnell, 1994; King and Schnell, 1998; Petersen and Ambus,
252 2006). The apparent first-order kinetics suggest that the rate of CH₄ uptake, and thus demand for O₂, was
253 proportional to the CH₄ concentration, which explains our observation that methane oxidation became more
254 affected by O₂ limitation at 10⁴ ppmv than at 10² ppmv CH₄ (Fig. 3).

255 The neutral to alkaline pH of slurry (Nielsen et al., 2010; Petersen and Ambus, 2006) suggests the presence
256 of free ammonia (NH₃), which is a competitive inhibitor for methane oxidation (Carlsen et al., 1991; Gulledge

257 and Schimel, 1998). In both the present (Table 3) and a previous study (Duan et al., 2013), inhibition by
258 ammonia was less at 10^4 compared to 10^2 ppmv CH_4 , suggesting that competitive inhibition is important for
259 effects of ammonia. The observations that NO_2^- was a more potent inhibitor of methane oxidation than NO_3^-
260 was also consistent with the report by Duan et al. (2013), where the concentration of NO_3^- resulting in 50%
261 inhibition was 100-fold higher than that of NO_2^- . It is likely that these N species both inhibit methane
262 oxidation *via* nitrite toxicity (Stein and Klotz, 2011).

263 We further tested interactions of N inhibition with O_2 availability by comparing relative activities. If there
264 were an interaction between O_2 and inorganic N, a given N amendment would result in different degree of
265 inhibition at 20% and 3% O_2 concentrations. At 10^2 ppmv initial CH_4 , 10 mM, but not 50 mM, NH_4^+ or NO_3^-
266 amendments caused a stronger inhibition at 3% than at 20% O_2 (Table 3), confirming an interaction between
267 N inhibition and O_2 limitation at low N concentrations. At 10^4 ppmv CH_4 , the fact that N amendments
268 generally inhibited methane oxidation at 20% O_2 but not further at 3% O_2 suggested that high-concentration
269 methane oxidation is more sensitive to O_2 limitation rather than to N inhibition.

270 Generally, as an essential substrate for aerobic methane oxidation, O_2 has a direct and immediate effect on
271 aerobic methanotrophic activity. On the other hand, the mechanism of N inhibition on methane oxidation is
272 much more complex and may include immediate toxicity to cell growth and enzyme synthesis, as well as
273 delayed influence on microbial community composition (Bodelier and Laanbroek, 2004). The mechanisms
274 behind interactions between controlling factors in surface crusts could not be explained with the data
275 presented here, and more research is needed to further elucidate this matter.

276 ***^{13}C PLFA Signatures for MOB***

277 Type I and Type II MOB produce unique membrane PLFAs, 16:1 ω 8 and 18:1 ω 8, respectively (Bodelier et al.,
278 2009; Hanson and Hanson, 1996). These signature PLFAs are not always present, or present only in low
279 amounts. However, there is also a general predominance of C_{16} and C_{18} PLFAs among Type I and Type II
280 MOB, respectively (Bodelier et al., 2009), and this has been used to evaluate sources of methane oxidation

281 by stable isotope probing (Qiu et al., 2008). In the present study, both types of MOB were active, with Type I
282 MOB dominating the methanotrophic activity especially where CH₄ availability was high (Fig. 4, b1 and b2).
283 Molecular analyses of microflora in other crust materials likewise suggested that Type I MOB dominated the
284 methanotrophic community in terms of both diversity and abundance (Duan et al., 2014).

285 Interestingly, the incorporation of ¹³C into PLFAs was high in surface crusts incubated with NO₂⁻ as compared
286 to those with other N species (Fig. 4, a3 and a4), especially considering that crusts incubated with NO₃⁻ had
287 over two-fold higher ¹³CH₄ uptake than NO₂⁻ treatments (data not shown). Roslev et al. (1997) reported that
288 NH₄⁺ decreased C conversion efficiency and increased respiration of C assimilated by MOB, which is
289 consistent with the low ¹³C recovery from NH₄⁺ treatments in this study. However, the mechanism by which
290 NO₃⁻ and NO₂⁻ could interfere with C assimilation for MOB remains unclear. Alternative pathways for CH₄
291 uptake may have contributed to the particularly high ¹³C assimilation in NO₂⁻ amended crusts. The
292 microorganism *Candidatus Methyloirabilis oxyfera* is able to couple anaerobic methane oxidation with
293 nitrite reduction (Ettwig et al., 2010), and may be widespread in natural environments (Ettwig et al., 2009;
294 Wang et al., 2012; Zhu et al., 2012). We recovered 10MeC_{16:0} in the present study, which is characteristic of
295 the lipid profile of *M. oxyfera* (Kool et al., 2012). Still, more concrete evidence, such as the recovery of
296 specific gene markers of *M. oxyfera* (Luesken et al., 2011), is needed to confirm the presence of *M. oxyfera*
297 in surface crusts. More recently, gammaproteobacterial (Type I) methanotrophs *Methylomonas denitrificans*
298 and *Methylomicrobium album* have been reported to be able to oxidize methane under hypoxia using
299 oxidized nitrogen as electron acceptor (Kits et al., 2015a; Kits et al., 2015b). The involvement of such a
300 process would be consistent with the high proportion of C₁₆ PLFAs recovered from NO₃⁻ and NO₂⁻
301 treatments, and the genera *Methylomonas* and *Methylomicrobium* are widespread (Knief, 2015) and were
302 indeed present in surface crusts (Duan et al., 2014). However, there was no direct evidence in the present
303 study to evaluate the presence of these specific strains.

In-situ O₂ and N₂O Distribution and Implications for CH₄ Oxidation

Oxygen distribution in surface crusts vary over short distances, as shown in this and other studies (Hansen et al., 2009; Nielsen et al., 2010). Despite this variation, the depth of O₂ penetration is generally shallow and was never more than 25% of the thickness of the crust (Nielsen et al., 2010). The restriction of significant O₂ penetration is likely due to surface crusts being a floating organic structure on top of liquid slurry, where the bottom of the crust is always saturated, while aerobic processes actively consume O₂ in upper layers. Nielsen et al. (2010) proposed that trapping of gases formed in the slurry could elevate the crust above the liquid slurry phase and, as a result, improve O₂ penetration. Measurements of O₂ at a fixed depth over 48 h did indicate gas pockets that lifted the crust, but they were intermittently deflated (Supplemental Fig. S3). Structural voids in the crusts, such as pores and crevices, could also provide access for O₂ to deeper parts of the crusts, and weather conditions such as precipitation and drought will influence O₂ permeability by altering the wetness of the crust (Hansen et al., 2009).

The present study observed N₂O levels as high as 100 µmol L⁻¹, which was far above the N₂O concentrations commonly found in other environments. For example, Baral et al. (2014) and Zhou et al. (2016) reported N₂O concentrations of < 5 µmol L⁻¹ near the soil surface. Careful examination of potential interferences to the microsensor is therefore warranted. Surface crusts may contain up to 300 µmol L⁻¹ of H₂S (Nielsen et al., 2010), and H₂S is known to affect the signal of N₂O microsensors. However, the microsensor used here is equipped with an alkaline oxygen guard which converts incoming H₂S to ionic forms, and this offers some protection to the cathode from H₂S (Andersen et al., 2001). With the same type of microsensor used in this study, Andersen et al. (2001) found a reduction in the sensitivity towards N₂O with increasing concentrations of H₂S up to 350 µmol L⁻¹, indicating that any H₂S interference would result in lower, not higher, N₂O readings. Therefore, the high N₂O concentrations observed were not likely to be a result of H₂S interference. In the above mentioned studies where soil N₂O concentrations were <5 µmol L⁻¹, the corresponding N₂O emissions were 3.6 and 25 µmol m⁻² h⁻¹ (Baral et al., 2014; Zhou et al., 2016). For livestock slurry with surface crusts, N₂O emissions as high as 393–1,429 µmol m⁻² h⁻¹ have been reported (Hansen et al., 2009;

329 Sommer et al., 2000). Considering the relationship between N₂O concentration and emission, N₂O
 330 concentrations up to 100 µmol L⁻¹ in surface crusts seem plausible.

331 Maximum N₂O accumulation occurred near the oxic-anoxic interface in most cases, and therefore both
 332 nitrification and denitrification were potential sources of N₂O (Braker and Conrad, 2011). Law et al. (2012)
 333 found a correlation between N₂O production and ammonia oxidation rate, possibly as a result of nitrifier
 334 denitrification to conserve O₂ or prevent NO₂⁻ toxicity (Lawton et al., 2013), and similar mechanisms could
 335 lead to N₂O accumulation via ammonia oxidation in surface crusts. At or below oxic-anoxic interfaces,
 336 incomplete heterotrophic denitrification was likely the main source of N₂O due to intolerance of N₂O
 337 reductase towards trace O₂ (Thomson et al., 2012). Nitrous oxide could also be released as a terminal
 338 product from methane oxidation coupled with partial denitrification by *M. denitrificans* (Kits et al., 2015b)
 339 and *M. album* (Kits et al., 2015a; Nyerges et al., 2010), which can be enabled by hypoxia developed at the
 340 oxic-anoxic interfaces. Denitrification could also act as a sink for NO₃⁻ or, particularly, NO₂⁻ which is a strong
 341 inhibitor of aerobic methane oxidation. The ubiquitous presence of N₂O in the surface crusts indicated active
 342 transformations of N species which could interfere with methane oxidation. Yet, as seen from the incubation
 343 experiments, the effect of N species on MOB may be either inhibitory or stimulatory depending on other
 344 factors. Moreover, Type I MOB utilizing NO₂⁻ and/or NO₃⁻ for methane oxidation are presumably more
 345 resilient to N inhibition than others (Zhu et al., 2016). In support of this, *M. denitrificans* and *M. album* have
 346 been shown to tolerate and grow under 10 mM NO₃⁻ (Kits et al., 2015b) and 2.5 mM NO₂⁻ (Nyerges et al.,
 347 2010), respectively. Also, Hu and Lu (2015) found that, while NH₄⁺ and NO₃⁻ both stimulated Type I MOB as
 348 determined from *pmoA* gene copy numbers, Type II MOB were inhibited by NH₄⁺ as concentrations
 349 increased.

350 Clearly, various physio-chemical and biological properties are involved in regulating methane oxidation in
 351 surface crust. Firstly, the heterogeneous structure of the crust adds complexity to the distribution of gases.
 352 Secondly, there are complex interactions between CH₄ and O₂ levels, and inorganic N species and
 353 concentration, with respect to methanotrophic activity. In parts of the crust where CH₄ availability is

relatively low, moderate O₂ limitation probably has little impact on aerobic methane oxidation activity, whereas N inhibition could be significant depending on N species and concentration. In contrast, in parts of the crust with high CH₄ availability, O₂ limitation is likely the main control of aerobic methane oxidation, and inhibition due to inorganic N is only important where O₂ is not limiting.

Conclusions

Microsensor measurements of *in situ* O₂ and N₂O profiles revealed shallow penetration of O₂ into slurry surface crusts and active N transformations around oxic-anoxic interfaces. Laboratory incubations suggested that O₂ availability was more important to high-concentration than low-concentration methane oxidation, and there were complex interactions between inorganic N and O₂ limitation. The incorporation of ¹³C from CH₄ into membrane PLFAs indicated that both Type I and Type II MOB were actively involved in methane oxidation, but with Type I MOB dominating the activity at high CH₄ concentrations. These observations together imply that manipulation of storage conditions to increase headspace CH₄ concentration, as proposed by Petersen and Miller (2006), could stimulate methane oxidation by Type I MOB in the upper parts of the crust where O₂ is non-limiting and mineral N availability low.

Acknowledgements

This study was partly funded by the Danish Agency for Science, Technology and Innovation. We thank Bodil Stensgaard for technical assistance, and Kristian Kristensen for statistical support.

References

- Acha, V., J. Alba and F. Thalasso. 2002. The absolute requirement for carbon dioxide for aerobic methane oxidation by a methanotrophic-heterotrophic soil community of bacteria. *Biotechnology Letters* 24: 675-679. doi:10.1023/a:1015265530501.
- Amaral, J.A., C. Archambault, S.R. Richards and R. Knowles. 1995. Denitrification associated with groups I and II methanotrophs in a gradient enrichment system. *FEMS Microbiology Ecology* 18: 289-298.
- Amaral, J.A. and R. Knowles. 1995. Growth of methanotrophs in methane and oxygen counter gradients. *FEMS Microbiology Letters* 126: 215-220. doi:10.1111/j.1574-6968.1995.tb07421.x.
- Andersen, K., T. Kjær and N.P. Revsbech. 2001. An oxygen insensitive microsensor for nitrous oxide. *Sensors and Actuators B: Chemical* 81: 42-48. doi:http://dx.doi.org/10.1016/S0925-4005(01)00924-8.
- Baral, K.R., A.G. Thomsen, J.E. Olesen and S.O. Petersen. 2014. Controls of nitrous oxide emission after simulated cattle urine deposition. *Agriculture, Ecosystems & Environment* 188: 103-110. doi:http://dx.doi.org/10.1016/j.agee.2014.02.029.
- Bodelier, P.L.E. 2011. Interactions between nitrogenous fertilizers and methane cycling in wetland and upland soils. *Current Opinion in Environmental Sustainability* 3: 379-388. doi:http://dx.doi.org/10.1016/j.cosust.2011.06.002.
- Bodelier, P.L.E., M.-J. Bar Gillisen, K. Hordijk, J.S. Sinninghe Damste, W.I.C. Rijpstra, J.A.J. Geenevasen, et al. 2009. A reanalysis of phospholipid fatty acids as ecological biomarkers for methanotrophic bacteria. *ISME J* 3: 606-617. doi:10.1038/ismej.2009.6.
- Bodelier, P.L.E. and H.J. Laanbroek. 2004. Nitrogen as a regulatory factor of methane oxidation in soils and sediments. *FEMS Microbiology Ecology* 47: 265-277. doi:10.1016/s0168-6496(03)00304-0.
- Bollmann, A. and R. Conrad. 1998. Influence of O₂ availability on NO and N₂O release by nitrification and denitrification in soils. *Global Change Biology* 4: 387-396. doi:10.1046/j.1365-2486.1998.00161.x.
- Bosse, U., P. Frenzel and R. Conrad. 1993. Inhibition of methane oxidation by ammonium in the surface layer of a littoral sediment. *FEMS Microbiology Ecology* 13: 123-134. doi:10.1016/0168-6496(93)90030-b.
- Braker, G. and R. Conrad. 2011. Chapter 2: Diversity, Structure, and Size of N₂O-Producing Microbial Communities in Soils – What Matters for Their Functioning? In: S. S. Allen I. Laskin and M. G. Geoffrey, editors, *Advances in Applied Microbiology*. Academic Press. p. 33-70.
- Broecker, W.S. and T.H. Peng. 1974. Gas exchange rates between air and sea. *Tellus* 26: 21-35. doi:10.1111/j.2153-3490.1974.tb01948.x.
- Cai, Z.C. and X.Y. Yan. 1999. Kinetic model for methane oxidation by paddy soil as affected by temperature, moisture and N addition. *Soil Biology & Biochemistry* 31: 715-725.
- Carlsen, H.N., L. Joergensen and H. Degn. 1991. Inhibition by ammonia of methane utilization in *Methylococcus capsulatus* (Bath). *Applied Microbiology and Biotechnology* 35: 124-127. doi:10.1007/bf00180649.
- Damgaard, L.R. and N.P. Revsbech. 1997. A Microscale Biosensor for Methane Containing Methanotrophic Bacteria and an Internal Oxygen Reservoir. *Analytical Chemistry* 69: 2262-2267. doi:10.1021/ac9611576.
- De Beer, D., A. Schramm, C.M. Santegoeds and M. Kuhl. 1997. A nitrite microsensor for profiling environmental biofilms. *Applied and Environmental Microbiology* 63: 973-977.
- De Beer, D., J.-P.R.A. Sweerts and J.C. van den Heuvel. 1991. Microelectrode measurement of ammonium profiles in freshwater sediments. *FEMS Microbiology Letters* 86: 1-6. doi:10.1111/j.1574-6968.1991.tb04789.x.

- De Visscher, A., D. Thomas, P. Boeckx and O. Van Cleemput. 1999. Methane Oxidation in Simulated Landfill Cover Soil Environments. *Environmental Science & Technology* 33: 1854-1859. doi:10.1021/es9900961.
- Duan, Y.-F., W. Abu Al-Soud, A. Brejnrod, S.J. Sørensen, L. Elsgaard, S.O. Petersen, et al. 2014. Methanotrophs, methanogens and microbial community structure in livestock slurry surface crusts. *Journal of Applied Microbiology* 117: 1066–1078. doi:10.1111/jam.12584.
- Duan, Y.-F., L. Elsgaard and S.O. Petersen. 2013. Inhibition of methane oxidation in a slurry surface crust by inorganic N: An incubation study. *Journal of Environmental Quality* 42: 507-515. doi:10.2134/jeq2012.0230.
- Dunfield, P. and R. Knowles. 1995. Kinetics of inhibition of methane oxidation by nitrate, nitrite, and ammonium in a humisol. *Applied and Environmental Microbiology* 61: 3129-3135.
- Ettwig, K.F., T.v. Alen, K.T.v.d. Pas-Schoonen, M.S.M. Jetten and M. Strous. 2009. Enrichment and molecular detection of denitrifying methanotrophic bacteria of the NC10 phylum. *Applied and Environmental Microbiology* 75: 3656-3662. doi:10.1128/aem.00067-09.
- Ettwig, K.F., M.K. Butler, D. Le Paslier, E. Pelletier, S. Mangelot, M.M.M. Kuypers, et al. 2010. Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* 464: 543-548. doi:10.1038/nature08883.
- Francesco, N.T., S. Mirella, R. Simone, F. Alessandro, F. Nuala and S. Pete. 2013. The FAOSTAT database of greenhouse gas emissions from agriculture. *Environmental Research Letters* 8: 015009.
- Graham, D.W., J.A. Chaudhary, R.S. Hanson and R.G. Arnold. 1993. Factors affecting competition between type I and type II methanotrophs in two-organism, continuous-flow reactors. *Microbial Ecology* 25: 1-17. doi:10.1007/bf00182126.
- Gulledge, J. and J.P. Schimel. 1998. Low-Concentration Kinetics of Atmospheric CH₄ Oxidation in Soil and Mechanism of NH₄⁺ Inhibition. *Applied and Environmental Microbiology* 64: 4291-4298.
- Hansen, R.R., D.A. Nielsen, A. Schramm, L.P. Nielsen, N.P. Revsbech and M.N. Hansen. 2009. Greenhouse Gas Microbiology in Wet and Dry Straw Crust Covering Pig Slurry. *Journal of Environmental Quality* 38: 1311-1319. doi:10.2134/jeq2008.0336.
- Hanson, R.S. and T.E. Hanson. 1996. Methanotrophic bacteria. *Microbiological Reviews* 60: 439-471.
- Hu, A. and Y. Lu. 2015. The differential effects of ammonium and nitrate on methanotrophs in rice field soil. *Soil Biology and Biochemistry* 85: 31-38. doi:http://dx.doi.org/10.1016/j.soilbio.2015.02.033.
- King, G.M. and S. Schnell. 1994. Ammonium and Nitrite Inhibition of Methane Oxidation by *Methylobacter albus* BG8 and *Methylosinus trichosporium* OB3b at Low Methane Concentrations. *Applied and Environmental Microbiology* 60: 3508-3513.
- King, G.M. and S. Schnell. 1998. Effects of Ammonium and Non-Ammonium Salt Additions on Methane Oxidation by *Methylosinus trichosporium* OB3b and Maine Forest Soils. *Applied and Environmental Microbiology* 64: 253-257.
- Kits, K.D., D.J. Campbell, A.R. Rosana and L.Y. Stein. 2015a. Diverse electron sources support denitrification under hypoxia in the obligate methanotroph *Methylobacterium albus* strain BG8. *Frontiers in Microbiology* 6: 1072. doi:10.3389/fmicb.2015.01072.
- Kits, K.D., M.G. Klotz and L.Y. Stein. 2015b. Methane oxidation coupled to nitrate reduction under hypoxia by the Gammaproteobacterium *Methylobacterium denitrificans*, sp. nov. type strain FJG1. *Environmental Microbiology* 17: 3219-3232. doi:10.1111/1462-2920.12772.
- Knief, C. 2015. Diversity and Habitat Preferences of Cultivated and Uncultivated Aerobic Methanotrophic Bacteria Evaluated Based on *pmoA* as Molecular Marker. *Frontiers in Microbiology* 6. doi:10.3389/fmicb.2015.01346.

- Kool, D.M., B. Zhu, W.I.C. Rijpstra, M.S.M. Jetten, K.F. Ettwig and J.S. Sinninghe Damsté. 2012. Rare Branched Fatty Acids Characterize the Lipid Composition of the Intra-Aerobic Methane Oxidizer "Candidatus Methyloirabilis oxyfera". *Applied and Environmental Microbiology* 78: 8650-8656. doi:10.1128/aem.02099-12.
- Larsen, L.H., N.P. Revsbech and S.J. Binnerup. 1996. A microsensor for nitrate based on immobilized denitrifying bacteria. *Applied and Environmental Microbiology* 62: 1248-1251.
- Law, Y., B.-J. Ni, P. Lant and Z. Yuan. 2012. N₂O production rate of an enriched ammonia-oxidising bacteria culture exponentially correlates to its ammonia oxidation rate. *Water Research* 46: 3409-3419. doi:10.1016/j.watres.2012.03.043.
- Lawton, T.J., K.E. Bowen, L.A. Sayavedra-Soto, D.J. Arp and A.C. Rosenzweig. 2013. Characterization of a Nitrite Reductase Involved in Nitrifier Denitrification. *Journal of Biological Chemistry* 288: 25575-25583. doi:10.1074/jbc.M113.484543.
- Liu, L. and T.L. Greaver. 2009. A review of nitrogen enrichment effects on three biogenic GHGs: the CO₂ sink may be largely offset by stimulated N₂O and CH₄ emission. *Ecology Letters* 12: 1103-1117. doi:10.1111/j.1461-0248.2009.01351.x.
- Luesken, F.A., B. Zhu, T.A. van Alen, M.K. Butler, M. Rodriguez Diaz, B. Song, et al. 2011. pmoA primers for detection of anaerobic methanotrophs. *Applied and Environmental Microbiology*. doi:10.1128/aem.02960-10.
- Melse, R.W. and A.W. van der Werf. 2005. Biofiltration for Mitigation of Methane Emission from Animal Husbandry. *Environmental Science & Technology* 39: 5460-5468. doi:10.1021/es048048q.
- Nielsen, D.A., L.P. Nielsen, A. Schramm and N.P. Revsbech. 2010. Oxygen Distribution and Potential Ammonia Oxidation in Floating, Liquid Manure Crusts. *J. Environ. Qual.* 39: 1813-1820. doi:10.2134/jeq2009.0382.
- Nielsen, D.A., A. Schramm, L.P. Nielsen and N.P. Revsbech. 2013. Seasonal Methane Oxidation Potential in Manure Crusts. *Applied and Environmental Microbiology* 79: 407-410. doi:10.1128/aem.02278-12.
- Nyerges, G., S.-K. Han and L.Y. Stein. 2010. Effects of Ammonium and Nitrite on Growth and Competitive Fitness of Cultivated Methanotrophic Bacteria. *Applied and Environmental Microbiology* 76: 5648-5651. doi:10.1128/aem.00747-10.
- Petersen, S. and P. Ambus. 2006. Methane Oxidation in Pig and Cattle Slurry Storages, and Effects of Surface Crust Moisture and Methane Availability. *Nutrient Cycling in Agroecosystems* 74: 1-11. doi:10.1007/s10705-005-3822-6.
- Petersen, S.O., B. Amon and A. Gattinger. 2005. Methane Oxidation in Slurry Storage Surface Crusts. *J. Environ. Qual.* 34: 455-461. doi:10.2134/jeq2005.0455.
- Petersen, S.O., P.S. Frohne and A.C. Kennedy. 2002. Dynamics of a Soil Microbial Community under Spring Wheat. *Soil Science Society of America Journal* 66: 826-833. doi:10.2136/sssaj2002.8260.
- Petersen, S.O. and D.N. Miller. 2006. Greenhouse gas mitigation by covers on livestock slurry tanks and lagoons? *Journal of the Science of Food and Agriculture* 86: 1407-1411. doi:10.1002/jsfa.2543.
- Petersen, S.O., M. Skov, P. Drøschler and A.P.S. Adamsen. 2009. Pilot Scale Facility To Determine Gaseous Emissions From Livestock Slurry During Storage. *J. Environ. Qual.* 38: 1560-1568. doi:10.2134/jeq2008.0376.
- Qiu, Q., M. Noll, W.-R. Abraham, Y. Lu and R. Conrad. 2008. Applying stable isotope probing of phospholipid fatty acids and rRNA in a Chinese rice field to study activity and composition of the methanotrophic bacterial communities in situ. *ISME J* 2: 602-614.

- Revsbech, N.P. 2005. Analysis of Microbial Communities with Electrochemical Microsensors and Microscale Biosensors. In: R. L. Jared, editor *Methods in Enzymology*. Academic Press. p. 147-166.
- Roslev, P., N. Iversen and K. Henriksen. 1997. Oxidation and assimilation of atmospheric methane by soil methane oxidizers. *Applied and Environmental Microbiology* 63: 874-880.
- Sommer, S., S. Petersen, P. Sørensen, H. Poulsen and H. Møller. 2007. Methane and carbon dioxide emissions and nitrogen turnover during liquid manure storage. *Nutrient Cycling in Agroecosystems* 78: 27-36. doi:10.1007/s10705-006-9072-4.
- Sommer, S.G., S.O. Petersen and H.T. Sørensen. 2000. Greenhouse Gas Emission from Stored Livestock Slurry. *Journal of Environmental Quality* 29: 744-751. doi:10.2134/jeq2000.00472425002900030009x.
- Stein, L.Y. and M.G. Klotz. 2011. Nitrifying and denitrifying pathways of methanotrophic bacteria. *Biochemical Society Transactions* 39: 1826-1831. doi:10.1042/bst20110712.
- Thomson, A.J., G. Giannopoulos, J. Pretty, E.M. Baggs and D.J. Richardson. 2012. Biological sources and sinks of nitrous oxide and strategies to mitigate emissions. *Philosophical Transactions of The Royal Society B* 367: 1157-1168. doi:10.1098/rstb.2011.0415.
- Urmann, K., E.S. Norina, M.H. Schroth and J. Zeyer. 2007. Methanotrophic activity in a diffusive methane/oxygen counter-gradient in an unsaturated porous medium. *Journal of Contaminant Hydrology* 94: 126-138. doi:10.1016/j.jconhyd.2007.05.006.
- Wang, Y., G. Zhu, H.R. Harhangi, B. Zhu, M.S.M. Jetten, C. Yin, et al. 2012. Co-occurrence and distribution of nitrite-dependent anaerobic ammonium and methane-oxidizing bacteria in a paddy soil. *FEMS Microbiology Letters* 336: 79-88. doi:10.1111/j.1574-6968.2012.02654.x.
- Wang, Z.-P. and P. Ineson. 2003. Methane oxidation in a temperate coniferous forest soil: effects of inorganic N. *Soil Biology and Biochemistry* 35: 427-433. doi:10.1016/s0038-0717(02)00294-8.
- Welte, C.U., O. Rasigraf, A. Vaksmaa, W. Versantvoort, A. Arshad, H.J.M. Op den Camp, et al. 2016. Nitrate- and nitrite-dependent anaerobic oxidation of methane. *Environmental Microbiology Reports* 8: 941-955. doi:10.1111/1758-2229.12487.
- Whittenbury, R., K.C. Phillips and J.F. Wilkinson. 1970. Enrichment, Isolation and Some Properties of Methane-utilizing Bacteria. *Journal of General Microbiology* 61: 205-218. doi:10.1099/00221287-61-2-205.
- Zhou, Z., X. Xu, Z. Bi, L. Li, B. Li and Z. Xiong. 2016. Soil concentration profiles and diffusion and emission of nitrous oxide influenced by the application of biochar in a rice-wheat annual rotation system. *Environmental Science and Pollution Research* 23: 7949-7961. doi:10.1007/s11356-015-5929-x.
- Zhu, B., G. van Dijk, C. Fritz, A.J.P. Smolders, A. Pol, M.S.M. Jetten, et al. 2012. Anaerobic Oxidation of Methane in a Minerotrophic Peatland: Enrichment of Nitrite-Dependent Methane-Oxidizing Bacteria. *Applied and Environmental Microbiology* 78: 8657-8665. doi:10.1128/aem.02102-12.
- Zhu, J., Q. Wang, M. Yuan, G.-Y.A. Tan, F. Sun, C. Wang, et al. 2016. Microbiology and potential applications of aerobic methane oxidation coupled to denitrification (AME-D) process: A review. *Water Research* 90: 203-215. doi:http://dx.doi.org/10.1016/j.watres.2015.12.020.

Figure 1: Profiles of oxygen (O₂, solid line, top X-axis) and nitrous oxide (N₂O, dashed line, bottom X-axis) profiles in a surface crust measured using microsensors at 20 cm intervals along the 200-cm diameter of a slurry storage tank. Distances of the sampling points from the edge of the tank are indicated on the bottom-right corner of each panel. Oxygen was traced to a depth of 30 mm in all cases; N₂O was traced to a depth of 30 mm at 20–80 cm and to a depth of 60 mm at 100–180 cm.

Figure 2: Dynamics of CH₄ concentrations during a 72-h incubation of slurry surface crusts with initial O₂ concentrations of 20, 5, 3, 2, 1, and 0%, and with initial CH₄ concentrations of 10² ppmv (a) and 10⁴ ppmv (b). Blank control contained 20% initial headspace O₂ and no crust material. Each point represents the mean of triplicate assays, and error bars show standard error.

Figure 3: First-order rate constants of potential CH_4 oxidation under different O_2 and CH_4 concentrations during the first 8 h of incubation. The bars show the mean of triplicate assays, and the error bars show standard error. For each CH_4 concentration, bars denoted with the same letter on top are not significantly different at $\alpha = 0.05$.

Figure 4: Total recovery of ¹³C in PLFA (expressed as peak area per gram fresh crust) derived from incubation of surface crusts with ¹³CH₄ (a1–a4), and proportions of ¹³C incorporation into PLFA belonging to C₁₆ or C₁₈ clusters (b1–b4) at different combinations of O₂, CH₄, and inorganic N availability. Data shown are means and standard errors of two replicates.

Table 1: Selected chemical properties of various livestock slurries and surface crusts from previous studies.

Sample	Surface Crust			Slurry	
	NH ₄ ⁺ μmol kg ⁻¹ WW*	NO ₃ ⁻ μmol kg ⁻¹ WW	NO ₂ ⁻ μmol kg ⁻¹ WW	pH	NH ₄ ⁺ mM
Cattle slurry surface crusts developed for 3 years (Duan et al., 2013)	116.3	370.9	n.a. [#]	n.a.	n.a.
Various cattle and swine slurries and surface crusts (Nielsen et al., 2010)	n.a.	7– 3,602	2–98,000	6.96–7.7	84–205
Surface crusts of various dryness (Hansen et al., 2009)	n.a.	290–35,000	80–1,990	7.7	175
Various cattle and swine slurries (Sommer et al., 2007)	n.a.	n.a.	n.a.	n.a.	79–257
Surface crusts (Petersen et al., 2006)	1,432–54,923	91–16,378	3–50	7.13–8.89	1.7–69
Digested cattle slurry (Clemens et al., 2006)	n.a.	n.a.	n.a.	7.4–7.8	85–127

* WW, wet weight.

[#] n.a., value not reported.

Table 2: Combinations of CH₄, O₂, and inorganic N amendments used in this study. The values indicate the number of replicates prepared. Treatments marked with asterisks (*) were used for analysis of ¹³C-labelled phosphate lipid fatty acids (PLFAs).

N Species and Concentrations		CH ₄ : 10 ² ppmv						10 ⁴ ppmv						
		O ₂ :	0%	1%	2%	3%	5%	20%	0%	1%	2%	3%	5%	20%
None			3	3*	3	3*	3	3*	3	3*	3	3*	3	3*
NH ₄ ⁺	10 mM					2		2				2		2
	50 mM					2		2				2*		2*
NO ₃ ⁻	10 mM					2		2				2		2
	50 mM					2		2				2*		2*
NO ₂ ⁻	1 mM					2		2				2*		2*

Table 3: First-order rate constants (h^{-1}) of CH_4 oxidation in slurry surface crust samples under different O_2 concentrations and N amendments. Numbers in parentheses indicate relative activity (the ratio of the treatment activity as compared to the activity of the N-free control). Under each CH_4 concentration, values followed by the same letter are not significantly different at $\alpha = 0.05$.

N Species and Concentrations		First-Order Rate Constants (h^{-1})			
		10^2 ppmv CH_4		10^4 ppmv CH_4	
		20% O_2	3% O_2	20% O_2	3% O_2
None		0.036 (1.00) a	0.037 (1.00) a	0.040 (1.00) b	0.016 (1.00) e
NH_4^+	10 mM	0.015 (0.42) d	0.010 (0.27) ef	0.031 (0.78) c	0.017 (1.06) e
	50 mM	0.007 (0.19) fg	0.005 (0.14) g	0.013 (0.33) e	0.005 (0.31) f
NO_3^-	10 mM	0.029 (0.81) b	0.022 (0.59) c	0.054 (1.35) a	0.022 (1.38) d
	50 mM	0.017 (0.47) d	0.013 (0.35) de	0.028 (0.70) c	0.021 (1.31) d
NO_2^-	1 mM	0.008 (0.22) fg	0.008 (0.22) fg	0.016 (0.40) e	0.017 (1.06) e

Methanotrophic activity in slurry surface crusts as influenced by CH₄, O₂, and inorganic N

Yun-Feng Duan ^{1*}, Sabine Reinsch ^{2#}, Per Ambus ^{2§}, Lars Elsgaard ¹, and Søren O. Petersen ¹

¹ Department of Agroecology, Aarhus University, Blichers Allé 20, DK-8830 Tjele, Denmark

² Department of Chemical and Biochemical Engineering, Technical University of Denmark, Frederiksborgvej 399, DK-4000 Roskilde, Denmark

* Corresponding Author: Yun-Feng Duan, Department of Agroecology, Aarhus University, Blichers Allé 20, DK-8830 Tjele, Denmark; E-mail: Kevin.YF.Duan@gmail.com.

Present address: Centre for Ecology and Hydrology, Deiniol Rd, Bangor Gwynedd, LL57 2 UW, UK

§ Present address: Department of Geosciences and Natural Resource Management, University of Copenhagen, Øster Voldgade 10, DK-1350 Copenhagen, Denmark

Figure S1: Custom-made mounting system for *in situ* measurement of O_2 and N_2O profiles using microsensors.

A 200-cm long aluminum bar with mm-scale distance marks (a) was installed on the inner rim of the storage tank, approximately 40 cm above the surface crust (b). A computer-controlled motorized micromanipulator (Unisense, Aarhus, Denmark) capable of vertical movement was mounted on a custom-made rack, which could be moved manually along the length of the aluminum bar (c). The microsensor was fixed to the micromanipulator using a rubber-lined clamp (d). When installed at the initial position for measurement, the tip of the microsensor was approximately 1 cm above the surface of the crust. During measurements, a cover was placed loosely over the storage tank to avoid heating of the surface crust by direct insolation.

When measurement at one sampling point was completed, the microsensor was retreated to initial position and temporarily detached from the micromanipulator to protect the tip from breaking during movement, and reinstalled after the micromanipulator had been moved to the next sampling point.

Figure S2: Diagram of the setup to prepare and distribute anoxic incubation medium.

A Duran bottle was modified to have three plastic tubes inserted through the cap: one as N₂ inlet, one as N₂ outlet, and the third as liquid outlet. The Duran bottle was filled with BS medium to ca. 70% of its volume and placed on an electric heater with the cap tightly secured. The BS medium was heated to the boiling point while bubbled with pure N₂ for at least 30 min, and then cooled to room temperature still under N₂ bubbling. Then, the N₂ outlet was closed and the liquid outlet valve was opened, and the entire tubing was flushed and filled with anoxic BS medium by the build-up of gas pressure inside the Duran bottle. When distributing BS medium to the serum bottle, a needle connected to pure N₂ flow was first inserted into the serum bottle through the rubber stopper and then, a second needle was inserted as N₂ outlet. The serum bottle was flushed with N₂ for 10 sec, and then a third needle connected to the three-port valve was inserted. Ports A and B of the three-port valve were opened while C was closed, and the syringe was filled with 20 mL incubation medium. Then, ports B and C were opened and A was closed and the incubation medium was injected from the syringe into the serum bottle while the bottle was being flushed by pure N₂.

Figure S3: Dynamics of O₂ concentration at a fixed depth in a surface crust over 48 hours. The storage tank was covered to eliminate wind effect but was passively ventilated. The microsensor was initially fixed in a position where the tip just touched the surface of the crust. Oxygen concentrations were measured every 10–15 min as described in the manuscript. The peaks in O₂ profile showed that the microsensor was alternately exposed to atmospheric O₂ and more anaerobic conditions. This indicated that the surface of the crust periodically rose up and moved down, likely due to inflation and deflation of gas pockets underneath the crust.

Table S1: Oxygen Consumption during CH₄ oxidation at 10,000 ppmv initial CH₄.

Initial State (time 0)		After 8 h		
Headspace O ₂ (%)	Dissolved O ₂ (μM)	CH ₄ Consumed (μM)	O ₂ Consumed (μM)	O ₂ Remaining (μM)
20	256.0	4.92	9.84	246.2
3	38.4	1.95	3.90	34.5
1	12.8	0.57	1.14	11.7

- Stoichiometry of CH₄ oxidation is:

where x is the fraction of carbon that is assimilated into biomass (CH₂O) (Urmann et al., 2007).

Therefore, CH₄ to O₂ ratio in CH₄ oxidation theoretically ranges between 1:1 (100% C assimilation) and 1:2 (no C assimilation). In this calculation, we assumed maximum O₂ consumption, i.e. a CH₄ to O₂ ratio of 1:2.

- Henry's Law constant (K) for O₂ and CH₄ are 1.28 and 1.34 mmol L⁻¹ atm⁻¹, respectively.
- Dissolved O₂ was calculated as: $K \times [\text{Headspace O}_2]$.
- Diffusivities of dissolved O₂ and CH₄ in water at 20°C are 2.06 and 1.75×10^{-5} cm² sec⁻¹, respectively (Broecker and Peng, 1974). Therefore, there's no diffusion limit of O₂ for CH₄ oxidation.