



## Effects of Different Lipophilized Ferulate Esters in Fish Oil-Enriched Milk: Partitioning, Interaction, Protein, and Lipid Oxidation

Qiu, Xujian; Jacobsen, Charlotte; Villeneuve, Pierre; Durand, Erwann; Sørensen, Ann-Dorit Moltke

*Published in:*  
Journal of Agricultural and Food Chemistry

*Link to article, DOI:*  
[10.1021/acs.jafc.7b02994](https://doi.org/10.1021/acs.jafc.7b02994)

*Publication date:*  
2017

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

*Citation (APA):*  
Qiu, X., Jacobsen, C., Villeneuve, P., Durand, E., & Sørensen, A-D. M. (2017). Effects of Different Lipophilized Ferulate Esters in Fish Oil-Enriched Milk: Partitioning, Interaction, Protein, and Lipid Oxidation. *Journal of Agricultural and Food Chemistry*, 65(43), 9496–9505. <https://doi.org/10.1021/acs.jafc.7b02994>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

**Effects of different lipophilized ferulate esters in fish oil enriched milk: Partitioning, Interaction, Protein and Lipid Oxidation**

|                               |   |
|-------------------------------|---|
| Journal:                      | <i>Journal of Agricultural and Food Chemistry</i>   |
| Manuscript ID                 | jf-2017-02994x.R1   |
| Manuscript Type:              | Article   |
| Date Submitted by the Author: | n/a   |
| Complete List of Authors:     | Qiu, xujian; Jimei University, Food science<br>Jacobsen, Charlotte; Technical University of Denmark, National food Institute<br>Villeneuve, Pierre; CIRAD<br>DURAND, Erwann; CIRAD, IATE<br>Sørensen, Ann-Dorit; Technical University of Denmark, National Food Institute, Division of Industrial Food Research |
|                               |   |

SCHOLARONE™  
Manuscripts

*Effects of different lipophilized ferulate esters in fish oil enriched milk:  
Partitioning, Interaction, Protein and Lipid Oxidation*

Xujian Qiu<sup>1,2</sup>, Charlotte Jacobsen<sup>2</sup>, Pierre Villeneuve<sup>3</sup>, Erwann Durand<sup>3</sup> and Ann-Dorit Moltke Sørensen<sup>2,\*</sup>

<sup>1</sup> College of Food and Biological Engineering, Xiamen Key Laboratory of Marine Functional Food, Jimei University, 43 Yindou Road, Xiamen 361021, Fujian, China

<sup>2</sup> Division of Food Technology, National Food Institute (DTU Food), Technical University of Denmark, DK 2800 Kgs Lyngby, Denmark

<sup>3</sup> CIRAD, UMR IATE, Montpellier, F-34398, France

\*Tel: +45 45252591. E-mail: adms@food.dtu.dk

1 **ABSTRACT:** Antioxidant effects of ferulic acid and lipophilized ferulate esters were investigated  
2 in fish oil enriched milk. Methyl ferulate (C1) and ethyl ferulate (C2) more efficiently prevented  
3 lipid oxidation than dodecyl ferulate (C12), followed by ferulic acid (C0). The combination of C1  
4 or C2 with C12 could have a “synergistic” effect indicated by peroxide value, hexanal, and 1-  
5 penten-3-ol analysis results. These antioxidants also showed protein oxidation inhibition effects.  
6 The most effective antioxidants (C1 and C2) had the highest concentration in the precipitate phase,  
7 but lowest concentration in aqueous phase which was opposite to the partitioning of C0. C12 had  
8 the highest concentration in the oil and emulsion phase. In particular, the interaction between  
9 ferulates esterified with short and medium alkyl chain lengths respectively could lead to their  
10 “synergistic” effects in fish oil enriched milk, which could be caused by the change of their  
11 partitioning or localization at the interface.

12 **KEY WORDS:** Ferulic acid, Ferulate ester, omega-3 PUFA, protein oxidation, lipid oxidation,  
13 antioxidants, antioxidant partitioning, phenolipids

## 14 INTRODUCTION

15 Fish oils are reported to have many health beneficial effects including reduced risk of  
16 cardiovascular and inflammatory diseases. This is attributed to the presence of long chain n-3  
17 polyunsaturated fatty acids (PUFAs).<sup>1,2</sup> It is thus of great interest to the food industry to enrich food  
18 with fish oil. However, fish oils that contain n-3 PUFA like eicosapentaenoic acid (EPA, 20:5) and  
19 docosahexanoic acid (DHA, 22:6) are highly prone to lipid oxidation. Lipid oxidation can cause  
20 development of compounds with an unpleasant off flavor in the product even at low concentrations  
21 (e.g. 1µg/kg). Moreover, lipid oxidation may also generate some oxidized compounds, which are  
22 potentially harmful to human health.<sup>3-5</sup> Therefore, the use of efficient antioxidants in food is  
23 required to limit oxidation of the healthy n-3 PUFAs and other compounds in the food product.

24 Currently, there is an increasing interest in using natural antioxidants such as phenolic acids in  
25 foods. However, phenolic acids could have limited effectiveness in oil-in-water (O/W) emulsions  
26 since they are generally hydrophilic and have relatively high polarity. Polar antioxidants are less  
27 efficient in O/W emulsions than apolar antioxidants according to the “polar paradox” theory put  
28 forwarded first by Porter<sup>6</sup> and Porter et al.<sup>7</sup> This was explained by the partitioning of the  
29 antioxidants into the different phases. Apolar antioxidants were closely located to the interface,  
30 where lipid oxidation is initiated, whereas polar antioxidants are mainly located in the aqueous  
31 phase. The polarity/hydrophilicity of phenolic acids has therefore been modified by esterification  
32 with fatty alcohols in order to change the partitioning of the antioxidant in the O/W emulsion  
33 system to get better antioxidant effects.<sup>8</sup> Interestingly, the lipophilized phenolic acids have a higher  
34 antioxidant activity in O/W emulsions with the increment of alkyl chain length but only to a certain  
35 limit termed critical chain length. With further increment of the chain length, the antioxidant effect  
36 was significantly decreased. Based on these results, the cut-off effect was introduced.<sup>9</sup> The

37 mechanism of the cut-off phenomenon is still not very clear despite the fact that there has been  
38 significant progress in this area recently.<sup>10</sup>

39 In particular, there are only a few studies on antioxidant efficacies of lipophilized phenolic acids in  
40 real food systems. A study on antioxidant effects of lipophilized caffeic acid has shown that methyl  
41 and butyl caffeates were most efficient in milk, whereas butyl, octyl and dodecyl caffeates were  
42 most efficient in mayonnaise, suggesting that the critical chain lengths of lipophilized phenolic  
43 compounds were different in different foods enriched with fish oil.<sup>11</sup> Moreover, a study on  
44 dihydrocaffeates and rutin esters in fish oil enriched milk showed that phenolipids with medium  
45 chain were more efficient although phenolipids with only two different alkyl chain lengths were  
46 evaluated.<sup>12</sup> In another study with lipophilized ferulic acid, methyl ferulate had the best antioxidant  
47 activity of all the ferulates and ferulic acid evaluated in fish oil enriched milk. Octyl and dodecyl  
48 ferulates were the least efficient compounds evaluated in this study where they acted as prooxidants  
49 in spite of their more apolar character than methyl ferulate.<sup>13</sup>

50 It is known that a combination of antioxidants can often provide synergistic antioxidant efficiency  
51 although the mechanism is not completely understood. To the best of our knowledge, there have  
52 been no studies performed on a combination of antioxidants of lipophilized ferulic acids with  
53 different alkyl chain lengths. Therefore, it is of interest to test the antioxidant efficiency of the  
54 mixed lipophilized ferulic acids. Based on former experiment with fish oil enriched milk<sup>13</sup>, the  
55 combination of ferulate esters with short alkyl chain length and medium alkyl chain length was  
56 selected. In addition, only few studies have evaluated the partitioning of antioxidants in fish oil  
57 enriched milk as well as the effect of phenolipids on protein oxidation. Thus, the first aim of this  
58 study was to evaluate the effects of ferulic acid, ferulate esters, and a combination of short and  
59 medium alkyl chain ferulate esters on the oxidative stability of fish oil enriched milk. Both lipid and

60 protein oxidation was assessed. The second aim was to compare the oxidative stability to the  
61 partitioning of ferulic acid and ferulates in fish oil enriched milk.

## 62 MATERIALS AND METHODS

63 **Materials.** Pasteurized milk with 0.4% and 1.5% fat content was purchased from a local  
64 supermarket. Cod liver fish oil was supplied by Maritex A/S (TINE BA, Sortland, Norway). The  
65 peroxide value (PV) of the fish oil was 0.65 meq peroxides/kg. The fish oil had 242 mg  $\alpha$ -  
66 tocopherol/kg, 119 mg  $\gamma$ -tocopherol/kg, 44 mg  $\delta$ -tocopherol/kg, respectively. The fatty acid  
67 composition of the oil was analyzed by GC as: C14:0, 3.8%; C16:0, 9.9%; C16:1 (n-7), 9.4%;  
68 C18:0, 2.0%; C18:1 (n-9), 17.2%; C18:1 (n-7), 4.8%; C18:2 (n-6), 2.0%; C18:3 (n-3), 2.8%;  
69 C18:4(n-3), 0.2%; C20:1, 13.2%; C20:5 (n-3) (EPA), 9.8%; C22:1(n-11), 5.8%; C22:5(n-3), 1.2%; C  
70 22:6 (n-3) (DHA), 11.9% (w/w).

71 Methyl (C1) and dodecyl (C12) ferulates used in the study were synthesized by the grafting  
72 respective alcohols to ferulic acid according to the procedure described by Sørensen et al.<sup>14</sup> Ferulic  
73 acid (C0) and ethyl ferulate (C2) were purchased from Sigma-Aldrich (Steinheim, Germany). All  
74 solvents used were of HPLC grade and purchased from Lab-Scan (Dublin, Ireland). Standards for  
75 quantification of volatile lipid oxidation products were from Sigma-Aldrich (Steinheim, Germany).

76 **Production of fish oil enriched milk with different antioxidants.** Milk with 1.5% and 0.4% fat  
77 content were mixed (6:5, w/w) to obtain a total fat content of 1%. Milk was then heated to 72 °C  
78 and kept for 15 sec. The selected temperature and time is known to get a better oxidative stability of  
79 the fish oil enriched milk according to a previous study.<sup>15</sup> Fish oil (0.5%, w/w) was then added to  
80 the milk while stirring manually. The fish oil enriched milk was homogenized with a two-valve  
81 homogenizer (Rannie, Albertslund, Denmark) with a pressure of 225 bar and recirculated for 3

82 times. To have enough fish oil enriched milk for all sample codes and sampling times, two batches  
83 (3 kg each batch) were prepared and then pooled.

84 The fish oil enriched milk was then divided into 1000 ml autoclaved bottles and the different  
85 methanolic antioxidant solutions (C0, C1, C2 and C12) were added to 4 bottles separately to make  
86 100  $\mu\text{M}$  final concentration. For the sample codes with combination of two ferulates, C1 (50  $\mu\text{M}$ ) or  
87 C2 (50  $\mu\text{M}$ ) was combined with C12 (50  $\mu\text{M}$ ) i.e. total antioxidant concentration 100  $\mu\text{M}$ . For the  
88 control (Con), only methanol was added in same amount as the antioxidant solutions. After gently  
89 shaking, the bulk samples of the different emulsions were then divided into 100 mL sterilized  
90 bottles for storage at 5 °C in darkness. On day 0, 3, 6, 9 and 13, one bottle of each code samples  
91 were taken out and divided into 50 ml brownbottles, flushed with nitrogen and stored at -40 °C for  
92 chemical analysis. On day 1 and day 13, droplet size of each sample code was measured without  
93 prior freezing of the samples. On day 1, 100 mL of each treatment except Con was taken out to  
94 determine the partitioning of ferulic acid and ferulate esters in the fish oil enriched milk.

## 95 **Methods of Analysis**

96 *Droplet size measurement.* A few drops of the sample were suspended in recirculating distilled  
97 water (2800 rpm, obscuration 13-15%, method: Fraunhofer) to measure the lipid droplet size in the  
98 fish oil enriched milk by a laser diffraction particle size analyzer (Mastersizer 2000, Malvern  
99 Instruments Ltd., Worcestershire, UK). Triplicate samples were measured.  $D_{3,2}$  was used to report  
100 the results.

101 *Lipid extraction by Bligh and Dyer method.* Lipid from fish oil enriched milk samples was extracted  
102 by Bligh and Dyer method.<sup>16</sup> Briefly, milk sample was homogenized with methanol, chloroform  
103 and water (1:2:2:1, w/v/v/v). After phase separation by centrifugation, the chloroform layer which



104 contained lipid was kept for analysis. For each treatment, duplicate lipid extractions were performed  
105 and analyzed for peroxide value (PV), fatty acid composition and tocopherols.

106 *PV analysis.* Lipid extracts from the samples were evaporated to dryness under nitrogen. PV  
107 analysis was performed on the same day of lipid extraction based on a method described by Shantha  
108 and Decker.<sup>17</sup> Results were reported as meq peroxides/kg lipids.

109 *Fatty acid composition analysis.* Lipid extracts were evaporated to dryness under nitrogen before  
110 analysis. 100  $\mu$ L internal standard C23:0 in heptane, 200  $\mu$ L heptane with BHT and 100  $\mu$ L toluene  
111 were added. Lipids were methylated with boron trifluoride reagent (20%) by a one-step procedure  
112 using a microwave (Multiwave 3000 SOLV, Anton Paar, Graz, Austria) with a 64MG5 rotor.  
113 Samples were heated for 5 min at 500 W and cooled down for 10 min. Then it was mixed with 1  
114 mL saturated NaCl solution and 0.7 mL heptane. The top layer was used for analysis by gas  
115 chromatography with flame ionization detection (GC-FID) (HP 5890A, Agilent Technology, Palo  
116 Alto, CA) according to AOCS official Method Ce 1b-89.<sup>18</sup> A DB-wax column (10 m $\times$ ID 0.1 mm  $\times$   
117 0.1  $\mu$ m film thickness, J&W Scientific, Folsom, CA) was used for analysis. The oven temperature  
118 program was set initially as 160  $^{\circ}$ C and increased gradually to 200  $^{\circ}$ C at 10.6  $^{\circ}$ C/min, kept at  
119 200  $^{\circ}$ C for 0.3 min, and increased to 220  $^{\circ}$ C at 10.6  $^{\circ}$ C/min, kept at 220  $^{\circ}$ C for 1 min, then  
120 increased to 240  $^{\circ}$ C at 10.6  $^{\circ}$ C/min and finally kept at 240  $^{\circ}$ C for 3.8 min. Injection volume is 0.2  
121  $\mu$ L with a split ratio of 1:50. Results were reported as percentages of total fatty acids.

122 *Tocopherol concentration determination.* Tocopherol concentration in samples were analyzed based  
123 on the AOCS official method Ce 8-89.<sup>19</sup> Briefly, lipid extracts were evaporated to dryness under  
124 nitrogen and dissolved in 1 mL heptane and directly analyzed by HPLC (Agilent 1100 series,  
125 Agilent Technology, Palo Alto, CA) with a fluorescence detector. The different tocopherols were  
126 separated on a silica column (Waters, 4.6 mm $\times$ 150 mm, 3  $\mu$ m). The mobile phase was isopropanol

127 in hexane (0.5:99.5, v/v). The excitation wavelength and the emission wavelength were set at 290  
128 nm and 330 nm respectively. Results were reported as  $\mu\text{g}$  tocopherol/g lipid.

129 *Volatile compound analysis.* A dynamic headspace gas chromatography mass spectrometry (GC-  
130 MS) method was used to analyze the volatile compounds of the milk samples. 4 g of milk samples  
131 with 0.5 mL of antifoam (Synperonic, 800  $\mu\text{L}/\text{mL}$  water) was put in a pear shaped bottle and heated  
132 in a water bath at 45  $^{\circ}\text{C}$  for 30 min. A nitrogen flow of approximately 150 mL/min was used to  
133 purge the milk sample in order to release the volatile compounds from the milk into a Tenax GR  
134 packed tube. Tenax GR tubes were put in an automatic thermal desorber (ATD-400, Perkin Elmer,  
135 Waltham, MA) to desorb the volatiles. Volatile compounds were then transferred automatically  
136 from ATD to a GC (Agilent 6890, Palo Alto, CA) with a mass selective detector (HP 5973). A  
137 DB1701 column (30 m, 0.25 mm  $\times$  1 mm film thickness, J&W Scientific, Folsom, CA) was used to  
138 separate the volatile compounds. The oven temperature was kept at 45  $^{\circ}\text{C}$  for 5 min, increased to  
139 55  $^{\circ}\text{C}$  at 1.5  $^{\circ}\text{C}/\text{min}$ , and increased from 55  $^{\circ}\text{C}$  to 90  $^{\circ}\text{C}$  at 2.5  $^{\circ}\text{C}/\text{min}$ , then increased to 220  $^{\circ}\text{C}$  at  
140 12  $^{\circ}\text{C}/\text{min}$  and finally kept at 220  $^{\circ}\text{C}$  for 4 min. Compounds were identified using the NIST data  
141 base and also confirmed with authentic standards. Volatile compounds in the samples were then  
142 quantified according to the external standard calibration curves using the ChemStation software. 30  
143 mg of mixed external standard solutions ranged from 0.1  $\mu\text{g}/\text{g}$  to 10  $\mu\text{g}/\text{g}$  was added to 4 g fresh  
144 milk (1.5% fat content) and also analyzed in the same way to prepare the calibration curves. Each  
145 sample was analyzed in triplicate and results were reported as ng volatile/g milk.

146 *Protein carbonyls analysis.* Protein oxidation in the milk samples was evaluated by carbonyl  
147 analysis based on the method by Levine et al.<sup>20</sup> and Fenaille et al.<sup>21</sup> Briefly, 50  $\mu\text{L}$  of milk was  
148 incubated with 0.5 mL 10 mM DNPH in 2 M HCl for 30 min in dark at room temperature. For the  
149 same milk sample a control was performed (0.5 mL 2 M HCl without DNPH). Milk proteins were  
150 precipitated with 10% (w/v) TCA (final concentration) and recovered by centrifugation. Protein

151 pellets were washed 3 times with 1 mL of ethanol/ethyl acetate 50:50 (v/v) to completely remove  
152 free DNPH reagent. The pellet was redissolved in 1 mL of 6 M guanidine hydrochloride (pH 2.3).  
153 Protein carbonyls and protein content were determined by UV at 370 nm and 280 nm respectively  
154 using a spectrophotometer. An extinction coefficient of  $2.2 \times 10^4 \text{ M}^{-1}\text{cm}^{-1}$  was used for the  
155 calculation of carbonyls. Results were expressed as nmol carbonyl/mg protein (n=3).

156 *Determination of the partitioning of ferulic acid and ferulates in fish oil enriched milk.* A simple  
157 ultracentrifuge procedure based on the method described by Jacobsen et al. with modification<sup>22</sup> was  
158 used to determine the partitioning of ferulic acid and ferulate esters in fish oil enriched milk  
159 samples. Triplicate samples of 20 mL were ultracentrifuged (Beckman Coulter Inc., Palo Alto, CA)  
160 for 2 h at 15 °C at 110,000 g. After centrifugation, samples were frozen at -18 °C to facilitate the  
161 separation of the obtained phases. There were 3 phases obtained after centrifugation: oil and  
162 emulsion (containing milk fat globule membrane), aqueous, and precipitate. Oil and emulsion phase  
163 was removed carefully by a sharp small spatula. Aqueous phase and precipitate could be poured out  
164 from the centrifuge tubes directly. Each phase was weighed and diluted with ethanol (1:4) and  
165 sonicated for 1 min except aqueous phase. Aqueous phase was directly filtered through a 0.45 µm  
166 PVDF syringe filter and analyzed. The mixed solution was then centrifuged and the supernatant was  
167 also filtered through the same type filter before HPLC analysis.

168 Ferulic acid and ferulate esters were determined by HPLC (Agilent 1100 series, Agilent Technology,  
169 Palo Alto, CA) at 328 nm based on the method described by Guyot et al.<sup>23</sup> with modification. An  
170 ODS-3 column (250\*4.6 mm, 5µm; Phenomenex, Torrance, CA) reversed phase column was used  
171 for the determination for all the samples. Elution was obtained by using a mobile phase of solvents  
172 A [water (pH 3)] and B [methanol/acetonitrile (1:1)]. Gradient elution was as follows: 95:5 (A/B)  
173 for 2 min then to 60:40 (A/B) over 3 min, to 0:100 (A/B) over 10 min and 0:100 (A/B) for 5 min,  
174 then reversion back to 95:5 (A/B) over 1 min, finally 95:5 (A/B) for 4 min. The flow rate was 0.9

175 mL/min. Standard solutions of ferulic acid and ferulates were analyzed under the same HPLC  
176 conditions to identify and quantify the ferulic acid and ferulates in each sample prepared from  
177 different phases.

178 **Statistical Analysis.** All data shown represent the mean value  $\pm$  standard deviation of duplicate or  
179 triplicate measurements. Data from the different quality parameters were subjected to one-way  
180 analysis of variance using SPSS software (version 19.0; SPSS Inc., Chicago, IL). Comparison of  
181 means was performed using a least squares difference (LSD) method ( $P < 0.05$ ). For PV or volatile  
182 compound results, we use the term “synergistic” effect of C1(C2) with C12 when there was no  
183 statistical difference between C1(C2)+C12 and C1(C2), but when the values of the combination  
184 were significantly lower than C12. Only when C1(C2)+C12 showed statistical significance when  
185 compared with C1(C2) and C12, they were compared with the expected values to determine the  
186 “synergistic” effects. For PV and volatile compound analysis results, expected C1(C2)+C12 were  
187 calculated by the average of C1(C2) with C12. This approach to evaluate synergistic effect is  
188 somewhat different from the commonly used approach where a synergistic effect is considered  
189 when the sum of the effects of individual compounds each added in a concentration of x and y is  
190 lower than the effect of the two compounds added together in a concentration of x+y. The effect is  
191 usually measured as % inhibition of the measured parameters. To highlight that a different approach  
192 has been used in this study the term synergistic is quoted.

## 193 **RESULTS AND DISCUSSION**

194 **Droplet size of the lipid in milk emulsion system.** Overall, the droplet size of the lipid in all the  
195 samples increased from an average of 0.81  $\mu\text{m}$  to 0.87  $\mu\text{m}$  during the whole storage period (data not  
196 shown). No creaming or flocculation was observed during 13 days of storage at 5 °C. The lipid  
197 droplet size of the fish oil enriched emulsion was slightly higher than in our previous studies.<sup>12,13</sup> It

198 may be related with the natural variation in the composition of milk used in the different studies.  
199 Particle size can range from 0.5 to 1.6  $\mu\text{m}$  in fish oil enriched milk samples depending on the  
200 homogenization temperature and pressure.<sup>24</sup> Particle size can influence the oxidation stability of  
201 emulsion as reported by Let et al.<sup>15</sup> This is not the focus of the present research and has been  
202 investigated before. Moreover, differences in droplet sizes were not significant and most likely did  
203 not affect lipid oxidation.

204 **Fatty acid composition.** EPA contents of all the samples on day 0 were from 3.54 - 3.70% and  
205 3.41 - 3.74% on day 13. DHA contents changed from 4.26 – 4.72% on day 0 to 4.13 – 4.45% on  
206 day 13 (data not shown). No significant reduction of EPA and DHA content was observed for  
207 samples between day 0 and day 13 except slight reduction for the C0 sample. EPA and DHA are  
208 prone to oxidation due to their highly unsaturated nature. In previous studies, the reduction could be  
209 significant or non-significant depending on the extent of lipid oxidation.<sup>11,12</sup>

210 **Change of PV in fish oil enriched milk emulsions.** In general, PV of the control sample increased  
211 dramatically from day 0 to day 6 and then maintained stable values (Figure 1). For both C0 and C12  
212 samples, PV increased gradually from day 0 to day 9 at a rate of 0.98 and 0.77 meq peroxides / kg  
213 lipid per day, respectively and then became steady. Control sample had significantly higher values  
214 than both C0 and C12 sample from day 3 till the end of the storage, while C0 had significantly  
215 higher PV than C12 after day 6 ( $P<0.05$ ). C1+C12 and C2+C12 had very similar slightly  
216 increasing PVs during the whole storage period, in agreement with the very similar PVs of C1 and  
217 C2. From day 6, the combination of C1 or C2 with C12 had significantly higher PVs than either C1  
218 or C2, but lower PVs than C12 and the expected combination values indicated as dashed lines in  
219 Figure 1 ( $P<0.05$ ). The lower PVs than expected values based on the individual treatments  
220 indicated a “synergistic” effect of the combination of ferulates with short and medium chain lengths.  
221 Further studies are required to confirm this preliminary finding on the “synergistic” effect and

222 synergistic effect with the use of other chain lengths or type of phenolipids. Furthermore, the  
223 synergistic effect should be investigated at other concentrations than that applied in the present  
224 study. C1 and C2 showed a very slight increase in PV throughout the storage period and had the  
225 lowest PV compared to all other treatments. However, the low PV did not necessarily indicate low  
226 lipid oxidation as the peroxides may be transformed into secondary oxidation products. The  
227 secondary oxidation products results presented next could provide a better profile of the lipid  
228 oxidation in all the treatments.

229 **Change of volatile secondary oxidation products during storage.** Secondary oxidation products  
230 are mainly derived from the decomposition of primary lipid hydroperoxides. Hexanal, 1-penten-3-  
231 one, 1-penten-3-ol, 2,4-heptadienal, 2-pentenal, 1-hexanol, 2,6-nonadienal were quantified. Figure 2  
232 shows the changes of hexanal, 1-penten-3-one, 1-penten-3-ol and 2,4-heptadienal. They represent  
233 the general development trend of the volatile secondary compounds quantified during storage. In  
234 addition, 1-penten-3-one, 1-penten-3-ol, 2,4-heptadienal were found to be related to the off flavor  
235 generated by oxidation of n-3 PUFAs.<sup>25</sup>

236 Hexanal are mainly from the oxidation of n-6 fatty acids and is a very important index of lipid  
237 oxidation. Both fish oil and milk fat contain significant amount of n-6 fatty acids, which are easily  
238 oxidized due to the unsaturated nature. Hexanal in all the samples reached the peak on day 6 and  
239 then gradually decreased or remained constant in concentration (Figure 2). Corresponding to the PV  
240 results, all the treatment had much lower concentrations than the control sample while C0 and C12  
241 had much higher concentrations than all other ferulate treatments after day 3 ( $P<0.05$ ). These  
242 results confirmed the strongest antioxidant effects of lipophilized ferulic acid with short alkyl chain  
243 lengths. In addition, there was no significant difference between C12 and C0 with regard to PV until  
244 day 9, however, C12 sample had significantly lower hexanal content than C0 sample starting from  
245 day 6 ( $P<0.05$ ). This finding suggested that C12 and other shorter lipophilized ferulic acid could

246 have stronger antioxidant effects by inhibiting not only primary oxidation but also the secondary  
247 oxidation reactions compared to ferulic acid. This is particular important as the volatile compounds  
248 due to oxidation could cause off flavor of the food products. On day 13, the ranking order of the  
249 concentration of hexanal was: Con<sup>a</sup>> C0<sup>b</sup>>C12<sup>c</sup>>C1+C12<sup>d</sup>=C2+C12<sup>d</sup>=C1<sup>d</sup>=C2<sup>d</sup>. The hexanal  
250 concentration in C1+C12 or C2+C12 was not significantly higher than C1 or C2, but significantly  
251 lower than C12, suggesting a “synergistic” effect of the combination of C1 or C2 with C12 on the  
252 reduction of hexanal content.

253 The content of 1-penten-3-one in all the samples also reached a peak concentration on day 6 except  
254 for C2 sample (Figure 3). In contrast to hexanal, 1-penten-3-one decreased sharply after day 6 and  
255 reached to approximately the same level for all the treatments at the end of the storage. The decline  
256 of 1-penten-3-one in fish oil enriched milk samples at later stage of storage period was also found in  
257 other similar studies.<sup>4,11,13</sup> C0 and C12 did not show lower concentration of 1-penten-3-one  
258 compared to the control until day 6 while the other treatments had significantly lower 1-penten-3-  
259 one concentration compared to the control starting from day 3 ( $P<0.05$ ). The biggest difference in  
260 1-penten-3-one concentrations among all treatments was shown on day 6. C2 had the lowest 1-  
261 penten-3-one concentration among all the treatments ( $P<0.05$ ). C1, C1+C12 and C2+C12 showed  
262 similar antioxidant effects ( $P>0.05$ ). The dramatic decline of 1-penten-3-one at the end of storage  
263 period was probably related to its reaction with other components in the milk sample.<sup>11</sup>

264 1-penten-3-ol is developed during the decomposition of hydroperoxides of the n-3 polyunsaturated  
265 fatty acids.<sup>26</sup> In contrast to 1-penten-3-one, 1-penten-3-ol increased steadily during the whole  
266 storage period after a lag phase of 3 days (Figure 4). Starting from day 6, the concentration of 1-  
267 penten-3-ol in both C2 and C1 had more than 50% reduction compared to the control. On day 13,  
268 the ranking order of the sample codes based on the concentration of 1-penten-3-ol was: Con<sup>a</sup> >  
269 C0<sup>b</sup>=C12<sup>b</sup>>C1+C12<sup>c</sup>=C2+C12<sup>c</sup>>C1<sup>d</sup>=C2<sup>d</sup>. Moreover, the concentration of 1-penten-3-ol in both

270 C1+C12 and C2+C12 were lower than their expected values, respectively ( $P<0.05$ ) (data not  
271 shown), indicating a “synergistic” effect of the combination of C1 or C2 with C12.

272 Corresponding to the increase in PV, a dramatic increase in 2,4-heptadienal for Con and C0 samples  
273 were observed after day 3, while a much slower increase was observed in all the ferulate treatments  
274 (Figure 5). At the end of storage, the ranking order of the sample codes based on the concentration  
275 of 2,4-heptadienal was:  $\text{Con}^a = \text{C0}^a > \text{C12}^b \geq \text{C1+C12}^{bc} = \text{C2+C12}^{bc} \geq \text{C1}^c = \text{C2}^c$ . A “synergistic” effect  
276 could not be determined due to the relatively high standard deviations.

277 Overall, for all the compounds quantified in the present study, C2 and C1 had the lowest  
278 concentration for all the volatile compounds followed by C2+C12, C1+C12, C12 and C0. Con had  
279 the highest concentration of the analyzed volatile compounds. This kind of difference was also  
280 observed in the PV results. These findings clearly demonstrated that lipophilization of ferulic acid  
281 with short alkyl chain lengths caused a dramatic increase of the antioxidant activity, but the  
282 antioxidant activity was sharply reduced with medium alkyl chain lengths in fish oil enriched milk.  
283 In addition, an indication of a “synergistic” antioxidant activity of ferulates with short and medium  
284 alkyl chain lengths were observed as shown in PV, hexanal and 1-penten-3-ol results.

285 **Change of tocopherol in the fish oil enriched milk samples.** Tocopherol of the milk samples  
286 mainly originated from the fish oil used in the study. The samples can be separated into three  
287 groups roughly based on the changing trend of  $\alpha$ -tocopherol contents (Figure 6). During the whole  
288 storage period, C1 and C2 treatments had the lowest  $\alpha$ -tocopherol reduction while  $\alpha$ -tocopherol  
289 levels in C1+C12 and C2+C12 were reduced moderately. C0, C12 and Con treatments had the  
290 highest reduction. There were significant differences in the content of  $\alpha$ -tocopherol among these 3  
291 groups starting from day 6 except C0 sample on day 6 ( $P<0.05$ ). In general, it showed that changes  
292 in  $\alpha$ -tocopherol levels were well correlated with the change in PV in the samples according to these



293 3 groups although there was some discrepancy within the highest reduction group (Con, C0, and  
294 C12). These findings were also in agreement with previous studies by Alemán et al.<sup>11</sup> and Sørensen  
295 et al.<sup>12</sup> There was not as high amount of  $\gamma$ -,  $\delta$ -tocopherol as  $\alpha$ -tocopherol in the samples. The  
296 reduction trend of  $\gamma$ - tocopherol was not as significant as for  $\alpha$ -tocopherol, and there was no obvious  
297 trend in the change of  $\delta$ -tocopherol (data not shown).

298 **Protein oxidation in fish oil enriched milk.** Changes in protein carbonyls in various milk samples  
299 are shown in Figure 7. Increase in carbonyls indicates protein oxidation during storage.<sup>27</sup> The  
300 amount of protein carbonyls in the control samples increased significantly during the storage from  
301 day 0 to day 13 ( $P<0.05$ ), corresponding to the dramatic increase of PV (Figure 1). The  
302 combination of C1 or C2 with C12 also showed inhibition effects for protein oxidation throughout  
303 the storage. Although C12 showed stronger lipid oxidation inhibition effects than C0, this was not  
304 the case for protein oxidation. C2 had the lowest carbonyls among all the treatments at the end of  
305 the storage ( $P<0.05$ ).

306 The protein oxidation observed by increase of carbonyls could be caused by the lipid oxidation  
307 products as these two reactions often occur simultaneously. Reactive oxygen species generated  
308 from lipid oxidation process can react with protein and cause protein oxidation.<sup>28</sup> C1 and C2  
309 showed strong inhibition of lipid oxidation and consequently protein oxidation. The contrast in lipid  
310 and protein oxidation results observed with C12 and C0 suggested that C0 could exhibit extra  
311 protection of protein from oxidation other than by reducing the lipid oxidation products. This could  
312 be related to the difference of hydrophobicity of C12 and C0 and thus their different partitioning in  
313 the milk sample as presented below. Further research such as loss of sulphhydryl content is needed to  
314 confirm the effects of lipophilized ferulate esters on protein oxidation. In addition, research on  
315 whether 1-pentent-3-one generated in the lipid oxidation process could react with protein could help  
316 to understand the mechanism of protein oxidation.

317 **Partitioning of lipophilized ferulates in the fish oil enriched milk.** Milk samples were physically  
318 separated into 3 different phases after ultracentrifugation: oil and emulsion, aqueous, and precipitate.  
319 The oil and emulsion phase was the top layer after ultracentrifugation and consisted of small  
320 amounts of oil and milk fat globule membrane. It was difficult to completely separate the oil from  
321 emulsion by the method used in this study, and this is the reason for the combination of them into  
322 one phase. The precipitate phase contained proteins both from the aqueous phase and the O/W  
323 interface layer. Approximately 61.9% of C12 partitioned in the oil and emulsion phase, followed by  
324 C2 (50.8%) and C1 (40.3%), while C0 (13.2%) had the lowest percentage in this phase ( $P<0.05$ )  
325 (Figure 8). This is in agreement with their corresponding hydrophobicity. In the aqueous phase, C0  
326 had the highest percentage (70%). Unexpectedly, C12 which is much more hydrophobic than either  
327 C1 or C2 had higher percentage in the aqueous phase than C1 or C2 ( $P<0.05$ ). However, this could  
328 be explained as some of C12 could have formed aggregates in the aqueous phase.<sup>29</sup> C1 and C2 with  
329 highest antioxidant effects had the lowest partitioning percentages in the aqueous phase. Some  
330 studies also found that antioxidants with high efficacies have low partitioning coefficients in the  
331 aqueous phase in simple O/W emulsions.<sup>30</sup> This study showed that this phenomenon could also  
332 happen in complex food systems such as fish oil enriched milk. Contrary to the aqueous phase, C1  
333 and C2 had relatively higher concentration in the precipitate phase than C12 and C0 ( $P<0.05$ ).  
334 Based on their antioxidant effects of these compounds, we may hypothesize that the higher  
335 partitioning of an antioxidant compound in the precipitate is related to a better antioxidant activity  
336 in fish oil enriched milk. Further research on other antioxidant phenolics to test this hypothesis is  
337 required. Protein plays an important role in milk as an emulsifier to stabilize the milk fat. The  
338 interaction of antioxidant with proteins and particularly those localized in the oil water interface  
339 could inhibit the lipid oxidation significantly, which is in agreement with the interfacial concept.<sup>31</sup>  
340 The combination of C1 or C2 with C12 seemed to change their relative partitioning behaviors. C12

341 decreased in the oil phase in both combination samples but increased in either the aqueous phase  
342 (C2+C12) or the precipitate phase (C1+C12) ( $P<0.05$ ). Both C1 and C2 had a slight decrease in the  
343 oil phase and a slightly higher increase in the protein phase in all the combination samples, which  
344 could increase the antioxidant effects of the combination and explain the “synergistic” effects  
345 observed by the antioxidant study. However, no statistical significances were found. An improved  
346 *in vivo* analysis method to determine the partitioning behavior among the oil, aqueous and the  
347 interface phases is required to provide a more clear picture.

348 **Localization of ferulic acid and lipophilized ferulates in fish oil enriched milk and the**  
349 **antioxidant activity mechanism.** Apart from the chemical properties of one particular antioxidant,  
350 its physical location in an emulsion is also important for its effectiveness.<sup>32</sup> C0 was the least  
351 effective with regard to lipid oxidation among all the antioxidants tested in this study. As it was  
352 reported that milk oil droplets were negatively charged,<sup>33</sup> this could be caused by the negative  
353 charge status of this compound in the milk. Consequently, C0 could be electrostatically repelled and  
354 have relatively lower concentration at the oil droplet surface than uncharged ferulates. In another  
355 study, researchers also showed that phenolic compounds with different electrical charge had an  
356 impact on their partitioning and the antioxidant activity in salmon oil-in-water emulsions.<sup>34</sup>

357 The difference of effectiveness among antioxidants is also probably due to the different  
358 hydrophobicity they have. The hydrophobicity of C12 is much higher than C1 or C2. Antioxidant  
359 with high hydrophobicity normally showed better antioxidant activity in O/W emulsions.<sup>7,31</sup>  
360 However, antioxidant activity of C1 or C2 was significantly higher than C12 from our observation  
361 which could be explained by the cut-off effect.<sup>9</sup> The critical chain length of a particular esterified  
362 phenolic acid differed in different food system.<sup>11</sup> There were currently three putative hypotheses to  
363 explain this cut-off phenomena: the “reduced mobility”, the “internalization”, and the “self-  
364 aggregation”, which was described in detail by Laguerre et al.<sup>9</sup> Figure 9 explains the assumed

365 localization of the ferulic acid and lipophilized ferulic acids in fish oil enriched milk based on the  
366 theory that lipid oxidation is initiated at the interface, measured lipid oxidation in the different  
367 sample codes and the partitioning study and how the combination of antioxidants could influence  
368 their localization. From this study, the “internalization” seemed to be the right mechanism to  
369 explain our results. C0 had the highest concentration in the aqueous phase which was showed in  
370 Figure 8 and illustrated in Figure 9A. C2 or C1 had the highest concentration in the precipitate and  
371 could have a relatively higher concentration in the interface due to their lower hydrophobicity  
372 compared to C12 (Figure 9B). A significant amount of C12 could be internalized inside the oil  
373 droplets due to its high hydrophobicity, which was indirectly confirmed by the high percentage of  
374 C12 detected in the oil and emulsion fraction (Figure 9C). Partitioning studies in simplified O/W  
375 emulsion system with rutin laurate and rutin palmitate showed that very low concentrations were  
376 detected in the aqueous phase of the different two or multiple phases systems (buffer / oil phases,  
377 buffer / emulsifier phases and emulsion).<sup>35</sup> This could also support our assumption for the location  
378 of C12 in this study. There is an equilibrium distribution of one particular antioxidant among the oil,  
379 interfacial and aqueous phases. It is assumed that the concentration of each compound in the  
380 interface decides the antioxidant activity because the oil droplet interface is where the lipid  
381 oxidation occurs.<sup>36</sup> The combination of C12 with C1 or C2 could disturb the equilibrium  
382 distribution of C1 or C2 between the oil and interface as also observed in other studies between  
383 tocopherol and caffeates in model emulsions.<sup>37</sup> The combination of C1 or C2 with C12 could lead  
384 to a relatively higher concentration of C1 or C2 in the interface than when C1 or C2 was added in  
385 the fish oil enriched milk emulsion alone (Figure 9D). The higher concentration of C1 or C2 at the  
386 interface can lead to better antioxidant effects, which could explain the indication of the  
387 “synergistic” antioxidant effect observed in this study. In another study, Panya et al.<sup>36</sup> showed that  
388 the equilibrium distribution of antioxidant compound in the oil, interface and aqueous phase could

389 be broken by adding a surfactant Tween-20 and the antioxidant activity was changed accordingly in  
390 O/W emulsions. The present study demonstrated that antioxidant-antioxidant interaction could  
391 influence their antioxidant activity in fish oil enriched milk, which could also be related to the  
392 change of partitioning coefficients of the antioxidants at the interface. The mechanism of this kind  
393 of interaction is still not clear. Challenges include the quantification of distribution of antioxidant  
394 compounds in the oil, interfacial, and aqueous phases *in vivo* in a complicated food system like fish  
395 oil enriched milk emulsion. In addition, there are also various compounds in a food system that  
396 could influence the partitioning of antioxidant in the interface. More research is needed to study the  
397 particular factors that could influence the localization of antioxidants and further clarify the  
398 “internalization” or “localization” mechanism.

399 Although partitioning of an antioxidant in the interface of an emulsion system is relevant for its  
400 antioxidant activity in inhibiting lipid oxidation, this could not apply to the protein oxidation. Lipid  
401 oxidation could originate from the interface and generate primary and secondary lipid oxidation  
402 products that react with protein and cause the induction of protein oxidation. An antioxidant such as  
403 C1 or C2 that localizes at the interface may inhibit lipid oxidation efficiently and thus protein  
404 oxidation. However, antioxidant like C0 which has high concentration in aqueous phase and  
405 consequently less efficient lipid oxidation activity could still inhibit protein oxidation. Further  
406 research on how C0 could retard the reaction between lipid oxidation products and proteins would  
407 help to explain the mechanism.

#### 408 **Funding**

409 The financial support provided by China Scholarship Council (No. 201608350049) for Xujian Qiu’s  
410 visit to DTU Food is acknowledged.

#### 411 **Notes**

412 The authors declare no competing financial interest.

413 **ACKNOWLEDGEMENTS**

414 We thank Maritex (subsidiary of TINE BA, Norway) for providing the fish oil to our research.

415 **REFERENCES**

- 416 (1) Calder, P. C. n-3 Fatty acids and cardiovascular disease: evidence explained and mechanisms  
417 explored. *Clin. Sci.* **2004**, *107* (1), 1–11.
- 418 (2) Calder, P. C. N-3 Fatty Acids, Inflammation and Immunity: New Mechanisms To Explain Old  
419 Actions. *Proc. Nutr. Soc.* **2013**, *72* (3), 326–336.
- 420 (3) Frankel, E. N. N. Lipid Oxidation. *Prog. Lipid Res.* **1980**, *19* (1–2), 488.
- 421 (4) Let, M. B.; Jacobsen, C.; Frankel, E. N.; Meyer, A. S. Oxidative flavour deterioration of fish oil  
422 enriched milk. *Eur. J. Lipid Sci. Technol.* **2003**, *105* (9), 518–528.
- 423 (5) Jacobsen, C.; Sørensen, A. -D.M.; Nielsen, N. S. Stabilization of omega-3 oils and enriched  
424 foods using antioxidants. In *Food Enrichment with Omega-3 Fatty Acids*; Jacobsen, C., Nielsen, N.  
425 S., Horn, A.F., Sørensen, A. -D.M., Eds.; Woodhead Publishing Limited: Cambridge, UK, 2013; pp  
426 130–149.
- 427 (6) Porter, W. L. Recent Trends in Food Applications of Antioxidants. In *Autoxidation in Food and*  
428 *Biological Systems*; Simic, M. G., Karel, M., Eds.; Springer US: Boston, MA, 1980; pp 295–365.
- 429 (7) Porter, W. L., Black, E. D., Drolet, A. M. Use of polyamide oxidative fluorescence test on lipid  
430 emulsions: contrast in relative effectiveness of antioxidants in bulk versus dispersed systems. *J.*  
431 *Agric. Food Chem.* **1989**, *37* (3), 615–624.
- 432 (8) Laguerre, M.; Bayrasy, C.; Lecomte, J.; Chabi, B.; Decker, E. A.; Wrutniak-Cabello, C.;  
433 Cabello, G.; Villeneuve, P. How to boost antioxidants by lipophilization? *Biochimie* **2013**, *95* (1),  
434 20–26.

- 435 (9) Laguerre, M.; Sørensen, A.-D. M.; Bayrasy, C.; Lecomte, J.; Jacobsen, C.; Decker, E. A.;  
436 Villeneuve, P. Role of hydrophobicity on antioxidant activity in lipid dispersions, From the polar  
437 paradox to the cut-off theory. In *Lipid Oxidation: Challenges in Food Systems*; Logan A., Nienaber  
438 U., Pan X., Eds.; AOCS Press: Urbana, IL, 2013; pp 261-295.
- 439 (10) Costa, M.; Losada-Barreiro, S.; Paiva-Martins, F.; Bravo-Diaz, C. Optimizing the efficiency of  
440 antioxidants in emulsions by lipophilization: tuning interfacial concentrations. *RSC Adv.* **2016**, *6*  
441 (94), 91483–91493.
- 442 (11) Alemán, M.; Bou, R.; Guardiola, F.; Durand, E.; Villeneuve, P.; Jacobsen, C.; Sørensen, A. -D.  
443 M. Antioxidative effect of lipophilized caffeic acid in fish oil enriched mayonnaise and milk. *Food*  
444 *Chem.* **2015**, *167*, 236–244.
- 445 (12) Sørensen, A.-D. M.; Petersen, L. K.; de Diego, S.; Nielsen, N. S.; Lue, B.-M.; Yang, Z.; Xu, X.;  
446 Jacobsen, C. The antioxidative effect of lipophilized rutin and dihydrocaffeic acid in fish oil  
447 enriched milk. *Eur. J. Lipid Sci. Technol.* **2012**, *114* (4), 434–445.
- 448 (13) Sørensen, A.-D. M.; Lyneborg, K. S.; Villeneuve, P.; Jacobsen, C. Alkyl chain length impacts  
449 the antioxidative effect of lipophilized ferulic acid in fish oil enriched milk. *J. Funct. Foods* **2015**,  
450 *18*, Part B, 959–967.
- 451 (14) Sørensen, A.-D. M.; Durand, E.; Laguerre, M.; Bayrasy, C.; Lecomte, J.; Villeneuve, P.;  
452 Jacobsen, C. Antioxidant Properties and Efficacies of Synthesized Alkyl Caffeates, Ferulates, and  
453 Coumarates. *J. Agric. Food Chem.* **2014**, *62* (52), 12553–12562.
- 454 (15) Let, M. B.; Jacobsen, C.; Sørensen, A.-D. M.; Meyer, A. S. Homogenization Conditions Affect  
455 the Oxidative Stability of Fish Oil Enriched Milk Emulsions: Lipid Oxidation. *J. Agric. Food Chem.*  
456 **2007**, *55* (5), 1773–1780.



- 457 (16) Bligh, E. G.; Dyer, W. J. A rapid method of total lipid extraction and purification. *Can. J.*  
458 *Biochem. Physiol.* **1959**, *37* (8), 911–917.
- 459 (17) Shantha, N. C.; Decker, E. A. Rapid, sensitive, iron-based spectrophotometric methods for  
460 determination of peroxide values of food lipids. *J. AOAC Int.* **1994**, *77* (2), 421–424.
- 461 (18) AOCS Official Method Ce 1b-89. *Fatty Acid Composition by GC. Marine Oils.* AOCS,  
462 Champaign, IL, 1998.
- 463 (19) AOCS Official Method Ce 8-89. *Determination of tocopherols and tocotrienols in vegetable*  
464 *oils and fats by HPLC.* AOCS, Champaign, IL, 1997.
- 465 (20) Levine, R. L.; Garland, D.; Oliver, C. N.; Amici, A.; Climent, I.; Lenz, A. G.; Ahn, B. W.;  
466 Shaltiel, S.; Stadtman, E. R. Determination of carbonyl content in oxidatively modified proteins.  
467 *Methods Enzymol.* **1990**, *186*, 464–478.
- 468 (21) Fenaille, F.; Parisod, V.; Visani, P.; Populaire, S.; Tabet, J. C.; Guy, P. A. Modifications of  
469 milk constituents during processing: A preliminary benchmarking study. *Int. Dairy J.* **2006**, *16* (7),  
470 728–739.
- 471 (22) Jacobsen, C.; Meyer, A. S.; Adler-Nissen, J. Oxidation mechanisms in real food emulsions:  
472 method for separation of mayonnaise by ultracentrifugation. *J. Food Lipids* **1998**, *5* (2), 87–101.
- 473 (23) Guyot, B.; Gueule, D.; Pina, M.; Farines, V.; Farines, M. Enzymatic synthesis of fatty esters in  
474 5-caffeoyl quinic acid. *Eur. J. Lipid Sci. Technol.* **2000**, 93–96.
- 475 (24) Sørensen, A. -D. M.; Baron, C. P.; Let, M. B.; Brüggemann, D. A.; Pedersen, L. R. L.;  
476 Jacobsen, C. Homogenization conditions affect the oxidative stability of fish oil enriched milk

- 477 emulsions: Oxidation linked to changes in protein composition at the oil-water interface. *J. Agric.*  
478 *Food Chem.* **2007**, *55* (5), 1781–1789.
- 479 (25) Venkateshwarlu, G.; Let, M. B.; Meyer, A. S.; Jacobsen, C. Chemical and Olfactometric  
480 Characterization of Volatile Flavor Compounds in a Fish Oil Enriched Milk Emulsion. *J. Agric.*  
481 *Food Chem.* **2004**, *52* (2), 311–317.
- 482 (26) Olsen, E.; Vogt, G.; Saarem, K.; Greibrokk, T.; Nilsson, A. Autoxidation of cod liver oil with  
483 tocopherol and ascorbyl palmitate. *J. Am. Oil Chem. Soc.* **2005**, *82* (2), 97–103.
- 484 (27) Soyer, A.; Özalp, B.; Dalmış, Ü.; Bilgin, V. Effects of freezing temperature and duration of  
485 frozen storage on lipid and protein oxidation in chicken meat. *Food Chem.* **2010**, *120* (4), 1025–  
486 1030.
- 487 (28) Heinecke, J. W.; Li, W.; Daehnke, H. L.; Goldstein, J. A. Dityrosine, a specific marker of  
488 oxidation, is synthesized by the myeloperoxidase-hydrogen peroxide system of human neutrophils  
489 and macrophages. *J. Biol. Chem.* **1993**, *268* (6), 4069–4077.
- 490 (29) Laguerre, M.; Bayrasy, C.; Lecomte, J.; Chabi, B.; Decker, E. A.; Wrutniak-Cabello, C.;  
491 Cabello, G.; Villeneuve, P. How to boost antioxidants by lipophilization? *Biochimie.* **2013**, *95*(1):  
492 20–26.
- 493 (30) Laguerre, M.; López Giraldo, L. J.; Lecomte, J.; Figueroa-Espinoza, M.-C.; Baréa, B.; Weiss,  
494 J.; Decker, E. A.; Villeneuve, P. Chain Length Affects Antioxidant Properties of Chlorogenate  
495 Esters in Emulsion: The Cutoff Theory Behind the Polar Paradox. *J. Agric. Food Chem.* **2009**, *57*  
496 (23), 11335–11342.
- 497 (31) Frankel, E. N.; Huang, S.-W.; Kanner, J.; German, J. B. Interfacial phenomena in the  
498 evaluation of antioxidants: bulk Oils vs emulsions. *J. Agric. Food Chem.* **1994**, *42* (5), 1054–1059.

499 (32) McClements, D.J.; Decker, E.A. Lipid oxidation in oil-in-water emulsions: impact of  
500 molecular environment on chemical reactions in heterogeneous food systems. *J. Food Sci.* **2000**, *65*  
501 (8), 1270–1282.

502 (33) Ménard, O.; Ahmad, S.; Rousseau, F.; Briard-Bion, V.; Gaucheron, F.; Lopez, C. Buffalo vs.  
503 cow milk fat globules: size distribution, zeta-potential, compositions in total fatty acids and in polar  
504 lipids from the milk fat globule membrane. *Food Chem.* **2010**, *120* (2), 544–551.

505 (34) Mei, L.; McClements, D. J.; Decker, E. A. Lipid oxidation in emulsions as affected by charge  
506 status of antioxidants and emulsion droplets. *J. Agric. Food Chem.* **1999**, *47* (6), 2267–2273.

507 (35) Lue, B.-M.; Sørensen, A.-D. M.; Jacobsen, C.; Guo, Z.; Xu, X. Antioxidant efficacies of rutin  
508 and rutin esters in bulk oil and oil-in-water emulsion. *Eur. J. Lipid Sci. Technol.* **2016**, *119*(4), 1–  
509 15.

510 (36) Panya, A.; Laguerre, M.; Bayrasy, C.; Lecomte, J.; Villeneuve, P.; McClements, D. J.; Decker,  
511 E. A. An Investigation of the Versatile Antioxidant Mechanisms of Action of Rosmarinate Alkyl  
512 Esters in Oil-in-Water Emulsions. *J. Agric. Food Chem.* **2012**, *60* (10), 2692–2700.

513 (37) Sørensen, A.-D. M.; Villeneuve, P.; Jacobsen, C. Alkyl caffeates as antioxidants in O/W  
514 emulsions: Impact of emulsifier type and endogenous tocopherols. *Eur. J. Lipid Sci. Technol.* **2017**,  
515 *119* (6), e201600276.

516

517

518

519

520 **Figure captions:**

521 Figure 1. Changes of PV in fish oil enriched milk samples with ferulic acid and different  
522 lipophilized ferulates during 13 days storage at 5 °C. Data points and error bars represent means ±  
523 standard deviations (n=2). Con, control; C0, ferulic acid; C1, methyl ferulate; C2, ethyl ferulate;  
524 C12, dodecyl ferulate.

525 Figure 2. Change of hexanal in fish oil enriched milk samples with ferulic acid and different  
526 lipophilized ferulates during 13 days storage at 5 °C. Data points and error bars represent means ±  
527 standard deviations (n=3). Con, control; C0, ferulic acid; C1, methyl ferulate; C2, ethyl ferulate;  
528 C12, dodecyl ferulate.

529 Figure 3. Change of 1-penten-3-one in fish oil enriched milk samples with ferulic acid and different  
530 lipophilized ferulates during 13 days storage at 5 °C. Data points and error bars represent means ±  
531 standard deviations (n=3). Con, control; C0, ferulic acid; C1, methyl ferulate; C2, ethyl ferulate;  
532 C12, dodecyl ferulate.

533 Figure 4. Change of 1-penten-3-ol in fish oil enriched milk samples with ferulic acid and different  
534 lipophilized ferulates during 13 days storage at 5 °C. Data points and error bars represent means ±  
535 standard deviations (n=3). Con, control; C0, ferulic acid; C1, methyl ferulate; C2, ethyl ferulate;  
536 C12, dodecyl ferulate.

537 Figure 5. Change of 2,4-hepatadienal in fish oil enriched milk samples with ferulic acid and  
538 different lipophilized ferulates during 13 days storage at 5 °C. Data points and error bars represent  
539 means ± standard deviations (n=3). Con, control; C0, ferulic acid; C1, methyl ferulate; C2, ethyl  
540 ferulate; C12, dodecyl ferulate.

541

542 Figure 6. Changes of  $\alpha$ -tocopherol in fish oil enriched milk samples with ferulic acid and different  
543 lipophilized ferulates during 13 days storage at 5 °C. Data points and error bars represent means  
544  $\pm$  standard deviations (n=2). Con, control; C0, ferulic acid; C1, methyl ferulate; C2, ethyl ferulate;  
545 C12, dodecyl ferulate.

546 Figure 7. Changes of carbonyl content in fish oil enriched milk samples with ferulic acid and  
547 different lipophilized ferulates during 13 days storage at 5 °C. Data points and error bars represent  
548 means  $\pm$  standard deviations (n=3). Con, control; C0, ferulic acid; C1, methyl ferulate; C2, ethyl  
549 ferulate; C12, dodecyl ferulate.

550 Figure 8. Partitioning of ferulic acid and lipophilized ferulates into 3 different phases in the milk  
551 samples: Oil and emulsion, aqueous and precipitate. Data points and error bars represent means  $\pm$   
552 standard deviations (n=3). C0, ferulic acid; C1, methyl ferulate; C2, ethyl ferulate; C12, dodecyl  
553 ferulate.

554 Figure 9. A schematic illustration of assumed localization and interaction of ferulic acid and  
555 different lipophilized ferulates in fish oil enriched milk: (A) ferulic acid is mainly localized at the  
556 aqueous phase; (B) ferulates with short alkyl chain lengths such as methyl ferulate and ethyl  
557 ferulate have high concentration at the interface; (C) Ferulates with medium alkyl chain lengths are  
558 highly hydrophobic and are mainly in the oil; (D) Interaction between ferulates with short and  
559 medium alkyl chain lengths can change the equilibrium distribution of ferulates with short alkyl  
560 chain length and increase its concentration at the interface.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6






Figure 7




Figure 8








Figure 9


## Graphic for table of contents































**Partitioning  
& Interaction**

**Protein &  
Lipid  
Oxidation**

