Vitamin D-fortified foods improve wintertime vitamin D status in women of Danish and Pakistani origin living in Denmark: a randomized controlled trial

Grønborg, Ida Marie; Tetens, Inge; Christensen, Tue; Andersen, Elisabeth Wreford; Jakobsen, Jette; Kiely, Mairead; Cashman, Kevin D.; Andersen, Rikke

Published in:
European Journal of Nutrition

Link to article, DOI:
[10.1007/s00394-019-01941-6](https://doi.org/10.1007/s00394-019-01941-6)

Publication date:
2020

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Grønborg, I. M., Tetens, I., Christensen, T., Andersen, E. W., Jakobsen, J., Kiely, M., Cashman, K. D., & Andersen, R. (2020). Vitamin D-fortified foods improve wintertime vitamin D status in women of Danish and Pakistani origin living in Denmark: a randomized controlled trial. *European Journal of Nutrition*, 59, 741–753. <https://doi.org/10.1007/s00394-019-01941-6>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Vitamin D fortified foods improve wintertime vitamin D status in women of Danish and Pakistani origin living in Denmark: A randomized controlled trial.

Ida M Grønberg¹; Inge Tetens^{1,2}; Tue Christensen¹; Elisabeth W Andersen^{3,4}; Jette Jakobsen¹; Mairead Kiely⁵; Kevin D Cashman^{5,6}; Rikke Andersen¹.

¹National Food Institute, Technical University of Denmark, Lyngby, Denmark; ²Vitality - Centre for good older lives, Department of Nutrition, Exercise and Sports, University of Copenhagen; ³Danish Cancer Society, Section for Statistics and Pharmacoepidemiology, Copenhagen, Denmark; ⁴Institute of Applied Mathematics and Computer Science, Technical University of Denmark, Lyngby, Denmark; ⁵Cork Centre for Vitamin D and Nutrition Research, School of Food and Nutritional Sciences and ⁶Department of Medicine, University College Cork, Cork, Ireland.

Corresponding author: Ida M Grønberg, Division of Nutrition, National Food Institute, Technical University of Denmark, Lyngby, Denmark. Phone: +4520997801. E-mail: idadamag@food.dtu.dk. **Orcid ID:** 0000-0001-7316-1073.

Running title: Food-based randomized controlled trial with vitamin D fortified foods

Registered at ClinicalTrials.gov with identifier: NCT02631629.

Financial support: This research was undertaken by the National Food Institute, Technical University of Denmark (DTU) and the study is a part of the European collaborative project “Food-based solutions for eradication of vitamin D deficiency and health promotion throughout the life cycle - ODIN”, which was funded by the European Commission (FP7/2007-2013) under grant agreement no. 613977 (ODIN) and by the National Food Institute, Technical University of Denmark. The cheese and yoghurt products were produced and provided free of charge by FrieslandCampina. The eggs were produced and provided free of cost by Hedegaard Agro, including chicken feed produced by Dava Foods, the crisp bread was produced by Smørum konditori (confectionary) with ingredients provided free of cost by Lantmännen cerialia and vitamin D₃ supplied by DSM Nutritional Products.

Abbreviations:

25(OH)D – Serum 25-hydroxyvitamin D

PTH – Parathyroid Hormone

RCT – Randomized Controlled trial

ODIN – Food-based solutions for optimal vitamin D nutrition and health through the life cycle

SD – Standard Deviation

CV – Coefficient of variance $((SD/mean)*100)$

UL – Tolerable Upper Intake Level

FFQ – Food Frequency Questionnaire

OPTIFORD – Towards a strategy for optimal vitamin D fortification

UCC – University College Cork

LC-MS/MS – Liquid chromatography-tandem mass spectrometry

VDSP – Vitamin D Standardization Program

ANOVA – Analysis of variance

ANCOVA – Analysis of co-variance

CI – Confidence Interval

ABSTRACT

Purpose: Low vitamin D status is prevalent worldwide. We aim to investigate the effect of vitamin D fortification on serum 25-hydroxyvitamin D (25(OH)D) concentration in women of Danish and Pakistani origin at risk of vitamin D deficiency.

Methods: A 12-week randomized, double-blinded, placebo-controlled intervention trial during winter time, designed to provide 20 µg vitamin D₃/day through fortified yoghurt, cheese, eggs and crisp bread and assess the change in serum 25(OH)D. Participants were 143 women of Danish and Pakistani origin, living in Denmark, randomized into four groups, stratified by ethnicity.

Results: Mean (SD) baseline 25(OH)D concentrations among women of Danish and Pakistani origin were 49.6 (18) and 46.9 (22) nmol/L, respectively ($P = 0.4$). While 9 % of Danish women had 25(OH)D < 30 nmol/L, the prevalence among women of Pakistani origin was 24 %. Median (IQR) vitamin D intake among Danish and Pakistani women at endpoint was 32.0 (27.0, 34.4) µg/d and 24.2 (19.2, 30.8) µg/d, respectively. Endpoint serum 25(OH)D increased in fortified groups to 77.8 (14) nmol/L among Danish women and 54.7 (18) nmol/L among women of Pakistani origin ($P < 0.01$). At endpoint, 0 % in the Danish fortified group and 3 % in the Pakistani fortified group had 25(OH)D < 30 nmol/L, compared with 23 % and 34 % in their respective control groups.

Conclusions: Vitamin D fortification of four different foods for 12-weeks during winter was effective in increasing serum 25(OH)D and reducing the prevalence of very low vitamin D status among women of Danish and Pakistani origin.

Keywords: Vitamin D; Food-based RCT; ODIN; women of Pakistani origin; women of Danish origin; fortified foods.

21 INTRODUCTION

22 Very low and low vitamin D status (reflected by serum 25-hydroxyvitamin D [25(OH)D] concentrations <
23 30 and < 50 nmol/L, respectively [1, 2]) are prevalent amongst the general population in countries of
24 northern latitudes [3], as well as being a more worldwide concern [4]. Notably, many immigrants (first
25 generation immigrants and descendants) living in Denmark, and other Nordic countries, are at a higher risk
26 of vitamin D deficiency compared to the native residents [3, 5–8]. Low vitamin D status has been linked with
27 risk of various chronic diseases, and in particular, with adverse bone health outcomes [1, 2, 9–11].

28 Denmark, located at 55° North, experiences a five-month long ‘vitamin D winter’, during which there is
29 increased emphasis on dietary supply of vitamin D [12]. However, food sources of vitamin D are sparse
30 (fish, meat, eggs, cheese) and their consumption patterns irregular [13]. Data from the most recent Danish
31 National Survey of Dietary Habits and Physical Activity (DANSDA 2011-13) shows that median intakes of
32 vitamin D are 3-4 µg/day (without supplement contribution) [13], considerably lower than the Average
33 Requirement of 7.5 µg/day for the Nordic region [10]. In Denmark, while vitamin D supplement use is only
34 recommended for populations at risk of deficiency; infants, pregnant women, elderly and individuals with
35 dark skin [14], the proportion in the general population taking a vitamin D supplement is relatively high, e.g.,
36 50-60% of the adult females [15, 16]. However, while vitamin D supplementation is an effective approach
37 for increasing vitamin D status in those individuals who consume them regularly, it has limitations as a
38 strategy at a population level due to a high proportion of non-users as well as concerns about potential risk of
39 excessive intake amongst high users [17].

40 Food-based strategies aiming at improving vitamin D intake and status across the population, such as vitamin
41 D food fortification, have been highlighted as being of high potential as a public health measure in Europe
42 [17–19]. Finland have been fortifying all fluid dairy products and margarine for more than 10 years and
43 consequently experienced increases of the Finnish population vitamin D status [20]. In countries such as
44 USA, Canada, food fortification with vitamin D have already been introduced using several different
45 fortification vehicles such as milk, orange juice and margarine [21]. In India solutions such as fortification of

46 commonly consumed staple foods are currently being investigated [22]. Furthermore, while many of the
47 environmental factors that contribute to the elevated risk of vitamin D deficiency in populations, such as
48 latitude, skin color, and cultural clothing practices are not modifiable, in contrast, dietary supply of vitamin
49 D is an important modifiable factor, again emphasizing the need for food-based strategies to offset low
50 intakes [23].

51 Vitamin D fortification of food has not yet been widely implemented and accepted in Denmark, despite
52 voluntary fortification of certain foods being permissible for over a decade [19, 24]. Low-dose fortification
53 of several foods rather than higher dose fortification in few foods has been proposed as an effective and safe
54 strategy on the population level [18, 25]. Yet, randomized controlled trial (RCT) data on the effectiveness
55 and safety of this approach, especially in at-risk population subgroups, is lacking despite the importance of
56 such data in informing food fortification policy. Thus, we aimed to investigate the effect of low-dose,
57 vitamin D₃ fortification of a number of commonly consumed foods (yoghurt, cheese, eggs and crisp bread)
58 on vitamin D status in a population of adult women of Danish and Pakistani origin at increased risk of
59 vitamin D deficiency, using a 12-week, winter-based RCT.

SUBJECTS AND METHODS

Study design

The *ODIN-FOOD* study was part of the European Commission-funded large scale collaborative ODIN (Food-based solutions for optimal vitamin D nutrition and health through the life cycle) Project. The *ODIN-FOOD* study was a three-month randomized double-blinded, placebo-controlled, intervention trial carried out during the winter months (January-March) of 2016, and enrolled a total of 143 women of Danish and Pakistani origin, aged between 18 and 50 years. Immigrant women in Denmark are considered an at risk group for vitamin D deficiency [5, 7]. The risk extends to the ethnic Danish female population in a modern work-life setting with a sedentary lifestyle, low UVB exposure and a low habitual vitamin D intake, particularly if they avoid or limit fish intake [13, 26]. The trial was performed during the winter months to minimize interference from UVB-induced cutaneous synthesis of vitamin D₃. Participants were randomized into four groups, stratified for ethnicity (two groups with women of Danish origin and two groups with women of Pakistani origin). The randomization sequence was generated by a researcher not involved in the project using block randomization with a block size of four within each ethnic group (Fig. 1). The subjects were given four different fortified foods aiming to contribute an additional 20 µg/day of vitamin D₃ or equivalent non-fortified foods (placebo). Participants were seen before the start of intervention and at the end of 3-month intervention, and at both visits non-fasting blood were drawn, anthropometrics, dietary vitamin D intake and muscle strength was measured. The main endpoint was the change in serum 25(OH)D concentration. Secondary endpoints included anthropometric measures and dietary intake of vitamin D.

The intervention was double-blinded, and the study foods were color and letter coded. The blinding was managed by a researcher not involved in the project. Both the participants and the researchers working within the project, during the intervention and subsequent analyses, were all blinded until the statistical analyses were completed.

82 Sample size of 143 women of two ethnicities was based on 90% power ($\alpha = 5\%$) to detect a change in
83 serum 25(OH)D concentration of 20 nmol/L in the treatment groups with a standard deviation (SD) of 23
84 nmol/L, and included a drop-out rate of 20 %.

85 *Subjects*

86 We recruited women of Pakistani origin (first generation immigrants or descendants) and women of Danish
87 origin from the Copenhagen area, city and suburbs (Denmark, 55°N). The recruitment was done by e-mail,
88 advertising, networking and interactions with local community groups, media, and social and cultural
89 initiatives in the Copenhagen area, as well as visiting local shops, libraries, mosques and women's societies.
90 Eligible women were invited to information meetings in which the study procedures were explained by a
91 researcher from the project group. Written informed consent was obtained from all participants on
92 enrolment. Inclusion criteria were a low consumption of fish and fish products (less than weekly), a low
93 frequency of use of vitamin D-containing supplements (less than weekly), no use of tanning facilities, no
94 planned sun-holiday (to a location more southerly than 47°N) between October 2015 and May 2016. There
95 was no upper limit on the vitamin D dose of the supplements as long as participants discontinued them
96 during the study period. Exclusion criteria were pregnancy and breastfeeding, menopause, non-Danish
97 speakers, serious diseases (cancer, server liver or kidney insufficiencies, sarcoidosis and other granulomatous
98 diseases) and medicines affecting vitamin D metabolism (steroids, antiepileptic, thyroid hormones,
99 bisphosphonates, estrogen).

100 *Fortification vehicles*

101 The intervention foods of choice were low-fat Milner cheese (gouda) and yoghurt (plain) produced and
102 provided free of cost by FrieslandCampina in the Netherlands, eggs (Livskraft) produced and provided free
103 of cost by Hedegaard in Denmark, and whole grain crisp bread produced by Smørum Konditori
104 (confectionary) using ingredients provided free of cost by Lantmännen Cerealia, Denmark, the vitamin D₃
105 for the crisp bread was supplied by DSM nutritional products, Switzerland. The study foods were chosen
106 because they are commonly consumed by both ethnic groups. Dietary calculations and pilot taste tests were

107 carried out to ensure the acceptance of the products. The taste tests were performed among 12 women of
108 Danish and Pakistani origin. The majority of the foods were low in fat and they are considered suitable
109 substitutions for the participants' habitual intake of these food products (Table 1). The participants were
110 given either placebo foods or vitamin D-fortified foods (aimed at providing approximately 20 µg/day of
111 additional vitamin D₃) [19] at no cost for the participants.

112 A daily dose of 20 µg, together with the contribution from habitual intake (~3-4 µg/d [6]), would be expected
113 to maintain winter serum 25(OH)D concentration above of 30 nmol/L in the vast majority of individuals [27]
114 and it allows for a large margin of safety in relation to the Tolerable Upper Intake Level (UL) of 100 µg/day
115 [19, 28]. The fortified and non-fortified foods have the same fat content and comparable content of nutrients,
116 except for the vitamin D concentration (Table 1). The packaging of the food products was identical, except
117 for color and letter coding added to distinguish the fortified from the placebo foods.

118 *Laboratory analysis of food samples*

119 The vitamin D₃/D₂ and 25-hydroxyvitamin D₃/D₂ content of food were analyzed at the National Food
120 Institute at the Technical University of Denmark using modifications of a sensitive liquid chromatography-
121 tandem mass spectrometry (LC-MS/MS), as described elsewhere [29]. The analysis was performed in a
122 laboratory accredited according to ISO17025, and has a limit of quantification at 0.01 µg/100 g, and a
123 precision < 10% (CV). For crisp bread and eggs, eight and six batches, respectively, were produced and all
124 were analyzed. Only one batch was produced for yoghurt and cheese and for this the stability was controlled.
125 Analysis confirmed there was no decrease of vitamin D during the 12 weeks of intervention. Total vitamin D
126 activity of each food was calculated as the vitamin D₃ content of the food plus the 25-hydroxyvitamin D₃
127 content x 2.5. The conversion factor of 2.5 was used as a conservative estimate based on factor of 5 as
128 currently used [30] and a factor of 1.5 recently obtained in an intervention study in Denmark [31].

129 *Intervention*

130 After the baseline visit, participants were each given food for two weeks (1 egg per day, 150 g (1 small pot)
131 yoghurt, 60 g (2 slices) of low fat Gouda cheese and 1 crisp bread per day). The participants were allowed to

freely plan how they distributed the provided foods over a week as long as they consumed the designated 7 eggs, 7 portions of yoghurt, 7 x 60 g cheese and 7 crisp breads per week. To maintain body weight, participants were advised to substitute some of their regularly consumed foods with the study foods. Fresh foods were distributed every two weeks. Adherence/compliance was monitored on a weekly basis by a user-friendly printed questionnaire in which the participant would mark each food item on a picture once consumed. Adherence was estimated individually by dividing the individual amount of food received by self-reported adherence, expressed as a percentage. Due to packaging, an individual would sometimes receive more than they needed for that period, and were asked not to consume the surplus after fulfilling their weekly amount, however, if they did consume the extra foods, this was accounted for. Thus, individual adherence could in some cases be > 100 %.

Measurements

Participants were examined twice at the National Food Institute at the Technical University of Denmark, including a baseline visit in January and an endpoint visit in April 2016. At each visit, a 40 mL non-fasting blood sample was obtained from each participant by an experienced phlebotomist or nurse.

Anthropometric measures were completed with participants wearing thin/light clothing, no shoes and after emptying their bladder. The measures were height (wall mounted stadiometer, seca, Hamburg, Germany) and weight, waist-hip circumference (standard tape measure) and body composition (Tanita BC 418 MA, Tokyo, Japan).

At baseline, participants completed two questionnaires, a general background questionnaire, assessing the health, lifestyle and sun habits, and other factors affecting the vitamin D status and a semi-quantitative Food Frequency Questionnaire (FFQ) estimating the average intake of vitamin D and calcium. The FFQ was a retrospective questionnaire asking about the participants' habitual dietary intake of vitamin D-rich foods over the 3 preceding months. The FFQ used in the *ODIN-FOOD* study was developed from existing questionnaires used in previous RCT's (OPTIFORD and VitmaD) at the National Food Institute [5, 24].

156 All questionnaires were self-administered, however to ensure that all participants completed the
157 questionnaire and to avoid misunderstandings, cultural as well as language-related, we set up a questionnaire
158 room, where all participants were introduced to the questionnaires and offered help when needed. The
159 questionnaires contained ethnic-specific questions regarding foods that contribute vitamin D and calcium.
160 All FFQ's were administered during the participant's first visit in January 2016. The questionnaire contained
161 the 8 food groups (Fish, meat, milk and milk products, egg, cheese, bread, fats and pulses) contributing to the
162 majority of dietary vitamin D (98 %) and calcium (71 %) [13]. Further questions were asked to estimate
163 intake of vitamin D-containing supplements prior to the study start. The questionnaire took between 30 and
164 50 minutes to complete and all participants completed the questionnaire.

165 Estimates of vitamin D intake were made from the data obtained from the FFQ matched to specific foods and
166 recipes made from the Danish food composition database (version 7) using the individually reported portion
167 sizes [32]. The total vitamin D intake for each participant prior to the study start was estimated as the sum of
168 dietary vitamin D and contribution from personal supplements, if consumed.

169 *Laboratory analysis of blood samples*

170 Serum concentrations of total 25(OH)D (i.e., 25(OH)D₂ plus 25(OH)D₃) of all serum samples (baseline and
171 endpoint visits) were measured at the Cork Centre for Vitamin D and Nutrition Research, University College
172 Cork (UCC), using the ODIN core LC-MS/MS analytical platform for serum 25(OH)D, described in detail
173 elsewhere [33]. The intra-assay CV of the method was < 5 % for all 25(OH)D metabolites, whilst the inter-
174 assay CV was < 6 %. The LC-MS/MS method at UCC is certified by the Center for Disease Control and
175 Prevention's *Vitamin D Standardization Certification Program*.

176 The biochemical analyses of serum calcium and PTH were performed at the University Hospital Aarhus.
177 Total serum calcium had an analysis precision of ± 0.032 mmol/L (standard deviation [SD]) at a
178 concentration of 2.161 mmol/L and ± 0.047 mmol/L (SD) at a concentration of 3.134 mmol/l. Serum PTH
179 had a precision of ± 0.2 pmol/L (SD) at a concentration of 2 pmol/L and ± 1.0 pmol/L (SD) at a
180 concentration of 10 pmol/l.

181 *Statistical analysis*

182 Results are shown as means and SDs unless otherwise specified. Descriptive statistics were generated for the
183 two ethnic groups at baseline, and the groups were compared using a two-sample *t* test when the data could
184 be assumed to be normal; otherwise using a non-parametric Kruskal Wallis test. Categorical variables were
185 compared using a Pearson's chi-square test.

186 Comparison of the daily vitamin D intake, serum 25(OH)D and serum PTH at baseline and endpoint across
187 the intervention groups were done by simple one way ANOVA and if significant differences were observed a
188 Tukey HSD test were performed in order to assess differences between groups. Analysis of covariance
189 (ANCOVA) was used to analyze the effect of the intervention on the outcome variable *change in vitamin D*
190 *status* (endpoint - baseline 25(OH)D). Two models were run – a minimal adequate model (Model 1) and a
191 maximal model (Model 2), which controlled for specific covariates that may influence the outcome. Model 1
192 included *baseline 25(OH)D* as a covariate and the factors *intervention group*, *ethnicity* and their interaction
193 allowing the effect of intervention to differ between the two groups of women. The interaction was tested for
194 statistical significance to see whether the model could be simplified. In Model 2, the covariates *Age* and *BMI*
195 at baseline were added to Model 1 construct as they are likely to have a strong association with the outcome
196 (serum 25(OH)D) [34]. The variables were chosen based on factors known to affect the change in vitamin D
197 status following fortification and variables associated with missing outcomes as well as the variables that
198 were significantly different between the two ethnic groups at baseline.

199 To check the model assumptions the standardized residuals of the final models were assessed for normality,
200 variance homogeneity and linearity.

201 Statistical analyses were performed using RStudio for Windows [35] (Version 1.1.414 – © 2009-2018
202 RStudio, Inc.) with a significance level of $\alpha = 0.05$.

203 RESULTS

204 A total of 143 women of Danish and Pakistani origin were randomly allocated to either vitamin D-fortified
205 foods or placebo (similar non-fortified foods), forming the four study groups. The completion rate of the
206 study was 89 % with the number of drop-outs at 16 in total for reasons as described within the Consort flow
207 diagram in Fig. 1. Drop-outs were equally distributed across the two ethnic groups ($P = 0.67$). In total, six
208 participants who finished the intervention were excluded from the analyses due to unplanned travels to
209 countries more southerly than 47° N during the study period and one was excluded as an outlier due to a
210 baseline serum 25(OH)D concentration > 125 nmol/L, which the Institute of Medicine (IOM) suggest is
211 potentially of concern, if sustained [1].

212 Adherence to consumption of the study foods was 92 % among participants of Danish origin and 73 %
213 among participants of Pakistani origin ($P < 0.05$). The study foods and the habitual dietary intake contributed
214 to a median (25th, 75th percentiles) daily intake of vitamin D of 32.0 (27.0, 34.4) $\mu\text{g/day}$ in the participants of
215 Danish origin randomized to the fortified food study group, and 24.2 (19.2, 30.8) $\mu\text{g/day}$ in the participants
216 of Pakistani origin randomized to the fortified food study group (Table 3). The chemical analysis of the crisp
217 bread showed a slightly higher concentration of vitamin D than expected due to a calculation error in the
218 production of the flour mix, whereas the eggs contained less vitamin D than expected due to lower vitamin D
219 (D_3 and 25(OH) D_3) doses in the chicken feed than expected, and this led to a slightly higher total daily
220 contribution of vitamin D from fortified foods (Table 1). The margin of safety was still sufficient (UL:100
221 $\mu\text{g/day}$) [19].

222 *Baseline characteristics of the participants*

223 There were no differences in baseline characteristics between the groups receiving vitamin D-fortified foods
224 or placebo foods within either ethnicity (data not shown); however, there were significant differences in
225 some baseline anthropometric characteristics between the two ethnic groups (Table 2). Women of Pakistani
226 origin had a significantly higher mean BMI and fat percentage compared to the women of Danish origin ($P =$
227 0.004 and $P < 0.001$, respectively). The intake of supplements containing only vitamin D was significantly

228 higher among the women of Pakistani origin compared to the women of Danish origin ($P = 0.001$). More
229 than 60 % of the women of Pakistani origin reported taking a supplement with vitamin D before study start.
230 The proportion of the participants using multivitamins was statistically similar in the two ethnic groups ($P >$
231 0.30) (Table 2). No significant differences ($P > 0.05$) were found in mean age between the two ethnic groups
232 (Table 2). The intake of vitamin D from dietary sources were significantly higher among the women of
233 Danish origin ($P < 0.05$), however both groups had very low intake.

234 Mean ($\pm$ SD) serum 25(OH)D₃ concentration at baseline was similar in women of Danish and Pakistani
235 origin (49.6 ± 18 v. 46.9 ± 22 nmol/L, respectively; $P = 0.80$), shown in Table 2. At baseline, none of the
236 participants, irrespective of ethnicity, had serum 25(OH)D < 10 nmol/L. The prevalence of 25(OH)D < 30
237 nmol/L and < 50 nmol/L was 9 and 50 % of the women of Danish origin and 24 and 32 % of the women of
238 Pakistani origin, respectively.

239 While serum 25(OH)D₂ was significantly higher in the women of Danish origin compared to women of
240 Pakistani origin (median [IQR] 2.0 [1.3-2.4] v. 1.0 [0.7-1.5] nmol/L; $P = 0.0001$), overall serum 25(OH)D₂
241 only represented 4%, on average, of serum total 25(OH)D in the entire group of women at baseline.

242 Baseline serum PTH was significantly higher among the women of Danish origin than of Pakistani origin (P
243 < 0.02), but all group values were within the reference range for normal PTH (Table 2). While serum PTH
244 was positively associated with serum 25(OH)D at baseline among the women of Pakistani origin ($P = 0.02$),
245 there was no significant ($P > 0.90$) association between these two variables in women of Danish origin (data
246 not shown).

247 *Effect of intervention*

248 The following analyses included 136 individuals following a complete case intention-to-treat approach. We
249 saw no significant differences in drop-outs between study groups ($P = 0.8$). There were no significant
250 changes in body weight (between baseline and endpoint) in either ethnic group ($P = 0.70$ and $P = 0.66$ for
251 Danish and Pakistani women, respectively) (data not shown).

252 The total vitamin D intake from the fortified foods during the intervention was higher among the Danish
253 women, compared to the women of Pakistani origin ($P = 0.004$), reflecting the lower adherence of the
254 women of Pakistani origin. The mean (SD) endpoint serum 25(OH)D concentration among the women of
255 Danish origin in the fortified food group was 77.8 (15) nmol/L, whereas among the women of Pakistani
256 origin in the fortified food group, it was significantly ($P < 0.01$) lower at 54.7 (18) nmol/L (Table 3). The
257 mean increase in serum 25(OH)D concentration from baseline to endpoint among the fortified food group
258 was higher ($P < 0.01$) in the women of Danish origin ($\Delta 26.4$ (16) nmol/L) compared to that in the women of
259 Pakistani origin ($\Delta 10.5$ (18) nmol/L). Serum 25(OH)D decreased by 2.8 (9) nmol/L in the Danish placebo
260 group and by 11.2 (12) nmol/L in the Pakistani placebo group over the 12 weeks of winter ($P = 0.02$). While
261 endpoint serum 25(OH)D₃ concentration was unaltered ($P = 0.09$) or significantly decreased ($P < 0.0001$) in
262 women of Danish and Pakistani origin in their respective placebo groups (mean ($\pm$ SD) change from pre- to
263 post-intervention: -2.8 ± 9.0 and -11.4 ± 12.5 nmol/L, respectively), it significantly increased in both Danish
264 and Pakistani women in the fortification groups ($+ 26.9 \pm 16.1$ nmol/L, $P < 0.0001$ and $+10.2 \pm 18.0$ nmol/L,
265 $P = 0.004$, respectively). Compared to pre-intervention, mean endpoint serum 25(OH)D₂ was unaltered in
266 women of either Danish or Pakistani origin, irrespective of placebo or fortification groups ($P > 0.11$ in all
267 cases; data not shown).

268 Following the intervention, none of the women of Danish origin in the fortified group had a serum 25(OH)D
269 concentration < 50 nmol/L. Among the women of Pakistani origin, 3 and 41 % of the fortified food group
270 had an endpoint serum 25(OH)D concentration < 30 and < 50 nmol/L, respectively (Table 3). In contrast,
271 vitamin D deficiency (serum 25(OH)D < 30) was evident in about a quarter of the Danish and a third of the
272 Pakistani women in the placebo groups at endpoint. Likewise, vitamin D insufficiency (serum 25(OH)D < 50)
273 was evident in about two-thirds and three-quarters of the Danish and Pakistani women in the placebo groups,
274 respectively, at endpoint (Table 4).

275 There was no significant increase in PTH concentration of the placebo groups (Danish and Pakistani)
276 following the intervention (0.21 and $P = 0.85$, respectively, t test). Likewise, there was no significant

277 decrease in PTH concentration of the fortified food groups (Danish and Pakistani) following the intervention
278 ($P = 0.40$ and 0.58 , respectively, t test).

279 *Analysis of covariance including factors influencing the change in serum 25(OH)D following the*
280 *intervention*

281 Based on the output of Model 1, compared to that of an equivalent woman in the non-fortified group, 12
282 weeks of intervention with vitamin D fortified foods (together with habitual diet, collectively supplying a
283 total of $30\text{ }\mu\text{g}$ vitamin D/day), resulted in an improvement of endpoint serum 25(OH)D of 31.1 nmol/L (25.0 ;
284 37.2) and 20.3 nmol/L (14.3 ; 26.3) in the women of Danish and Pakistani origin, respectively (Table 5). The
285 intervention effect is significantly higher in the Danish group compared to the Pakistani group ($P = 0.008$).

286 In terms of other factors which affected the response of serum 25(OH)D to intervention, the ANCOVA
287 models showed that baseline 25(OH)D status and BMI had a negative effect on the change in serum
288 25(OH)D following the intervention, however the effect of BMI was very small (Model 2). Model 2 was also
289 run where body weight was substituted for BMI but this did not affect the findings majorly (e.g., for body
290 weight at baseline, Effect = -0.18 , P value = 0.04) (data not shown). According to the model, participants
291 with e.g. a higher baseline serum 25(OH)D had an expected lower increase in serum 25(OH)D following the
292 intervention (Table 5).

293 **DISCUSSION**

294 The present study demonstrated that fortification of four different foods for 12 weeks during winter (January
295 through to March) was effective in increasing serum 25(OH)D among Danish and Pakistani women and
296 reduced the prevalence of low and very low vitamin D status during the winter months. The Danish fortified
297 group increased significantly more than the fortified Pakistani group. To our knowledge few previous studies
298 have assessed the effects of vitamin D food fortification with two different ethnic groups [36] and none in
299 Denmark, the results of our study may be of importance for the planning of fortification policies in countries
300 with multi-ethnic populations and no vitamin D fortification program.

301 The increase in serum 25(OH)D of the fortified participants was mainly dependent on baseline serum
302 25(OH)D and ethnicity in accordance with the linear models. However, we saw significant difference in the
303 intake of vitamin D from the fortified foods provided in the study due to a lower adherence in the women of
304 Pakistani origin compared to women of Danish origin. Generally, the results show a successful preventative
305 effect of an intervention with vitamin D-fortified foods in a country without mandatory fortification in a
306 population of women at risk of deficiency, as well as an important effect in terms of maintaining serum
307 25(OH)D > 50 nmol/L (reflective of sufficiency) in all participants of Danish origin and in 60 % of the
308 participants of Pakistani origin.

309 These results are in line with the findings of an 11-year Finnish follow-up study which showed a positive
310 effect of an implementation of a national voluntary vitamin D fortification policy. The study revealed an
311 average increase in vitamin D status of 20 nmol/L (95% CI: 19, 21) among supplement non-users, although
312 no records of ethnic background was obtained [20].

313 Several studies have investigated fortification of a single food, but only few have experience from low dose
314 fortification of several foods [17, 24]. We decided on a low-dose approach in which the daily dose of vitamin
315 D was spread out into several (four) foods in order to ensure the effectiveness and safety in a population
316 approach, based on previous experience [24, 25]. Our study participants were of Danish and Pakistani origin,
317 therefore we made special considerations when choosing the fortification vehicles since acceptance of the
318 foods in both groups was of high importance. As an example of our considerations we did not use milk as a

319 fortification vehicle despite its use in previous Nordic studies [20, 24, 25], since the prevalence of lactose
320 intolerance among individuals of Pakistani origin is > 60 % [37]. Instead we chose plain yoghurt. Yoghurt
321 has lower concentrations of lactose, and more importantly it contains the enzyme β -galactosidase which, by
322 intra-intestinal digestion, enables lactose intolerant people to absorb more lactose [38]. Yoghurt is thus better
323 tolerated by lactose intolerant people, compared to milk. Additionally, it can be used in breakfast, snacks and
324 sweet or savory cooking.

325

326 All the study foods were provided with no cost for the participants in portions that fit into a healthy diet. The
327 foods were all low in fat and energy. We instructed the participants to substitute their normal diet with food
328 from the study, so that they would remain approximately isocaloric throughout the study. This was
329 successful and we saw no changes in the body composition (BMI) of the participants following the
330 intervention.

331

332 The effect of intervention with fortified foods on vitamin D status was significantly different in the two
333 ethnic groups. This ethnic difference may in part be explained by the differences in adherence with the study
334 foods as adherence while high (73%) was significantly lower in the Pakistani women. Assessing the
335 distribution of baseline serum 25(OH)D it was evident that despite a similar mean vitamin D status, the IQR
336 revealed a much longer tail among the women of Pakistani origin which was evident as per the higher
337 percentage < 30 nmol/L. In the statistical models ethnicity was significant and there could be some ethnic-
338 specific differences in the serum 25(OH)D response that stem from genetic variation of vitamin D
339 modulating genes. Analyses of Single Nucleotide Polymorphism (SNP) data collected in this study, but not
340 yet studied, may explain parts of the found ethnic specific intervention effect.

341 We found that the baseline serum 25(OH)D among the women of Pakistani origin was higher than expected
342 (mean $\approx$ 50 nmol/L), when compared with data from a previous Danish study which included only
343 participants of Pakistani origin living in Denmark. In that study, conducted in 2002, a low vitamin D status
344 was reported for both girls and women of Pakistani origin (median serum 25(OH)D was 10.9 and 12.0

345 nmol/L, respectively) [5]. It should be noted, however, that the earlier study by Andersen *et al.* [5] was
346 performed prior to the initiation of the Vitamin D Standardization Program (VDSP) and this may affect the
347 comparability of serum 25(OH)D data. More importantly, several factors may have affected the change in
348 vitamin D status over the course of the last 15+ years. For example, it may partly be a result of the strong
349 focus on vitamin D in the general public as well as public health actions initiated by the Danish health
350 authorities between 2005 and 2010, targeting the general population, ethnic minorities as well as health
351 professionals [39, 40].

352

353 An additional result of the mentioned public health actions may be an increased intake of dietary
354 supplements containing vitamin D. In our study we saw a relatively high self-reported intake of vitamin D
355 from supplements, importantly, participants taking supplements prior to the study start were asked to stop
356 their supplement routine for the duration of the intervention study. The prior intake of vitamin D-containing
357 supplements was adjusted for by ensuring that baseline serum 25(OH)D concentration was an input in all of
358 the linear models. We did not intend to perform a screening serum 25(OH)D test prior to study start due to
359 ethical reasons as well as the time constraint present in this vitamin D RCT's due to the relatively short study
360 period of a winter only study.

361 ***Strengths and limitations***

362 This study was a real-life based design in which the participants would incorporate the study foods by
363 substitution of their habitual intake of similar foods. We encouraged the participants to consume the same
364 amount of the study foods every day to create a habit and incorporate the foods into their normal diet,
365 however, we allowed for the participants to include foods from several days in one meal, permitting people
366 to consume e.g. 4 eggs and 60 g cheese in one meal (e.g. omelet). This approach was chosen to mimic a real-
367 life situation in order for the results to be used in a public health setting as well as to strengthen the study by
368 increasing adherence and decreasing drop-out rate. The study achieved a good power and a low drop-out rate
369 for this study population.

370 In Denmark we have an increasingly multiethnic society. Research studies involving two or more ethnicities
371 are therefore highly needed in order to give suitable health advice and ensure equality in health of all citizens

372 in Denmark. The trial was carried out in winter time when there is no cutaneous production of vitamin D,
373 this lowers the risk of vitamin D contribution from UV sources which would confound the intake-status
374 relationship.

375 The intake of dietary vitamin D was very low, compared to the Danish National Survey of Dietary Habits
376 and Physical Activity (DANSDA 2011-13) [13], however, the participants were recruited based on their low
377 intake of fish, so this was not surprising. The dietary calcium intake recorded in the FFQ was also low,
378 especially considering the habitual dietary intake of calcium in Denmark which is on average approximately
379 1000 mg/day, among women [13]. The calcium intake data suggest that the study participants are in the
380 lower 25th percentile of the distribution intake range of age matched women in the general Danish
381 population [13]. One methodological explanation may be that the questionnaire, which was focused on
382 dietary vitamin D, was not designed to capture more than about 70 % of dietary calcium.

383 The chemical analysis of the crisp bread showed a slightly higher concentration than expected due to a
384 calculation error in the production of the flour mix, whereas the eggs contained less vitamin D than expected
385 due to lower vitamin D (D_3 and $25(OH)D_3$) doses in the chicken feed than expected, and this affected the
386 total daily dose from fortified foods of approximately 30 $\mu\text{g/day}$ (Table 1).

387 *Perspectives*

388 Results of the present study may be applicable for designing fortification policies prior to implementing them
389 in modern multiethnic population groups in Denmark. Though the road to implementation of a national
390 fortification program may be country specific, the overall goal is the same, improving vitamin D status of the
391 population, with a minimal risk of toxicity [41]. Data from countries such as Finland, having already
392 implemented widespread systematic vitamin D fortification, show that fortification is effective in terms of
393 increasing vitamin D status on a population level [41]. Future studies may go further into the topic of vitamin
394 D fortification at Northern latitude and effects of genetic variation in different ethnic populations complying
395 with the VDSP when analyzing the serum $25(OH)D$ concentration.

396 **CONCLUSION**

397 Vitamin D fortification of 30 µg/day, provided in four different foods, for 12 weeks during winter was
398 effective in increasing vitamin D status and preventing vitamin D deficiency in women of Danish and
399 Pakistani origin living in Denmark. Women of Pakistani origin had a lower response to the intervention that
400 did women of Danish origin. Adherence was lower among the women of Pakistani origin compared to the
401 women of Danish origin.

402 **Acknowledgements**

403 We thank the following individuals for their help and assistance with this trial: Majken Ege, Karin Hess Ygil,
404 Dorte L Korsbech, Erika Baig for assisting with the measurements, dietary intake calculations and FFQ
405 work, Bashy Quraishy, Raza Mustafa, Saima Raza, Shahnaz Qureshi and Shais Anaïs for introducing us to
406 the Pakistani societies in Copenhagen, Denmark. We would also like to thank Dr. George Hull and Ms.
407 Kirsten Dowling at University College Cork for their contribution to the analysis of serum 25(OH)D within
408 the ODIN project. We thank our industrial partners FrieslandCampina, Hedegaard, Lantmännen Cerealia and
409 DSM nutritional products for delivering the foods used in the trial.

410 **Contributors:** IMG and RA collected the data, TC managed intake data, JJ analyzed the vitamin D content
411 of the food, KC oversaw the analysis of serum 25(OH)D, IMG undertook the statistical analyses and wrote
412 this paper. EWA assisted with the statistical analyses. RA, IT, KC and MK designed the study. All
413 contributed to the manuscript.

414 **Ethical standards:**

415 Written informed consent was obtained from all participants on enrolment. The study protocol was approved
416 by the local ethical committee (protocol no. H-15008276) and registered at ClinicalTrials.gov with identifier:
417 NCT02631629. The study was carried out in accordance with the Declaration of Helsinki.

418 **Conflicts of interest:**

419 None of the authors had conflicts of interest. The industry partners had no influence on the design of the
420 study, the interpretation of the results or the writing of this manuscript.

421

REFERENCES

- 422 1. Ross AC, Taylor CL, Yaktine AL, Del Valle HB (2011) Calcium and Vitamin D. Institute of
423 Medicine, the National Academic Press, Washington D.C.
- 424 2. EFSA Panel on Dietetic Products Nutrition and Allergies (NDA) (2016) Scientific Opinion on
425 Dietary Reference Values for vitamin D. EFSA J 1–179. doi: 10.2903/j.efsa.2016.NNN
- 426 3. Cashman KD, Dowling KG, Skrabakova Z, et al (2016) Vitamin D deficiency in Europe: pandemic?
427 Am J Clin Nutr 103:1033–1044. doi: 10.3945/ajcn.115.120873.
- 428 4. Holick M, Chen T (2008) Vitamin D deficiency: a worldwide problem with health consequences. Am
429 J Clin Nutr 87:1080–1086.
- 430 5. Andersen R, Mølgaard C, Skovgaard LT, et al (2008) Pakistani immigrant children and adults in
431 Denmark have severely low vitamin D status. Eur J Clin Nutr 62:625–634. doi:
432 10.1038/sj.ejcn.1602753
- 433 6. Andersson Å, Björk A, Kristiansson P, Johansson G (2013) Vitamin D intake and status in immigrant
434 and native Swedish women: a study at a primary health care centre located at 60 °N in Sweden. Food
435 Nutr Res 1:1–8.
- 436 7. Meyer HE, Falch JA, Sjøgaard AJ, Haug E (2004) Vitamin D deficiency and secondary
437 hyperparathyroidism and the association with bone mineral density in persons with Pakistani and
438 Norwegian background living in Oslo, Norway, The Oslo Health Study. Bone 412– 417. doi:
439 10.1016/j.bone.2004.04.003
- 440 8. Cashman KD, Dowling KG, Škrabáková Z, et al (2015) Standardizing serum 25-hydroxyVitamin D
441 data from four Nordic population samples using the Vitamin D Standardization Program protocols:
442 Shedding new light on Vitamin D status in Nordic individuals. Scand J Clin Lab Invest 75:549–561.
443 doi: 10.3109/00365513.2015.1057898
- 444 9. Holick MF (2006) Resurrection of vitamin D deficiency and rickets. J Clin Invest 116:2062–2072.
445 doi: 10.1172/JCI29449
- 446 10. Nordic council of Ministers (2014) Nordic Nutrition Recommendations, 5th ed. København
- 447 11. SACN (Scientific Advisory Committee on Nutrition) (2016) Vitamin D and Health. 116–130.
- 448 12. Brot C, Vestergaard P, Kolthoff N, et al (2001) Vitamin D status and its adequacy in healthy Danish
449 perimenopausal women: relationships to dietary intake, sun exposure and serum parathyroid
450 hormone. Br J Nutr 86:97–103. doi: 10.1079/BJN2001345
- 451 13. Pedersen AN, Christensen T, Matthiessen J, et al (2015) Danskernes kostvaner 2011-2013 [Danish
452 Dietary survey]. Søborg
- 453 14. Danish National Health Authority (2010) Vitamin D: Prevention of deficiency.
- 454 15. Knudsen VK (2014) Danskernes forbrug af kosttilskud (The use of dietary supplements in Denmark).
455 e-paper Natl. Food Institute, DTU
- 456 16. Tetens I, Biltoft-Jensen A, Spagner C, et al (2011) Intake of micronutrients among Danish adult users
457 and non-users of dietary supplements. Food Nutr Res 55:1–8. doi: 10.3402/fnr.v55i0.7153
- 458 17. Cashman KD, Kiely M (2016) Tackling inadequate vitamin D intakes within the population:

fortification of dairy products with vitamin D may not be enough. *Endocrine* 51:38–46. doi: 10.1007/s12020-015-0711-x

18. Kiely M, Black LJ (2012) Dietary strategies to maintain adequacy of circulating 25-hydroxyvitamin D concentrations. *Scand J Clin Lab Invest Suppl* 243:14–23. doi: 10.3109/00365513.2012.681893
19. Grønborg IM, Tetens I, Ege M, et al (2018) Modelling of adequate and safe vitamin D intake in Danish women using different fortification and supplementation scenarios to inform fortification policies. *Eur J Nutr* 1–6. doi: 10.1007/s00394-017-1586-9
20. Jääskeläinen T, Itkonen ST, Lundqvist A, et al (2017) The positive impact of general Vitamin D food fortification policy on Vitamin D status in a representative adult Finnish population: Evidence from an 11-y follow-up based on standardized 25-hydroxyVitamin D data. *Am J Clin Nutr* 105:1512–1520. doi: 10.3945/ajcn.116.151415
21. Calvo MS, Whiting SJ, Barton CN (2004) Vitamin D fortification in the United States and Canada : current status and data needs 1 – 4 LAWFUL ADDITION OF VITAMIN D TO FOODS IN. *Am J Clin Nutr* 80:1710–1716. doi: 10.1093/ajcn/80.6.1710S
22. Ritu G, Gupta A, Gupta A (2014) Fortification of foods with vitamin D in India. *Nutrients* 6:3601–3623. doi: 10.3390/nu6093601
23. O'Neill CM, Kazantzidis A, Kiely M, et al (2017) A predictive model of serum 25-hydroxyvitamin D in UK white as well as black and Asian minority ethnic population groups for application in food fortification strategy development towards vitamin D deficiency prevention. *J Steroid Biochem Mol Biol* 173:245–252. doi: 10.1016/j.jsbmb.2016.09.010
24. Madsen K, Rasmussen L, Andersen R, et al (2013) Randomized controlled trial of the effects of vitamin D–fortified milk and bread on serum 25-hydroxyvitamin D concentrations in families in Denmark during winter: the VitmaD study. *Am J Clin Nutr* 98:374–382. doi: 10.3945/ajcn.113.059469.1
25. Hirvonen T, Sinkko H, Valsta L, et al (2007) Development of a model for optimal food fortification: vitamin D among adults in Finland. *Eur J Nutr* 46:264–70. doi: 10.1007/s00394-007-0660-0
26. Thuesen B, Husemoen L, Fenger M, et al (2012) Determinants of vitamin D status in a general population of Danish adults. *Bone* 50:605–610. doi: 10.1016/j.bone.2011.12.016
27. Cashman KD, Ritz C, Kiely M (2017) Improved dietary guidelines for vitamin D: Application of individual participant data (IPD)-level meta-regression analyses. *Nutrients* 9:1–17.
28. EFSA Panel on Dietetic Products Nutrition and Allergies (NDA) (2012) Scientific Opinion on the Tolerable Upper Intake Level of vitamin D. *EFSA J* 10:1–45. doi: 10.2903/j.efsa.2012.2813.
29. Burild A, Frandsen HL, Poulsen M, Jakobsen J (2014) Quantification of physiological levels of vitamin D3 and 25-hydroxyvitamin D3 in porcine fat and liver in subgram sample sizes. *J Sep Sci* 37:2659–63. doi: 10.1002/jssc.201400548
30. National Food Institute DTU (2015) Fooddata. <http://frida.fooddata.dk>. Accessed 8 Jan 2018
31. Jakobsen J, Wreford Andersen EA, Christensen T, et al (2018) Vitamin D vitamers affect vitamin D status differently in young healthy males. *Nutrients* 10 (1):12.
32. National Food Institute DTU (2009) The Danish food composition database (version 7). <http://www.foodcomp.dk/v7/>. Accessed 4 Mar 2018

499 33. Cashman KD, Kiely M, Kinsella M, et al (2013) Evaluation of vitamin D standardization program
500 protocols for standardizing serum 25-hydroxyvitamin D data: A case study of the program's potential
501 for national nutrition and health surveys. *Am J Clin Nutr* 97:1235–1242. doi:
502 10.3945/ajcn.112.057182

503 34. Bischof MG, Heinze G, Vierhapper H (2006) Vitamin D status and its relation to age and body mass
504 index. *Horm Res* 66:211–215. doi: 10.1159/000094932

505 35. R Core Team (2017) R: A language and environment for statistical computing. R Foundation for
506 Statistical Computing.

507 36. Tripkovic L, Wilson LR, Hart K, et al (2017) Daily supplementation with 15 µg vitamin D 2
508 compared with vitamin D 3 to increase wintertime 25-hydroxyvitamin D status in healthy South
509 Asian and white European women: a 12-wk randomized, placebo-controlled food-fortification trial.
510 *Am J Clin Nutr* 106:481–490. doi: 10.3945/ajcn.116.138693

511 37. Ahmad M, Flatzb G (1984) Prevalence of Primary Adult Lactose Malabsorption in Pakistan. *Hum*
512 *Hered* 34:69–75. doi: 10.1007/BF00270566

513 38. Kolars JC, Levitt MD, Aouji M, Savaiano DA (1984) Yogurt — An Autodigesting Source of Lactose.
514 *N Engl J Med* 5:1–3. doi: 10.1056/NEJM198401053100101

515 39. Danish National Health Authority (2005) Do you get enough vitamin D? - Important if you have dark
516 skin or wears covered clothing. 0–11.

517 40. Danish National Health Authority (2009) Make sure that your child's bones are healthy and strong.

518 41. Pilz S, März W, Cashman KD, et al (2018) Rationale and plan for vitamin D food fortification: A
519 review and guidance paper. *Front Endocrinol (Lausanne)* 9:1–16. doi: 10.3389/fendo.2018.00373

520

Fig. 1
 Consort flow diagram of the number of participants enrolled, randomized, completed and analyzed in the *ODIN-FOOD* study. Consort, Consolidated standards of Reporting Trials. DK: Women of Danish origin, Pa: Women of Pakistani origin

530 **Table 1**
531 Nutritional composition, portion size, energy and mean total vitamin D contribution from the
532 fortified and placebo foods

Food product	Macro nutrients (g/100g) ¹	Kcal/100g ²	Daily portion size (g/day)	Kcal/day	Contribution from fortified product (µg/day) to:		Contribution from placebo product (µg/day) to:	
					Vitamin D ₃ ³	25(OH)D ₃ ⁴	Vitamin D ₃ ³	25(OH)D ₃ ⁴
Yoghurt		70	150	105	2.00 (0.03) ⁵	< 0.025	< 0.01	< 0.025
Fat	3.5							
CHO	6.6							
Protein	2.9							
Cheese		272	60	163	8.30 (0.27)	0.07 (0.01)	0.03 (0.01)	0.07 (0.01)
Fat	19							
CHO	-							
Protein	30							
Eggs		141	54	76	0.18 (0.17)	0.68 (0.43)	0.81 (0.06)	0.54 (0.04)
Fat	9.9							
CHO	0.8							
Protein	12.6							
Crisp bread		460	8.9	40	18.5 (2.6)	< 0.025	< 0.025	< 0.025
Fat	20							
CHO	58							
Protein	10							
Total		943	-	384	29.0 (2.6)	≈ 0.7	≈ 0.8	≈ 0.6

533 ¹Macro nutrient composition for the specific products given by the suppliers.

534 ²Fortified and placebo foods had similar energy content, only shown once.

535 ³Analysed vitamin D₃ concentrations of fortified and placebo foods. Contents of vitamin D₂ in cheese and eggs
536 in fortified and placebo products would additional account for <0.03 µg/day.

537 ⁴Analysed 25(OH)D concentration, all 25(OH)D₃ values are multiplied with a factor 2.5. Contents of 25-
538 hydroxyvitamin D₂ in cheese in fortified and placebo products would additional account for <0.04 µg/day.

539 ⁵Mean (SD) for all such values.

Table 2Baseline characteristics of participants by study group and ethnicity¹

	Danish			Pakistani		
	Total (n=66)	Placebo (n=35)	Fortified (n=31)	Total (n=70)	Placebo (n=37)	Fortified (n=33)
Born in Denmark (%)	99	100	99	62	70	54
Mean Age (y)	33 (11)	34 (11)	32 (11)	36 (9)	36 (9)	36 (10)
Mean weight in kg (SD)	68 (13)	70 (15)	67 (11)	70 (12)	68 (13)	70 (12)
Mean BMI (kg/m ²)	24 (5)	25 (5)	24 (4)	27 (5)*	27 (5)	27 (5)
Mean fat percentage (%)	31 (8)	33 (8)	30 (7)	37 (6)*	37 (6)	38 (6)
Mean serum 25(OH)D (nmol/L)	49.6 (18)	46.2 (19)	53.3 (17)	46.9 (22)	49.0 (23)	44.5 (21)
< 9.9 nmol/L, n (%)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
≥ 10 - < 29.9 nmol/L, n (%)	6 (9)	5 (14)	1 (3)	17 (24)	8 (22)	9 (27)
≥ 30 - < 49.9 nmol/L, n (%)	33 (50)	20 (57)	13 (42)	22 (32)	11 (30)	11 (33)
≥ 50 nmol/L, n (%)	27 (41)	10 (29)	17 (55)	31 (44)	18 (48)	13 (40)
Mean PTH (pmol/L)	5.0 (1.7)	5.3 (1.9)	4.9 (1.6)	4.4 (1.8)*	4.2 (1.8)	4.5 (1.8)
Median vitamin D intake from the diet (µg/day)	1.5 (1.0 ; 2.0)	1.5 (1.0; 2.0)	1.7 (1.0; 2.1)	1.1 (1.0; 2.0)*	1.1 (0.8; 1.4)	1.0 (0.7; 1.7)
Total median vitamin D intake from vitamin D suppl. and multivitamin suppl. (µg/day)	2.9 (1.8; 9.0)	7.9 (4.3; 16.0)	2.0 (1.0; 3.7)	13 (6.8; 29.3)*	13 (7.9; 29.3)	13 (5.5; 22.2)
Median dietary calcium intake (mg/day)	465 (324; 688)	487 (309; 685)	442 (331; 692)	441 (260; 589)	422 (262; 547)	465 (259; 634)
Vitamin D supplements (%)	15	17	13	61 [¶]	65	58
Multivitamin supplements (%)	24	17	32	30	32	27

Smokers (%)	14	14	13	4	3	6
Alcohol drinkers (%)	86	89	84	1.4¶	3	0
Wearing hijab (%)	0	0	0	44¶	43	45

¹Means and SD unless otherwise specified. If non-normally distributed, medians and 25th and 75th percentiles. No variables were transformed for analysis.

*Means significantly different from women of Danish origin; Unpaired *t* test, *P* <0.05

¶Percentage significantly different from women of Danish origin; Pearson's chi² test, *P* <0.05

Table 3Total vitamin D intake, and serum 25(OH)D and PTH concentration at baseline and endpoint in each of the four study groups¹

	Danish		Pakistani		<i>P</i> ⁵
	Placebo (n = 35)	Fortified (n = 31)	Placebo (n = 37)	Fortified (n = 33)	
Total vitamin D Intake ² , µg/d	1.5 (1.0, 2.0) ^{3a}	32.0 (27.0, 34.4) ^b	1.1 (0.8, 1.4) ^a	24.2 (19.2, 30.8) ^c	<0.0001
Serum 25(OH)D, nmol/L:					
Baseline	46.2 (19) ⁴	53.3 (17)	49.0 (23)	44.5 (21)	0.31
Endpoint	44.0 (17) ^a	77.8 (14) ^b	36.5 (16) ^a	54.7 (18) ^c	<0.0001
Change	-2.8 (9) ^a	26.4 (16) ^b	-11.2 (12) ^a	10.5 (18) ^c	<0.0001
Serum PTH, pmol/L:					
Baseline	5.3 (1.9)	4.9 (1.6)	4.2 (1.8)	4.5 (1.8)	0.07
Endpoint	4.7 (1.4)	4.4 (1.8)	4.3 (1.8)	4.3 (1.5)	0.97
Change	-0.54 (9.0)	-0.42 (16)	-0.002 (12)	-0.27 (18)	0.60

¹25(OH)D = 25-hydroxyvitamin D; PTH, parathyroid hormone.²Total vitamin D intake during study; placebo groups had only dietary intake, fortified groups had diet plus study fortified foods.³Median (25th, 75th percentiles).⁴Means (SD), all such values.⁵P values for baseline comparisons by intervention group were determined with the use of a simple one-way ANOVA, followed by a Tukey HSD test.^{a,b,c}Different superscript letters represent significant (*P*<0.01) differences in group means for endpoint total vitamin D intake, serum 25(OH)D and change in serum 25(OH)D.

Table 4

Number and percentage of women with serum 25(OH)D below 30 and 50 at baseline and endpoint in each of the four study groups

	Danish		Pakistani	
	Placebo	Fortified	Placebo	Fortified
Serum 25(OH)D < 30 nmol/L:	n (%)		n (%)	
Baseline	5 (14)	1 (3)	8 (22)	9 (27)
Endpoint	7 (23)	0 (0)	11(34)	1 (3)
Serum 25(OH)D < 50 nmol/L:				
Baseline	25 (71)	14 (45)	19 (51)	20 (60)
Endpoint	20 (65)	0 (0)	25 (78)	11 (41)

Table 5

Analysis of covariance models exploring the intervention effects on serum 25(OH)D response

Coefficients	Model 1 ¹ (minimal)			Model 2 ² (maximal)		
	Effect	95 % CI	<i>P</i>	Effect	95 % CI	<i>P</i>
Intercept ³	16.5	(9.76; 23.2)	<0.0001 (***)	33.8	(19.6; 48.1)	<0.0001 (***)
Baseline 25(OH)D	-0.41	(-0.52; -0.30)	<0.0001 (***)	-0.42	(-0.53; -0.30)	<0.0001 (***)
Ethnicity:						
Danish	-	-	-	-	-	-
Pakistani	-8.41	(-13.9; -2.14)	0.008 (**)	-6.12	(-12.14; -0.07)	0.05 (*)
Danish*Fortified	31.1	(25.0; 37.2)	<0.0001 (***)	33.8	(19.6; 48.1)	<0.0001 (***)
Pakistani*Fortified	20.3	(14.3; 26.3)	<0.0001 (***)	27.7	(12.2; 43.3)	<0.0001 (***)
BMI at baseline				-0.64	(-1.16; -0.12)	0.02 (*)
Age				-0.05	(0.006; 0.20)	0.69

¹Model 1 (minimal): change in vitamin D status ~ Baseline status + intervention group + (ethnicity*intervention group), adjusted R²= 65%.

²Model 2 (maximal): change in vitamin D status ~ Baseline status + intervention group + (ethnicity*intervention group) + BMI + Age, adjusted R²= 67%.

³The reference group is included in the intercept: Danish ethnicity and placebo study group.