Enzyme kinetics of fungal glucuronoyl esterases on natural lignin-carbohydrate complexes

Mosbech, Caroline; Holck, Jesper; Meyer, Anne; Agger, Jane Wittrup

Published in:
Applied Microbiology and Biotechnology

Link to article, DOI:
[10.1007/s00253-019-09797-w](https://doi.org/10.1007/s00253-019-09797-w)

Publication date:
2019

Document Version
Other version

[Link back to DTU Orbit](#)

Citation (APA):
Mosbech, C., Holck, J., Meyer, A., & Agger, J. W. (2019). Enzyme kinetics of fungal glucuronoyl esterases on natural lignin-carbohydrate complexes. *Applied Microbiology and Biotechnology*, 103(10), 4065-4075.
<https://doi.org/10.1007/s00253-019-09797-w>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Online Resources

Enzyme Kinetics of Fungal Glucuronoyl Esterases on Natural Lignin-Carbohydrate Complexes

Applied Microbiology and Biotechnology

Caroline Mosbech, Jesper Holck, Anne Meyer, Jane Wittrup Agger

Protein Chemistry and Enzyme Technology

Department of Biotechnology and Biomedicine, Technical University of Denmark, Søltøfts Plads 221, DK-2800 Kgs. Lyngby

Corresponding author:

Jane Wittrup Agger: jaag@dtu.dk

Contents

Online Resource 1: Codon optimized DNA sequences.....	2
Online Resource 2: Cloning vector and SDS gel.....	5
Online Resource 3: Assay flow diagram	7
Online Resource 4: Ions for quantification.....	8
Online Resource 5: Sequence alignment.....	9
Online Resource 6: Percent Identity Matrix (PIM).	10
Online Resource 7: <i>Cu</i> GE Temperature Stability.....	11

Online Resource 1: Codon optimized DNA sequences

Armillaria fuscipes, AfGE

>GAATTCGGTTGTTCTGATTTGCCATCTAACATTCCATTGACTCCTAATTCTATGTTGCCAGATCCTTTTACTTTCTTGAAT
GGTACTGCTGTTACTTCTTGTCTGCTTTTACTTGTAGACAAGCTGAAATTTCTGCTTTGTTGCAACAATTTCGAG
TTGGGTCAATTGCCACCTAGACCTGAATCTGTTGTTGGTACTTTGTCTGGTAACACTTTGACTATTAATGTTGCT
CATTCTGGTAGATCTATCTCTTTTACTGTTGGTATTCAAAAGCCAACCTGGTTCTGGTACTGGTCCATTTCTGCT
TTGATTGTTTTGGGTGGTTTCTTGTCTTTGCCTCAACCACCTGGTGTGCTATCTTGTCTTTCAACAACGATGAT
ATTGCTGCTCAAAACAATGCTCAATCTAGAGGTCAGGGTAAATTTTCACTTTGTACGGTTCTAACCACCCAGCT
GGTGTCTTGTATGGCTTGGGCTTGGGGTGTGCTAGAAATTATGGATGCTTTGGAGCAAGTTAACTCTAACATCGAT
TTGACTAGAGTTGGTGTTACTGGTTGTTCTAGAAACGGTAAAGGTGCTTTGGTTGCTGGTGCTTTTGAGCCAAGA
ATTGCTTTGACTATTCTCAAGAATCTGGTCTGGTGGTGCTGGTTGTTGGAGAATTTCTGATGCTATGTTGGCT
TCTGGTGTCTACTCAAACCTGCTTCTGAAATTGTTCAAGAGAACGTTTGGTTTTCTCCTGCTTTTAATACTTTT
GCTCAATCTGGTGTTAACCATTGCCTTTGATCATCACATGTTGTCTGCTTTGATTGCTCCAAGACCTTTGTTG
GTTATTGATAACACTGGTATCGATTGGTTGGGTCCAGAGTCTGTTTGGGGTTGTCAAAGACTGCTTCTTTGGTT
TGGAAAGCTTTGGGTGTTCTGATAGAAATGGGTGTTTCTCAAGTTGGTAACCATGCTCACTGTGCTTTTCCAAC
CAAGAAAAGCCTGATTGGATGCTTTCGTTAACAGATTCTTGTGGGTCAAGGTGGTGTTAACACTTCTGGTATT
GTTAAGACTGATGGTGCTAATGGTTTGGGTTTCGTTGATTCTAGATGGGTGGTTGGAAAGTTCAAACCTGGCT
GGTGTCGAC

Thielavia terrestris, TtGE

>GAATTCGCTAGACAAGGTGGTGCTGGTAACGCTACTATTCAATGTGCTGCTATTCCATCTCCTTTTCCAACCTGGCAACAA
TTGCCTCAACAATCTACTTTGCCTGATCCATTCTTGCCTTTGAAGTACACTACTCCATCTGATGCTCAAAATATT
ATGGCTGGTAAAGGTCAAGGTAGAGTTCAAACCTCTGATGAATGGTATCAATGTAGACAACCAGAAAATTTGCAA
ATGTTGCAAGAGTACCAATATGGTTACTATCCTGATCATTCTCAAGAAAAGGTTGAGGCTACTAGATCTGGTACT
ACTTTGAACATTGCTGTTACTGCTGGAGATAAGACTGGTCTTTTAAAGCTTCTTTCATTTGCCATCTGGTGCT
TCTGCTTCTAAACCTGCTCCAGTTGTTATTAACATTGGTGGTATGCAAAATCAACCTTACTTGTCTGCTGGTATC
GCTATCGCTCAATTCGATTACACTTCTGTTGCTCCAGATTCTAACTCTAAGACTGGTGCTTTCTGGTCTATCTAC
AATGGTAGAGATATTGGTGTTTTGACTGCTTGGGCTTGGGGTTCCACAGAGTTTTGGATGCTATTAATATGACT
GTTCCAGAAATTGATGCTACTAGAGTTGGTGTTACTGGTTGTTCTAGATTGGGTAAAGGTGCTTTGGCTGCTGGT
TTGTTGATAAGAGAATCACTTTGACTATGCCTATGTCTTCTGGTGTTCAAGGTATGGGTCCATACAGATACTAT
ACTATGTCTGGTCAAGGTGAAAACCTGGAAAACCTCTAAACAAGGTGCTCCTTGGTGGACTGATTCTACTTTGGGT
ACTTTTGTTAACCATTGCTGAGAATTTGCCTTATGATGCTCACACTATTGCTGCTGCTATTGCTCCAAGAGCTTTG
GTTATTGATCAAGGTACTGGAGATCAATTCATAACTCTAAGGCTACTGCTGTTATTATCTACCCAGCTGCTAAA
TTGGTTTATGATTGGTTGGGTGTTGGAGATCAAATTGCTATGTCTGTTAGATCTGGTGGTCATTGTGATATGTCT
GGTTACACTTCTGTTTTGCCTTACGTTCAAAGATTTTCTTTGGTACTCCAACCTAACAAAGATTACAACCTTTG
GGTCTTATGTTTCTCCTATTACTACTGCTTATCCATGGGCTACTGCTACTCCACCTTCTAAGGCTGCT
GTCGAC

Punctularia strigosonata, PsGE

>GAATCCAATCTACTTGTGATACTCCATCTACTATTGCTGGTTATTCTAACGCTATGTTGCCAGATCCTTTCAATCCCT
AATGGTACTAGAGTTACTACTGTTGATGAATGGAAGTCTAGAAAGATCTCAAATTGCTGCTTTGGTTCAAGGTTAC
GAGGCTGGTACTTTGCCACCTAAGCCATCTTCTGTTACTGGTTCTTTTCTGGTAACACTTTGACTGTAAACGTT
AACACGGTGGTCTTCTATCTCTTCGCTAACACTATCACTTACCCTTCTGGTACTGCTCCAGCTGGTGGTTGG
CCTTTGTTGATTGCTTATGATGGTGGTCTATTCCAGTTCCTTCTGGTGTGCTGTTTTGAACTACAACATCGGT
AACATCGCTCAACAACTGATACTACTTCTAGAGGTGTTGGTTTGTGTTTACAACCTGTACGGTAAAACGCTACT
GCTTCTGCTATGACTGCTTGGGTTTGGGGTTTGTCTAGAATCATCGATGTTTTGGAAGTACTCCAGCTGCTATG
GTTAACACTGCTAAGTTGGCTGTTACTGGTGTCTAGAGATGGTAAAGGTGCTTTGATGGCTGGTGCCTTCGAA
CCTAGAATTGCTTTGACTATTGCTCAAGAGTCTGGTGTGGTGGAGATGCTTGTGGAGAATTACTAAAACGAA
GGAGATGAGGGTGTGGTGTCAACCATCTACTGAAATTGTTCAAGAGAACGTTTGGTCTCTACTAACTTCAAC
AACTACGTTTCTAACTGAATGTTTTGCCATATGATCATCACTTGTGGCTGCTATGATTGCTCCTAGAGCTTTG
TTGTCTTACGGTAATACTGATTATGCTTGGTGTCTCCTATGTCTTCTTGGGGTGTATGAATGCTGCTCATACT
GTTTGGCAAGCTTTGGGTGTTCCACAAAACACGGTTTTGTTGAGGTGGTGGTCATGCTCACTGTGCTTCCCT
ACTTCTTTGGTTCAAGTTTGGATGCTTCTTTGATACTTTCTTGTGGGTCAAACGTTTCTACTGATTACTTC
TCTACTAACACGTTTTCGGTTCTGCTCAATGGGTTCCAGCTAACTTCATTAATTGGACTGCTCCTGCTTTGACT
GGTCTTCTACTTCTTCTTCTACTTCTGTTATTTCTTCTATTACTCCAACCTCCTGTTCCAACACTTCTCCTACT
ACTGTTCCAACACTTCTGCTCCTGCTCCAACCTGGTGTCCAGTTGCTAAATGGGGTCAATGTGGTGGTATTGGT
TGGACTGGTCTACTACTTGTACTTCTGGTACTACTTGTACTGTTTTGAACGCTTACTATTCTCAATGTTTG
GTCGAC

Cerrena unicolor, CuGE

>GAATCCAAGCATCCGCTCCTCAATGGGGTCAATGTGGTGGTATCGGTTGGACTGGTCCT
ACTGCCTGCCCTAGTGGTTGGGCTGTGAGCAATTGAACGCTTACTATTCTCAATGTTG
CAAGGTGCTGCTCCTGCTCCAGCTAGAACTACTGCTGCTCCACCTCCACCTCCAGCTACT
ACTGCTGCTCCTCCACCTCCAACACTTCTGCTCCTACTGGTCTTCTCCAGTTGCTGGT
GCTTGTGGTGTCTATTGCTTCTACTGTTCCAAACTATAACAATGCTAAGTTGCCTGATCCA
TTCACTTTGCTAATGGTACTGCTTTGAGAACTAAAGCTGATTGGTCTTGTAGAAGAGCT
GAAATTTCTGCTTTGATTCAAAACTACGAGGCTGGTACTTTGCCTCCAAAGCCTCCAGTT
GTTACTGCTTCTTTTCTAAATCTGGTAACACTGGTACTTTGGCTATTACTGCTGGTTTG
TCTAATTCTCAAACATTAAATTCTCTCCTACTATCTCTTACCCATCTGGTACTCCTCCA
GCTAATGGTTGGCCATTGATTATTGCTTATGAAGGTGGTCTATTCTTCTTCCAGCTGGT
GTTGCTACTTTGACTTACTCTAACTCTGATATGGCTCAACAAAATTCTGCTTCTTCCAGA
GGTCAAGGTTTGTGTTTACCAATTGTATGGTCTACTCATTCTGCTTCTGCTATGACTGCT
TGGGTTTGGGGTGTGTTCCAGAATTATTGATGCTTTGGAGATGACTCCTACTGCTCAAATT
AATACTCAAAGAATCGGTGTTACTGGTTGTTCCAGAGATGGTAAAGGTGCTTTGATGGCT
GGTGTCTTTCGAAGAGAGAATTGCTTTGACTATTCCACAAGAATCTGGTTCTGGTGGAGAT
GCTTGTGGAGATTGTCTAAGTACGAGATCGATAACGGTAACCAAGTTCAAGATGCTGTT
GAAATTGTTGGAGAGAACGTTTGGTCTCTACTAACTTCAACAACTACGTTCAAAAGTTG
CCTACTGTTCCAGAAGATCATCACTTGTGGCTGCTATGGTTGCTCCTAGAGCTATGATC
TCTTCGAGAACTGATTATTTGTGGTGTCTCCAATGTCTTCTTTCGGTTGTATGACT

GCTGCTCATACTGTTTGGCAAGGTTTGGGTATTGCTGATTCTCACGGTTTTGCTCAAGTT
GGTGGTCATGCTCACTGTGCTTGGCCTTCTTCTTTGACTCCACAATTGAACGCTTTTATT
AACAGATTCTTGTTGGATCAATCTGCTACTACTAATGTTTTCACTACTAACAATCAATTC
GGAAAGGTTCAGTGGAATGCCGCAAATTGGATTACTTGGACTACCCCTACTCTTACTGGT
CTAGA

Online Resource 2: Cloning vector and SDS gel

Graphic representation of the pPICZalphaA vector delivered by Genscript holding the CE15 genes between the *EcoRI* and *SalI* restriction sites

(pPICZalpha-A vector (2019) Genscript <https://www.genscript.com/gsfiles/vector-map/yeast/pPICZalpha-A.pdf?1897614803> Accessed March 12th 2019)

SDS gel of the expressed enzymes after purification. Loaded in the following order with theoretical molecular weight given in bracket; *CuGE* (48 kDa), *AfuGE* (40 kDa), *PsGE* (47 kDa), *TtGE* (42 kDa). The four enzymes are all glycosylated resulting in the smear and the increase in size compared to the theoretical weight.

Online Resource 3: Assay flow diagram

Schematic overview of assay procedure for the kinetic study. Varying substrate concentrations of LRP with a fixed enzyme concentration. Assay conducted in eppendorph tubes in thermomixer. 50 µL samples taken every 5 minutes. Reaction stopped after initial removal of substrate by centrifugation and heat inactivation at 100 °C for 10 min. GH10 endo-xylanase added to the reaction supernatant to decrease the size of the products released by CE15 enzymes. Acetylation removed from the products by alkaline treatment to simplify the products profile. Characterization and quantification of products done by LC-MS.

Online Resource 4: Ions for quantification.

Complete list of adducts used for quantification of enzyme reaction products and their corresponding retention times.

Analyte	m/z [M+Na] ⁺	m/z [M+NH ₄] ⁺	Retention time (min), peak max
MeGlcAXyl	363	358	7.6
MeGlcAXyl ₂	495	490	9.8
MeGlcAXyl ₃	627	622	12.6
Reduced MeGlcAXyl ₃	629	624	11.4

Online Resource 5: Sequence alignment

Alignment of segments of the amino acid sequences from CE15 characterized in this study. Conserved GCSRXC motif with a catalytic nucleophile serine marked with red curly bracket. Catalytic glutamic acid and histidine marked with red arrow. *TtGE* belonging to PPR group 8 holds a serine at the position of the glutamic acid. Alignment and sequence logo made with CLC Workbench 7.5.

Online Resource 6: Percent Identity Matrix (PIM).

Percent query coverage given in brackets. Color scale from light to dark orange indicate percent sequence identity from the lowest value 28% in light orange to the highest value 100% in dark orange. Values obtained using BLAST (<https://blast.ncbi.nlm.nih.gov>).

	<i>Tt</i> GE	<i>Afu</i> GE	<i>Ps</i> GE	<i>Cu</i> GE
<i>Tt</i> GE	100 (100)	35 (84)	31(79)	28 (96)
<i>Afu</i> GE	35 (84)	100 (100)	51 (99)	49 (99)
<i>Ps</i> GE	31 (77)	51 (89)	100 (100)	65 (88)
<i>Cu</i> GE	28 (98)	49 (99)	65 (97)	100 (100)

Online Resource 7: *CuGE* Temperature Stability.

Thermostability of *CuGE* obtained by incubation at specific temperatures for 48 hours and followed by activity measurement on Me-MeGlcA (left graph). % conversion calculated relative to activity with *CuGE* not pre-exposed to incubation. Right graph: A zoom of the first 6 hours of incubation. The study was conducted in triplicates. The kinetic assays (Fig 2) were conducted for 20 min at 50 °C. Under these conditions *CuGE* still retained around 85 % of its relative activity. After 25 hours at 50 °C approx. 50% relative activity remained.