



Size selection of Antarctic krill (*Euphausia superba*) in a commercial codend and trawl body

Herrmann, Bent; Krag, Ludvig Ahm; Krafft, Bjørn A.

Published in:
Fisheries Research

Link to article, DOI:
[10.1016/j.fishres.2018.05.028](https://doi.org/10.1016/j.fishres.2018.05.028)

Publication date:
2018

[Link back to DTU Orbit](#)

Citation (APA):
Herrmann, B., Krag, L. A., & Krafft, B. A. (2018). Size selection of Antarctic krill (*Euphausia superba*) in a commercial codend and trawl body. *Fisheries Research*, 207, 49-54.
<https://doi.org/10.1016/j.fishres.2018.05.028>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Research

Manuscript Draft

Manuscript Number: FISH8081R1

Title: Size selection of Antarctic krill (*Euphausia superba*) in a commercial codend and trawl body

Article Type: Research Paper

Keywords: zooplankton fishery; size selectivity; sequential selectivity process

Corresponding Author: Dr. bent herrmann, Ph.D

Corresponding Author's Institution: SINTEF Fisheries and Aquaculture

First Author: bent herrmann, Ph.D

Order of Authors: bent herrmann, Ph.D; Ludvig A Krag, PhD; Bjørn A Krafft, PhD

Manuscript Region of Origin: DENMARK

1 **Size selection of Antarctic krill (*Euphausia superba*) in a commercial codend and trawl**
2 **body**

3 Bent Herrmann^{1,a*}, Ludvig A. Krag^{2,a}, Bjørn A. Krafft^{3,a}

4 ¹SINTEF Fisheries and Aquaculture, Fishing Gear Technology, Hirtshals, Denmark

5 ²DTU Aqua, Technical University of Denmark, Hirtshals, Denmark

6 ³Institute of Marine Research, Bergen, Norway

7 ^aThese authors have equal authorship

8 *: Corresponding author. Tel.: +45 98 94 43 00

9 E-mail address: Bent.Herrmann@sintef.no (B. Herrmann).

10 **Keywords:** zooplankton fishery, size selectivity, sequential selectivity process

11 **Abstract**

12 During fishing, many fish species are able to avoid the net walls of the trawl body and so the
13 majority of size selection occurs in the codend of the net. Antarctic krill (*Euphausia superba*)
14 are regarded as true planktonic organisms passively drifting with currents, but they also
15 display self-locomotion by active swimming. There is a lack of knowledge regarding the
16 behavior of krill during the fishing process, and extrapolating results obtained for other
17 species to krill is of limited value. In the case of krill, it is largely unknown to what extent the
18 codend versus the trawl body contributes to the size selection process. The current study aims
19 to quantify the size selection of krill in a commercially applied codend during experimental
20 fishing. Combining these results with a model for full trawl size selectivity it was possible to
21 provide an insight to the size selection process in the trawl body. Specifically, the study
22 applied a two-step approach by first estimating the size selectivity of a commercial codend
23 and second used the codend size selectivity obtained in this study to estimate the trawl body

24 size selectivity of a commercial trawl based on entire trawl-selectivity obtained in a previous
25 study. The results of this two-step analysis revealed that the trawl body contributes
26 significantly to the total size selection process, demonstrating that size selectivity of Antarctic
27 krill in commercial trawls is affected by both the trawl body and the codend.

28 **1. Introduction**

29 Several fish species avoid the netting of trawls during capture (Wardle, 1993) and so the
30 majority of size selection for those species occurs in the codend of the trawl (Wileman et al.,
31 1996). Other species, such as smaller invertebrates, may display a different pattern of
32 behavior. For example, prawns tend to display a more limited response to trawl stimuli
33 (Lochhead, 1961; Newland & Chapman, 1989) and size selection resembles more of a sieving
34 process in which individuals may meet the trawl netting frequently and with a more random
35 orientation. Polet (2000) found that it was mainly the rounded lateral part of the net belly that
36 was responsible for size selectivity for Crangon shrimps (*Crangon crangon*). Antarctic krill
37 (*Euphausia superba*) are generally regarded as true planktonic organisms that drift with the
38 currents, however they also display the ability to move horizontally and vertically in the water
39 column, by swimming at higher speeds for limited periods of time (Marr, 1962; Kanda et al.
40 1982). Krag et al. (2014) speculated if size selection may occur throughout the entire trawl
41 body when harvesting Antarctic krill.

42
43 Size selectivity results and underwater video recordings indicate that Antarctic krill escape
44 through the mesh head first, at an angle perpendicular to the netting wall (Krag et al., 2014).
45 This suggests that individual krill are either able to orientate themselves optimally in relation
46 to the net mesh to facilitate their escape or, alternatively, their escape is a random process, where
47 frequent contact with the trawl netting will result in some krill meeting the netting at an
48 optimal orientation for escape by chance. Recent trawl designs in the fishing industry also

49 support these mechanisms: Traditional net designs in the krill fishery comprised midwater
50 trawls (Budzinski et al., 1985) with large openings (e.g. 60x50m) and large meshes near the
51 mouth of the net with a successive reduction in size towards the small meshed codend. More
52 recent designs comprise small mouthed (20x20m), low-tapered trawls with small meshes
53 throughout the length of the trawl body (Bakketeig et. al, 2017). Detailed knowledge of the
54 selection processes operating in fishing gear is important both in terms of understanding catch
55 efficiency and gaining a better insight into ecosystem based management practices (Krafft et
56 al., 2016).

57

58 Krag et al. (2014) assessed the selectivity of a full commercial trawl. However, it is unknown
59 whether their results represented size selection over the full trawl body, with krill having
60 multiple random contacts with the mesh in the trawl body, eventually resulting in escape, or
61 they were due to the fact that krill are very effective at orientating themselves towards the
62 meshes at an angle that facilitates escape in the codend. Therefore, it is unknown to what
63 extent trawl body and codend each contribute to the size selection in the trawl. If the majority
64 of size selection occurs in the codend, management of size selection in the krill fishery would
65 only require changes in codend design. However, if the trawl body is important, adjusting the
66 gear selectivity would require changes to other parts of the trawl . Therefore, it is important to
67 quantify size selection in commercial codends and trawl bodies . The current study aimed to
68 provide data to bridge this knowledge gap. Specifically, the main objectives were:

- 69 - To quantify size selection in a commercial krill trawl codend.
- 70 - To investigate to what extent size selection of krill in commercial trawls is attributed
- 71 to the codend and the main trawl body.

72 **2. Materials and Methods**

73 To obtain the objects described above, the study applied a two-step approach: i)
74 estimating the size selectivity of a commercial codend (sections 2.1 and 2.2); and
75 ii) used the codend size selectivity obtained in this study to estimate the trawl
76 body size selectivity of a commercial trawl based on entire trawl-selectivity
77 obtained in a previous study under the assumption that the codend selectivity in
78 both studies is similar (sections 2.2 and 2.3).

79 *2.1 Sea trials and gear specifications*

80 To quantify the size selection process that occur in the codend, a survey trawl with a codend
81 of commercial mesh size was used. The codend was surrounded by a small-meshed cover to
82 collect codend escapees. The trawling was carried out off the coast of the South Orkney
83 Islands (60°35'S, 45°30'W) in January and February 2014 and 2015, using the
84 Norwegian commercial ramp trawlers FV *Saga Sea* (96m, 6000 hp) in 2014, and the FV
85 *Juvel* (99.5 m, 8158 hp) in 2015. A 30 m long small mesh survey trawl ('Macroplankton
86 trawl') was used (see Krafft et al., 2010; 2016; Krafft & Krag, 2015), with a 6 × 6 m
87 mouth and 7 mm netting from the trawl mouth to the end of the last tapered section. The
88 trawl body and cover were supported by an outer 200 mm protection net (single 3mm PE twine).
89 The codend was 5 m long (stretched) with four similar panels joined into four selvages.
90 Each codend panel was 270 meshes wide forward and 96 meshes wide at the codline
91 following a 3N2B cutting rate. The codend was about 440 meshes in circumference
92 where the codend was closed and made of 16 mm (nominal; 15.4 mm measured)
93 diamond mesh PA netting. The actual mesh size was obtained by placing a small sample
94 of the codend netting on a flatbed scanner with no tension in the netting together with a
95 measuring unit to determine the precise mesh size. Individual meshes in the picture were
96 analysed in FISHSELECT software tool (Herrmann et al., 2009) using the built-in image


97 analysis function, and mesh size was assessed following the procedures described in
98 Sistiaga et al. (2011). Standard mesh measuring methods using the OMEGA measuring
99 gauge (Fonteyne, 2005), which are applied for larger mesh sizes, could not be used in this
100 study because the measuring jaws are too large for the small mesh sizes used in the krill
101 fishery.

102 A 26.5 m long cover comprised of 7 mm mesh was mounted to the codend to collect
103 escaping individuals. To prevent the cover net from masking the codend, two aluminium
104 hoops (4 m diameter) were used (Fig 1). The cover had a zipper to facilitate easy access
105 to the codend catch. The trawl was towed at speeds of approximately 2.5 knots as used in
106 the commercial fishery.

107 When a trawl was landed on deck, a random subsample of krill from both the codend
108 and the cover was taken. The length of the krill in the subsamples were measured from
109 the anterior margin of the eye to the tip of the telson excluding the setae,
110 following Marr (1962). The catch data was sorted into 1 mm wide length classes
111 with count numbers quantifying the number of krill belonging to each length class
112 from the codend and cover catch, respectively. The total catch and the subsample
113 were weighed for both cover and codend in all hauls.

114

115 Fig. 1: Covered codend sampling system used to collect krill codend escapees and retainers.



116

117

118 2.2 Analysis of data from sea trials to estimate codend size selectivity

119 Data was pooled from different hauls in order to estimate average size selection over hauls
120 $r_{av}(l, \mathbf{v})$ (Herrmann et al., 2012), where \mathbf{v} is a vector consisting of the parameters of the size
121 selectivity model and l is the length of the krill. The purpose of this analysis is to estimate the
122 values of the parameters \mathbf{v} that make the experimental data (averaged over hauls) most likely
123 to be observed, assuming that the selectivity model is able to describe the data sufficiently
124 well. Therefore, expression (1) was minimized with respect to parameters \mathbf{v} , which is
125 equivalent to maximizing the likelihood for the observed data in form of the length-dependent
126 number of krill retained in the codend (nR_{jl}) versus those escaping to the cover (nE_{jl}):

$$127 - \sum_{j=1}^k \sum_l \left\{ \frac{nR_{jl}}{qR_j} \times \ln(r_{av}(l, \mathbf{v})) + \frac{nE_{jl}}{qE_j} \times \ln(1.0 - r_{av}(l, \mathbf{v})) \right\} \quad (1)$$

128 The outer summation in (1) is over k hauls conducted and the inner summation is over length
129 classes l . qR_j and qE_j are the sampling factors for the fraction of krill length measured in the
130 codend and cover, respectively.

131 Four different models were chosen as basic candidates to describe $r_{av}(l, \mathbf{v})$: Logit, Probit,
132 Gompertz and Richard (Wileman et al., 1996). The first three models are fully described by
133 the two selection parameters L50 (length of krill with 50% probability of being retained) and
134 SR (difference in length between krill with 25% and 75% probability of being retained,
135 respectively). The Richard model requires one additional parameter ($1/\delta$) that describes the
136 asymmetry of the curve. The formulas for the four selection models, together with additional
137 information, can be found in Wileman et al. (1996). In addition to the four classical size
138 selection models (Logit, Probit, Gompertz, Richard), which assume that all individual krill
139 entering the codend are subject to the same size selection process, we also considered one

140 additional model that we refer to as the double logistic model DLogit (Herrmann et al., 2016).
 141 The Dlogit model is constructed by assuming that a fraction C_1 of krill entering the codend
 142 will be subject to one logistic size selection process with parameters $L50_1$ and SR_1 while the
 143 remaining fraction $(1.0 - C_1)$ will be subject to an additional logistic size selection process but
 144 with parameters $L50_2$ and SR_2 . The rationale behind considering the DLogit model for the
 145 codend size selection of krill is the expectation that the selection process may constitute more
 146 than one process. Therefore, a total of five models were considered for $r_{av}(l, \nu)$:

147 $r_{av}(l, \nu) =$

$$\left\{ \begin{array}{l}
 \textit{Logit}(l, L50, SR) \\
 \textit{Probit}(l, L50, SR) \\
 \textit{Gompertz}(l, L50, SR) \\
 \textit{Richard}(l, L50, SR, 1/\delta) \\
 \textit{DLogit}(l, C_1, L50_1, SR_1, L50_2, SR_2) = C_1 \times \textit{Logit}(l, L50_1, SR_1) + (1.0 - C_1) \times \textit{Logit}(l, L50_2, SR_2)
 \end{array} \right.$$

149 (2)

150 Each of the five models were fitted in (1). Selection of the best model of the five considered
 151 in (2) was carried out by comparing the AIC values for the model fit in (1). The selected
 152 model is the one with the lowest AIC value (Akaike, 1974). Evaluating the ability of a model
 153 to describe the data sufficiently is based on calculating the corresponding p -value, which
 154 expresses the likelihood of obtaining at least as big a discrepancy between the fitted model
 155 and the observed experimental data as would be expected by coincidence. Therefore, for the
 156 fitted model to be a candidate to model the size selection data, this p -value should not be
 157 below 0.05 (Wileman et al., 1996). In the case of a poor fit statistic (p -value < 0.05), the
 158 residuals were inspected to determine whether the result was due to structural problems when
 159 modeling the experimental data using the different selection curves or if it was due to
 160 overdispersion in the data (Wileman et al., 1996).

161 Once the specific size selection model was identified, bootstrapping was applied to estimate
162 the confidence limits for the average size selection. We applied the software tool SELNET
163 (Herrmann et al., 2012) for size selection analysis and utilized the double bootstrap method
164 implemented in this tool to obtain confidence limits for the size selection curve and the
165 corresponding parameters. This bootstrapping approach is identical to the one described in
166 Millar (1993) and takes both within-haul and between-haul variation into consideration. Each
167 of the 1000 bootstrap repetitions conducted resulted in a “pooled” set of data which was
168 analyzed using the identified selection model. The bootstrap results were used to estimate the
169 Efron percentile 95% confidence limits for the selection curve and its parameters (Herrmann
170 et al., 2012).

171 *2.3 Assessing contribution to full trawl size selectivity from trawl body*

172 The commercial trawl used by Krag et al. (2014) was a four panel *Omega 7* krill trawl
173 having a 400m² mouth opening (20 *20m) and a total length of about 220m. The trawl
174 was supported by an outer netting ranging from 400mm in 2*6mm PE in the mouth area
175 to 144mm in 2*4mm PE in the codend. 20 N-cut in-liner sections in 16mm PA netting
176 were sequentially attached from the mouth of the trawl to the codend. These in-liners
177 were only attached in the forward end and there was about 1m overlap between in-liner
178 sections. The codend was about 50m long having about 2000 meshes in circumference.
179 The entire codend section was supported by an arrangement of roundstraps and lastridge
180 ropes to provide strength to the section. The codend used during the experimental fishing in
181 this study was made of the exact same netting as used in both the codend and the trawl body
182 in the trials reported in Krag et al. (2014). This means that the two diamond mesh codends are
183 identical with respect to at least two of the most important factors, mesh size and twine
184 properties, for determining codend size selectivity (O'Neill & Herrmann, 2007). For fish
185 trawls number of meshes in codend circumference have been found to influence size selection

186 in diamond mesh codends by affecting the openness of the meshes (Herrmann et al., 2007;
187 O'Neill and Herrmann, 2007; O'Neill et al., 2008; Wienbeck et al., 2011; Tokaç et al., 2016).
188 However, for the small mesh krill codends we expect that the water flow acting on the netting
189 will keep the meshes open and therefore lowering the potential influence of number of meshes
190 in circumference on the codend size selection of krill. Therefore, despite not all codend design
191 factor are identical, including number of meshes incircumference, we assume for explorative
192 purposes that the two codends would have approximately similar size selectivity. Considering
193 that the codend was attached to a small meshed survey trawl in the current study and to a
194 commercial trawl in the study by Krag et al. (2014) we could interpret the difference in size
195 selection between the experiments to be mainly due to size selection in the commercial trawl
196 body as opposed to the codend. Therefore, any significantly higher retention probabilities for
197 the size selection curve in the current study in comparison to the full trawl and codend size
198 selectivity curve of Krag et al. (2014) are assumed to be caused by size selection in the
199 commercial trawl body in Krag et al. (2014).

200 If we look at the size selection of the whole net from Krag et al. (2014) $r_{total}(l)$ as a
201 sequential process we get:

$$\begin{aligned} r_{total}(l) &= r_{body}(l) \times r_{codend}(l) \\ &\quad \downarrow \\ r_{body}(l) &= \frac{r_{total}(l)}{r_{codend}(l)} \end{aligned} \quad (2)$$

203 Where $r_{body}(l)$ is the size selectivity in the main trawl body and $r_{total}(l)$ is the full trawl size
204 selectivity from Krag et al. (2014).

205 By using (2) and $r_{total}(l)$ from Krag et al. (2014) and the estimate for $r_{codend}(l)$ from the
206 dataset in this study, an estimate for $r_{body}(l)$ for the commercial trawl applied by Krag et al.
207 (2014) was obtained. 95% confidence intervals for $r_{body}(l)$ are based on the two bootstrap

208 populations of results (1000 bootstrap repetitions in each) from $r_{codend}(l)$ in the current study
209 and $r_{total}(l)$ from Krag et al. (2014), respectively. As these values were obtained
210 independently, a new bootstrap population of results for $r_{body}(l)$ was created using:

$$211 \quad r_{body}(l)_i = \frac{r_{total}(l)_i}{r_{codend}(l)_i} \quad i \in [1 \dots 1000] \quad (3)$$

212 Where i denotes the bootstrap repetition index. As the sampling was random and independent
213 for the two groups of results (the current study and Krag et al. (2014)) it is valid to generate
214 the bootstrap population of results for the ratio based on (3) using two independently
215 generated bootstrap files (Moore et al., 2003). Based on the bootstrap population we can
216 obtain Efron 95% percentile confidence limits for $r_{body}(l)$ as described above. This analysis
217 was conducted using the analysis tool SELNET.

218

219 *2.4 Ratio of release form codend and trawl body to full trawl*

220 To quantify the length dependent release potential of the codend and the trawl body relative to
221 that of the complete trawl the following length dependent release ratios were calculated:

$$222 \quad \begin{aligned} e_{codend}(l) &= \frac{1.0 - r_{codend}(l)}{1.0 - r_{total}(l)} \\ e_{body}(l) &= \frac{1.0 - r_{body}(l)}{1.0 - r_{total}(l)} \end{aligned} \quad (4)$$

223 In (4) the estimated $r_{codend}(l)$ and $r_{body}(l)$ as described in the previous two sections are
224 used, in addition to $r_{total}(l)$ from Krag et al. (2014). Efron percentile 95% confidence
225 intervals for $e_{codend}(l)$ and $e_{body}(l)$ were obtained by creating a new bootstrap file
226 following the approach described for $r_{body}(l)$ in the last section.

227 **3. Results**

228 *3.1 Codend size selection obtained from sea trials conducted in this study*

229 A total of eight valid hauls were carried out during the sea trials in 2014/2015. Table 1
230 summarizes the catch data from these hauls. Fishing was based on acoustic registrations of
231 krill swarms resulting in relatively short towing times ranging from 13 to 57 minutes
232 (Table 1).

233

234 Table 1: Catch data and haul information. Haul 1 and 2 are from the 2014 cruise while the
235 remaining hauls are from the 2015 cruise.*: from time the gear is at fishing depth until it is on
236 deck again.

Haul ID (j)	Number of length measurements from codend (nR_j)	Number of length measurements from cover (nE_j)	Sampling factor for codend (qR_j)	Sampling factor for cover (qE_j)	Catch in codend (kg)	Catch in cover (kg)	Towing duration (min)*	Maximum towing depth (m)
1	332	292	0.0015	0.0050	108	22	13	60
2	481	270	0.0053	0.0450	61	3.5	19	111
3	246	88	0.0137	0.0534	10	0.5	34	155
4	237	40	0.1155	0.2780	1	0.05	47	160
5	225	345	0.0016	0.0198	58	6	43	123
6	249	345	0.0019	0.0222	50	7	27	155
7	326	322	0.0180	0.2050	9	0.5	33	98
8	414	442	0.0018	0.0086	15	0.25	57	106

237

238 Length measurements were obtained for a total of 4654 krill during the cruises and these data
239 form the basis for the analysis of codend size selection.

240

241

242 Each of the five size selection models considered (section 2.2) were fitted to the pooled size
243 selection data. Table 2 shows the AIC values for the fit of each model to the experimental
244 data and it is clear that average size selectivity was best described by the DLogit model.

245 Therefore the Dlogit model is selected to represent the codend size selection (Fig. 2) it is


246 Table 2: AIC values for models. The model with lowest AIC value is highlighted in bold.

Model	AIC value
Logit	807872.17
Probit	808023.37
Gompertz	807795.25
Richard	807797.31
DLogit	807050.66

247

248

249 Fig. 2: On the top plot fit of the DLogit size selection model (black curve) to the
 250 experimental retention rates (white diamond marks). The grey curve represents the raised
 251 codend catch from the eight valid hauls and the black broken curve represents the raised cover
 252 catch. The bottom plot shows the deviance residuals for the fit of the DLogit model to the
 253 experimental data.



254

255

256 The fact that the DLogit model provided the best fit could indicate that size selection in a
 257 diamond mesh codend involves more than one size selection process, which is potentially
 258 caused by krill having few contacts with the mesh that facilitate escape in the codend
 259 (Frandsen et al., 2010; Herrmann et al., 2016). The two sets of selection parameters ($L50_1$,
 260 SR_1) and ($L50_2$, SR_2) can be interpreted as the selection parameters to represent the two
 261 different selection processes accounted for by the DLogit model (Table 3). The difference in
 262 values for $L50_1$ and $L50_2$ estimated at respectively 32.55 mm and 25.02 mm indicate a
 263 considerable difference in those two selection processes. The p -value < 0.05 could indicate
 264 problems describing the experimental data, but as the deviation between experimental rates
 265 and the fitted curve as the deviance residual plot (Fig. 2) did not show any systematic patterns
 266 as only few consecutive residual values was found to have same sign. Therefore, it was
 267 assumed that the low p -value was caused by overdispersion in the data probably resulting
 268 from working with subsampled and data pooled over hauls. Based on this, it was assumed that
 269 the DLogit model can be applied to describe the size selection of krill in the codend.

270 Table 3: Selection parameters and corresponding fit statistics for DLogit modelling of codend
 271 selectivity data. Values in () represent 95% confidence limits.


L50 (mm)	26.04 (13.82-29.19)
SR (mm)	7.07 (1.65-27.19)
C_1	0.4361 (0.0346-0.6889)
$L50_1$ (mm)	32.55 (28.17-50.00)
SR_1 (mm)	12.73 (1.00-50.00)
$L50_2$ (mm)	25.02 (16.87-33.18)
SR_2 (mm)	2.69 (1.00-26.35)
Deviance	213.75
DOF	31
P-value	<0.0001

272

273 *3.2 Comparison with full trawl selectivity from former study and predicting trawl* 274 *body size selection for trawl in the former study*

275 The estimated codend size selectivity curve was compared with the full trawl selectivity curve
 276 obtained by Krag et al. (2014) (Fig. 3).

278 Fig. 3: Size selectivity for: full trawl, codend and trawl body. Top: Comparison of size
 279 selectivity curves for the codend in the current study (black curve) and for the full trawl by
 280 Krag et al. (2014) (grey curve). Bottom: Predicted size selection curve for the trawl body in
 281 the commercial trawl applied by Krag et al. (2014). Broken curves represent 95% confidence
 282 bands.



283

284

285 From Fig. 3 it is clear that the codend retains significantly higher proportions of krill between
 286 27 and 33 mm in comparison to the full trawl (Krag et al., 2014). As it is assumed that codend

287 size selection was similar in both studies, it is likely that this difference is caused by size
288 selection processes in the trawl body in the commercial trawl applied by Krag et al. (2014).
289 For larger krill (37-50 mm) the codend size selection curve is estimated to have a slightly
290 lower retention rate than the full trawl, which violates the assumption that the two codends
291 have similar size selection. However, the confidence intervals of the two curves clearly
292 overlap for krill of these sizes and therefore this result is not a violation of the assumption
293 regarding similar codend size selection. Based on the size selection curves for the codend and
294 the full trawl (Fig. 3, top), size selection in the trawl body for the commercial trawl applied by
295 Krag et al. (2014) was predicted based on the method described in section 2.3 (Fig. 3,
296 bottom).


297

298

299 From Fig. 3 it was predicted that the trawl body enables release of krill up to about 37 mm in
300 length because the size selection curve first reach full retention above that size. Considering
301 the confidence bands, significant size selectivity for krill ranging from 23-33 mm is predicted.
302 The predicted trawl body release efficiency is high for krill up to 30 mm in length with less
303 than 25% retained, demonstrating a considerable size selection process in the trawl body of
304 the commercial trawl. For krill approximately 28 mm long, the upper confidence limit for the
305 size selection curve is below 50%, demonstrating that more than 50% of krill at that size
306 entering the trawl will be released through the trawl body. The contributions of both the trawl
307 body and the codend in size selection for the commercial trawl can be further illustrated by
308 quantifying the length dependent fraction of the full trawl escape that can be obtained by the
309 trawl body and codend provided from a standalone deployment. This is obtained by the
310 method described in section 2.4, with results shown in Fig. 5.

311

312 Fig. 5: Fraction of full trawl krill escape rate obtainable for the trawl body alone (top) and
313 codend (bottom). Broken curves represent 95% confidence bands.



314

315

316

317 From Fig. 5 it is predicted that more than 80% of the full trawl escape rate can be obtained in
318 the trawl body for krill up to 30 mm in length. For some sizes of krill, the fraction is very high
319 with the lower significance limit above the 50% fraction (value above 0.5). In contrast, for the
320 codend the upper limit for the release fraction does not exceed 75% for sizes of krill between
321 27 and 33 mm in length. The results in Fig. 5 clearly depict the potential contribution of both
322 the trawl body and the codend in total krill release through the meshes of the commercial
323 trawl.

324

325 **4. Discussion**

326 Detailed quantification of the size selection of both the codend and the trawl body is essential
327 to estimate escape mortality, and total removal by the fishery, for the optimization of gear
328 design and the technical regulation of a fishery. In this study, the covered codend method was
329 used to investigate size selectivity for Antarctic krill using a 16 mm diamond mesh codend.
330 Codend selectivity was best described by the double logistic model, indicating that more than
331 one process affects codend size selectivity. It is possible that only a small fraction of krill
332 meet the codend mesh at an optimal orientation for escape and so a double logistic model is
333 necessary to describe size selection in the codend, as opposed to a single logistic for the full
334 trawl, as in Krag et al. (2014).

335 By combining new codend size selection results obtained within this study with results for full
336 trawl size selectivity obtained in a former study, this study provided an insight into the size
337 selection process in the main trawl body of the commercial trawl, contributing to an
338 understanding of full trawl size selectivity.

339 This analysis demonstrates that the trawl body contributes significantly to the size selection
340 process and that size selectivity of Antarctic krill is affected by the trawl body of commercial
341 trawls and by the attached codend. Conclusions from this study are based on the assumption
342 that the codend in the current study provides similar size selectivity for krill as the one used in
343 the trials described by Krag et al. (2014). The same type of netting was used for both
344 experiments, but it is possible that different fishing conditions could affect the predicted size
345 selectivity. However, we expect the potential maximum difference in codend size selection is
346 well within the confidence bands obtained in this study and thus is reflected in the
347 uncertainties for the trawl body size selectivity.

348 The results for trawl body size selectivity demonstrate considerable size selection for krill <32
349 mm using commercial 16 mm mesh. Therefore, this study has shown that commercial trawl
350 bodies in krill-fishery can generally contribute to size selectivity. Nevertheless, a number of
351 parameters (e.g. tapering of body) will influence the specific selectivity. Therefore, the
352 specific findings about size selectivity of trawl body are not general, but an example for this
353 specific gear used in Krag et al. (2014). Other trawl designs might have different selectivity.
354 In this respect, it is important to mention that some commercial krill trawl designs include
355 “flapper-panels”, which prevent “stickers” and increase net avoidance (active or passive),
356 enhancing transportation towards the codend (Bakketeig et. al, 2017). With such flappers
357 mounted, the size selectivity in the trawl body could potentially be considerably lower than
358 that estimated in Krag et al. (2104).

359 The current study found that for krill, size selectivity occurs across the entire trawl. This is
360 different to what is observed for most fish species, but it is in keeping with results from
361 fisheries targeting smaller crustaceans (e.g. Polet, 2000). The results of the current study
362 revealed that a substantial fraction of size selectivity for Antarctic krill occurred in the trawl
363 body ahead of codend. Such findings can be incorporated into fisheries management, where
364 technical regulations should consider the entire trawl and not just the codend section.

365 **Acknowledgements**

366 The Institute of Marine Research in Norway and The Norwegian Research Council provided
367 financial support (grant number 243619) for this study. We thank the two Norwegian fishing
368 companies Rimfrost AS and Aker Biomarine AS, for providing their vessels and crew for the
369 collection of data and thus making this field experiment possible. Finally, we want to express
370 our gratitude to the anonymous reviewers for their valuable comments during the review
371 process which helped improving this manuscript greatly.

372

373 **References**

374 Akaike, H., 1974. A new look at the statistical model identification. IEEE Trans. Autom.
375 Control 19, 716–723.

376 Bakketeig, I.E., Hauge M., Kvamme, C. (Editors), 2017. Havforskningsrapporten 2017.

377 Fisken og havet, særnr. 1–2017 (in Norwegian). ISBN: 0802 0620.

378 Budzinski, E., Bykowski, P., Dutkiewics, D., 1985. Possibilities of processing and marketing
379 of products made from Antarctic krill. FAO Fisheries Technical Paper No. 268. Rome: FAO.
380 46p.

381 Frandsen, R., Herrmann, B., Madsen, N., 2010. A simulation-based attempt to quantify the
382 morphological component of size selectivity of *Nephrops norvegicus* in trawl codends. Fish.
383 Res. 101, 156-167.

384 Herrmann, B., Priour, D., Krag, L.A., 2007. Simulation-based study of the combined effect on
385 codend size selection for round fish of turning mesh 90 degree and of reducing the number of
386 meshes in the circumference. Fish. Res. 84, 222–232.

387 Herrmann, B., Krag, L.A., Frandsen, R.P., Madsen, N., Lundgren, B., Staehr, K.J., 2009.
388 Prediction of selectivity from morphological conditions: methodology and a case study on cod
389 (*Gadus morhua*). Fish. Res. 97, 59–71.

390 Herrmann, B., Sistiaga, M., Nielsen, K.N., Larsen, R.B., 2012. Understanding the size
391 selectivity of redfish (*Sebastes* spp.) in North Atlantic trawl codends. J. Northw. Atl. Fish.
392 Sci., 44, 1-13.

393 Herrmann, B., Krag, L.A., Feekings, J., Noack, T., 2016. Understanding and predicting size
394 selection in diamond mesh codends for Danish seining: a study based on sea trials and
395 computer simulations. *Marine and Coastal Fisheries* 8, 277-291.

396 Kanda, K., Takagi, K., Seki, Y. 1982. Movement of larger swarms of Antarctic krill
397 *Euphausia superba* off Enderby Land during 1976–1977 season. *J Tokyo Univ Fish* 68:24–
398 42.

399 Krafft, B.A., Krag, L.A., Engås, A., Nordrum, S., Bruheim, I., Herrmann, B., 2016.
400 Quantifying the escape mortality of trawl caught Antarctic krill (*Euphausia superba*).
401 PlosONE 11(9):e0162311 doi:10.1371/journal.pone.0162311.

402 Krafft, B.A., Krag, L.A., 2015. Assessment of mortality of Antarctic krill (*Euphausia*
403 *superba*) escaping from a trawl. *Fisheries Research* 170:102-105.

404 Krafft, B.A., Melle, W., Knutsen, T., Bagøien, E., Broms, C., Ellertsen, B., Siegel, V., 2010.
405 Distribution and demography of Antarctic krill in the Southeast Atlantic sector of the
406 Southern Ocean during the austral summer 2008. *Polar Biology* 33:957-968

407 Krag, L.A., Herrmann, B., Iversen, S., Engås, A., Nordrum, S., Krafft, B.A., 2014. Size
408 selection of Antarctic krill (*Euphausia superba*) in trawls. *PloS ONE* 9(8): e102168.
409 doi:10.1371/journal.pone.0102168.

410 Lochhead, J.H., 1961. Locomotion. In: Waterman T.H. (Ed), *The physiology of the*
411 *Crustacea*. Academic Press, New York, pp 313–356.

412 Marr, J. 1962. The natural history and geography of the Antarctic krill *Euphausia superba*.
413 *Discov Rep* 32:33–46

414 Millar, R. B. 1993. Incorporation of between-haul variation using bootstrapping and
415 nonparametric estimation of selection curves. Fisheries Bulletin 91, 564-572.

416 Moore, D.S, McCabe, G.P., Duckworth, W.M., Sclove, S.L, 2003. Practice of Business
417 Statistics using data for decisions. Published by W. H. Freeman. ISBN 10: 0716757230 /
418 ISBN 13: 9780716757238.

419 Newland, P.L., Chapman, C.J., 1989. The swimming and orientation behaviour of the Norway
420 lobster, *Nephrops norvegicus* (L), in relation to trawling. Fisheries Research 8, 63–80.

421 O’Neill, F.G., Herrmann, B., 2007. PRESEMO- a predictive model of codend selectivity- a
422 tool for fisheries managers. ICES Journal of Marine Science 64, 1558-1568.

423 O’Neill, F.G., Graham, N., Kynoch, R.J., Ferro, R.S.T., Kunzlik, P.A., Fryer, R.J., 2008. The
424 effect of varying cod-end circumference, inserting a ‘flexi-grid’ or inserting a Bacoma type
425 panel on the selectivity of North Sea haddock and saithe. Fish.Res. 94, 175–183

426 Polet, H., 2000. Codend and whole trawl selectivity of a shrimp beam trawl used in the North
427 Sea. Fisheries Research, 48, 167-183.

428 Sistiaga, M., Herrmann, B., Nielsen, K.N., Larsen, R.B., 2011. Understanding limits to cod
429 and haddock separation using size selectivity in a multi-species trawl fishery: an application
430 of FISHSELECT. Can. J. Fish. Aquat. Sci. 68, 927–940.

431 Tokaç, A., Herrmann, B., Gökçe, G., Krag, L. A., Nezhade, D. S., Lök, A., Kaykaç, H.,
432 Aydın, C., Ulaş, A., 2016. Understanding the size selectivity of red mullet (*Mullus barbatus*)
433 in Mediterranean trawl codends: A study based on fish morphology. Fish. Res. 174, 81–93.

434 Wardle, C.S., 1993. Fish behaviour and fishing gear. In: Behaviour of teleost fishes, 2nd
435 edition. Pitcher TJ (Ed). Chapman and Hall, London. pp 609–644.

436 Wienbeck, H., Herrmann, B., Moderhak, W., Stepputtis, D, 2011. Effect of netting direction
437 and number of meshes around on size selection in the codend for Baltic cod (*Gadus morhua*).
438 Fisheries Research 109: 80-88.

439 Wileman, D., Ferro, R.S.T., Fonteyne, R., Millar, R.B. (Eds.), 1996. Manual of Methods of
440 Measuring the Selectivity of Towed Fishing Gears. , ICES Coop. Res. Rep., No. 215, 126 pp.