Protective effect of in-feed specific IgM towards *Yersinia ruckeri* in rainbow trout

Chettri, Jiwan Kumar; Al-Jubury, Azmi; Hansen, Marie B.; Lihme, Allan; Dalsgaard, Inger; Buchmann, Kurt; Heegaard, Peter M. H.

Published in:
Fish and Shellfish Immunology

Link to article, DOI:
[10.1016/j.fsi.2019.08.024](https://doi.org/10.1016/j.fsi.2019.08.024)

Publication date:
2019

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Chettri, J. K., Al-Jubury, A., Hansen, M. B., Lihme, A., Dalsgaard, I., Buchmann, K., & Heegaard, P. M. H. (2019). Protective effect of in-feed specific IgM towards *Yersinia ruckeri* in rainbow trout. *Fish and Shellfish Immunology*, 93, 934-939. <https://doi.org/10.1016/j.fsi.2019.08.024>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Journal Pre-proof

Protective effect of in-feed specific IgM towards *Yersinia ruckeri* in rainbow trout

Jiwan K. Chettri, Azmi Al-Jubury, Marie B. Hansen, Allan Lihme, Inger Dalsgaard, Kurt Buchmann, Peter M.H. Heegaard

PII: S1050-4648(19)30824-1

DOI: <https://doi.org/10.1016/j.fsi.2019.08.024>

Reference: YFSIM 6366

To appear in: *Fish and Shellfish Immunology*

Received Date: 22 February 2019

Revised Date: 8 August 2019

Accepted Date: 9 August 2019

Please cite this article as: Chettri JK, Al-Jubury A, Hansen MB, Lihme A, Dalsgaard I, Buchmann K, Heegaard PMH, Protective effect of in-feed specific IgM towards *Yersinia ruckeri* in rainbow trout, *Fish and Shellfish Immunology* (2019), doi: <https://doi.org/10.1016/j.fsi.2019.08.024>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier Ltd.

Protective effect of in-feed specific IgM towards *Yersinia ruckeri* in rainbow trout

Jiwan K. Chettri^{ab}, Azmi Al-Jubury^b, Marie B. Hansen^c, Allan Lihme^c, Inger Dalsgaard^a, Kurt Buchmann^b and Peter M. H. Heegaard^a

^aNational Veterinary Institute, Technical University of Denmark, Frederiksberg C, Denmark

^bDepartment of Veterinary and Animal Science, Faculty of Health and Medical Sciences, University of Copenhagen, Frederiksberg C, Denmark

^c Lihme Protein Solutions, Kongens Lyngby, Denmark

Abstract

Tightened regulations and an environmentally friendly approaches in fish production have greatly reduced the use of antibiotics but green solutions are continuously being explored. The use of functional feed may have a potential in the aquaculture sector in securing biomass and minimizing the loss from disease. In the present study, we tested the concept that blood from the fish slaughterhouse can be used for mass purification of specific antibodies which subsequently can be used for feeding fish and thereby confer protection against diseases. IgM was purified from serum from *Yersinia ruckeri* vaccinated rainbow trout and an IgM sandwich ELISA was developed for quantification of rainbow trout IgM. The purified IgM was encapsulated in alginate microparticles and top-coated in fish feed. IgM re-extracted from the alginate microparticles was shown to retain high reactivity towards *Y. ruckeri* antigens indicating that its bioactivity remained intact after encapsulation. IgM release from the alginate microparticles was only observed at high pH (pH 8.2) and minimal at low pH, indicating protection of IgM at low pH in the fish stomach during passage. In a feeding – challenge experiment (feeding 1 week before *Y. ruckeri* challenge and for two weeks following challenge), a statistically non-significant 10% lower mortality was observed in the high dose (400 µg IgM/fish/day fed over 3 weeks) group.

Introduction

The annual growth of the aquaculture sector has exceeded 7.8% in the 1990-2010 period [1] making it the fastest growing food producing sector globally. Intensive fish farming systems face a number

of challenges, including increased risk of disease outbreaks with associated environmental and animal welfare issues, but immunoprophylactic control measures may reduce disease problems. Vaccination is generally applied when fish are immune-competent and have well-developed immune organs but vaccine-induced protection may not always be achievable when considering young and not fully immune-competent stages. Alternative methods of protection, such as passive immunization using specific antibodies, may confer protection to young and vulnerable fish. A series of studies have documented that vaccine-induced protection is in many cases directly related to production of specific antibodies [2-4] which in rainbow trout comprise IgM, IgT and IgD where IgM represents the systemic immunoglobulin, whereas IgT and IgD contribute to protection of mucosal surfaces [5-7]. Passive immunization of fish against bacterial pathogens, such as *Vibrio anguillarum*, *V. vulnificus* and *Streptococcus* sp., has been demonstrated to confer protection in larger fish [8-10]. Oral administration of specific antibodies may be a solution for small fish however the conditions (low pH, proteolytic activity) in the stomach represent a problem. Digestive degradation of antigens by low pH and proteolytic conditions and antigens or proteins should be protected in such an environment. The use of a low cost biodegradable polymer, such as alginate, may offer resistance to proteolysis without being immunogenic and toxic. Alginate is one of the most used polymers for microencapsulation [11] and the alginate from brown algae has been successfully used for oral vaccination of Atlantic salmon, *Salmon salar* against *Y. ruckeri* [12]. Recently, it was shown that the feeding of weaning piglets with IgG purified from pig slaughterhouse blood significantly reduced disease and shedding of pathogenic bacteria, and at the same time maintained ileal microbial diversity, suggesting its applicability in controlling post-weaning diarrhea in piglets [13]. The present study was conducted to test a parallel concept in fish for controlling early stage diseases by making use of fish slaughterhouse blood containing specific immunoglobulins against different pathogens obtained through active vaccination or through

environmental exposure during the grow-out phase. To test this concept, IgM with high activity against *Y. ruckeri* was purified from fish vaccinated with *Y. ruckeri* and encapsulated in alginate microparticles. The IgM containing alginate microparticles were then top-coated in fish feed and fed to rainbow trout fry to evaluate the protective effect of orally administered immunoglobulin against subsequent *Y. ruckeri* challenge by immersion exposure.

2. Materials and Methods

2.1. Vaccination

Rainbow trout (N=35, total body weight 0.5- 0.6 kg) were vaccinated with an experimental vaccine containing formalin-killed whole cell *Y. ruckeri* bacterin (biotype 1 and 2) [14] in combination with Freund's incomplete adjuvant (FIA) (vaccine:adjuvant, (vol:vol), 1:1; water-in-oil emulsion). The final adjuvanted vaccine contained 3×10^{10} CFU/ml and was administered by intraperitoneal injection (IP: 0.1 ml). Booster vaccination (IP) was applied three weeks (315 degree days) post primary vaccination and blood samples were taken 7 weeks (735 degree days) post primary vaccination. Fish were kept at 15°C and daily fed with pelleted feed (2% of biomass). Blood samples from unvaccinated fish (N= 50; 1.5 – 2.5 kg) were used as a control.

2.2. Blood sampling

Fish were euthanized by MS222 (ethyl 3-aminobenzoate methanesulfonate; Sigma-Aldrich A5040) immersion (concentration 100 mg/l) and blood samples from 35 immunized and 10 non-immunized fish by caudal vein puncture (BD Vacutainer®). Serum was recovered by centrifugation ($3000 \times g$) for 10 min at 4 °C and kept at -80°C for later analysis.

2.3. ELISA

Specific ELISA to measure antibody titer

Serum samples were analysed for *Y. ruckeri* specific antibody titers using ELISA [2, 15]. In brief: microtiter plates (flat-bottom 96-well plates, MaxiSorp™, Nunc) were coated with sonicated

bacterial lysate at 5 µg/ml protein concentration (*Y. ruckeri* O1 biotype 2) in bicarbonate coating buffer (Sigma-Aldrich Cat. No. C3041, Denmark) and incubated overnight in a refrigerator. Non-specific binding sites were hereafter blocked with 2% bovine serum albumin (BSA, Sigma-Aldrich, A4503) in PBS with 0.1% Tween 20 for 1 h at room temperature followed by four times washing in PBS-T (PBS with 0.1% Tween 20) and plates were stored at -20°C until further use. Serum samples in duplicate were incubated overnight at 4°C in antigen coated plates, followed by washing and incubation with mouse anti-salmonid Ig (AbD Serotec, Dusseldorf, Germany) (dilution:1:500) and HRP (horseradish peroxidase) conjugated rabbit anti-mouse IgG (AbD Serotec Dusseldorf, Germany) (dilution1:500) for 1 h at room temperature. After every step, plates were washed 4 times in PBS-T. To optimize a working dilution, two representative samples from each group were run in 10 fold dilution series (1/50 to 1/500 000) and two serum dilutions (where the non-specific binding was minimal) were chosen for the final analysis. As a final step plates were incubated with substrate TMB (tetramethylbenzidine) PLUS (AbD Serotec, Dusseldorf, Germany) for 5-10 min and the reaction was stopped by adding 1N HCL whereafter optical density (OD) was measured at 450 nm using an Epoch Spectrophotometer (BioTek Instruments, Inc., Winooski, USA).

2.4. Preparation of IgM loaded alginate particles

Alginate encapsulation of IgM molecules was performed by a previously described method [16]. Briefly, purified concentrated IgM (91.8 mg/ml) was combined with 630 mg medium viscosity alginate from brown algae (Sigma-Aldrich, A2033) dissolved in 20 ml distilled water and mixed in a magnetic stirrer for 20 min at room temperature. The aqueous phase (alginate containing IgM) was then added slowly to an oil phase containing 37 ml octane and 3 ml of Span-80 (continuous stirring with a handheld pitched-blade homogenizer at 300 rpm). Then, 3 ml of Tween-80 was added and the emulsification procedure was continued for 1 h at room temperature at 500 rpm

whereafter 7.7 ml of 8% aqueous solution of calcium chloride was added dropwise to facilitate gelation of microparticles through ionic crosslinking by calcium cations. The mixture was stirred for 1 h at room temperature at 200 rpm. In order to harden the microparticles, 10 mL isopropyl alcohol was added and allowed to stir at room temperature for 30 mins. The microparticles were collected by centrifuging at 250 g for 10 mins and collected microparticles were washed twice with isopropyl alcohol. The final product was washed in distilled water (2x), lyophilized overnight and stored at 4°C until use. Using this procedure, four types of IgM-loaded alginate microparticles were prepared: (i) IgM from *Y. ruckeri* vaccinated fish (50 mg, 250 mg and 500 mg); (ii) IgM from unvaccinated fish (250 mg) and (iii) only plain alginate microparticles (no IgM encapsulation).

2.5. Evaluation of stability of alginate encapsulated IgM at different pH and determination of bioactivity

The release of IgM from alginate encapsulation was tested by incubating 20 mg of encapsulated microparticles in 1 ml solution at different pH (pH = 2.7, 5, 7, 7.7, 8.2 and 10, respectively (glycine, sodium acetate, PBS, milliQ water, EDTA, sodium bicarbonate and sodium phosphate) for 4 days at room temperature or at 4°C. Hereafter, 100 µl of the incubated solution was used for running the *Y. ruckeri* specific ELISA to analyse for release of bio-active IgM. A total of 32.5 µl of the incubated solution was used for running SDS-PAGE and Western blot to check if the IgM had been degraded during the process.

2.6. Coating of fish feed with IgM-loaded alginate microparticles

Fish feed (commercial 2 mm pellets, Aller Aqua A/S) was top-coated with alginate microparticles and sealed with vegetable oil. The encapsulated microparticles containing different amounts of IgM (50 mg, 250 mg and 500 mg) were added to 50 g of feed and mixed by gentle stirring while continuously being sprayed with a fine oil spray. In a similar way, 250 mg of anti-*Yersinia* IgM as

well as oxolinic acid (1.25 g/kg feed; Sigma-Aldrich O0240000) was directly top-coated with vegetable oil. All coated feed were stored at 4°C.

2.7. Feeding and challenge experiment

Rainbow trout (*Oncorhynchus mykiss*) fry produced in a re-circulated pathogen-free facility (Salmon hatchery Bornholm, Nexø, Denmark) were transported to our experimental facility at the University of Copenhagen, Denmark. Fish were acclimatized in our facility for 1 week at 15°C and distributed in the experimental tanks. A total of eight groups in duplicate (average body weight 2 g; 25 fish/tank; 16 tanks with 20 L volume) were included in the experiment. Three control groups in duplicate were included: One group fed with ordinary feed coated with plain alginate (this group was not given infection and served as uninfected control; group 1). A second group fed the same way but exposed to *Y. ruckeri* infection (infected control; group 2). The third control group was given alginate encapsulated IgM supplemented feed purified from the unvaccinated fish (250 mg IgM per 50 g feed or 200 µg IgM/fish/day) and exposed to infection (group 3). The three treatment groups were fed different levels of anti-*Yersinia ruckeri* encapsulated immunoglobulin: 50, 250 and 500 mg IgM per 50 g feed (corresponding to 40 µg IgM/fish/day, 200 µg IgM/fish/day and 400 µg IgM/fish/day; group 4, group 5 and group 6, respectively). The seventh group (group 7) of fish was fed with anti-*Yersinia ruckeri* immunoglobulin directly oil coated on feed (250 mg IgM per 50 g feed corresponding to 200 µg IgM/fish/day). Finally, group 8 was fed with antibiotic (oxolinic acid: 1.25 g/kg feed; Sigma-Aldrich O0240000) coated feed (positive control). The experimental groups are detailed in Table 1.

Fish was fed 2% of body weight (40 mg feed/fish/day) with different codes of feed comprising 40 µg IgM/fish/day, 200 µg IgM/fish/day and 400 µg IgM/fish/day. Fish were fed once in the morning by slowly adding feed to individual tanks ensuring total consumption of the added feed. After 1 week of feeding with experimental feed, fish were exposed to experimental infection: Water from

each tank was lowered to 5 L and fish were exposed (duration 4 h) to a 48 h culture of *Y. ruckeri* (100415-1/4) with a final bacterial concentration of 1.2×10^8 CFU/ml by adding 50 ml of culture broth. Hereafter clean water was added to the tank reaching a water volume 20 L. On the following day, water was exchanged with fresh tap water to remove bacteria from the tank. In the uninfected group control (Group 1, Table 1), sterile LB medium was added (sham infection) and treated otherwise as infected groups. The fish were continuously fed with experimental feed for 2 weeks after exposure whereafter the experiment was terminated. Fish were closely observed following the infection, any fish swimming abruptly, swimming at the surface with asphyxia and showing severe sign of disease were euthanized in a high dose of MS222 (ethyl 3-aminobenzoate methanesulfonate; Sigma-Aldrich A5040) and recorded as mortality. Bacteria from freshly euthanized fish (samples from head kidney, spleen and brain) were re-isolated on blood agar plates for confirmation of specific cause of death.

3. Statistical analysis

Mortality data were analyzed by Kaplan-Meier survival analysis and are presented as cumulative mortality (%). Two-tailed unpaired t-test with Welch's correction was applied for testing ELISA data. For determining the concentration of IgM, a standard curve was plotted and analyzed with non-linear regression (curve fit) with Sigmoidal four points logistic (4PL) and the concentration of unknown samples was interpolated based on the standard curve (appendix II). The data were considered significant when $P \leq 0.05$.

4. Results

4.1. ELISA

Fish vaccinated with *Y. ruckeri* antigens showed a significantly higher pathogen-specific antibody level (although with a considerable variation) than the unvaccinated control group (Fig. 1).

4.2. IgM loaded alginate microparticles

Alginate microparticles loaded with anti-*Yersinia* antibody were exposed to various pH conditions and IgM release measured using a sandwich ELISA designed to quantify IgM concentration (Appendix I & II). The highest IgM release at alkaline pH 8.2 (sodium bicarbonate), a lower release at pH 10 (sodium phosphate 0.1M) but no or minimal release of IgM at low or neutral pH. The recovered IgM from alginate beads showed a high bioactivity (binding to *Y. ruckeri*) antigens and no loss of bioactivity due to the gelation process was detected (Fig. 3). This was further confirmed by SDS-PAGE (Fig. 4-left) and Western blot (Fig. 4-right) showing that the IgM remained intact after the gelation process.

4.3. Feeding and challenge experiment

Fish were fed 2% of body weight per day corresponding to specific IgM feeding levels of 40 µg IgM/fish/day (50 mg), 200 µg IgM/fish/day (250 mg) and 400 µg IgM/fish/day (500 mg). Fish were fed IgM encapsulated feed for 1 week before challenge with *Y. ruckeri* and during the course of infection. The control IgM (from un-vaccinated fish) were offered to fish as 200 µg IgM/fish/day (250 mg). No protective effect of feeding with specific IgM on the survival from lethal challenge with *Y. ruckeri* was observed (Fig. 5). The challenge bacterium was re-isolated from all the euthanized fish and was confirmed by MALDI-TOF MS (matrix-assisted laser desorption ionization-time of flight mass spectrometry). Only the group fed with oxolinic acid (antibiotic) showed a significantly lower mortality. Rainbow trout offered feed with the highest anti-*Yersinia* IgM concentration showed a 10% lower mortality ($p>0.05$) than the infected control group and delay in onset of mortality was observed in this group. Fish fed un-specific IgM also showed a trend for a lower mortality. An enhancing effect of alginate encapsulation was noted when comparing mortality patterns between the IgM (only top-coated with oil) and IgM encapsulated in alginate microparticles (250 mg IgM) (Fig. 5).

5. Discussion

193 Fish possess an adaptive immune system with an ability to mount a specific antibody response
194 against pathogens. However, early life stages (fish larvae and fry), have not yet developed a fully
195 functional immune response and rely to a wide extent on innate immune molecules and maternal
196 transfer of vital immune factors (complement factors, SAA, serine proteases, and immunoglobulin)
197 for survival and well-being of young fish [17, 18]. Transfer of immunoglobulin from mother to
198 offspring occurs in fish [19-22] and this vertical transmission confers protection [23] which frames
199 the importance of securing immunized brood fish for generation of healthy offspring [24]. Passive
200 immunization (transfer of immune serum to naive fish by injection) has shown to enhance
201 protection for more than 2 months post administration against vibriosis caused by *Vibrio*
202 *anguillarum* [20]. Passive immunization by use of oral administration of fish serum containing
203 specific immunoglobulin from immunized fish would be a less laborious and less costly method.
204 However, preliminary trials could not document enhanced survival in rainbow trout fingerlings
205 following challenge with *Vibrio anguillarum* but the lack of protection could be due to proteolysis
206 of antibodies in the gastro-intestinal tract of salmon [25]. We evaluated the effect of alginate-
207 encapsulation to overcome the degradation in the gut as intact antibodies, protected against
208 proteolysis, could confer short-term immunity until the young fish is fully immune-competent and
209 able to respond to vaccination. This approach would be equally beneficial in protecting larger fish
210 or high-value fish which are close to harvesting but are at high risk to disease outbreak and for
211 which active vaccination is not an option. In the present study, blood harvested from vaccinated fish
212 with high antibody titers against *Y. ruckeri* was used for the production of alginate encapsulated
213 IgM microparticles which subsequently were applied for the feeding of rainbow trout fry. In order
214 to optimize precision in the study we further developed and validated a sandwich ELISA for
215 quantification of IgM from rainbow trout using a mouse monoclonal anti-salmon antibody [26, 27]
216 as both catching and detection antibody (supplementary material II). Oral active immunization

applying alginate encapsulated bacterial cells was previously demonstrated to be effective in boosting immune response and increasing protection of salmon [12, 28] which suggests that alginate encapsulation is a useful technology for delivery of cells, proteins and low molecular weight drugs [29]. In the present study alginate from brown algae was used for encapsulation of purified IgM from vaccinated or un-vaccinated fish. The *in vitro* incubation experiment suggests that IgM is protected from degradation at acidic conditions as there was no release of IgM at pH 2 and merely partial discharge of IgM occurred at higher pH (maximum at pH 8.2 and minor release at pH 10). Analysis of gut samples would have shed more light on the quantity of specific IgM delivered into the gut and its intestinal release, but was not performed in the present study. However, previous studies suggest that alginate microcapsules are retained in the gut lumen or found in the intestinal epithelium within 1.5 h and in the lamina propria 6 h after oral administration [16]. We showed that the bio-activity of IgM remained intact as re-harvested IgM from alginate microparticles showed a strong binding towards *Y. ruckeri* antigen as tested by pathogen-specific ELISA. This indicates that there is no deleterious effect of the emulsification and gelation process on the activity of anti-*Yersinia* IgM which supports previous studies showing that the immunogenicity of antigen or protein are not affected by the process [12, 16].

Our *in vivo* studies (feeding a diet containing anti-*Yersinia ruckeri* IgM) did not show a significantly enhanced survival of fish challenged with *Y. ruckeri* but a tendency towards protection (10%) was observed for the fish group fed the highest IgM dose (400 µg/fish/day). The bacterial dose used for challenge may have been too high compared to the administered amount of IgM and a lower challenge dose may allow a better segregation between differently IgM-dosed groups. Future studies should therefore elucidate if increased IgM levels in feed and/or a lowered infection pressure result in elevated protection. It may be speculated that IgM reaching the gut in feed may bind to and inactivate a pathogen in the lumen or within the intestinal wall and/or that

elevated IgM levels at bacterial entry portals (anal opening, intestine, stomach, gills, lateral line, epidermis, dorsal fins and olfactory bulb [30, 31] may delay or prevent invasion. As IgM is a high molecular weight protein its uptake from the gut to the circulation is unlikely unless M-cells assist the process but such a scenario should be further investigated.

Acknowledgments

Authors would like to thank fish keepers at the Bornholm Salmon Hatchery and at the fish keeping facility at the Technical University of Denmark. The technical assistance of Henriette Vorsholt, Kári Karbech Mouritsen and productive discussions with Lone Madsen, Chris Juul Hedegaard and Prof. Niels Lorenzen are highly appreciated. The project was funded of the Green Development and Demonstration Programme (GUDP) under “IMMFEED: Immunoglobulin for Fish Production- combating infections without antibiotics” project no. 34009-14-0795.

References

- [1] M. Troell, R.L. Naylor, M. Metian, M. Beveridge, P.H. Tyedmers, C. Folke, K.J. Arrow, S. Barrett, A.S. Crepin, P.R. Ehrlich, A. Gren, N. Kautsky, S.A. Levin, K. Nyborg, H. Osterblom, S. Polasky, M. Scheffer, B.H. Walker, T. Xepapadeas, A. de Zeeuw, Does aquaculture add resilience to the global food system?, *Proc Natl Acad Sci U S A* 111(37) (2014) 13257-63.
- [2] J.K. Chettri, S. Deshmukh, L. Holten-Andersen, R.M. Jafaar, I. Dalsgaard, K. Buchmann, Comparative evaluation of administration methods for a vaccine protecting rainbow trout against *Yersinia ruckeri* O1 biotype 2 infections, *Vet Immunol Immunopathol* 154(1-2) (2013) 42-7.
- [3] S. Deshmukh, P.W. Kania, J.K. Chettri, J. Skov, A.M. Bojesen, I. Dalsgaard, K. Buchmann, Insight from Molecular, Pathological, and Immunohistochemical Studies on Cellular and Humoral Mechanisms Responsible for Vaccine-Induced Protection of Rainbow Trout against *Yersinia ruckeri*, *Clinical and vaccine immunology : CVI* 20(10) (2013) 1623-41.
- [4] K. Rømer Villumsen, I. Dalsgaard, L. Holten-Andersen, M.K. Raida, Potential Role of Specific Antibodies as Important Vaccine Induced Protective Mechanism against *Aeromonas salmonicida* in Rainbow Trout, *PLoS ONE* 7(10) (2012) e46733.
- [5] I. Hordvik, Immunoglobulin isotypes in Atlantic salmon, *Salmo salar*, *Biomolecules* 5(1) (2015) 166-77.
- [6] Y.-A. Zhang, I. Salinas, J. Li, D. Parra, S. Bjork, Z. Xu, S.E. LaPatra, J. Bartholomew, J.O. Sunyer, IgT, a primitive immunoglobulin class specialized in mucosal immunity, *Nature Immunology* 11(9) (2010) 827-835.
- [7] J. Skov, J.K. Chettri, R.M. Jaafar, P.W. Kania, I. Dalsgaard, K. Buchmann, Effects of soluble immunostimulants on mucosal immune responses in rainbow trout immersion-vaccinated against *Yersinia ruckeri*, *Aquaculture* 492 (2018) 237-246.
- [8] M. Akhlaghi, Passive immunisation of fish against vibriosis, comparison of intraperitoneal, oral and immersion routes, *Aquaculture* 180(3-4) (1999) 191-205.

- [9] M. Akhlaghi, B.L. Munday, R.J. Whittington, Comparison of passive and active immunization of fish against streptococcosis (enterococcosis), *Journal of Fish Diseases* 19(3) (1996) 251-258.
- [10] B.R. LaFrentz, C.A. Shoemaker, Passive transfer of serum from tilapia vaccinated with a *Vibrio vulnificus* vaccine provides protection from specific pathogen challenge, *Aquaculture* 442 (2015) 16-20.
- [11] L. Chan, H. Lee, P. Heng, Production of alginate microspheres by internal gelation using an emulsification method, *International journal of pharmaceutics* 242(1-2) (2002) 259-62.
- [12] B. Ghosh, T.D. Nguyen, P.B.B. Crosbie, B.F. Nowak, A.R. Bridle, Oral vaccination of first-feeding Atlantic salmon, *Salmo salar* L., confers greater protection against yersiniosis than immersion vaccination, *Vaccine* 34(5) (2016) 599-608.
- [13] C.J. Hedegaard, M.L. Strube, M.B. Hansen, B.K. Lindved, A. Lihme, M. Boye, P.M.H. Heegaard, Natural Pig Plasma Immunoglobulins Have Anti-Bacterial Effects: Potential for Use as Feed Supplement for Treatment of Intestinal Infections in Pigs, *PLOS ONE* 11(1) (2016) e0147373.
- [14] R.M. Jaafar, J.K. Chettri, I. Dalsgaard, A. Al-Jubury, P.W. Kania, J. Skov, K. Buchmann, Effects of adjuvant Montanide™ ISA 763 A VG in rainbow trout injection vaccinated against *Yersinia ruckeri*, *Fish & Shellfish Immunology* 47(2) (2015) 797-806.
- [15] J.K. Chettri, J. Skov, R.M. Jaafar, B. Krossøy, P.W. Kania, I. Dalsgaard, K. Buchmann, Comparative evaluation of infection methods and environmental factors on challenge success: *Aeromonas salmonicida* infection in vaccinated rainbow trout, *Fish & Shellfish Immunology* 44(2) (2015) 485-495.
- [16] B. Ghosh, B.F. Nowak, A.R. Bridle, Alginate Microencapsulation for Oral Immunisation of Finfish: Release Characteristics, Ex Vivo Intestinal Uptake and In Vivo Administration in Atlantic Salmon, *Salmo salar* L, *Marine biotechnology* (New York, N.Y.) 17(6) (2015) 841-53.
- [17] M. Lovoll, T. Kilvik, H. Boshra, J. Bogwald, J.O. Sunyer, R.A. Dalmo, Maternal transfer of complement components C3-1, C3-3, C3-4, C4, C5, C7, Bf, and Df to offspring in rainbow trout (*Oncorhynchus mykiss*), *Immunogenetics* 58(2-3) (2006) 168-79.
- [18] P. Swain, S.K. Nayak, Role of maternally derived immunity in fish, *Fish & Shellfish Immunology* 27(2) (2009) 89-99.
- [19] A. Hanif, V. Bakopoulos, G.J. Dimitriadis, Maternal transfer of humoral specific and non-specific immune parameters to sea bream (*Sparus aurata*) larvae, *Fish & Shellfish Immunology* 17(5) (2004) 411-435.
- [20] L.W. Harrell, H.M. Etlinger, H.O. Hodgins, Humoral factors important in resistance of salmonid fish to bacterial disease. I. Serum antibody protection of rainbow trout (*Salmo gairdneri*) against vibriosis, *Aquaculture* 6(3) (1975) 211-219.
- [21] A. Mor, R.R. Avtalion, Transfer of antibody activity from immunized mother to embryo in tilapias, *Journal of Fish Biology* 37(2) (1990) 249-255.
- [22] Y.M. Sin, K.H. Ling, T.J. Lam, Passive transfer of protective immunity against ichthyophthiriasis from vaccinated mother to fry in tilapias, *Oreochromis aureus*, *Aquaculture* 120(3) (1994) 229-237.
- [23] P. Swain, S. Dash, J. Bal, P. Routray, P.K. Sahoo, S.K. Sahoo, S. Saurabh, S.D. Gupta, P.K. Meher, Passive transfer of maternal antibodies and their existence in eggs, larvae and fry of Indian major carp, *Labeo rohita* (Ham.), *Fish & Shellfish Immunology* 20(4) (2006) 519-527.
- [24] S. Oshima, J. Hata, C. Segawa, S. Yamashita, Mother to fry, successful transfer of immunity against infectious haematopoietic necrosis virus infection in rainbow trout, *The Journal of general virology* 77 (Pt 10) (1996) 2441-5.
- [25] M.E. Nielsen, K. Buchmann, Effects of Oral Administration of Specific Antibodies to Rainbow Trout *Oncorhynchus mykiss*, *Journal of the World Aquaculture Society* 34(1) (2003) 11-17.
- [26] J.K. Chettri, M.K. Raida, P.W. Kania, K. Buchmann, Differential immune response of rainbow trout (*Oncorhynchus mykiss*) at early developmental stages (larvae and fry) against the bacterial pathogen *Yersinia ruckeri*, *Developmental & Comparative Immunology* 36(2) (2012) 463-474.
- [27] M.M. Olsen, P.W. Kania, R.D. Heinecke, K. Skjoedt, K.J. Rasmussen, K. Buchmann, Cellular and humoral factors involved in the response of rainbow trout gills to *Ichthyophthirius multifiliis* infections: Molecular and immunohistochemical studies, *Fish & Shellfish Immunology* 30(3) (2011) 859-869.

- 325 [28] J.L. Romalde, A. Luzardo-Alvárez, C. Ravelo, A.E. Toranzo, J. Blanco-Méndez, Oral immunization using
326 alginate microparticles as a useful strategy for booster vaccination against fish lactococcosis, *Aquaculture*
327 236(1–4) (2004) 119-129.
- 328 [29] K.Y. Lee, D.J. Mooney, Alginate: Properties and biomedical applications, *Progress in Polymer Science*
329 37(1) (2012) 106-126.
- 330 [30] U. Khimmakthong, S. Deshmukh, J.K. Chettri, A.M. Bojesen, P.W. Kania, I. Dalsgaard, K. Buchmann,
331 Tissue specific uptake of inactivated and live *Yersinia ruckeri* in rainbow trout (*Oncorhynchus mykiss*):
332 visualization by immunohistochemistry and in situ hybridization, *Microbial pathogenesis* 59-60 (2013) 33-
333 41.
- 334 [31] M. Ohtani, K.R. Villumsen, E.O. Koppang, M.K. Raida, Global 3D Imaging of *Yersinia ruckeri* Bacterin
335 Uptake in Rainbow Trout Fry, *PLOS ONE* 10(2) (2015) e0117263.

Table 1. The experimental set-up used for testing the protective effect of feeding *Yersinia ruckeri* specific IgM encapsulated in alginate microparticles. Each group consisted of 50 fish divided into duplicate (25 fish/tank) and fish were fed with medicated feed for 1 week before challenge infection and 2 weeks during the infection period. The water level was lowered to 5 L during the challenge infection and fish was exposed to 1.2×10^8 CFU/ml concentration of bacteria soup for 4 h. The experiment was conducted at 15°C for 3 weeks and moribund fish were euthanized using a high dose of MS222.

Group no.	Groups	Feed coated with:	IgM concentration	No. of fish/group	Challenge infection Bath exposure for 4 h
1	Uninfected control	Plain alginate microparticles	0	50	No
2	Infected control	Plain alginate microparticles	0	50	1.2×10^8 CFU/ml
3	Un-specific IgM	Alginate microparticles with un-specific IgM	200 µg/fish/day	50	1.2×10^8 CFU/ml
4	Anti- <i>Yersinia</i> IgM	Alginate microparticles with anti- <i>Yersinia</i> IgM	40 µg/fish/day	50	1.2×10^8 CFU/ml
5	Anti- <i>Yersinia</i> IgM	Alginate microparticles with anti- <i>Yersinia</i> IgM	200 µg/fish/day	50	1.2×10^8 CFU/ml
6	Anti- <i>Yersinia</i> IgM	Alginate microparticles with anti- <i>Yersinia</i> IgM	400 µg/fish/day	50	1.2×10^8 CFU/ml
7	Anti- <i>Yersinia</i> IgM	Feed oil coated with anti- <i>Yersinia</i> IgM	200 µg/fish/day	50	1.2×10^8 CFU/ml
8	Oxolinic acid	Feed oil coated with Oxolinic acid (1.25 g/kg)	0	50	1.2×10^8 CFU/ml

Figure 1. *Yersinia ruckeri* specific antibody titer from vaccinated and unvaccinated fish. Fish were vaccinated with *Y. ruckeri* 3×10^{10} CFU/ml (containing both biotype 1 and 2) in Freund's incomplete adjuvant (1:1). ELISA plates were coated with sonicated lysate of *Y. ruckeri* (5 µg/ml protein concentration) and tested for antibody titer. Each symbol represents a single fish and asterisk (*) on the top indicates a significantly different ($P \leq 0.05$) from the unvaccinated control group.

Figure 2. The sandwich ELISA developed for quantification of rainbow trout IgM. ELISA plates were coated with monoclonal antibody against rainbow trout IgM (1/1000 dilution) and detection antibody was based on the same antibody with biotinylation. Pooled samples from vaccinated or unvaccinated fish were run in dilutions and calculation of IgM was performed based on the standard purification. Asterisk sign (*) on the top indicates a significantly different ($P \leq 0.05$) from the control group.

Figure 3. Binding of specific antibody against *Yersinia ruckeri* after elution from alginate microparticles. The IgM encapsulated alginate microparticles were incubated with sodium bicarbonate pH 8.2 for 4 days and the re-extracted IgM (100 µl) was used for testing bio-activity in pathogen-specific ELISA (plate coated with sonicated lysate of *Y. ruckeri* (5 µg/ml protein concentration)).

Figure 4. Quality of specific antibody against *Yersinia ruckeri* following elution from alginate microparticles. The IgM encapsulated alginate microparticles were incubated with sodium bicarbonate pH 8.2 for 4 days and the re-extracted IgM was used for running SDS-PAGE (left) and Western Blot (right) to test if encapsulation process leads to any changes in protein integrity and to test the release of encapsulated IgM at different pH.

Figure 5. Cumulative mortality of fish after challenge infection with *Yersinia ruckeri* biotype 2. Fish were fed with a diet containing anti-*Yersinia ruckeri* antibody (IgM) for three weeks (1-week pre-feeding before challenge and 2 weeks feeding during the infection). A total of eight groups in duplicate was included in the experiment. Three control groups were included in the study where one group was fed with ordinary feed coated with plain alginate (this group was not given infection and served as uninfected control) and the second group was given the same feed type as group 1 but this group was given infection (infected control). The third control group was given alginate encapsulated IgM supplemented feed purified from the unvaccinated fish and infection was applied. Three groups were given immunoglobulin in feed (three dosages). The seventh group of fish was fed with anti-*Yersinia ruckeri* immunoglobulin directly oil coated on feed. And the last group (8) was fed with antibiotic (oxolinic acid: 1.25 g/kg feed) coated feed (positive control). The experimental groups are detailed in table 1. For infection, Fish were exposed to 48 h culture of *Y. ruckeri* (100415-1/4) bacterial soup for 4 h with a final bacterial concentration of 1.2×10^8 CFU/ml.

Highlights

- *Yersinia ruckeri* specific IgM was purified and encapsulated in alginate microparticles
- Bio-activity and integrity of encapsulated IgM remain intact
- IgM encapsulated microparticles were top-coated in fish feed
- Feeding and challenge experiment was performed to evaluate protection
- Highest dose of IgM fed group showed 10% lower mortality indicating its positive effect.