



From biogas to biorefinery: Modelling green chemical production in a full-scale anaerobic digester

Junicke, Helena; Flores-Alsina, Xavier; Pinto, Tiago; Gernaey, Krist V.

Publication date:
2019

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Junicke, H., Flores-Alsina, X., Pinto, T., & Gernaey, K. V. (2019). *From biogas to biorefinery: Modelling green chemical production in a full-scale anaerobic digester*. Abstract from 10th IWA Symposium on Modelling and Integrated Assessment, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Conference title:

10th IWA Symposium on Modelling and Integrated Assessment (Watermatex)

Conference scope:

International

Conference date:

1-4 September 2019

Conference place:

Copenhagen, DK

Conference URL:

<https://www.conferencemanager.dk/watermatex2019/home.html>

Talk title:

From biogas to biorefinery: Modelling green chemical production in a full-scale anaerobic digester

Authors:

Helena Junicke, Xavier Flores-Alsina, Tiago Pinto, Krist V. Gernaey

Abstract:

Industrial waste streams are a versatile feedstock for biorefinery processes. In this work, we study the upgrading of such feedstock into butanol, a valuable chemical and promising biofuel. The Anaerobic Digestion Model No. 1 (ADM1) was extended with butanol-forming reactions, thermodynamics and butanol gas-liquid mass transfer. Continuous butanol production was modelled in a full-scale anaerobic digester. ADM1 without thermodynamic constraints overestimates butanol productivity by 121% as compared to the thermodynamics based implementation. The highest butanol productivity (2.6 MWh.d⁻¹) was obtained at pH 5 with a COD recovery of 12%. Results lead us to propose a two-stage biorefinery process for butanol/methane production.