



Listeria monocytogenes control in cold smoked salmon using natural and sodium free preservatives

Heintz, Eelco; Dalgaard, Paw; van Lent, Henkjan ; Eliassen, Michael; Vega, Leonardo

Published in:
Journal of Food Protection

Publication date:
2019

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Heintz, E., Dalgaard, P., van Lent, H., Eliassen, M., & Vega, L. (2019). Listeria monocytogenes control in cold smoked salmon using natural and sodium free preservatives. *Journal of Food Protection*, 82(Suppl A).

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

***Listeria monocytogenes* Control in Cold Smoked Salmon using Natural and Sodium Free Preservatives.**

Heintz, E.¹ Dalgaard, P.²

¹ Niacet b.v., Papesteeg 91, 4006 WC Tiel, The Netherlands.

² National Food Institute (DTU Food), Technical University of Denmark Lyngby, Denmark.

Abstract:

Listeria monocytogenes is an opportunistic pathogen associated with over 1000 illnesses and high mortality rates each year in the US (ref. CDC). *L. monocytogenes* contamination and growth is a major concern for chilled cold smoked salmon (CSS). In response to WHO recommendation salt is being reduced in food products, which results in an even bigger challenge to control *L. monocytogenes* growth. This leads to an increasing demand for highly effective sodium free preservatives.

Organic acids and particularly acetic acid/acetates are well known for their inhibitive activity on *Listeria*, but consumers have negative perceptions towards synthetic-derived additives and increasingly prefer natural alternatives. Neutralized vinegar is a natural alternative for acetates and vinegar based products are developed in order to serve the modern consumer with a natural based powerful ingredient to increase the safety of low sodium CSS.

Methods:

A Danish CSS processor in collaboration with DTU Food used a combination of salting with injection brining followed by treatments on surface of filets in order to have optimal distribution of additives. Main product characteristics of CSS were 3 % water phase NaCl and pH of 6.1-6.3 (Control). Other treatments included lactate/diacetate (60% solution), 2.5%; Provian K (solid acetate/diacetate), 0.25%, 0.5% and 0.75%; Provian NDV (solid vinegar), 0.3%, 0.6%, 0.9%. For each treatment sliced products were inoculated with approximately 2 log CFU/g of a *L. monocytogenes*-cocktail. Inoculated slices were vacuum packaged and stored for 50 days at 4°C and 7°C. At each time of sampling during storage *L. monocytogenes* were enumerated in triplicate by surface plating on PALCAM agar.

Results:

Growth of *L. monocytogenes* to 7-8 log CFU/g was observed during 50 days on the CSS control treatment at both 4°C and 7°C. With an increase of 1 log CFU/g reached after one week for this treatment. All other treatments showed reduced growth of *L. monocytogenes* and 0.75% Provian K and 0.9% Provian NDV both reduced growth to less than a 2 log CFU/g increase in concentrations during 50 days at 4°C and 7°C.

Significance:

This research demonstrated the possibility to increase safety of cold smoked salmon using preservatives that meet current food trends, like sodium reduction and natural origin.