Predicting the effect of salt on heat tolerance of *Listeria monocytogenes* in meat and fish products

Abdallas, Somaya; Al-Hilali, Iman; Hansen, Tina Beck

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Abdallas, S., Al-Hilali, I., & Hansen, T. B. (2019). *Predicting the effect of salt on heat tolerance of Listeria monocytogenes in meat and fish products*. Poster session presented at 11th International Conference on Predictive Modelling in Food , Braganza, Portugal.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Predicting the effect of salt on heat tolerance of *Listeria monocytogenes* in meat and fish products

Somaya Abdalas, Iman Al-Hilali and Tina Beck Hansen (tibha@food.dtu.dk)
DTU Food, Division for Microbiology and Production, Research Group for Food Microbiology and Hygiene

Listeria monocytogenes is a potentially fatal foodborne pathogen that can be found in various food products, including meats. It can tolerate adverse conditions such as high salt concentrations, thus, the elimination of the organism is difficult in food processing. This study was conducted to enable prediction of the effect of salt on heat tolerance of *L. monocytogenes* in food. The obtained model enables food processors to design proper thermal processes to eliminate *L. monocytogenes* in meat products to ensure safety and prevent foodborne listeriosis.

Figure 1. Overview of modelling procedure.

Figure 3. Validation – predicted values versus literature values of $\log_{10}(3D)$ of *Listeria monocytogenes* heated in meat and fish products.

Methodology

The work involved minced pork, chicken and salmon with different water phase salt (WPS) concentrations from 0 to 5.6 % inoculated with late-stationary phase *L. monocytogenes* cultures from three strains associated with fish, meat and industrial environment. Samples were vacuum-packaged in sterile bags, immersed in water bath, and held at constant temperatures from 57 to 65 °C.

Modelling

Heat tolerance was defined as the time to achieve 3 decimal reductions, *i.e.* 3D-values, by using both log-linear and non-linear-regression models (Figure 1). Development of a secondary predictive model describing the combined effect of temperature and WPS was carried out.

Temp. °C	WPS %	3D log-linear (min)	3D GInaFIT (min)	3D F-test (min)
57	0	41	46	43
57	3.4	108	97	94
57	4.9	151	154	144
60	0	20	28	20
60	3.4	22	23	23
60	4.9	27	32	29
65	0	3.8	4.4	4.4
65	3.4	7.1	5.3	6.9
65	4.9	8.6	8.2	8.7
RSS – $\log_{10}(3D)$		0.323	0.375	0.257

Table 1. Heat tolerance of *Listeria monocytogenes* in pork products estimated using three approaches. Average of two trials.

Figure 2. Secondary model - fitted values versus observed values of $\log_{10}(3D)$ for *Listeria monocytogenes* heated in meat and fish products. Diagonal line: perfect fit.

The mathematical structure of the model was adopted from Li *et al.* (Food Control 73:406, 2017):

$$\log_{10}(3D) = a + b \times \text{Temp} + c \times \text{Temp} \times \text{WPS}$$

Three different approaches i) Log-linear, ii) GInaFIT and iii) F-test were applied for selection of the 3D-values to be included as responses in the development of the model (Figure 1).

The F-test approach resulted in best fit (Table 1) with $a=11.8476$, $b=-0.1771$ and $c=0.00142$. Goodness-of-fit is shown in Figure 2.

Validation

The secondary model was validated using 86 external literature values from six different sources (Figure 3). WPS from 0 to 6 %, temperatures from 57.5 to 65.0 °C and salmon roe, minced meat (beef and pork) and pork slurry were represented.

The bias and accuracy factors were 2.92 and 3.14, respectively, when D-values were used as response. When $\log_{10}(3D)$ -values were used, the bias factor was 1.78 and the accuracy factor was 1.81.