Biomass composition of *Arthrospira platensis* during cultivation on industrial process water and harvesting

Ljubic, Anita; Safafar, Hamed; Holdt, Susan Løvstad; Jacobsen, Charlotte

Published in:
Journal of Applied Phycology

Link to article, DOI:
[10.1007/s10811-017-1332-y](https://doi.org/10.1007/s10811-017-1332-y)

Publication date:
2018

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Ljubic, A., Safafar, H., Holdt, S. L., & Jacobsen, C. (2018). Biomass composition of *Arthrospira platensis* during cultivation on industrial process water and harvesting. *Journal of Applied Phycology*, 30(2), 943–954.
<https://doi.org/10.1007/s10811-017-1332-y>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 Biomass composition of *Arthrospira platensis* during cultivation on industrial 2 process water and harvesting

3

4 Anita Ljubic^{1*}, Hamed Safafar², Susan L. Holdt¹ and Charlotte Jacobsen¹

5 ¹Technical University of Denmark, National Food Institute, Kemitorvet, Building 202, Kongens Lyngby 2800,
6 Denmark; suho@food.dtu.dk (0000-0002-6888-782X); chja@food.dtu.dk (0000-0003-3540-9669)

7 ²BioMar A/S, Mylius-Erichsensvej 35, 7330 Brande, Denmark; hamsa@biomar.dk (0000-0002-8538-7456)

8 * Author for correspondence: e-mail: aniljub@food.dtu.dk (0000-0001-9370-8830), phone: +4581611094

9

10 **Acknowledgments:** The authors wish to acknowledge Inge Holmberg for her technical assistance. We would like to
11 thank Kalundborg WWTP and Novozymes A/S for providing access to industrial process water.

12 **Conflicts of Interest:** The authors declare no conflict of interest.

13

14 Abstract

15 Microalgae have the ability to utilize nutrients from wastewater and use it for biomass production.
16 The effluent from a biogas process was tested as a nutrient source for blue-green microalga
17 *Arthrospira platensis* cultivation and compared with conventional synthetic medium. Cultivation
18 was carried out in four different concentrations of industrial process water (25%, 50%, 75% and
19 100%). The biomass was then harvested by microfiltration and centrifugation followed by freeze
20 drying. Variations in biomass composition were studied, in order to investigate effects of industrial
21 process water on *A. platensis* over 30 days of cultivation. Applied harvesting techniques were
22 evaluated for their effect on physiochemical properties of the biomass. *Arthrospira platensis* was
23 able to grow in all tested wastewater concentrations except 100%, however, increase of wastewater
24 concentration in medium resulted in a decreased growth rate. Partial substitution of synthetic
25 Zarrouk medium with 25% of wastewater showed no adverse effect on chemical composition of the
26 biomass including high protein content (45-58% dry weight) and favourable fatty acids composition
27 (42-45% PUFAs of total fatty acids). Evaluation by optical microscopy showed that microfiltration
28 caused cell rupture at the moderate level while centrifugation had more severe effect on *A.*
29 *platensis*. Effect of centrifugal forces and shear stress on *A. platensis* cells was confirmed by

detecting lower lipid content in samples after applying both microfiltration and centrifugation due to cell content leakage.

Keywords: Spirulina, wastewater, biomass composition, cross-flow microfiltration, centrifugation, cell rupture.

1. INTRODUCTION

Microalgae are a group of microscopic organisms with a broad phylogenetic diversity. They are a potential sustainable source of feedstock that can be harnessed for commercial use, due to their high photosynthetic activity, short growth cycle and low land area requirements compared to terrestrial plants (Mata et al. 2010). During the last 30 years, the microalgal industry has grown significantly. The first large-scale cultivation facilities were established as the potential solution for the food and feed shortage. Nowadays, different species of microalgae, mainly from the genus *Chlorella*, *Dunaliella* and *Arthrospira*, are used for mass cultivation with a wide range of applications such as bioenergy production, food and feed supplements and pharmaceutical products (Priyadarshani and Rath 2012; Milledge 2011).

Arthrospira platensis, traditionally known as Spirulina, is a photosynthetic, prokaryotic, blue green microalgae. Characteristic morphological features of *A. platensis* are the spiral shape of multicellular cylindrical filaments in an open helix with the length from 0.3-1.0 mm. It is considered to be sensitive towards external stress, due to the relatively large size and fragile cellulose-free cell wall, and therefore external stress is found to facilitate extraction of essential compounds. Typical chemical composition of *A. platensis* is 55-70% proteins, 15-25% carbohydrates and 4-7% lipids (Ali and Saleh 2012). It serves as a rich source of essential nutrients such as vitamins, minerals, proteins including all the essential amino acids, pigments and polyunsaturated fatty acids (PUFAs). Due to the high content of biological active compounds, it has found applications in human and animal nutrition, cosmetics, high-value molecules production for pharmaceuticals etc. (Priyadarshani and Rath 2012).

Despite the large potential and wide range of applications, industrial production of microalgal biomass often meets economical challenges. However, several approaches have been considered for production process optimization and thereby a reduction of production costs (El-Sheekh et al. 2016; Barros et al. 2015). Using wastewater as complete or partial substitution of synthetic growth

61 medium can reduce production costs and fresh water requirement (Abdel-Raouf et al. 2015).
62 Bioremediation by microalgae is particularly effective due to their ability to assimilate nutrients and
63 to convert light energy into valuable biomass. Thus, applying wastewater as cultivation medium
64 presents environmentally beneficial sources of carbon, nitrogen and phosphorous for microalgae
65 growth (El-Sheekh et al. 2016). Efficient bioremediation of *A. platensis* was reported previously in
66 several studies (Markou et al. 2016; Mezzomo et al. 2010; Phang et al. 2000). In the present study
67 industrial process water (ICW) was acquired by methanogenic conversion of organic compounds to
68 methane, carbon dioxide and an effluent with relatively high ammonia content. The anaerobic
69 digestion process was carried out in an anaerobic sludge tower reactor with internal circulation
70 (ICT), therefore, the effluent is called IC water (ICW). The growth performance of several
71 microalgae species was previously tested on ICW where some of the tested species showed ability
72 to grow even on 100% ICW (Safafar 2017). In general, microalgae use nitrogen in amino acid,
73 protein and pigment syntheses and it is utilized in the form of nitrate (NO^{-3}), nitrite (NO^{-2}) or
74 ammonia (NH_3). However, as long as NH_3 is available most of the microalgal cells will not utilize
75 other nitrogen sources because assimilation of any other nitrogen form requires energy consumption
76 for reduction (Chaiklahan et al. 2010).

77 Another optimizing approach relates to downstream processes due to their high share of the total
78 production costs. Harvesting presents a major challenge due to the small size of microalgae cells
79 and low density of the cultures. An efficient optimized harvesting process should be suitable for a
80 variety of microalgae strains resulting in high biomass concentrations while requiring low
81 operational costs for energy and maintenance (Barros et al. 2015). Among many harvesting
82 techniques, centrifugation is the most energy-intensive harvesting method. Regardless of that, due
83 to its high separation efficiency and ability to harvest a large majority of microalgae, it is the most
84 commonly used harvesting method in lab-scale and large-scale microalgal plant systems (Dassey
85 and Theegala 2013). In order to decrease energy demand, microalgae can be pre-concentrated
86 before centrifugation. Membrane filtration presents an optimal pre-treatment process, where
87 biomass can be pre-concentrated 5-10 times (Bilad et al. 2014). It represents closed harvesting
88 system, which is commonly needed for production of high valued products (e.g. omega-3-fatty
89 acids, pigments). However, it has previously been reported that exposure of microalgal cells to high
90 gravitational forces during centrifugation and providing sufficient shear at the membrane surface
91 during filtration can lead to structural cell damage (Safafar 2017; Xu et al. 2015; Bilad et al. 2014).
92 Whether this happens will to a large degree depend on the applied force level and the specific

features of the microalgal strain. Unwanted cell disruption liberates the cell content, which can greatly reduce the shelf life and nutrition value of the microalgal biomass.

The aim of this study was to investigate the effects of different concentrations of ICW during cultivation on biomass production and nutritional composition of *A. platensis*. This study evaluated the optimal ICW concentration for obtaining maximum biomass production and favourable biomass composition (proteins, lipids, fatty acids, pigments and amino acids). Furthermore, the applied two-step harvesting process (cross-flow microfiltration and centrifugation) was tested for its effect on the changes in the biomass as a result of external stress exposure with an aim of investigating potential for developing large scale harvest process after cultivation in photo bioreactors.

2. MATERIALS AND METHODS

2.1. Microalgal cultivation and growth measurements

Arthrospira platensis (SAG 85.79) was obtained from SAG (Sammlung von Algenkulturen der Universität Göttingen) culture collection of algae in Germany. The inoculum was prepared in synthetic Zarrouk medium (ZM; Zarrouk 1966) with the following composition: 16.8 g L⁻¹ NaHCO₃, 2.5 g L⁻¹ NaNO₃, 0.5 g L⁻¹ KH₂PO₄, 1.0 g L⁻¹ K₂SO₄, 1.0 g L⁻¹ NaCl, 40 mg L⁻¹ CaCl₂, 80 mg L⁻¹ Na₂EDTA, 200 mg L⁻¹ MgSO₄·7H₂O, 10 g L⁻¹ FeSO₄·7H₂O and 1.0 mL of trace elements stock solutions: 2.86 g L⁻¹ H₃BO₃, 20 mg L⁻¹ (NH₄)₆Mo₇O₂₄, 1.8 g L⁻¹ MnCl₂·4H₂O, 80 mg L⁻¹ CuSO₄ and 220 mg L⁻¹ ZnSO₄·7H₂O. Industrial process water (ICW) used for the experiments was obtained from Novozyme's plant (Kalundborg, Denmark) and it was filtered by cross-flow microfiltration (0.2 µm pore size) prior use. Chemical composition of ICW is presented in Table 1.

The cultivation experiments were carried out in two sets using 1 and 5 L GS Schott bottles. First set of experiments (preliminary experiments) included 12 days cultivation of *A. platensis* in four different concentrations of ICW (25%, 50%, 75% and 100% (all concentrations were prepared in ZM)) and 100% Zarrouk medium. Working volume was 1 L including 100 ml of inoculum.

The second set of experiments was designed based on the growth curves acquired from the first set of experiments. Two best growing cultures were recultivated in 5 L bottles in order to collect samples during cultivation period. Therefore, the second set of experiments included cultivation in 25% ICW and 100% ZM in duplicates over 31 days with a working volume of 5 L. Both ICW and ZM were autoclaved before use at 120 °C for 45 min. The cultures were aerated with a mixture of carbon dioxide (5%) and sterile air. Cultivation temperature range was 25±1 °C and pH range

125 9.5±5. pH monitoring was performed by Milwaukee MC-122-pH controller (Milwaukee
126 Electronics, Szeged, Hungary) equipped with a solenoid valve to control CO₂ addition. Provided
127 light intensity was 100 μmol photons m² s⁻¹ (measured on the outer side of the bottle) under
128 fluorescent lamp illumination (Green-line A/S, Maribo, Denmark) during 14/10 hours of light/dark
129 cycle.

130 Growth was monitored daily by taking 5 mL culture sample and measuring optical density at 670
131 nm. Dry matter was determined on a daily basis by taking liquid samples, weighing and drying in
132 oven at 100 °C until dryness, kept in desiccator, cooled down to room temperature and weighted.
133 The specific growth rate was calculated using the following formulae (Guillard and Ryther 1962):

$$\mu = \frac{\ln \frac{x_1}{x_0}}{t_1 - t_0} \quad (1)$$

135 where x₁ and x₀ were biomass concentrations (g L⁻¹) at t₁ and t₀ cultivation days, respectively.

137 2.3. Analytical methods

138 For chemical analysis 0.4 L of culture was sampled. In order to achieve biomass separation,
139 samples were centrifuged at 11 000 x g and the biomass was washed by deionized water. The
140 resulting biomass was freeze dried and stored at -20 °C until analysis.

141 The protein concentration in the microalgal samples (approximately 1 g of dried microalgal
142 biomass) was estimated using the Dumas method for the quantitative determination of nitrogen in
143 different matrices (Elementar, Mt. Laurel, NJ, USA). The following steps were automated including
144 sample combustion in a chamber at a high temperature (900 °C) in the presence of oxygen.
145 Conversion of the estimated total nitrogen content to the crude protein was done using conversion
146 factor 6.35 for *A.platensis* reported by Safi et al. (2013). Protein concentration is reported as % of
147 dry biomass.

148 The amino acid composition was analyzed using Phenomenex EZ:faastTM amino acid analysis kit
149 (Phenomenex Inc. CA, USA). Approximately 30 mg of dry microalgal biomass was weighted in
150 microwave glass vials. Samples were hydrolyzed in 6 M HCl in a microwave oven (Microwave
151 3000 SOLV, Anton Paar, Ashland, VA, USA) for 60 minutes at 110 °C. Amino acid derivatization
152 was done by following clean-up steps in order to remove matrix interference. Liquid
153 chromatography with Agilent 1100 series LC/MSD Trap mass spectrometry (Agilent technologies,
154 Santa Clara, CA, USA) was used for amino acid composition determination. Applied column was
155 EZ:fastTM LC-MS column (250 x 3.0 mm, Phenomenex, Torrance, CA, USA). Amino acids are
156 reported as μg/g of dry biomass.

Lipids were extracted from approximately 1 g of dried microalgal biomass by subsequent addition of methanol, chloroform and water while stirring as described by Bligh & Dyer (1959), but with a reduced amount of solvent. Samples were centrifuged at 1400 x g for 10 min, in order to separate methanol/water phase from chloroform/oil phase. Bligh & Dyer extracts were used for analyses of oil content, fatty acids and tocopherols. For determining total lipid content around 15 g of extract was weighted in beakers and left overnight in a fume hood in order to evaporate chloroform. Lipid content was calculated using the equation below (2).

$$\% \text{ lipid} = \frac{g(\text{lipid}) * 41 g * 100}{(g(\text{extract}) - g(\text{lipid}) * g(\text{sample}))} \quad (2)$$

Fatty acid profile was determined according to a slightly modified FAME method based on the American Oil Chemist's Society (AOCS) official method Ce 1i-07 (Firestone 2009). Approximately 5 g of Bligh & Dyer extract was weighted in methylation glass tube. Extracts were evaporated under the stream of nitrogen until they were completely dry. A mixture containing 100 µL of internal standard solution, 200 µL of heptane with BHT and 100 µL of toluene was added. Samples were methylated in a microwave oven (Microwave 3000 SOLV, Anton Paar, Ashland, VA, USA) for 10 min. at 100 °C at 500 watts. 0.7 mL of heptane with BHT and 1 mL of saturated salt water (NaCl) were added. The separated upper phase (heptane) was transferred into vials and analyzed by gas chromatography (HP-5890 A, Agilent Technologies, Santa Clara, CA, USA). Fatty acid methyl esters were separated by the GC column Agilent DB wax 127-7012 (10 µm x 100 µm x 0.1 µm; Agilent technologies, Santa Clara, CA, USA). Standard mix of fatty acids methyl esters (Sigma, St. Louis, MO, USA) was used for identification of individual fatty acid. Fatty acids were quantified as in area % of total fatty acids.

Pigment analysis was done by the method described by Safafar et al. (2015). Extraction was carried out on approximately 0.05 g of dried algal biomass by methanol containing BHT in a sonication bath (Branson Ultrasonics, Danbury, CA, USA) for 15 min at 5±2 °C. Pigments analysis was performed by HPLC using Agilent 1100 Liquid Chromatograph with DAD. Separation was carried out on a Zorbax Eclipse C8 column 150 mm x 46 mm x 3.5 µm (Phenomenex Inc. CA, USA). The mobile phase was a mixture of 70% methanol + 30% of 0.028 M tertiary butyl ammonium acetate in water and methanol at a flow rate of 1.1 mL min⁻¹ with total acquisition time of 40 min. Identification of peaks was performed using DHI pigment standard mix (DHI LAB Products, Horsholm, Denmark). Detection of chlorophylls and carotenoids was done at 660 nm and 440 nm,

188 respectively, and for internal standard (BHT) at 280 nm. Pigments are reported as $\mu\text{g g}^{-1}$ of dry
189 biomass.

190

191 **2.4. Harvest - microfiltration and centrifugation**

192 Harvest of microalgal biomass was carried out after 30 days of cultivation (cultivation conditions
193 described in section 2.1), in a batch mode by tangential flow filtration using a silicon carbide
194 membrane with pore size 3 μm ($\varnothing$ 25 x 305 mm, $\varnothing$ 3 mm channel; Liqtech A/S, Denmark). The
195 laboratory scale filtration unit (Liqtech A/S, Denmark) had a capacity of 20-50 L h^{-1} , and 5 L of
196 algal biomass was used during the filtration process at a constant pressure of 1.0 ± 0.2 bar and a
197 temperature of 20 ± 3 $^{\circ}\text{C}$. Harvested microalgal biomass was recycled by the pump (Watson-
198 Marlow 604 U/R, Falmouth, Cornwall, UK) back to the feed tank, so that the concentration of the
199 feed increased, with processing time. A schematic drawing of the used batch mode microfiltration
200 process is shown in Fig. 1. Calculation of flux, filtration rate and concentration ratio was done by
201 the following formulae (EPA 2005):

$$202 \quad J = \frac{Q_p}{A_m} \quad (3)$$

203 J = flux ($\text{L h}^{-1}\text{m}^{-2}$); Q_p = filtrate flow rate through membrane (L h^{-1}); A_m = surface area of
204 membrane (m^2)

$$205 \quad J_{avg} = J_0 - 0.33 (J_0 - J_f) \quad (4)$$

206 J_{avg} = average flux rate; J_0 = initial flux; J_f = final flux

207

$$208 \quad CF = \frac{C_f}{C_0} \quad (5)$$

209 CF = concentration ratio; C_f = final concentration of a given solute (g L^{-1}); C_0 = initial
210 concentration of the solute (g L^{-1}).

211

212 A bench scale centrifuge (refrigerated centrifuge, IEC Centra-GP8R, Buckinghamshire, England)
213 was used for the biomass up-concentration after applying microfiltration. Complete separation was
214 accomplished at 11 000 x g for 10 min. The supernatant was discarded and microalgal biomass was
215 freeze-dried. Protein, lipid and pigment content, as well as the fatty acid composition were
216 determined in order to evaluate changes in the biomass composition (same procedures as describes
217 above). In addition, samples were evaluated by optical microscopy.

218

2.5. Statistical analysis

Cultivation experiments in 5 L bottles were carried out in 2 biological replicates (second set of experiments). All compositional analyses were performed on the samples from 5 L and repeated two times. The results are given as the mean ($\pm$ standard deviation). Analysis of variance (two-way ANOVA) was used to evaluate the effect of time and growth media on chemical composition of biomass. Data have met the assumption of normality and homogeneity of variance. Tukey's post hoc test was used to detect significant differences between groups where p values <0.05 were considered significant. The Statistica v. 13.2 software (Dell Inc.,Tulsa, OK, USA) was used for all statistical analyses.

3. RESULTS

3.1. Growth characteristics of *A. platensis*

Arthrospira platensis biomass concentration was continuously increasing in three tested ICW concentrations (25%, 50%, 75%) over the 12 days of cultivation (Fig. 2). However, it showed no ability for growing on 100% ICW. Biomass increment curves indicate that the increase in concentration of ICW resulted in lower biomass accumulation. The highest biomass increment was detected in 100% ZM. Standard curve of concordance between optical density and dry matter showed strong correlation ($R^2=0.99$; not shown) of *A. platensis* cultivated in 25% ICW and 100% ZM in experiment 2 (Fig. 3). Specific growth rate (μ) of *A. platensis* was $0.098 \pm 0.002 \text{ day}^{-1}$ and $0.089 \pm 0.005 \text{ day}^{-1}$ cultivated in 100% ZM and 25% ICW, respectively. No significant difference ($p > 0.05$), in biomass accumulation was observed between the cultures in 100% ZM and 25 % ICW during first 20 days of cultivation, after which *A. platensis* cultivated in 100% ZM grew significantly faster.

3.2. Biomass composition

Total protein and lipid concentration of *A. platensis* were not significantly affected ($p > 0.05$) by growth medium or cultivation time (Fig. 4 and 5). Protein concentration varied from 44-52% and 50-58% of dry biomass for *A. platensis* cultivated in 100% ZM and 25% ICW, respectively. Hultberg et al. (2016) tested protein content of *A. platensis* cultivated in Zarrouk medium (54-66%) and effluent-based medium (60-66%) where they reported no significant differences between the cultures grown in different medium, which corresponds to our study. Lipid concentration of *A.*

252 *platensis* varied from 4-6% of dry biomass for both cultures and similar values were reported in the
253 few previous studies with *A. platensis* (Baunillo et al. 2012; Colla et al. 2007). The amino acid
254 composition of *A. platensis* grown on different growth media is shown in Fig. 6a,b. Principal amino
255 acids included leucine, valine, glutamine and alanine, which is in agreement with the study reported
256 by Hultberg et al. (2016). The amino acid compositions were similar in *A. platensis* cultivated in
257 100% ZM and 25% ICW with no significant differences ($p > 0.05$). Tryptophan was not detected in
258 any of the cultures and it assumed that it was due to the fact that it is destroyed during the
259 hydrolysis. The current study confirms the presence of all essential amino acids with particularly
260 high contents of valine, 13-17%, and leucine, 11-17% of total amino acids. Significantly higher
261 contents ($p < 0.05$), of essential amino acids, compared to non-essential amino acids, were present
262 in algal biomass, however no significant difference between the proportions of essential and non-
263 essential amino acids, when comparing the two growth media (Table 3).

264 *Arthrospira platensis* is rich in a γ -linolenic acid (18:3 (n-6)) from omega-6 family. Other major
265 fatty acids in *A. platensis* are palmitic acid (16:0) and essential linoleic acid (18:2 (n-6)), and the
266 latter a precursor for synthesis of other polyunsaturated fatty acids. In the current study, these fatty
267 acids constitute 78-83% of total fatty acids. There was no significant differences ($p > 0.05$), in fatty
268 acid composition between different treatments (Table 4a,b), which agrees with previous findings on
269 *A. platensis* cultivated in ZM and anaerobic digestate effluent (Hultberg et al. 2016).

270 The pigment composition of *A. platensis* includes phycobiliproteins, chlorophylls and carotenoids.
271 In this study chlorophylls and carotenoids were analyzed (Fig. 7a,b), which are rarely reported for
272 *Arthrospira* sp. in contrast to the most abundant phycobiliproteins. Both treatments were showing
273 similar pigment composition pattern with no significant differences ($p > 0.05$). Slightly higher,
274 although not significantly, concentration of pigments was observed in *A. platensis* cultivated in
275 100% ZM. Concentrations of pigments in *A. platensis* increased over the 30 days of cultivation,
276 which positively correlate with the biomass production. The highest estimated concentration of
277 chlorophyll a was $4.099 \pm 67 \mu\text{g g}^{-1}$, followed by carotenoids: zeaxanthin ($1.465 \pm 531 \mu\text{g g}^{-1}$), β -
278 carotene ($1.745 \pm 132 \mu\text{g g}^{-1}$) and astaxanthin ($433 \pm 59 \mu\text{g g}^{-1}$).

279 3.3 Harvest - microfiltration and centrifugation

280 Microfiltration showed to be an efficient harvesting method with no membrane fouling during the
281 biomass separation. *Arthrospira platensis* cell size is relatively big, which favours separation

282 efficiency. Average flux (J_{avg}) was $405.3 \text{ L m}^{-2} \text{ h}^{-1}$ with a concentration ratio (CF) of 5.16, which is
283 above average compared to other microalgal species (*Monodopsis subterranea*, *Nannochloropsis*
284 *salina*, *Chlorella vulgaris*) harvested by cross-flow microfiltration under the same conditions
285 (Safafar 2017).

286 Exposure of *A. platensis* to transmembrane pressure during microfiltration, and high gravitational
287 and shear forces during centrifugation may cause structural cell damage (Xu et al. 2015; Bilad et al.
288 2014). The fresh culture of *A. platensis* (Picture 1a) showed no cell damage, while after applying
289 harvesting processes cell rupture was detected (Picture 1b,c). Biomass, analyzed after
290 microfiltration step, showed moderate cell damage, while applying microfiltration followed by
291 centrifugation, resulted in severe cell damage. Microscopy showed (circled points) indicators of
292 possible cell rupture such as presence of exopolysaccharides (EPS) and cellular fragments. EPS are
293 polymers composed of sugar residues that are secreted by microalgae into the surrounding
294 environment under stress conditions such as external pressure during harvest.

295 Protein, lipid and pigment composition were determined in order to test effects of applied
296 harvesting techniques on changes in *A. platensis* biomass. Significantly lower ($p < 0.05$), protein,
297 lipid and chlorophyll content were detected in samples experiencing both microfiltration and
298 centrifugation compared to only microfiltration (Table 5), which indicates more severe cell rupture
299 and leakage of the cell content.

300

301 4. DISCUSSION

302 4.1. Growth and chemical composition

303 *Arthrospira platensis* grows optimally under alkaline conditions, which means it requires use of
304 inorganic carbon from bicarbonate. Uptake of this inorganic carbon is possible due to the well-
305 developed carbon concentrating mechanism (CCM) in *A. platensis* (Klanchui et al., 2017). ZM
306 contains 16.8 g L^{-1} of sodium bicarbonate whereas ICW has no bicarbonate present. This could be
307 the reason for no growth detected in 100% ICW (Fig. 2). Furthermore, sodium and chloride ions
308 affect the osmotic pressure of the solution. Therefore osmolarity factor may contribute significantly
309 to the growth of *A. platensis*. Sodium concentration in ICW was significantly lower as compared to
310 ZM, which indicates that diluting ZM by ICW will result in salinity reduction. Lowering salinity
311 level may contribute to growth inhibition. In addition, ICW contains cyanide, which has toxic effect
312 on microalga, which could be another possible reason for growth inhibition.

313 Shorter adaptation period (lag phase) for *A. platensis* cultivated in lower levels of ICW and in ZM
 314 can be attributed also to a relatively low initial pH 8.1 of ICW compared to 9.3 in ZM (Fig. 2). Kim
 315 et al. (2007) reported the highest biomass accumulation of *A. platensis* at pH 9.5, which was later
 316 confirmed by Soundarapandian and Vasanthi (2008). The proposed explanation was that optimal
 317 activity of all the enzymes needed for photosynthesis and respiration is lower at below-optimal pH.
 318 Leema et al. (2010) reported 0.23 day^{-1} for *A. platensis* grown under similar conditions as in the
 319 current study. Observed deviations in these values and ones determined in this study are most likely
 320 result of the light limitation during cultivation. Growth rate of *A. platensis* strongly depends on the
 321 photosynthesis capacity, which depends on the light availability. In addition, high cell densities
 322 cause mutual shading and increase in turbidity of the culture, which will cause lower photosynthetic
 323 activity (Wondraczek et al. 2013). In a study by Leema et al. (2010), cultivation was carried out in
 324 500 mL bottles and higher μ was reported when comparing to 5 L glass bottle used in the current
 325 study. This can be confirmed by comparing biomass increment curves in Fig. 2 and 3, where OD_{670}
 326 is higher for *A. platensis* cultivated in 1 L bottle (Fig. 2) compared to *A. platensis* cultivated in 5 L
 327 bottle (Fig. 3) for the same time period. Light is a major factor influencing also pigment synthesis in
 328 microalgae. When microalgae are cultivated under a constant light intensity, the light intensity per
 329 cell will decrease due to the increase in cell density. Therefore, mutual shading will lead to the
 330 increase in pigment concentration as a result of competition for light harvesting (Myers and Kratz
 331 1955), which was confirmed by this study. Moreover, the increase of the total chlorophyll content
 332 during cultivation period indicates that the growth medium was nitrogen sufficient, otherwise
 333 chlorophyll content would decrease as a result of nitrogen limitation for the synthesis (Cohen 1997).
 334 In general, the chemical composition of ZM had higher concentration of inorganic carbon
 335 (NaHCO_3) available for *A. platensis* growth, which is most likely the main reason for increased
 336 biomass accumulation after day 20 (Fig. 3). Another important difference between ICW and ZM is
 337 nitrogen source, ammonia (NH_3) and nitrate (NO_3^-), respectively. Ammonia concentration is known
 338 to be a critical factor for *A. platensis* biomass accumulation and even though it presents the most
 339 preferable chemical form of nitrogen available to microalgae, high ammonia concentration has
 340 inhibitory effect on the growth of *A. platensis*. Ogbonna et al. (2000) reported complete inhibition
 341 of *A. platensis* by 400 mg L^{-1} of ammonia in the growth medium. Markou et al. (2014) suggested
 342 that low biomass densities are more susceptible to ammonia inhibition compared to high densities.
 343 In contrast, higher biomass densities assimilate ammonia rapidly as a response to inhibition. In our
 344 study concentration of ammonia in 100% ICW was 150 mg L^{-1} and initial biomass density was low,

345 which resulted in complete growth inhibition. However, *A. platensis* cultivated in 75% ICW, where
 346 ammonia concentration was 112.5 mg L⁻¹ showed biomass increment after 6 days of adaptation. In
 347 contrast, 100% ZM contained nitrate as nitrogen source with no ammonia and gave higher biomass
 348 yields compared to ZM diluted with ICW. Depleted amount of nitrogen present in the growth
 349 medium was estimated (data not shown) based on the chemical composition of *A. platensis*
 350 biomass. The data suggest that there was no nitrogen starvation in any of the two cultures.
 351 Protein synthesis is greatly affected by nitrogen level in the growth medium, which was confirmed
 352 by Sassano et al. (2010) who reported significant changes in protein content of *A. platensis* affected
 353 by nitrogen availability in the growth medium. Sufficient levels of nitrogen, means that carbon
 354 provided through the photosynthesis, will be used in protein synthesis (Hu, 2013). During starvation
 355 phase, when nitrogen limitation is present, lipids or carbohydrate will start accumulating. It is
 356 known that microalgae lipid accumulation can be enhanced by nitrogen limitation. However, it was
 357 also suggested that higher cultivation temperature (30-35 °C) may increase lipid accumulation in
 358 *A. platensis* (Markou et al. 2016; Colla et al. 2007). The fact, that there was no significant decrease
 359 in protein concentration or increase in lipid concentration over time (Fig. 4 and 5), suggest that *A.*
 360 *platensis* had sufficient nitrogen level in both experiments, and did not reach the stationary phase
 361 during 31 days of cultivation. Furthermore, it is expected that decrease in protein content would be
 362 observed first in culture grown on medium with 25% ICW, due to the lower nitrogen content
 363 compared to 100% ZM. In addition, sodium nitrate (NaNO₃), which is the nitrogen form available
 364 in ZM, was found to be the most preferable nitrogen form for *A. platensis* utilization (Costa et al.
 365 2001). *Arthrospira platensis* is a well-known rich source of high quality protein, which refers to the
 366 quantity of essential amino acids and high digestibility for animal and human organisms (Becker,
 367 2007). It can compete, due to its favourable amino acid composition and concentration, with other
 368 plant proteins such as soybean, where the content of essential amino acids is lower per unit of mass
 369 (Becker, 2007).
 370 *Arthrospira platensis* is a known commercial producer of γ -linolenic acid (18:3 (n-6)), which is a
 371 highly valuable fatty acid from the omega-6 family. The content of this fatty acid can reach up to
 372 30% of total fatty acids (Muhling et al. 2005). Saturated fatty acids start to accumulate, when the
 373 nitrogen concentration in the growth medium is limited (Hu et al. 2008). This finding confirms
 374 again that there was no nutrient limitation during the cultivation period for *A. platensis* in this study.
 375 However, there was a significant increase in the proportion of linoleic acid, and a decrease in γ -
 376 linolenic acid during the cultivation period. This trend indicates that there is a correlation between

the syntheses of these fatty acids. Thus, a possible transformation of tri-unsaturated fatty acid into di-unsaturated is suggested. Similar transformations have already been proven for some fatty acids in the process called retro-conversion, such as docosahexaenoic acid (22:6) transformed to eicosapentaenoic acid (20:5) and docosapentaenoic acid (22:5) transformed to arachidonic acid (20:4) in mammals. Also, it is suggested that microscopic organisms of phylum *Rotifers* and organisms of genus *Artemia* have the same ability (Barclay and Zeller 1996). This has not previously been reported in algal organism. However, it cannot be ruled out that retro-conversion is a process that can occur in fatty acids metabolism of other organisms than suggested by literature. In general, *Arthrospira* sp. has higher content of γ -linolenic acid compared to the content of linoleic acid (Muhling et al. 2005). However, Muhling et al. (2005) reported the fatty acid composition of 35 *Arthrospira* strains and observed different proportions of linoleic acid and γ -linolenic acid in *A. platensis* (strain SAG 85.79) compared to other *A. platensis* strains, where strain SAG 85.79 had considerably higher content of linoleic acid. Therefore, fatty acids composition reported in our study is in agreement with the study by Muhling et al. (2005).

391

392 **4.2. Harvesting**

In general, during harvesting the membrane is gradually being fouled during microfiltration, due to clogging of pores by small particles. Source of fouling can be the cell content released, due to the cell wall damage, or extracellular polymeric substances (exopolysaccharides) extracted by the cell under stress conditions (Rossi et al. 2008). Therefore, flux is being reduced over time as a result of gradual fouling (Fig. 8). If the cell rupture was severe during microfiltration, complete membrane fouling would occur fast and separation efficiency (average flux and concentration ratio) would decrease. According to Xu et al. (2015) high centrifugal forces can cause microalgae cell rupture, with a loss up to 40% of total lipid content. The level of disruption depends on the toughness and shape of the microalgae cell, as well as the hydrodynamic forces applied to the microalgae (Xu et al. 2015). According to Safafar (2017) larger average cell size will result in higher degree of the leakage, which is in agreement with the results of this study.

Chlorophyll content was higher before centrifugation and decreased after (Table 5). By damaging cell wall, content of the cell will lack the protection layer, which can result in pigment degradation (Hosikian et al. 2010). However, the content of β -carotene significantly increased after centrifugation. Carotenoids accumulate in the chloroplasts and recover in the cell at high centrifugation forces. Therefore, cell rupture can have both positive and negative consequences,

409 depending on the final product. If the aim is to achieve a high extractability of different compounds
410 from the cell, already broken cell walls will favour the extraction process.

411

412 **5. CONCLUSION**

413 *Arthrospira platensis* was able to grow on different dilutions of ICW, which offers potential low-
414 cost source of nutrients for microalgal growth. Partial substitution of synthetic medium with
415 wastewater can potentially reduce microalgal production costs and reduce fresh water requirements.
416 Growth was not detected in 100% ICW due to the non-optimal environment for growth of *A.*
417 *platensis*. Optimal concentration of ICW for obtaining high growth rate with no adverse effect on
418 the biomass composition was shown to be 25% dilution in synthetic medium. This study suggests *A.*
419 *platensis* as a potential species for wastewater treatment. Further research is needed in order to
420 investigate the efficiency of the nutrient removal from industrial process water, as well as testing of
421 possible toxic compound concentrating in the biomass. Furthermore, different environmental stress
422 factors could be tested in combination with ICW, in order to increase production of valuable
423 compounds such as protein, PUFAs and carotenoids.

424 Harvesting methods should be adjusted for specific microalgal species due to their wide diversity. A
425 suitable process needs to be applied, in order to preserve quality of the end product. Microfiltration
426 was demonstrated to be an efficient method for biomass separation with moderate cell rupture, as a
427 result of the filtration shear. However, economically it still cannot compete with standard passive
428 screen filtration commonly used for filamentous and large cell size specie. Also, microfiltration by
429 itself may not be sufficient for harvesting, in case it is necessary to further up-concentrate the
430 microalgal biomass before drying, which then requires additional up-concentrating steps.
431 Centrifugation was shown not to be a suitable harvesting (or up-concentration) method for
432 *A. platensis* due to severe cell damage, followed by cell content leakage. Effects of centrifugal force
433 on *A. platensis* were confirmed by changes in biomass composition including lower protein, lipid
434 and chlorophyll content after centrifugation.

435

436

437

References

- Abdel-Raouf N, Al-Homaidan AA, Ibraheem IBM (2012) Microalgae and wastewater treatment. *Saudi J Biol Sci* 19:257-275.
- Ali SK, Saleh AM (2012) Spirulina: an overview. *Int J Pharm Pharm Sci* 4(3), 9-15.
- Barclay W, Zeller S (1996) Nutritional enhancement of n-3 and n-6 fatty acids in Rotifers and *Artemia Nauplii* by feeding spray-dried *Schizochytrium* sp. *J World Aquacult Soc* 27: 314–322.
- Barros AI, Gonçalves AL, Simões M, Pires JCM (2015) Harvesting techniques applied to microalgae: A review. *Renew Sust Energ Rev* 41:1489–1500.
- Baunillo KE, Tan RS, Barros HR, Luque R (2012) Investigations in microalgal oil production from *Arthrospira platensis*: towards more sustainable biodiesel production. *RCS Advances* 2:11267-11272.
- Becker EV (2007) Micro-algae as a source of protein. *Biotechnol Adv* 25(2):207-210.
- Bilad MR, Arafat HA, Vankelecom IFJ (2014) Membrane technology in microalgae cultivation and harvesting: A review. *Biotechnol Adv* 32:1283–1300.
- Bligh EG, Dyer WJ (1959) A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 37:911-917.
- Chaiklahan R, Nattayaporn C, Wipawan S, Kalyanee P, Boosya B (2010) Cultivation of *Spirulina platensis* using pig wastewater in a semi-continuous process. *J Microbiol Biotechnol* 20:609–614.
- Cohen Z (1997) The chemicals of Spirulina. In: Vonshak A. (ed) *Spirulina platensis (Arthrospira): Physiology, cell biology and biotechnology*. Taylor and Francis, London, pp 175–204.
- Colla LM, Oliveira RC, Reichert C, Costa JAV (2007) Production of biomass and nutraceutical compounds by *Spirulina platensis* under different temperature and nitrogen regimes. *Bioresour Technol* 98:1489–93.
- Costa JAV, Colla LM, Filho PD (2003) *Spirulina platensis* - growth in open raceway ponds using fresh water supplemented with carbon, nitrogen and metal ions. doi: 10.1515/znc-2003-1-214
- Dassey AJ, Theegala CS (2013) Harvesting economics and strategies using centrifugation for cost effective separation of microalgae cells for biodiesel applications. *Bioresour Technol* 128:241-245.
- El-Sheekh MM, Farghl AA, Galal HR, Bayoumi HS (2016) Bioremediation of different types of polluted water using microalgae. *Rend Fis Acc Lincei* 27:401-410.
- EPA (2005) Membrane filtration guidance manual. *Environ Prot* <https://www.epa.gov>; searched on 23 March 2017
- Firestone D.A.O.A.C. (2009) Official Methods and Recommended Practices of the AOCS, 5th ed.; American Oil Chemist's Society: Urbana, IL, USA.
- Guillard RRL, Ryther JH (1962) Studies on marine planktonic diatoms I. *Cyclotella nana* Hustedt and *Detonula confervacea* (Cleve) Gran *Can J Micro* 8: 229-239.
- Hosikian A, Lim S, Halim R, Danquah M.K. (2010) Chlorophyll extraction from microalgae: A review on the process engineering aspects. *Int J Chem Eng* doi:10.1155/2010/391632
- Hu Q (2013) Environmental effects on cell composition. In *Handbook of microalgal culture: Applied phycology biotechnology*. Wiley-Blackwell: Hoboken, New York, pp 114–122.
- Hultberg M, Lind O, Birgersson G, Asp H (2016) Use of the effluent from biogas production for cultivation of *Spirulina*. *Bioprocess Biosyst Eng* doi:10.1007/s00449-016-1726-2
- Kim CJ, Jung YH, Oh HM (2007) Factors indicating culture status during cultivation of *Spirulina (Arthrospira) platensis*. *J Microbiol* 45(2):122-127.
- Klanchui A, Cheevadhanarak S, Prommeenate P (2017) Exploring components of the CO₂-concentrating mechanism in alkaliphilic cyanobacteria through genome-based analysis. *Computat Struct Biotechnol J* 15:340-350.
- Leema MJT, Kirubakaran R, Vinithkumar NV, Dheenan PS, Karthikayulu S (2010) High value pigment production from *Arthrospira (Spirulina) platensis* cultured in seawater. *Bioresur Technol* 101:9221-9227.
- Markou G, Georgakakis D (2011) Cultivation of filamentous cyanobacteria (blue-green algae) in agro-industrial wastes and wastewaters: A review. *Appl Energ* 88:3389–3401.
- Markou G, Ionomou D, Muylaert K (2016) Applying raw poultry litter leachate for the cultivation of *Arthrospira platensis* and *Chlorella vulgaris*. *Algal Res* 13:79-84.

Markou G, Vandamme D, Muylaert K (2014) Ammonia inhibition on *Arthrospira platensis* in relation to the initial biomass density and pH. *Bioresour Technol* 166:259–265.

Mata TM, Martins AA, Caetano NS (2010) Microalgae for biodiesel production and other applications: A review. *Renew Sust Rev*, 14:217-232.

Mezzomo N, Saggiorato AG, Siebert R, Tatsch PO, Lago MC, Hemkemeier M, Costa JAV, Bertolin TE, Colla LM (2010) Cultivation of microalgae *Spirulina platensis* (*Arthrospira platensis*) from biological treatment of swine wastewater. *Cienc Tecnol Aliment* 30(1): 173-178.

Milledge JJ (2011) Commercial application of microalgae other than as biofuels: a brief review. *Rev Environ Sci Biotechnol* doi:10.1007/s11157-010-9214-7

Muhling M, Belay A, Whitton BA (2005) Variation in fatty acid composition of *Arthrospira* (*Spirulina*) strains. *J Appl Phycol* 17:137–146.

Myers J, Kratz WA (1955) Relations between pigment content and photosynthetic characteristics in a blue-green alga. *J gen Physiol* 29:11-22.

Ogbonna JC, Yoshizawa H, Tanaka H (2000) Treatment of high strength organic wastewater by a mixed culture of photosynthetic microorganisms. *J Appl Phycol* 12:277–84.

Phang SM, Miah MS, Yeoh BG, Hashim MA (2000) *Spirulina* cultivation in digested sago starch factory wastewater. *J Appl Phycol* 12:395–400.

Priyadarshani I, Rath B (2012) Commercial and industrial applications of microalgae – A review. *J Algal Biomass Utln* 3(4):89–100.

Rossi N, Derouiniot-Chaplain MP, Legentilhomme P, Petit I (2008) *Arthrospira platensis* harvesting with membranes: Fouling phenomenon with limiting and critical flux. *Bioresour Technol* 99:6162–6167.

Safafar H (2017) Microalgae biomass as an alternative resource for fishmeal and fish oil in the production of fish feed. PhD thesis. Technical University of Denmark, Denmark 34-93 pp

Safafar H, van Wagenen J, Møller P, Jacobsen C (2015) Carotenoids, phenolic compounds and tocopherols contribute to the antioxidative properties of some microalgae species grown on industrial wastewater. *Mar Drugs* 13:7339–7356.

Safi C, Charton M, Pignolet O, Silvestre F, Vaca-Garcia C, Pierre-Yves P (2013) Influence of microalgae cell wall characteristics on protein extractability and determination of nitrogen-to-protein conversion factors. *J Appl Phycol* doi: 10.1007/s10811-012-9886-1

Sassano CEN, Gioielli LA, Ferreira LS, Rodrigues MS, Sato S, Converti A (2010) Evaluation of the composition of continuously cultivated *Arthrospira* (*Spirulina*) *platensis* using ammonium chloride as nitrogen source. *Biomass Bioenerg* 34:1732–1738.

Soundarapandian P, Vasanthi B (2008). Effects of chemical parameters on *Spirulina platensis* biomass production: optimized method for phycocyanin extraction. *Internat J Zool Res* 4(1):1-11.

Wondraczek L, Batentschuk M, Schmidt MA, Borchardt R, Scheiner S, Seemann B, Schweizer P, Brabec CJ (2013) Solar spectral conversion for improving the photosynthetic activity in algae reactors. *Nat Commun* doi:10.1038/ncomms3047

Xu Y, Milledge JJ, Abubakar A, Swamy RAR, Bailey D, Harvey PJ (2015) Effects of centrifugal stress on cell disruption and glycerol leakage from *Dunaliella salina*. *Microalgae Biotechnol* 1:1–8.

Zarrouk C (1966) Contribution a l'étude du cyanophycée. Influence de divers facteurs physiques et chimiques sur la croissance et la photosynthèse de *Spirulina maxima*. PhD thesis, Setch et Gardner, Paris

538 FIGURES AND TABLES:

539

540 **Table 1.** Chemical composition of industrial process water

Item	Unit	Amount
pH	-	8.1
Alkalinity	mmol L ⁻¹	62.6
Suspended solids	mg L ⁻¹	20
Ammonia + ammonium-N	mg L ⁻¹	150
Nitrite + nitrate	mg L ⁻¹	<0.1
Total nitrogen	mg L ⁻¹	190
Total phosphorous	mg L ⁻¹	11
Sulphate	mg L ⁻¹	3.6
Total cyanide	µg L ⁻¹	2.5
EDTA	mg L ⁻¹	<0.5
Sodium (Na)	mg L ⁻¹	1500
Cadmium (Cd)	µg L ⁻¹	<0.05
Copper (Cu)	µg L ⁻¹	3.4
Iron (Fe)	mg L ⁻¹	0.23
Cobalt (Co)	µg L ⁻¹	<0.5

541

542 **Table 2.** Concentration (mg/L) of nitrogen (N) and phosphorous (P) of growth mediums

Growth media	Total N	Total P
100% ZM	412	114
75% ZM + 25% ICW	356	91
50% ZM + 50% ICW	301	62
25% ZM + 75% ICW	245	37
100% ICW	190	11

Fig. 1 Schematic drawing of the microfiltration process. The pump is forcing the microalgal suspension through the microfiltration unit, where feed is passing across the filter membrane at positive pressure relative to the permeate side. Material, that is smaller in size than the membrane pore size, passes through the membrane as permeate, while the rest is retained on the inner side of the membrane as retentate and subsequently, collected back in the bottle

Fig. 2 Effect of different concentrations of ICW (25, 50, 75 and 100%) on growth of *A. platensis* during 12 days cultivation period (n=1)

Fig. 3 Effect of 25% ICW on growth of *A. platensis* during 31 days cultivation period. The results are presented as the means of $n = 4$ measurements from two biological replicates; error bars represent standard deviation

Fig. 4 Protein concentration of *A. platensis* cultivated on 100% ZM and 25% ICW over 30 days. The results are presented as the means of $n = 4$ measurements from two biological replicates; error bars represent standard deviation.

Table 3. Concentration of essential and non-essential amino acids in *A. platensis* cultivated in 100% ZM and 25% ICW. The results are presented as the means $\pm$ standard deviation of $n = 4$ measurements from two biological replicates.

Experiment	Essential amino acids	Non-essential amino acids
100% ZM	56.53 $\pm$ 2.80%	43.47 $\pm$ 2.20%
75% ZM + 25% ICW	57.83 $\pm$ 0.30%	42.17 $\pm$ 0.20%

Fig. 5 Lipid concentration of *A. platensis* cultivated on 100% ZM and 25% ICW over 30 days. The results are presented as the means of $n = 4$ measurements from two biological replicates; error bars represent standard deviation

Table 4. Fatty acid composition of *A. platensis* at day 1, 10, 20 and 30 of the 30 day cultivation: (a) in 100% ZM; (b) in 25% ICW. Concentrations are expressed as % of total fatty acid. The results are presented as the means of $n = 4$ measurements from two biological replicates. Different letters in the same row represent significant differences ($p < 0.05$)

(a)

Fatty acid	Day 1	Day 10	Day 20	Day 30
14:0	2.28 ± 0.12	2.16 ± 0.16	2.17 ± 0.10	1.77 ± 0.22
14:1	0.72 ± 0.00	1.68 ± 0.29	1.41 ± 0.28	1.32 ± 0.87
16:0	37.97 ± 1.95 ^a	40.31 ± 1.62 ^a	40.96 ± 0.70 ^a	40.55 ± 1.30 ^a
16:1 (n-7)	5.43 ± 0.86 ^a	3.43 ± 0.18 ^b	3.13 ± 0.16 ^b	3.62 ± 0.07 ^b
16:2 (n-4)	0.37 ± 0.09	0.20 ± 0.02	0.22 ± 0.03	0.16 ± 0.02
16:3 (n-4)	0.18 ± 0.05	0.17 ± 0.01	0.17 ± 0.01	0.12 ± 0.01
17:0	0.10 ± 0.15	0.22 ± 0.01	0.30 ± 0.00	0.33 ± 0.02
16:4 (n-1)	0.06 ± 0.09	0.16 ± 0.02	0.18 ± 0.00	0.18 ± 0.02
18:0	1.57 ± 0.07	1.65 ± 0.04	1.37 ± 0.11	1.06 ± 0.10
18:1 (n-9)	3.21 ± 0.35 ^a	4.79 ± 0.06 ^b	5.20 ± 0.33 ^b	5.92 ± 1.42 ^b
18:1 (n-7)	2.47 ± 0.17 ^a	2.69 ± 0.47 ^a	1.93 ± 0.14 ^a	2.61 ± 0.65 ^a
18:2 (n-6)	20.94 ± 0.93 ^a	23.39 ± 0.11 ^b	27.19 ± 0.86 ^c	28.16 ± 0.47 ^c
18:3 (n-6)	21.50 ± 1.65 ^a	17.23 ± 0.08 ^b	14.03 ± 0.40 ^c	12.58 ± 0.77 ^c
18:3 (n-3)	0.05 ± 0.08	0.04 ± 0.06	0.07 ± 0.02	0.08 ± 0.01
18:4 (n-3)	0.08 ± 0.11	0.03 ± 0.05	0.05 ± 0.01	0.07 ± 0.00
20:1 (n-11)+(n-9)	0.57 ± 0.52	0.08 ± 0.12	0.08 ± 0.05	0.02 ± 0.01
20:2 (n-6)	0.17 ± 0.00	0.31 ± 0.01	0.38 ± 0.02	0.41 ± 0.04
20:3 (n-6)	0.29 ± 0.03	0.40 ± 0.05	0.53 ± 0.04	0.57 ± 0.27
20:5 (n-3)	0.48 ± 0.11	0.22 ± 0.19	0.13 ± 0.12	0.14 ± 0.00
22:1 (n-11)	0.37 ± 0.29	0.21 ± 0.14	0.08 ± 0.11	0.00 ± 0.00

22:5 (n-3)	0.51 ± 0.09	0.36 ± 0.03	0.35 ± 0.08	0.14 ± 0.04
22:6 (n-3)	0.40 ± 0.56	0.09 ± 0.06	0.04 ± 0.06	0.13 ± 0.00
Σ SAFA	41.92 ± 2.29 ^a	44.34 ± 1.84 ^a	44.79 ± 0.91 ^a	43.71 ± 1.26 ^a
Σ UFA	57.97 ± 5.98 ^a	55.49 ± 1.93 ^a	55.16 ± 2.73 ^a	56.22 ± 3.18 ^a
Σ PUFA	45.03 ± 3.78 ^a	42.61 ± 0.67 ^a	43.34 ± 1.65 ^a	42.74 ± 1.77 ^a

(b)

Fatty acid	Day 1	Day 10	Day 20	Day 30
14:0	1.89 ± 0.08	2.15 ± 0.16	1.29 ± 0.70	1.42 ± 0.75
14:1	0.44 ± 0.02	1.57 ± 0.28	1.95 ± 0.51	2.72 ± 0.87
16:0	38.02 ± 1.49 ^a	38.23 ± 0.14 ^a	38.30 ± 1.28 ^a	37.12 ± 1.30 ^a
16:1 (n-7)	6.55 ± 0.82 ^a	4.85 ± 0.44 ^b	4.15 ± 0.11 ^b	4.25 ± 0.07 ^b
16:2 (n-4)	0.31 ± 0.03	0.31 ± 0.06	0.26 ± 0.01	0.24 ± 0.02
16:3 (n-4)	0.12 ± 0.00	0.17 ± 0.01	0.16 ± 0.00	0.15 ± 0.01
17:0	0.00 ± 0.00	0.24 ± 0.00	0.25 ± 0.03	0.28 ± 0.02
16:4 (n-1)	0.22 ± 0.03	0.21 ± 0.03	0.21 ± 0.03	0.22 ± 0.02
18:0	1.09 ± 0.01	1.38 ± 0.02	1.46 ± 0.19	1.42 ± 0.10
18:1 (n-9)	3.23 ± 0.41 ^a	3.28 ± 0.47 ^a	4.44 ± 1.05 ^a	4.52 ± 1.42 ^a
18:1 (n-7)	5.51 ± 0.39 ^a	5.14 ± 3.24 ^a	3.95 ± 1.42 ^a	4.64 ± 0.65 ^a
18:2 (n-6)	22.04 ± 2.44 ^a	22.14 ± 0.13 ^a	25.38 ± 0.04 ^b	26.21 ± 0.47 ^b
18:3 (n-6)	18.98 ± 2.28 ^a	18.59 ± 1.87 ^a	16.24 ± 0.22 ^a	14.78 ± 0.77 ^b
18:3 (n-3)	0.19 ± 0.18	0.02 ± 0.03	0.10 ± 0.03	0.08 ± 0.01
18:4 (n-3)	0.06 ± 0.02	0.02 ± 0.03	0.07 ± 0.00	0.09 ± 0.00
20:1 (n-11)+(n-9)	0.11 ± 0.03	0.22 ± 0.31	0.03 ± 0.04	0.05 ± 0.01
20:2 (n-6)	0.12 ± 0.00	0.25 ± 0.04	0.39 ± 0.07	0.41 ± 0.04
20:3 (n-6)	0.14 ± 0.02	0.28 ± 0.13	0.72 ± 0.37	0.73 ± 0.27
20:5 (n-3)	0.29 ± 0.07	0.26 ± 0.08	0.25 ± 0.02	0.23 ± 0.00
22:1 (n-11)	0.08 ± 0.00	0.02 ± 0.03	0.00 ± 0.00	0.00 ± 0.00
22:5 (n-3)	0.31 ± 0.06	0.23 ± 0.12	0.34 ± 0.01	0.34 ± 0.04
22:6 (n-3)	0.08 ± 0.01	0.07 ± 0.02	0.00 ± 0.00	0.00 ± 0.00
Σ SAFA	41.00 ± 1.58 ^a	41.99 ± 0.34 ^a	41.30 ± 2.19 ^a	40.24 ± 2.16 ^a
Σ UFA	58.80 ± 6.79 ^a	57.62 ± 7.31 ^a	58.64 ± 3.93 ^a	59.64 ± 4.68 ^a
Σ PUFA	42.87 ± 5.13 ^a	42.55 ± 2.54 ^a	44.13 ± 0.80 ^a	43.47 ± 1.66 ^a

SAFA = saturated fatty acids; UFA = unsaturated fatty acids; PUFA = polyunsaturated fatty acids

(a)

(b)

Fig. 6 Amino acid composition of *A. platensis* cultivated in medium containing: (a) 100% ZM; (b) 25% ICW. The results are presented as the means of $n = 4$ measurements from two biological replicates; error bars represent standard deviation

(a)

(b)

Fig. 7 Pigment composition and concentration ($\mu\text{g/g}$) of *A. platensis* cultivated in medium containing: (a) 100% ZM; (b) 25% ICW. The results are presented as the means of $n = 4$ measurements from two biological replicates; error bars represent standard deviation

Fig. 8 Harvest performance (flux) for *A. platensis* harvested by SiC membrane with $3 \mu\text{m}$ pore size

(a)

(b)

(c)

Picture 1. Optical microscopy (100x) of *A. platensis*: (a) Fresh culture with no cell rupture detectable; (b) Culture after microfiltration with moderate cell rupture detectable; (c) Culture after microfiltration followed by centrifugation with severe cell rupture detectable. Circles indicating mixture of material associated with cellular fragments and EPS

613 **Table 5.** Changes in biomass composition of *A. platensis* after microfiltration (MF) and
614 centrifugation (CF). Values are expressed as mean $\pm$ standard deviation of n = 4 measurements from
615 two replicates

	Protein (% of DW)	Lipid (% of DW)	Chlorophyll ($\mu\text{g/g DW}$)	Carotenoids ($\mu\text{g/g DW}$)
MF	56.70 $\pm$ 2.97	6.06 $\pm$ 0.15	3634.60 $\pm$ 189. 49	1010.07 $\pm$ 334.79
MF + CF	43.69 $\pm$ 2.83	4.40 $\pm$ 0.73	2405.61 $\pm$ 823.29	2229.86 $\pm$ 468.34

616

617