Risk assessment of contaminants of emerging concern in the context of wastewater reuse for irrigation: An integrated modelling approach

Delli Compagni, Riccardo ; Gabrielli, Marco; Polesel, Fabio; Turolla, Andrea; Trapp, Stefan; Vezzaro, Luca; Antonelli, Manuela

Published in:
Chemosphere

Link to article, DOI:
[10.1016/j.chemosphere.2019.125185](https://doi.org/10.1016/j.chemosphere.2019.125185)

Publication date:
2020

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Delli Compagni, R., Gabrielli, M., Polesel, F., Turolla, A., Trapp, S., Vezzaro, L., & Antonelli, M. (2020). Risk assessment of contaminants of emerging concern in the context of wastewater reuse for irrigation: An integrated modelling approach. *Chemosphere*, 242, [125185]. <https://doi.org/10.1016/j.chemosphere.2019.125185>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Risk assessment of contaminants of emerging concern in the context of
wastewater reuse for irrigation: An integrated modelling approach**

Riccardo Delli Compagni¹, Marco Gabrielli¹, Fabio Polesel^{2,3}, Andrea Turolla¹, Stefan Trapp², Luca
Vezzaro², Manuela Antonelli^{1*}

¹ Politecnico di Milano, Department of Civil and Environmental Engineering (DICA), Piazza
Leonardo da Vinci 32, 20133 Milano, Italy

² DTU Environment, Technical University of Denmark, Bygningstorvet, Building 115, 2800
Kongens Lyngby, Denmark

³ DHI A/S, Agern Allé 5, 2970 Hørsholm, Denmark

*Corresponding author: manuela.antonelli@polimi.it

14 **Abstract:**

15 Direct reuse of reclaimed wastewater (RWW) in agriculture has recently received increasing
16 attention as a possible solution to water scarcity. The presence of contaminants of emerging concern
17 (CECs) in RWW can be critical, as these chemicals can be uptaken in irrigated crops and eventually
18 ingested during food consumption.

19 In the present study, an integrated model was developed to predict the fate of CECs in water reuse
20 systems where RWW is used for edible crops irrigation. The model was applied to a case study
21 where RWW (originating from a municipal wastewater treatment plant) is discharged into a water
22 channel, with subsequent irrigation of silage maize, rice, wheat and ryegrass. Environmental and
23 human health risks were assessed for 13 CECs, selected based on their chemical and hazard
24 characteristics. Predicted CEC concentrations in the channel showed good agreement with available
25 measurements, indicating potential ecotoxicity of some CECs (estrogens and biocides) due to their
26 limited attenuation. Plant uptake predictions were in good agreement with existing literature data,
27 indicating higher uptake in leaves and roots than fruits. Notably, high uncertainties were shown for
28 weakly acidic CECs, possibly due to degradation in soil and pH variations inside plants. The human
29 health risk due to the ingestion of wheat and rice was assessed using the threshold of toxicological
30 concern and the hazard quotient. Both approaches predicted negligible risk for most CECs, while
31 sulfamethoxazole and 17 α -ethinylestradiol exhibited the highest risk for consumers. Alternative
32 scenarios were evaluated to identify possible risk minimization strategies (e.g., adoption of a more
33 efficient irrigation system).

34

35 **Keywords:** dietary intake; micropollutants; model-based risk assessment; plant uptake

36

37 **Declarations of interest:** none

38 **1. Introduction**

39 Water scarcity induces monetary and job losses throughout all continents (Asano, 2002; Ding et al.,
40 2011) and the situation is expected to worsen due to population growth, increase of water use per
41 capita, climate change and other stress factors (IPCC, 2014; Gosling and Arnell, 2016). Following
42 circular economy principles, the reuse of reclaimed wastewater (RWW) originating from urban
43 wastewater treatment plants (WWTPs) is identified as one of the main measures to alleviate fresh
44 water depletion (COM/2015/614). This can be a valid alternative to water supply for agricultural
45 irrigation, the largest source of freshwater consumption worldwide (COM/2015/614). This approach
46 has recently been promoted by the European Union, which also proposed minimum requirements
47 for a safe reuse (COM/2018/337). This document acknowledges the need of assessing the risk
48 (where relevant) associated to contaminants of emerging concern (CECs), whose threat for
49 environment and human health is well recognized (Daughton and Ternes, 1999; Monteiro and
50 Boxall, 2010).

51 Existing WWTPs are typically not designed for removal of CECs (e.g. pharmaceuticals, estrogens,
52 biocides), which are generally released from households into sewer systems (Luo et al., 2014;
53 Castiglioni et al., 2018). Hence, some CECs persist in RWW and can reach agricultural fields
54 through irrigation, leading to accumulation in soil (Durán-Alvarez et al., 2009; Gibson et al., 2010)
55 and contamination of groundwater (Siemens et al., 2008; Lesser et al., 2018). Irrigated crops can
56 also accumulate CECs in roots (Miller et al., 2016) and translocate them towards edible plant
57 organs such as leaves and fruits (Goldstein et al., 2014; Christou et al., 2017). Consequently,
58 potential exposure of humans and organisms living in environmental recipients to CECs requires a
59 careful assessment, and adverse effects can be quantified based on the estimated exposure
60 concentrations (Piña et al., 2018). Besides being discontinuous in time and subject to uncertainty
61 (due to e.g. analytics), empirical measurements of environmental CEC levels are site-specific and
62 do not allow extrapolating contamination levels to other RWW reuse systems. In this case,

63 modeling tools can contribute with valuable complementary information (e.g. filling temporal gaps
64 between measurements), and account for the influence of site-specific conditions. The use of
65 complex modelling tools, especially for integrated systems, can be challenged by parameter
66 identifiability (Voinov and Shugart, 2013; Bach et al., 2014) and limited input data availability.
67 These sources of uncertainty can be addressed by using statistical analysis methods (e.g., sensitivity
68 and uncertainty analysis), allowing the identification of important model factors and the
69 quantification of results uncertainty. So far, integrated models have shown the ability of evaluating
70 the impacts of discharges from storm- and wastewater system on the chemical status of surface
71 water (De Keyser et al., 2010). However, a dynamic chemical fate model capable of estimating
72 CEC concentrations (and thereby risk) in different environmental compartments (surface water,
73 soil, crops) under different water management and reuse scenarios is currently in high need.

74 The objective of this work was to develop a flexible integrated model to predict CEC fate in
75 different types of RWW reuse systems, irrespective of size and complexity, and to assess
76 environmental and human health risks. The model was: (i) applied to a real RWW reuse scenario,
77 comprising a discharge channel and cultivation of four different types of crops (silage maize, rice,
78 wheat and ryegrass); (ii) verified with site-specific measurements, where available, and literature
79 data in different compartments; (iii) used to estimate the environmental risk associated to CEC
80 occurrence in surface water, and human health risk following the ingestion of irrigated crops; (iv)
81 used to evaluate alternative risk management and minimization scenarios, which were compared to
82 the present situation and to a worst-case scenario (characterized by the absence of a WWTP).

83

84 **2. Materials and methods**

85 **2.1 Fate model development and evaluation**

86 A generic RWW reuse system was considered to include a surface water body, receiving RWW
87 discharges, and cultivated fields with irrigation. The river water quality model extended with CEC
88 fate processes (from the IUWS_MP library) (Vezzaro et al., 2014) and the coupled soil-plant model
89 (CSPM) (Trapp and Matthies, 1998; Trapp, 2017) were identified as the state-of-the-art models
90 (section 2.1.1) to describe CECs fate in the generic system. These models were extended (section
91 2.1.2) to increase their applicability and coupled (section 2.1.3). A sensitivity analysis (section
92 2.1.4) was also performed to identify the most significant parameters in the developed model.

93

94 **2.1.1 State of the art models**

95 The IUWS_MP library (Vezzaro et al., 2014) utilizes a conceptual approach to simulate river water
96 quality, where water transport is simulated by a series of continuously stirred tanks reactors
97 (CSTRs). Each CSTR consists of two compartments (bulk water and sediments), in which fate
98 processes (abiotic and biotic degradation, sorption to suspended solids and colloids) occur.

99 The CSPM includes (i) a tipping bucket model to describe the movement of water and dissolved
100 CECs through soil and (ii) a numerical four-compartment model to simulate CEC uptake and
101 translocation through xylem and phloem flows, in roots, stem, leaves and fruits. Partitioning
102 between plant tissues and soil, and xylem and phloem flows is described through a detailed
103 intracellular model (Trapp, 2004; Trapp and Horobin, 2005; Trapp, 2009; Trapp 2017).

104

105 **2.1.2 Model extensions**

106 Many CECs are ionized at environmental pH, which affects their partitioning behavior (Franco et
107 al., 2010). Hence, the equations described in Franco and Trapp (2008) were included in the
108 IUWS_MP library to describe sorption of ionized monovalent acidic and basic compounds.

109 In the CSPM, phloem flow introduced to improve the dynamics of the internal plant circulation,
110 through which weakly acidic CECs translocate from leaves through stem to fruits and roots
111 (Gonzalez-Garcia et al., 2019; Kleier & Hsu, 1996).
112 Model structures were further modified to accommodate for temporal and spatial dynamics of
113 environmental conditions (e.g., air, soil and water temperature, water and soil pH, sunlight intensity,
114 etc.), which can substantially affect CEC fate (Kunkel and Radke, 2011; Matamoros and Rodriguez,
115 2017). Moreover, the CSPM was also modified to simulate complex types of irrigational systems, in
116 which water can partially submerge the cultivations (e.g. rice). Particularly, the field capacity of the
117 first soil layer was adjusted to simulate the saturation effect introduced by a standing water layer.
118 The reader is referred to the Supplementary Material (SM) for a detailed description of the
119 extensions of the IUWS_MP library and of CSPM model.

120

121 **2.1.3 Model coupling and software selection**

122 Predicted CEC concentrations were obtained twice a day (noon and midnight) from the river model
123 (implemented in WEST 2014®, DHI A/S, Denmark) to consider high and low peaks, and input to
124 the CSPM (implemented in Microsoft Excel), which provided daily outputs.

125

126 **2.1.4 Identification of influential parameters**

127 A sensitivity analysis (SA) was performed to identify: (i) the most influential model parameters for
128 concentration predictions and (ii) non-sensitive model parameters uncertainties, which could be
129 fixed to a default value without affecting model predictions uncertainties. As for the river model, a
130 one-at-a-time (OAT) approach (Frey and Patil, 2002) was performed by varying each parameter
131 $\pm 5\%$ from its default value. A total of 52 parameters were investigated. The variation in predicted
132 concentrations was calculated as reported in Equation 1:

$$Deviation = \frac{1}{n} \sum_i^n \frac{y_i^{p,max} - y_i^{p,min}}{y_i^{p,default}} \quad (\text{Eq. 1})$$

where $y_i^{p,max}$, $y_i^{p,min}$, $y_i^{p,default}$ are the predicted concentrations obtained with the maximum, minimum and default values of the parameter p for i^{th} time step; n is the number of time steps in the simulation. Three CECs (i.e. diclofenac, triclosan and carbamazepine, see section 2.2.1) were chosen as benchmark for different physicochemical properties. Parameters influencing predicted concentrations were also assessed for four conventional pollutants (ammonia, nitrite, nitrate and phosphorus).

As for the SA of CSPM, partial rank correlation coefficients (PRCCs) (Saltelli et al., 1993) were calculated through the software Crystal Ball 11.1.2 (Oracle®). First, 1000 runs were conducted by varying separately plant and soil parameters. The parameters presenting |PRCCs| above 0.1 were selected and included into a second SA of 10000 runs. This time, parameters with |PRCCs| exceeding $\sqrt{1/n}$ (with n being the number of parameters varied in each simulation) were identified as relevant. The validity of the PRCCs was checked through the R^2 between the two sets of ranks (Sin et al., 2011). The second SA was carried out both with and without degradation in soil to better highlight its sensitivity on model variations.

2.2 Model testing

The developed model was tested on an existing Italian case study, where RWW from a municipal WWTP (flow = $3.5 \text{ m}^3 \text{ s}^{-1}$) is the only source providing water through a surface water channel to subsequently irrigate crop fields. The channel extends for about 12 km and provides irrigation water for 90 farms with more than 40 km^2 of crop fields (Pizza, 2014). Cultivated crops include winter wheat and rice, which are sold for human food consumption; and silage maize and ryegrass that are used for animal feed production. Irrigation systems included surface irrigation, the dominant form

155 of irrigation in southern European countries (Masseroni et al., 2017) for all the crops but rice, which
156 is grown in paddies. In the area, crop rotation is a standard procedure and is performed each season.

157

158 **2.2.1 Chemicals**

159 Thirteen CECs of different classes of use were assessed: clarithromycin (CLA), sulfamethoxazole
160 (SMX), diclofenac (DCF), ibuprofen (IBU), paracetamol (PAR), carbamazepine (CBZ), furosemide
161 (FUR), 17- α ethinylestradiol (EE2), 17- β estradiol (E2), estrone (E1), perfluorooctanoic acid
162 (PFOA), perfluorooctane sulfonate (PFOS) and triclosan (TCS). These CECs were selected from an
163 initial group of 80 CECs, for which measurements were available at the outlet of the WWTP
164 (Castiglioni et al., 2018a, 2018b). Specifically, two criteria were used: i) selected CECs had to
165 cover a broad range of physicochemical properties in terms of: air/water partitioning coefficient
166 (K_{aw}), normalized organic carbon partitioning coefficients (K_{oc}), and ionization state at
167 environmental pH; ii) selected CECs had to belong to different CEC classes (e.g. antibiotics,
168 hormones, etc.) and to possibly exceed the predicted no-effect concentration (PNEC).
169 Physicochemical properties (Table S2) and degradation rates (Tables S3-S15) of the selected CECs
170 were collected from literature (see SM for sources) and/or retrieved from QSAR software
171 (Advanced Chemistry Development Inc., 2015; National Food Institute, Technical University of
172 Denmark, 2018).

173

174 **2.2.2 System conceptualization**

175 The RWW reuse system was conceptualized as shown in Figure 1. The channel was described by
176 the river modules from the IUWS_MP library, resulting in 11 in-series CSTRs, each one
177 approximately 500 m long. This length was selected in agreement with Benedetti et al. (2004) as a
178 compromise between the need of simulating temporal dynamics in a realistic way and the required
179 computational effort. Within the channel, nine equidistant locations were selected, each having

180 irrigation water withdrawal that was proportional to the required water demand to harvest the
181 closest crops. The water demand ($\text{m}^3 \text{d}^{-1}$) was defined as the product of the water consumption to
182 harvest one unit of land and total land extension (Brouwer et al., 1992). Crop types and
183 corresponding extensions were derived from georeferenced maps of the area
184 (www.geoportale.regione.lombardia.it) and elaborated with QGIS 2.18. Periods of water
185 withdrawal and crop rotation were simulated according to the irrigation calendars (Figure 1b) based
186 on agronomic information of the area (Nelli and Sodi, 2007; Provincia di Milano, 2007; Borrelli et
187 al., 2014; Moretti et al., 2015). Two water withdrawal scenarios were considered: (i) scenario *MR*,
188 where silage maize and ryegrass were grown at the same time of rice and ryegrass and, (ii) scenario
189 *MW*, where silage maize and ryegrass were grown at the same time of winter wheat. Moreover,
190 different irrigation systems were simulated depending on crop types: (i) a monthly irrigation water
191 pulse for silage maize, ryegrass and winter wheat and, (ii) a continuous irrigation for rice. Three
192 types of soils were found predominantly in the study area, with more than one crop per type of soil
193 (see Table S18). For this reason, crop simulations were repeated for different soils. The simulated
194 soil structure was also adapted to match the identified stratigraphy for a soil depth of 1 m, splitting
195 (if needed) the widest horizons over two layers.

196

Figure 1. Outline of the wastewater reuse system (a) and irrigation calendars (b) for the two crop rotation scenarios.

2.2.3 Model input and parameters

Concentrations of conventional pollutants in RWW (i.e. ammonia, nitrite, nitrate and phosphorus) represented a 24-hour time-proportional composite sample and were retrieved from the laboratory measurements conducted by WWTP staff. CEC concentrations in RWW were obtained from previous measurements at the outlet of the WWTP (Castiglioni et al., 2018a, 2018b) and assumed constant over the year. Such assumption can be held valid for CECs (e.g., DCF and CBZ) that show low to no seasonal fluctuations in usage (Sui et al., 2011). For antibiotics such as CLA, for which seasonal fluctuations were observed (McArdell et al., 2003), measurements may reflect a worst-case scenario as campaigns were conducted during winter time. Environmental conditions (Tables S16-17) were directly obtained from public databases or indirectly estimated (e.g., potential evaporation, soil temperature, bulk density and water retention properties). Crop-specific data (Table S19) were collected from literature preferring, where available, local sources. Soil initial conditions (CECs

213 concentrations and water content) were assumed equal to the pseudo-steady state reached using the
214 median CECs. For more details, the reader is referred to SM.

215

216 **2.2.4 Uncertainty propagation**

217 The uncertainty associated to CEC concentrations in RWW and transformation rates (e.g., aerobic
218 degradation in water, photodegradation and aerobic/anoxic degradation in sediments) were
219 propagated to the predicted concentration in the surface water channel through a Monte-Carlo based
220 Uncertainty Analysis (UA). Uniform, triangular or empirical parameter distributions were chosen
221 based on literature values (Tables S3-S15). For every CEC, 2000 runs were performed.

222 Resulting uncertainties in the predicted CEC concentrations along the river were then propagated,
223 together with soil degradation rates and influential crop parameters, to CSPM predictions.
224 Specifically, triangular distributions were considered for soil degradation rates and CEC
225 concentrations. Crop parameters were considered as uniformly distributed with a variation of $\pm 10\%$
226 from literature data (Felle, 2001). Attention was paid to avoid parameter combinations that may
227 lead to non-realistic scenarios (Steinwand et al., 2001). 2000 latin-hypercube Monte Carlo runs
228 were performed for each CEC and soil type (Table S18). Physicochemical properties (Table S2) of
229 perfluorinated compounds exhibited a high degree of uncertainty due to analytical limitations
230 (Wang et al., 2011), and were thus included in UA. More details on UA are provided in SM.

231

232 **2.3 Model objectives**

233 The three main objectives of the developed integrated model were:

234 *I – Fate predictions and assessment of model performance:* long term simulations (≥ 1 year) were
235 performed to assess CECs fate in surface water, irrigated crops and groundwater. Assessment of
236 model performance was carried out by comparing model predictions with site-specific
237 measurements, where available (as for DCF) and/or literature data. For the surface water channel,

238 further verification was made by deriving first-order attenuation rates by fitting predicted median
239 concentrations at different locations and comparing to experimental literature data. In this case,
240 overall attenuation derives from the combination of different biotic and abiotic degradation
241 processes.

242 As for the uptake in crops, the bioconcentration factor (BCF, $\text{g}_{\text{dw}}^{-1} \text{g}_{\text{dw}}^{-1}$) was calculated as
243 $C_{\text{organ}}/C_{\text{soil}}$, where C_{organ} and C_{soil} are the predicted dry weight concentrations in a given plant organ
244 (root, stem, leaf, fruit) and in the first soil layer. There is currently limited consensus on calculation
245 of BCFs since different approaches have been considered with respect to C_{soil} values (Polesel et al.,
246 2015). For this reason, both the simulated maximum and median soil concentrations were used for
247 the BCF calculation, corresponding to BCF_{max} and $\text{BCF}_{\text{median}}$, respectively. Calculated BCF was
248 compared with literature (when not explicitly reported, empirical BCFs were derived from reported
249 concentration data) except for CLA, which showed limited plant uptake (Limmer and Burken,
250 2014; Lamshoeft et al., 2018). Predicted leaching of CECs into groundwater was also compared
251 with measurements from the nearby area.

252

253 *II – Ecological and human health risk assessment:* environmental risk assessment was carried out
254 through the risk quotients (RQs) (Hernando et al., 2006; Kuzmanovic et al., 2013). RQs were
255 calculated for each CEC as $PEC/PNEC$, where PEC is the predicted environmental concentration
256 and $PNEC$ the predicted no-effect concentration. $PNECs$ were obtained from literature (Table S1)
257 and $PECs$ were the predicted concentrations at the end of the surface water channel; specifically,
258 maximum, median and minimum concentrations were predicted to obtain RQ_{max} , $\text{RQ}_{\text{median}}$ and
259 RQ_{min} , respectively. Two alert thresholds of RQ above 0.1 and 1 were identified as medium and
260 high risk, respectively (Hernando et al., 2006). Monthly frequency of exceedance above the alert
261 thresholds was calculated as additional risk indicator.

262 The human health risk associated to dietary intake was calculated with two different approaches: the
263 threshold of toxicological concern (TTC) (Kroes et al., 2004; Malchi et al., 2014) and the hazard
264 quotient (HQ) (Prosser et al., 2014b; Prosser and Sbiley, 2015). TTC values were obtained through
265 the Cramer classes decision tree implemented in the Toxtree v3.1 software (Patlewicz et al., 2008)
266 and used to calculate the crop ingestion required for posing a risk as $(BW \text{ TTC}) / C_{fw}$, where BW
267 (kg) is the reference body weight (EFSA, 2012) and C_{fw} ($\mu\text{g}/\text{kg}_{dw}$) is the CEC concentration in the
268 edible part, based on fresh weight. These values were then compared with typical Italian food
269 consumption rates for infants and adults (Leclercq et al., 2009). HQs were calculated as the ratio
270 between the estimated daily intake (EDI) and the admissible daily intake (ADI), which is the
271 amount of CECs that can be consumed daily over a person's lifespan without evocating an adverse
272 effect. The level of concern was set to 0.1 as dietary ingestion represents a single pathway of
273 exposure (Prosser and Sibley, 2015). Hazard Index (HI) was also calculated as the sum of HQs to
274 provide a conservative first-tier estimate of the mixture risk (Prosser and Sibley, 2015; Evans et al.,
275 2015). Median and 97.5% percentile were derived for HQ, TTC and HI values from UA
276 propagation. Further details on the two methods are given in SM.

277 III – *Management scenarios analysis*

278 Two agronomic management strategies were simulated to assess their potential for human health
279 risk mitigation compared to the existing surface irrigation:

- 280 (i) the adoption of sprinklers, which require a lower investment cost than drip irrigation, and
281 are a more efficient irrigation system than surface irrigation (i.e. irrigation efficiency
282 increases from 60% to 75% according to Brouwer et al., 1989);
- 283 (ii) one week extension of the period without irrigation before harvest (also known as crop-
284 drying stage). Waiting periods before harvest are a commonly required health safety
285 measure in case of biosolids application (Prosser et al., 2014b).

286 The benefits in terms of human health risk reduction of the existing WWTP, although not
287 specifically designed to remove CECs, were simulated in a specific scenario, where untreated
288 wastewater was directly used for irrigation. Specifically, CEC concentrations measured in WWTP
289 influent (Castiglioni et al., 2018a) were used as input to the river model and first order attenuation
290 rates (calculated beforehand) were used to estimate *PECs* at the end of the channel. Based on *PECs*
291 as input, the CSPM was then used to predict concentrations in fruits, allowing to estimate dietary
292 intake and health risk for infants as shown earlier.

293 **3. Results and discussion**

294 **3.1 Sensitivity analysis**

295 Among the 52 parameters of the river model in the IUWS_MP library, 12 parameters (Figure S2)
296 had an impact on predicted CEC concentrations. The fraction of CEC sorbed onto total suspended
297 solids (TSS) showed higher sensitivity if compared to the dissolved fraction sorbed to colloids,
298 mainly due to sedimentation parameters, in agreement with De Schepper et al. (2012). As to
299 conventional pollutants, ammonium, nitrite and nitrate were impacted by microbial growth
300 parameters, in agreement with Deksissa et al. (2004), while phosphorus was influenced by the
301 sorption rate and sedimentation parameters (Figure S2).

302 For the CSPM model, variations of CECs concentrations were influenced by several soil and plants
303 parameters (Figure S3), in agreement with previous modeling studies (Trapp, 2015) and
304 experimental data (Chen et al., 2013; Goldstein et al., 2014; Dodgen et al., 2015; Lamshoeft et al.,
305 2018). Notably, the effect of the Arrhenius temperature correction coefficient on soil concentrations
306 ($|\text{PRCC}| > 0.9$) out-weighed the effect of the other parameters, as seen before (Legind et al., 2011).
307 Predicted concentrations of the weakly acidic DCF showed considerable sensitivity to internal plant
308 pH, which was further assessed.

309

310 **3.2 Fate analysis**

311 **3.2.1 Surface water channel**

312 Verification of IUWS_MP for conventional pollutants was adequate for the intended purpose,
313 showing predicted concentrations in agreement with the measurements (≤ 0.75 log units) and being
314 able to follow seasonal trends (Figure S4).

315 Along the water channel, model predictions (Figure 2 and Figures S5-14) showed that all the
316 investigated CECs were almost entirely found as dissolved in the water phase ($\sim 98\%$ of the load
317 released from the WWTP). The only exception was CLA, for which the fractions on colloids and

318 TSS were, on average, equal to 6% and 17% of the total mass (Figure 2c). Most of the simulated
319 CECs have lower sorption affinity than CLA ($\log K_{ow} = 3.2$), or they were negatively charged at the
320 measured pH (i.e. electrical repulsion). These results were in good agreements with the
321 measurements in the Ebro river (Ferreira da Silva et al., 2011), where most of the CECs were found
322 as predominantly dissolved (84%, 87%, 95%, 95%, 95% of the total mass for DCF, CLA, PAR,
323 CBZ and IBU, respectively). The width of concentration prediction bounds (about ± 200 % of the
324 median) was mainly explained by the propagation of the uncertainties associated to the initial
325 concentrations of CECs in RWW and by degradation rates. Low variation in predicted
326 concentrations (~ 0.3 log units) resulted from the propagation of uncertainties of conventional
327 pollutant parameters.

328 The temporal dynamics of predicted concentrations followed different daily and/or seasonal
329 patterns depending on the simulated CEC. For example, photodegradation caused an important
330 reduction in concentrations for DCF (52-89%) and FUR (25-87%) between night and day time, in
331 agreement with the findings of Hanamoto et al. (2013). Predicted DCF concentrations during day
332 time were also confirmed by the available measurements (Figure 2a). Photodegradation and
333 biodegradation also caused seasonal patterns with the lowest concentrations during June, July and
334 August (Figure 2a and 2b), when temperature and sunlight irradiation reached their maximum
335 values (22-25 °C and up to 900 W m⁻², respectively). A considerable reduction was observed in
336 summer months, as compared to December, for DCF (30-41%; summer median: 204-244 ng L⁻¹),
337 and FUR (49-72%; summer median: 49.4-90.6 ng L⁻¹). Moreover, when rice was cultivated, median
338 concentrations of DCF and FUR were 16% and 67% lower than when winter wheat was cultivated
339 (orange and red lines in Figure 2). In fact, a high water withdrawal for rice cultivation led to a
340 reduction of the water level and thus enhancing photodegradation and sedimentation since rates of
341 these two processes are inversely proportional to the water depth (Schwarzenbach, 2003).

342 Due to the low tendency to photodegrade (median $k_{\text{pho, nearsurf}} \leq 23.1 \text{ d}^{-1}$) and the overall slow
 343 biodegradation in water (median $k_{\text{bio, water}} \leq 0.32 \text{ d}^{-1}$), the other CECs showed smaller daily
 344 fluctuations (intra-day variation: $\leq 10 \%$), seasonal trends (reduction in summer months compared
 345 to December: $\leq 11\%$) and variations due to water withdrawal (concentration decrease during rice
 346 cultivation: $\leq 4\%$) (see Figures S5-S14).

347 **Figure 2.** Annual variation (from January 1st to December 31st) of daytime DCF (a), FUR (b), CLA
 348 (c) concentrations at the end of the channel.

350

351 Predicted attenuation rates were in good agreement with literature data (Figure S15), with slight
352 underestimation (≤ 1 order of magnitude). Differences might be due to the biofilm growth on
353 submerged vegetation and sediment surface (Winkler et al., 2001; Sabaliunas et al., 2003; Kunkel
354 and Radke, 2011; Writer et al., 2013), that enhances attenuation processes with respect to
355 biodegradability processes accounted by our model.

356 Median attenuation rates varied throughout the year up to one order of magnitude due to different
357 environmental conditions (temperature, sunlight intensity and water depth), with predicted
358 maximum and minimum rates during July and December, respectively. Due to low CLA
359 degradation (median $k_{\text{pho, nearsurf}} = 0.17 \text{ d}^{-1}$, median $k_{\text{bio, water}} = 3.85\text{E-}^{02} \text{ d}^{-1}$), CLA attenuation rate
360 variations were mostly caused by sedimentation, which was enhanced by the lower summer water
361 depth, in agreement with Castiglioni et al. (2006). Similar seasonal variations of the attenuation
362 rates were also observed by Matamoros et al. (2017) and Labadie and Budzinski (2005).

363

364 3.2.2 Crops

365 In Figure 3a we report BCF_{max} predictions for three CECs in different plant compartments, where
366 the maximum simulated soil concentration is used for normalization (see Figures S16-S19 for the
367 BCF_{max} calculated with maximum soil concentrations for all the other CECs). Figure 3b-c presents
368 $\text{BCF}_{\text{median}}$ values for fruits and leaves of the investigated crops based on the simulated median soil
369 concentration, given the larger data availability for plausibility check (see Figures S16-S19 for roots
370 and stems $\text{BCF}_{\text{median}}$). Box-plots were obtained by aggregating BCFs based on crop type (i.e.
371 maize, rice, ryegrass and wheat), and resulting ranges (i.e. width of the whiskers) included both
372 variability (of CEC concentrations in the irrigation water, soil pH and soil organic content SOM)
373 and uncertainty of model input.

374 Within crops (Figure 3a), predicted results for weakly acidic CECs, such as DCF, PFOA, FUR and
375 IBU, mainly show accumulation in roots, which can be partly attributed to phloem translocation.
376 For these CECs, a considerable BCF variation (more than factor 100) was generally obtained. This
377 could be attributed to pH variability within the plant cell organs (Vreugdenhil and Koot-Gronsveld,
378 1989; Felle, 2001), which can strongly affect speciation of ionizable CECs and, consequently, the
379 extent of CECs adsorption and translocation within the plant (Trapp, 2004). Moreover, large BCF
380 variations for PFOA and PFOS were also associated to $\log K_{ow}$ and K_{HSA} uncertainty, which affect
381 in-plant translocation and in-soil bioavailability (Trapp, 2009). TCS also showed maximum
382 accumulation in roots, but mainly due to its high lipophilicity, which limits partitioning into xylem
383 (Hsu et al., 1990). Conversely, neutral and hydrophilic CECs, such as CBZ, translocated upwards
384 through the xylem to the leaves (Collins et al., 2006; Miller et al., 2016), where they showed the
385 highest accumulation. In this case, variations of BCF were mainly caused by SOM variability,
386 which influenced sorption of CECs in neutral form, such as estrogens (E1, E2 and EE2), TCS, PAR
387 and CBZ (Goldstein et al., 2014), affecting bioavailability and uptake. SMX presented a more
388 uniform accumulation among the plant organs due to both high phloem and xylem mobility (Kleier
389 and Hsu, 1996; Goldstein et al., 2014). Regardless of the type of CEC, rice was generally the crop
390 showing the highest uptake (Figure 3a), given that the irrigation system constantly floods the crop
391 over the irrigational period, thus enhancing CECs loading. Conversely, the other crops were
392 intermittently irrigated, and degradation and growth dilution lower uptake between irrigation
393 events.

394 Box-plot medians (Figure 3b-c) were generally close (within 1 order of magnitude) to
395 measurements, while for estrogens and PAR few to no measurements were available. Conversely,
396 CBZ, TCS and SMX (having the highest data availability) showed measured values overall lower
397 than predicted BCF_{median} . This may be due to metabolization within the plant compartments, which
398 has been experimentally observed for CBZ in leaves (Riemenschneider et al., 2016; Goldstein et al.,

399 2014; Malchi et al., 2014), for TCS in roots (Macherius et al., 2012) and for SMX (Dodgen et al.,
 400 2013; Bircher et al., 2015; He et al., 2017; Li et al., 2018). In the model, metabolization within
 401 plants was assumed negligible since no kinetic data are available within the existing literature to
 402 account for such process. The approach used with respect to in-plant metabolism represents a
 403 realistic worst-case scenario. Moreover, in-plant metabolism does not always lead to degradation of
 404 the chemical but may lead to the formation of (i) conjugated metabolites that are prone to be
 405 transformed back to their parent analogue (Macherius et al., 2012); (ii) other metabolites that may
 406 be as hazardous as their respective parent (e.g., the case of carbamazepine metabolites; Malchi et
 407 al., 2015).

408 BCF calculation was highly affected by in-soil degradation, since it increases the differences
 409 between maximum and median soil concentration (hence, the difference between BCF_{max} and
 410 BCF_{median}). In particular, BCF_{median} variation for estrogens was the most affected (≥ 2 orders of
 411 magnitude) by in-soil degradation rates. The effect was more pronounced for maize, as the longer
 412 time between irrigation events allows for larger variations of the median soil concentration than for
 413 other crops. Overall, these results further confirm the need for a standard methodology for BCF
 414 calculation to allow for inter-study comparison of empirical results.

415 The model also predicted mass loss above 10^{-4} mg m⁻² season⁻¹ from soil to groundwater for CBZ,
 416 SMX, IBU, PFOS and PFOA (Figure S20), in agreement with the high GW concentrations
 417 measured in the area (Castiglioni et al., 2018b). Losses to groundwater were also predicted for DCF
 418 and FUR, while measurements were below detection limits, possibly due to further degradation
 419 outside the model boundary. Castiglioni et al. (2018b) also found traces of CLA and TCS in the
 420 groundwater, in disagreement with model predictions that showed almost complete accumulation in
 421 the surface soil layers, as confirmed by Chen et al. (2013) and Xu et al. (2009). The chemicals
 422 might stem from other sources (e.g. house imhoff tanks located in rural area) or reach groundwater
 423 by macropore transport.

Figure 3. (a) Bioconcentration factors (BCFs, g_{dw}/g_{dw}) for CBZ, IBU and SMX in different crops and soil types. BCFs were calculated with respect to maximum soil concentrations (COLOR FIGURE). (b, c) Fruit and leaf BCFs (g_{dw}/g_{dw}) for all evaluated CECs in silage maize, winter wheat and rice in different soil types, and comparison with literature values (lit.). BCFs were calculated with respect to median soil concentrations. BioS indicates experimental studies conducted with the use of biosolids. Red borders indicate the use of a LOD value in the calculations. Refs: Bizkarguenaga et al. (2016), Blaine et al. (2013), Carter et al. (2014), Christou et al. (2017), Fu et al. (2016), Goldstein et al. (2014), Hurtado et al. (2017), Malchi et al. (2014), Mordechay et al. (2018), Navarro et al. (2017), Pannu et al. (2012), Shenker et al. (2011), Wen et al. (2014), Winker et al. (2010), Wu et al. (2010, 2012), Yager et al. (2014), Yoo et al. (2011).

432 3.3 Risk assessment

433 3.3.1 Ecological risk

434 Three estrogens (E1, E2 and EE2), two antibiotics (SMX and CLA), one non-steroidal anti-
435 inflammatory drug (IBU) and one antiseptic (TCS) showed RQ_{median} above the alert thresholds at
436 least once over the 1-year simulation. EE2 displayed the highest risk both in terms of magnitude
437 and frequency of exceedance, with a daily median concentration always above the PNEC (RQ_{median}
438 ~ 27), and a maximum concentration during winter corresponding to a RQ_{max} of 46. Predicted
439 RQ_{median} for E2, CLA and TCS were approximately 1.5, with an average monthly frequency of
440 exceeding the corresponding PNECs of 0.95, 0.79 and 0.77, respectively. IBU, E1 and SMX
441 showed RQ_{median} below 1, although a non-negligible ecological risk (RQ_{max} ranging between 0.15
442 and 0.45) was shown. Other investigated CECs had an RQ below 0.1, showing low or negligible
443 ecological risk (see Table S22). These results, in good agreement with what found by Riva et al.
444 (2019) in the surrounding area, highlight that certain CECs may negatively affect aquatic organisms
445 at a medium distance (approximately 12 km) from the WWTP.

446

447 3.3.2 Human health risk from dietary uptake

448 The estimated consumption of rice and wheat required to exceed TTCs (Table S23) showed
449 unrealistic values (>2 kg/d per person) for all the CECs but SMX. For this compound a
450 consumption of 10 g/d and 50 g/d (considering the 97.5th rice concentration percentile) might imply
451 potential genotoxic effects (Malchi et al., 2015) on infants and adults, respectively. Conversely,
452 SMX had an HQ below the alert threshold of 0.1 in accordance with the work of Christou et al.
453 (2017). Consequently, experimental measurements should be performed in rice to clarify a possible
454 risk related to SMX. The calculated negligible human health risk is in good agreement with the
455 findings of Malchi et al. (2014) and Riemenschneider et al. (2016).

456 As for the predicted risk by the HQ approach (Table 1), results indicated negligible risk for all the
457 CECs, although an HQ (infants) of 0.3 was calculated for EE2. A comparison with the TTC could
458 not be done, since this latter approach is not valid for classes of compounds such as estrones (Kroes
459 et al., 2004).

460 To conclude, the HI value (sum of all individual HQs, see Table 1) was below 1, showing no
461 potential risk for human health from the investigated CECs. Moreover, the risk from indirect
462 exposure (possible biotransfer of CECs from forage to milk and beef and, subsequently to humans)
463 was not quantified since available empirical models (Travis and Arms, 1988; Hendriks et al., 2007)
464 failed to properly describe fate of ionisable compounds and were not validated for the CECs here
465 considered. Mechanistic models (Rosenbaum et al., 2009; Trapp et al., 2008) predict
466 bioaccumulation in milk and meat solely for chemicals with strong partitioning into lipids, which is
467 not the case for the investigated compounds.

468

469

470 **Table 1.** HQs and HIs for crops irrigated with water withdrawn at the end of the channel. P97.5
 471 stands for values calculated with the 97.5th percentile CEC concentration.

ECs	Rice		Wheat		Total	
	Median	P97.5	Median	P97.5	Median	P97.5
Infants						
SMX	2.94E-03	2.03E-02	1.86E-04	1.58E-03	3.12E-03	2.19E-02
DCF	2.66E-06	6.35E-04	6.14E-08	1.26E-05	2.72E-06	6.48E-04
IBU	1.12E-09	5.27E-07	1.21E-10	5.94E-08	1.24E-09	5.86E-07
PAR	7.12E-07	1.05E-06	5.64E-08	9.08E-08	7.69E-07	1.14E-06
CBZ	2.62E-03	3.91E-03	2.21E-03	2.80E-03	4.84E-03	6.71E-03
FUR	5.37E-08	8.49E-05	1.25E-07	9.82E-05	1.79E-07	1.83E-04
EE2	2.77E-02	2.57E-01	1.95E-03	3.85E-02	2.97E-02	2.95E-01
E2	1.07E-05	2.53E-05	8.72E-07	2.02E-06	1.15E-05	2.73E-05
E1	5.23E-04	4.82E-03	4.06E-05	7.80E-04	5.63E-04	5.60E-03
PFOS	6.98E-06	2.75E-04	5.44E-06	1.24E-04	1.24E-05	4.00E-04
PFOA	2.23E-08	1.22E-04	1.17E-06	2.66E-04	1.19E-06	3.88E-04
TCS	6.19E-09	1.79E-08	2.57E-09	1.04E-08	8.76E-09	2.83E-08
Hazard Index					3.82E-02	3.31E-01
Adults						
SMX	4.45E-04	3.08E-03	8.88E-05	7.52E-04	5.34E-04	3.83E-03
DCF	4.03E-07	9.63E-05	2.93E-08	6.03E-06	4.33E-07	1.02E-04
IBU	1.70E-10	7.99E-08	2.93E-08	6.03E-06	2.27E-10	1.80E-07
PAR	1.08E-07	1.59E-07	2.69E-08	4.33E-08	1.35E-07	2.20E-07
CBZ	3.98E-04	5.93E-04	1.06E-03	1.33E-03	1.45E-03	1.93E-03
FUR	8.14E-09	1.29E-05	5.96E-08	4.69E-05	6.78E-08	5.97E-05
EE2	4.20E-03	3.89E-02	9.32E-04	1.84E-02	5.13E-03	5.73E-02
E2	1.62E-06	3.84E-06	4.16E-07	9.63E-07	2.03E-06	4.80E-06
E1	7.93E-05	7.30E-04	1.94E-05	3.72E-04	9.86E-05	1.10E-03
PFOS	1.06E-06	1.31E-04	2.60E-06	5.94E-05	3.65E-06	1.90E-04
PFOA	3.38E-09	1.85E-05	5.58E-07	1.27E-04	5.62E-07	1.46E-04
TCS	9.38E-10	2.71E-09	1.23E-09	4.96E-09	2.17E-09	7.67E-09
Hazard Index					7.22E-03	6.47E-02

472

473 3.4 Risk management strategies

474 The adoption of a more efficient irrigation system (i.e. sprinklers) led to a calculated reduction of
 475 the human health risk due to the lower CECs load that reach the crops. Particularly, median HQs for

476 infants decreased by 25% for EE2 and SMX. In the model simulations, direct deposition of
477 chemicals on leaf surfaces due to sprinkling has not been considered and may lead to an additional
478 uptake of chemicals via this pathway, while significant uptake into grains via surface deposition is
479 unlikely. However, weak acids could reach the grains by phloem transport. We are not aware of
480 studies addressing this process. On the other hand, the adoption of a crop-drying stage before rice
481 harvest did not show any substantial reduction ($< 2\%$) in the median rice concentrations of EE2
482 (from $3.71 \cdot 10^{-4}$ ng/g_{dw} to $3.65 \cdot 10^{-4}$ ng/g_{dw}) and SMX (from $5.21E^{-01}$ ng/g_{dw} to $5.12E^{-01}$ ng/g_{dw}). In
483 fact, at the later stages of crop growth the transpiration rate is minimum (Hidayati et al., 2016),
484 consequently resulting in limiting the predicted CECs uptake during this stage. To best of our
485 knowledge no experimental studies compared the CECs uptake under different irrigation systems
486 preventing further verification of the model results.

487 Although not explicitly designed to remove CECs, the existing WWTP showed to be crucial for
488 guaranteeing human health safety. In the scenario where the WWTP was absent, only few CECs
489 (PFOS, PFOA, DCF, FUR and CBZ) had a HQ median risk below 0.1, while it was substantially
490 above 1 for all the other CECs. For example, HQ_{median} of TCS increased from 1.43 to 429, reaching
491 in the worst-case scenario an HQ_{max} of 842. Predicted rice concentrations increased up to 3 orders
492 of magnitude, especially for TCS, IBU, E1 and PAR, which are efficiently removed ($>99\%$) by the
493 WWTP (Castiglioni et al., 2018b). These results were in agreement with the work of Goldstein et al.
494 (2014) and Wu et al. (2014), in which higher CECs concentrations in TWW led to higher
495 accumulation in crops. The risk for infants associated to SMX and PAR (considering the 97.5th
496 percentile of the concentration in rice) showed a required ingestion to exceed TTC lower than 1 g d^{-1}
497 and the HQs of E1, EE2 and SMX were 0.34, 0.26 and 0.33, respectively. Median and 97.5th
498 percentile for infants HIs increased up to 8 times, reaching values of 0.27 and 0.95. respectively.
499 These results further indicate that irrigation with untreated wastewater requires specific attention
500 when considering the risk posed by CECs.

501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523

4. Conclusions

This study proposed a new integrated model for simulating the fate of different contaminants of emerging concern under dynamic conditions when reclaimed wastewater is reused for irrigation purposes. By extending and integrating existing fate models, the model showed capability and flexibility in describing the fate of 13 contaminants, covering a wide range of physicochemical properties, across different compartments and over long time intervals. Model predictions for contaminant occurrence in irrigation water and crops were generally verified with measured data, thus allowing for the evaluation of ecological and human health risk posed by wastewater discharge and reuse. Although dependent on the approach used, sulfamethoxazole and 17 α -ethinylestradiol were associated to potential health risk from dietary intake of irrigated crops, particularly rice. While the existing wastewater treatment plant contributes significantly to reduce the overall risk, further reductions can be obtained by adopting more efficient irrigation practices. Overall the proposed integrated approach represents a valuable decision support tool for assessing the safety of reclaimed wastewater reuse practices within the circular economy, particularly when considering recent developments in the regulation (EC, 2018).

Acknowledgments

The authors would like to acknowledge the COST Action ES1403 NEREUS “New and emerging challenges and opportunities in wastewater reuse”, supported by COST (European Cooperation in Science and Technology, www.cost.eu), for enabling the collaboration among the authors of the paper.

524 **References**

- 525 Asano, T. (2002). Water from (waste)water - the dependable water resource. *Water Science and*
526 *Technology*, 45(8):23-33.
- 527 Bach, P. M., Rauch, W., Mikkelsen, P. S., McCarthy, D. T., and Deletic, A. (2014). A critical
528 review of integrated urban water modelling: Urban drainage and beyond. *Environmental*
529 *Modelling and Software*, 54(4):88-107.
- 530 Benedetti, L., Meirlaen, J., Sforzi, F., Facchi, A., Gandolfi, C., & Vanrolleghem, P. A. (2004).
531 Dynamic integrated modelling: a case study on the river Lambro. *Novatech 2004*, 1–8.
- 532 Borrelli, L., Castelli, F., Ceotto, E., Cabassi, G., & Tomasoni, C. (2014). Maize grain and silage
533 yield and yield stability in a long-term cropping system experiment in northern Italy. *European*
534 *Journal of Agronomy*, 55, 12–19.
- 535 Brouwer, C., Hoevenaars, J.P.M. & Van Bosch, B.E., (1992) *Scheme Irrigation Water Needs and*
536 *Supply*. Rome, Italy: FAO.
- 537 Brouwer, C., Prins, K., & Heibloem, M. (1989). *Irrigation Water Management: Irrigation*
538 *Scheduling*. Rome, Italy: FAO.
- 539 Castiglioni, S., Bagnati, R., Fanelli, R., Pomati, F., Calamari, D., & Zuccato, E. (2006). Removal of
540 pharmaceuticals in sewage treatment plants in Italy. *Environmental Science and Technology*,
541 40(1), 357–363.
- 542 Castiglioni, S., Davoli, E., Riva, F., Palmiotto, M., Camporini, P., Manenti, A., & Zuccato, E.
543 (2018a). Data on occurrence and fate of emerging contaminants in a urbanised area. *Data in*
544 *Brief*, 17, 533–543.
- 545 Castiglioni, S., Davoli, E., Riva, F., Palmiotto, M., Camporini, P., Manenti, A., & Zuccato, E.
546 (2018b). Mass balance of emerging contaminants in the water cycle of a highly urbanized and
547 industrialized area of Italy. *Water Research*, 131:287-298.

548 Chen, W., Xu, J., Lu, S., Jiao, W., Wu, L., & Chang, A. C. (2013). Fates and transport of PPCPs in
549 soil receiving reclaimed water irrigation. *Chemosphere*, 93(10), 2621–2630.

550 Christou, A., Agüera, A., Bayona, J.M., Cytryn, E., Fotopoulos, V., Lambropoulou, D., Manaia,
551 C.M., Michael, C., Revitt, M., Schröder, P., Fatta-Kassinos, D. (2017). The potential
552 implications of reclaimed wastewater reuse for irrigation on the agricultural environment: The
553 knowns and unknowns of the fate of antibiotics and antibiotic resistant bacteria and resistance
554 genes – A review. *Water Res.* 123:448-467.

555 Collins, C., Fryer, M., and Grosso, A. (2006). Plant uptake of non-ionic organic chemicals.
556 *Environmental Science and Technology*, 40(1):45-52.

557 COM (2015) 614 Closing the loop- An EU action plan for the Circular Economy. Communication
558 from the Commission to the European Parliament, the Council, the European Economic and
559 Social Committee and the Committee of the Regions. European Commission, Brussels, Belgium.

560 COM (2018) 337 Proposal for a REGULATION OF THE EUROPEAN PARLIAMENT AND OF
561 THE COUNCIL on minimum requirements for water reuse. Proposal from the Commission to
562 the European Parliament, the Council, the European Economic and Social Committee and the
563 Committee of the Regions. European Commission, Brussels, Belgium.

564 Daughton, C. G. and Ternes, T. A. (1999). Pharmaceuticals and personal care products in the
565 environment: agents of subtle change? *Environmental Health Perspectives*, 107(suppl 6):907-
566 938.

567 De Keyser, W., Gevaert, V., Verdonck, F., Nopens, I., De Baets, B., Vanrolleghem, P.A.,
568 Mikkelsen, P.S., Benedetti, L., (2010). Combining multimedia models with integrated urban
569 water system models for micropollutants. *Water Sci. Technol.* 62, 1614–1622.

570 De Schepper, V. C. J., Holvoet, K. M. A., Benedetti, L., Seuntjens, P., & Vanrolleghem, P. A.
571 (2012). Extension of the River Water Quality Model No. 1 with the fate of pesticides. *Journal of*
572 *Hydroinformatics*, 14(1), 48–64.

573 Deksis, T., Meirlaen, J., Ashton, P. J., & Vanrolleghem, P. A. (2004). Simplifying dynamic river
574 water quality modelling: A case study of inorganic nitrogen dynamics in the Crocodile River
575 (South Africa). *Water, Air, and Soil Pollution*, 155(1–4), 303–320.

576 Ding, Y., Hayes, M. J., and Widhalm, M. (2011). Measuring economic impacts of drought: A
577 review and discussion. *Disaster Prevention and Management*, 20(4):434-446.

578 Dodgen, L. K., Ueda, A., Wu, X., Parker, D. R., and Gan, J. (2015). Effect of transpiration on plant
579 accumulation and translocation of PPCP/EDCs. *Environmental Pollution*, 198:144-153.

580 Dudley, S., Sun, C., Jiang, J., & Gan, J. (2018). Metabolism of sulfamethoxazole in *Arabidopsis*
581 *thaliana* cells and cucumber seedlings. *Environmental Pollution*, 242, 1748–1757.

582 Durán-Alvarez, J. C., Becerril-Bravo, E., Castro, V. S., Jiménez, B., and Gibson, R. (2009). The
583 analysis of a group of acidic pharmaceuticals, carbamazepine, and potential endocrine disrupting
584 compounds in wastewater irrigated soils by gas chromatography-mass spectrometry. *Talanta*,
585 78(3):1159-1166.

586 EC (European Commission), 2018. Proposal for a Regulation of the European Parliament and of the
587 Council on minimum requirements for water reuse. Brussels, 28th May 2018.

588 EFSA Scientific Committee. (2012). Guidance on selected default values to be used by the EFSA
589 Scientific Committee, Scientific Panels and Units in the absence of actual measured data. *EFSA*
590 *Journal*, 10(3), 2579.

591 Evans, R. M., Scholze, M., & Kortenkamp, A. (2015). Examining the feasibility of mixture risk
592 assessment: A case study using a tiered approach with data of 67 pesticides from the Joint
593 FAO/WHO Meeting on Pesticide Residues (JMPR). *Food and Chemical Toxicology*, 84, 260–
594 269.

595 Felle, H. H. (2001). pH: Signal and messenger in plant cells. *Plant Biology*, 3(6), 577–591.

596 Ferreira da Silva, B., Jelic, A., López-Serna, R., Mozeto, A. A., Petrovic, M., & Barceló, D. (2011).
 597 Occurrence and distribution of pharmaceuticals in surface water, suspended solids and sediments
 598 of the Ebro river basin, Spain. *Chemosphere*, 85(8), 1331–1339.

599 Franco, A., Struijs, J., Gouin, T., & Price, O. R. (2013). Evolution of the sewage treatment plant
 600 model SimpleTreat: Applicability domain and data requirements. *Integrated Environmental*
 601 *Assessment and Management*, 9(4), 560–568.

602 Franco, A., Ferranti, A., Davidsen, C., & Trapp, S. (2010). An unexpected challenge: Ionizable
 603 compounds in the REACH chemical space. *International Journal of Life Cycle Assessment*,
 604 15(4), 321–325.

605 Franco, A., & Trapp, S. (2008). Estimation of the soil-water partition coefficient normalized to
 606 organic carbon for ionizable organic chemicals. *Environ. Toxicol. Chem.* 27, 1995-2004.

607 Frey, C. H., & Patil, S. R. (2002). Identification and Review of Sensitivity Analysis Methods. *Risk*
 608 *Analysis*, 22(3), 553–578.

609 Gibson, R., Durán-Alvarez, J. C., Estrada, K. L., Chávez, A., and Jiménez Cisneros, B. (2010).
 610 Accumulation and leaching potential of some pharmaceuticals and potential endocrine disruptors
 611 in soils irrigated with wastewater in the Tula Valley, Mexico. *Chemosphere*, 81(11):1437-1445.

612 Goldstein, M., Shenker, M., and Chefetz, B. (2014). Insights into the Uptake Processes of
 613 Wastewater-Borne Pharmaceuticals by Vegetables. *Environmental Science and Technology*,
 614 48:5593-5600.

615 Gonzalez-Garcia, M., Fernandez-Lopez, C., Polesel, F., Trapp, S. (2019). Predicting the uptake of
 616 emerging organic contaminants in vegetables irrigated with treated wastewater – Implications for
 617 food safety assessment. *Environ. Res.* 172, 175-181.

618 Gosling, S. N. and Arnell, N. W. (2016). A global assessment of the impact of climate change on
 619 water scarcity. *Climatic Change*, 134(3), 371-385.

620 Hanamoto, S., Nakada, N., Yamashita, N., & Tanaka, H. (2013). Modeling the photochemical
621 attenuation of down-the-drain chemicals during river transport by stochastic methods and field
622 measurements of pharmaceuticals and personal care products. *Environmental Science and*
623 *Technology*, 47(23), 13571–13577.

624 Hendriks, A. J., Smítková, H., & Huijbregts, M. A. J. (2007). A new twist on an old regression:
625 Transfer of chemicals to beef and milk in human and ecological risk assessment. *Chemosphere*,
626 70(1), 46–56.

627 Hernando, M. D., Mezcuá, M., Fernández-Alba, A. R., & Barceló, D. (2006). Environmental risk
628 assessment of pharmaceutical residues in wastewater effluents, surface waters and sediments.
629 *Talanta*, 69(2 SPEC. ISS.), 334–342.

630 Hidayati, N., Triadiati, & Anas, I. (2016). Photosynthesis and Transpiration Rates of Rice
631 Cultivated Under the System of Rice Intensification and the Effects on Growth and Yield.
632 *HAYATI Journal of Biosciences*, 23(2), 67–72.

633 Hsu, F. C., Marxmiller, R. L., & Yang, A. Y. S. (1990). Study of Root Uptake and Xylem
634 Translocation of Cinmethylin and Related Compounds in Detopped Soybean Roots Using a
635 Pressure Chamber Technique. *Plant Physiology*, 93(4), 1573–1578.

636 IPCC (2014). *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and*
637 *Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the*
638 *Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United*
639 *Kingdom and New York, NY, USA.*

640 Kleier, D. A., & Hsu, F. C. (1996). Phloem Mobility of Xenobiotics. VII. The Design of Phloem
641 Systemic Pesticides. *Weed Science*, 44(3), 749–756.

642 Kroes, R., Renwick, A. ., Cheeseman, M., Kleiner, J., Mangelsdorf, I., Piersma, A., ... Würtzen, G.
643 (2004). Structure-based thresholds of toxicological concern (TTC): guidance for application to
644 substances present at low levels in the diet. *Food and Chemical Toxicology*, 42(1), 65–83.

645 Kunkel, U., & Radke, M. (2011). Reactive tracer test to evaluate the fate of pharmaceuticals in
646 rivers. *Environmental Science and Technology*, 45(15), 6296–6302.

647 Kuzmanovic, M., Banjac, Z., Ginebreda, A., Petrovic, M., & Barcelo, D. (2013). Prioritization:
648 Selection of Environmentally Occurring Pharmaceuticals to Be Monitored. In M. Petrovic, D.
649 Barcelo, & S. Pérez (Eds.), *Comprehensive Analytical Chemistry* (2nd ed., Vol. 62, pp. 71–90).

650 Labadie, P., & Budzinski, H. (2005). Determination of steroidal hormone profiles along the Jalle
651 d'Eysines River (near Bordeaux, France). *Environmental Science and Technology*, 39(14),
652 5113–5120.

653 Lamshoeft, M., Gao, Z., Ressler, H., Schriever, C., Sur, R., Sweeney, P., ... Reitz, M. U. (2018).
654 Evaluation of a novel test design to determine uptake of chemicals by plant roots. *Science of the*
655 *Total Environment*, 613–614, 10–19.

656 Lazarova, V. and Asano, T. (2013). Milestones in water reuse: main challenges, keys to success and
657 trends of development. An overview. In Lazarova, V., Asano, T., Bahri, A., and Anderson, J.,
658 editors, “Milestones in Water Reuse”, pages 1-22.

659 Leclercq, C., Arcella, D., Piccinelli, R., Sette, S., & Le Donne, C. (2009). The Italian National Food
660 Consumption Survey INRAN-SCAI 2005–06: main results in terms of food consumption. *Public*
661 *Health Nutrition*, 12(12), 2504–2532.

662 Legind, C. N., Kennedy, C. M., Rein, A., Snyder, N., & Trapp, S. (2011). Dynamic plant uptake
663 model applied for drip irrigation of an insecticide to pepper fruit plants. *Pest Management*
664 *Science*, 67(5), 521–527.

665 Lesser, L. E., Mora, A., Moreau, C., Mahlnecht, J., Hernández-Antonio, A., Ramirez, A. I., and
666 Barrios-Piña, H. (2018). Survey of 218 organic contaminants in groundwater derived from the
667 world's largest untreated wastewater irrigation system: Mezquital Valley, Mexico. *Chemosphere*,
668 198:510-521.

669 Li, Y., Chuang, Y. H., Sallach, J. B., Zhang, W., Boyd, S. A., & Li, H. (2018). Potential metabolism
 670 of pharmaceuticals in radish: Comparison of in vivo and in vitro exposure. *Environmental*
 671 *Pollution*, 242, 962–969.

672 Limmer, M. A., & Burken, J. G. (2014). Plant Translocation of Organic Compounds: Molecular and
 673 Physicochemical Predictors. *Environmental Science and Technology Letters*, 1(2), 156–161.

674 Luo, Y., Guo, W., Ngo, H. H., Nghiem, L. D., Hai, F. I., Zhang, J., Liang, S., Wang, X. C. (2014).
 675 A review on the occurrence of micropollutants in the aquatic environment and their fate and
 676 removal during wastewater treatment. *Science of the Total Environment*, 473-474:619-641.

677 Macherius, A., Eggen, T., Lorenz, W. G., Reemtsma, T., Winkler, U., & Moeder, M. (2012).
 678 Uptake of Galaxolide, Tonalide, and Triclosan by Carrot, Barley, and Meadow Fescue Plants.
 679 *Journal of Agricultural and Food Chemistry*, 60(32), 7785–7791.

680 Malchi, T., Maor, Y., & Chefetz, B. (2015). Comments on “Human health risk assessment of
 681 pharmaceuticals and personal care products in plant tissue due to biosolids and manure
 682 amendments, and wastewater irrigation.” *Environment International*, 82, 110–112.

683 Malchi, T., Maor, Y., Tadmor, G., Shenker, M., & Chefetz, B. (2014). Irrigation of Root Vegetables
 684 with Treated Wastewater: Evaluating Uptake of Pharmaceuticals and the Associated Human
 685 Health Risks. *Environmental Science & Technology*, 48(16), 9325–9333.

686 Masseroni, D., Ricart, S., de Cartagena, F. R., Monserrat, J., Gonçalves, J. M., de Lima, I., ...
 687 Gandolfi, C. (2017). Prospects for improving gravity-fed surface irrigation systems in
 688 mediterranean european contexts. *Water (Switzerland)*, 9(1), 20.

689 Miller, E. L., Nason, S. L., Karthikeyan, K. G., and Pedersen, J. A. (2016). Root Uptake of
 690 Pharmaceuticals and Personal Care Product Ingredients. *Environmental Science and*
 691 *Technology*, 50(2):525-541.

692 Matamoros, V., & Rodríguez, Y. (2017). Influence of seasonality and vegetation on the attenuation
 693 of emerging contaminants in wastewater effluent-dominated streams. A preliminary study.
 694 Chemosphere, 186, 269–277.

695 McArdell, C. S., Molnar, E., Suter, M. J. F., & Giger, W. (2003). Occurrence and Fate of Macrolide
 696 Antibiotics in Wastewater Treatment Plants and in the Glatt Valley Watershed, Switzerland.
 697 Environmental Science and Technology, 37(24), 5479–5486.

698 Monteiro, S. C. and Boxall, A. B. (2010). Occurrence and Fate of Human Pharmaceuticals in the
 699 Environment. In Whitacre, D. M., editor, “Reviews of Environmental Contamination and
 700 Toxicology”, volume 202, pages 54-154. Springer, New York, NY.

701 Moretti, B., Grignani, C., Bechini, L., & Sacco, D. (2015). Efficacia e convenienza delle catch crop.
 702 L’informatore Agrario, (36), 52–56.

703 National Food Institute, Technical University of Denmark (2018). Danish (Q)SAR Database. URL:
 704 <http://qsar.food.dtu.dk>. Last accessed: 23-03-2019.

705 Nelli, E. and Sodi, F. (2007). Riso - *Oryza sativa* L. - Atlante delle coltivazioni erbacee - Piante
 706 industriali. URL: <http://www.agraria.org/coltivazionierbacee/riso.htm>. Last accessed: 5-10-2018.

707 Patlewicz, G., Jeliaskova, N., Safford, R. J., Worth, A. P., & Aleksiev, B. (2008). An evaluation of
 708 the implementation of the Cramer classification scheme in the Toxtree software. SAR and QSAR
 709 in Environmental Research, 19(5–6), 495–524.

710 Piña, B., Bayona, J.M., Christou, A., Fatta-Kassinos, D., Guillon, E., Lambropoulou, D., Michael,
 711 C., Polesel, F., Sayen, S. (2018). On the contribution of reclaimed wastewater irrigation to the
 712 potential exposure of humans to antibiotics, antibiotic resistant bacteria and antibiotic resistance
 713 genes – NEREUS COST Action ES1403 position paper. Journal of Environmental Chemical
 714 Engineering.

715 Pizza, F. (2014). Agricultural reuse of treated wastewater : the case of Milano-Nosedo municipal
 716 wastewater treatment plant. In D. Santoro, Trojan Technologies, & Western University (Eds.),

717 Wastewater and Biosolids Treatment and Reuse: Bridging Modeling and Experimental Studies.
 718 ECI Symposium Series.

719 Polesel, F., Plósz, B.G., Trapp, S. (2015). From consumption to harvest: Environmental fate
 720 prediction of excreted ionizable trace organic chemicals. *Water Research* 84, 85-98.

721 Prosser, R. S., Lissemore, L., Topp, E., & Sibley, P. K. (2014a). Bioaccumulation of triclosan and
 722 triclocarban in plants grown in soils amended with municipal dewatered biosolids.
 723 *Environmental Toxicology and Chemistry*, 33(5), 975–984.

724 Prosser, R. S., & Sibley, P. K. (2015). Human health risk assessment of pharmaceuticals and
 725 personal care products in plant tissue due to biosolids and manure amendments, and wastewater
 726 irrigation. *Environment International*, 75, 223–233.

727 Prosser, R. S., Trapp, S., & Sibley, P. K. (2014b). Modeling Uptake of Selected Pharmaceuticals
 728 and Personal Care Products into Food Crops from Biosolids-Amended Soil. *Environmental*
 729 *Science & Technology*, 48(19), 11397–11404.

730 Provincia di Milano (2007). Piano di Settore Agricolo. Relazione Generale.

731 Riemenschneider, C., Al-Raggad, M., Moeder, M., Seiwert, B., Salameh, E., & Reemtsma, T.
 732 (2016). Pharmaceuticals, Their Metabolites, and Other Polar Pollutants in Field-Grown
 733 Vegetables Irrigated with Treated Municipal Wastewater. *Journal of Agricultural and Food*
 734 *Chemistry*, 64(29), 5784–5792.

735 Riva, F., Zuccato, E., Davoli, E., Fattore, E., & Castiglioni, S. (2019). Risk assessment of a mixture
 736 of emerging contaminants in surface water in a highly urbanized area in Italy. *Journal of*
 737 *Hazardous Materials*, 361, 103–110.

738 Rosenbaum, R. K., Mckone, T. E., & Jolliet, O. (2009). CKow: A dynamic model for chemical
 739 transfer to meat and milk. *Environmental Science and Technology*, 43(21), 8191–8198.

740 Sabaliunas, D., Webb, S. F., Hauk, A., Jacob, M., and Eckhoff, W. S. (2003). Environmental fate of
 741 Triclosan in the River Aire Basin, UK. *Water Research*, 37(13):3145–3154.

742 Saltelli, A., Andres, T. H., & Homma, T. (1993). Sensitivity analysis of model output: An
 743 investigation of new techniques. *Computational Statistics & Data Analysis*, 15(2), 211–238.

744 Schwarzenbach, R. P., Gschwend, P. M., & Imboden, D. M. (2003). *Environmental Organic*
 745 *Chemistry* (Second Edition).

746 Siemens, J., Huscsek, G., Siebe, C., and Kaupenjohann, M. (2008). Concentrations and mobility of
 747 human pharmaceuticals in the world's largest wastewater irrigation system, Mexico City-
 748 Mezquital Valley. *Water Research*, 42(8-9):2124-2134.

749 Sin, G., Gernaey, K. V., Neumann, M. B., van Loosdrecht, M. C. M., & Gujer, W. (2011). Global
 750 sensitivity analysis in wastewater treatment plant model applications: Prioritizing sources of
 751 uncertainty. *Water Research*, 45(2), 639–651.

752 Steinwand, A. L., Harrington, R. F., & Groeneveld, D. P. (2001). Transpiration coefficients for
 753 three Great Basin shrubs. *Journal of Arid Environments*, 49(3), 555–567.

754 Sui, Q., Huang, J., Deng, S., Chen, W., & Yu, G. (2011). Seasonal variation in the occurrence and
 755 removal of pharmaceuticals and personal care products in different biological wastewater
 756 treatment processes. *Environmental Science and Technology*, 45(8), 3341–3348.

757 Trapp, S. (2017). Coupled soil-plant uptake model for monovalent ionics. First release June 2017 at
 758 https://homepage.env.dtu.dk/stt/2017Release_Plant_Model/index.htm

759 Trapp, S. (2015). Calibration of a plant uptake model with plant- and site-specific data for uptake of
 760 chlorinated organic compounds into radish. *Environmental Science and Technology*, 49(1), 395–
 761 402.

762 Trapp S. (2009) Bioaccumulation of Polar and Ionizable Compounds in Plants. In: Devillers J. (eds)
 763 *Ecotoxicology Modeling. Emerging Topics in Ecotoxicology (Principles, Approaches and*
 764 *Perspectives)*, vol 2. Springer, Boston, MA

765 Trapp S, Ma L Bomholtz, Legind CN. (2008). Coupled mother-child model for bioaccumulation of
 766 POPs in nursing infants. *Environmental Pollution* 156, 90-98.

767 Trapp, S. (2004). Plant Uptake and Transport Models for Neutral and Ionic Chemicals.
 768 Environmental Science and Pollution Research, 11(1), 33–39.

769 Trapp, S., & Horobin, R. W. (2005). A predictive model for the selective accumulation of chemicals
 770 in tumor cells. European Biophysics Journal, 34(7), 959–966.

771 Trapp, S., & Matthies, M. (1998). Chemodynamics and Environmental Modelling. An introduction.
 772 New York, NY: Springer-Verlag.

773 Travis, C. C., & Arms, A. D. (1988). Bioconcentration of Organics in Beef, Milk, and Vegetation.
 774 Environmental Science and Technology, 22(3), 271–274.

775 Vezzaro, L., Benedetti, L., Gevaert, V., De Keyser, W., Verdonck, F., De Baets, B., ... Mikkelsen,
 776 P. S. (2014). A model library for dynamic transport and fate of micropollutants in integrated
 777 urban wastewater and stormwater systems. Environmental Modelling and Software, 53, 98–111.

778 Voinov, A., Shugart, H.H., 2013. “Integronsters”, integral and integrated modeling. Environ.
 779 Model. Softw. 39, 149–158.

780 Vreugdenhil, D., & Koot-Gronsveld, E. A. M. (1989). Measurements of pH, Sucrose and
 781 Potassium-Ions in the Phloem Sap of Castor Bean (*Ricinus-Communis*) Plants. Physiologia
 782 Plantarum, 77, 385–388.

783 Wang, Z., MacLeod, M., Cousins, I. T., Scheringer, M., & Hungerbühler, K. (2011). Using
 784 COSMOtherm to predict physicochemical properties of poly- and perfluorinated alkyl
 785 substances (PFASs). Environmental Chemistry, 8(4), 389–398.

786 Winkler, M., Lawrence, J. R., & Neu, T. R. (2001). Selective degradation of ibuprofen and clofibric
 787 acid in two model river biofilm systems. Water Research, 35(13), 3197–3205.

788 Writer, J. H., Antweiler, R. C., Ferrer, I., Ryan, J. N., & Thurman, E. M. (2013). In-stream
 789 attenuation of neuro-active pharmaceuticals and their metabolites. Environmental Science and
 790 Technology, 47(17), 9781–9790.

791 Wu, X., Conkle, J. L., Ernst, F., & Gan, J. (2014). Treated wastewater irrigation: Uptake of
792 pharmaceutical and personal care products by common vegetables under field conditions.
793 Environmental Science and Technology, 48(19), 11286–11293.

794 Xu, J., Wu, L., & Chang, A. C. (2009). Degradation and adsorption of selected pharmaceuticals and
795 personal care products (PPCPs) in agricultural soils. Chemosphere, 77(10), 1299–1305.

Risk assessment of contaminants of emerging concern in the context of wastewater reuse for irrigation: An integrated modelling approach

Supplementary Material

Number of pages: 32;

Number of Tables: 24;

Number of Figures: 20;

Riccardo Delli Compagni¹, Marco Gabrielli¹, Fabio Polesel^{2,3}, Andrea Turolla¹, Stefan Trapp², Luca Vezzaro², Manuela Antonelli^{1*}

¹ Politecnico di Milano, Department of Civil and Environmental Engineering (DICA), Piazza Leonardo da Vinci 32, 20133 Milano, Italy

² DTU Environment, Technical University of Denmark, Bygningstorvet, Building 115, 2800 Kongens Lyngby, Denmark

³ DHI A/S, Agern Allé 5, 2970 Hørsholm, Denmark

*Corresponding author: manuela.antonelli@polimi.it

Index

SI.1 Models extension	3
SI.1.1 IUWS_MP	3
SI.1.2 Coupled soil-plant model (CSPM)	3
SI.2 Chemicals	4
SI.2.1 CECs physicochemical properties and degradation rates	4
SI.3 Model parameters and model inputs	12
SI.3.1 Soil properties	13
SI.3.2 Plant parameters	14
SI.4 Uncertainty propagation	16
SI.5 Sensitivity analyses (SA)	19
SI.6 Fate analysis	20
SI.6.1 Traditional pollutants	20
SI.6.2 Uncertainty of CECs concentrations in canal	21
SI.6.3 Attenuation along the canal	24
SI.6.4 Bioconcentration factors	25
SI.7 Risk assessment	30
SI.7.1 Environmental risk – Results	30
SI.7.2 Human health risk – Methods	31
SI.7.3 Human health risk – Results	31
SI.8 References	33

SI.1 Models extension

SI.1.1 IUWS_MP

CECs are often ionized at environmentally relevant pH (Franco et al., 2010). While sorption of neutral chemicals occurs due to Van der Waals forces, in the presence of ions, the process is dominated by the stronger electrical forces. Hence, neutral molecules, anions and cations present different sorption behaviors which are furthermore affected by the degree of ionic speciation (Franco and Trapp, 2008). The Koc formula was updated accordingly to Franco et al. (2013) as shown in Eq. S1.

$$Koc_{acid} = \Phi_n 10^{0.54 \log Kow + 1.11} + \Phi_i 10^{0.11 \log Kow + 1.54} \quad (\text{Eq. S1})$$

$$Koc_{base} = 10^{\log Dow + 2.78}$$

The sorption of the charged and neutral fractions of monovalent chemicals is described as a function of Kow or Dow, pK_a and, in case of acids, also the solution pH. Φ_n and Φ_i are respectively the neutral and anionic CECs fractions calculated at pH_{opt} = pH - 0.6. The range of applicability is pK_a < 10 for monovalent acids and pK_a > 4 for monovalent bases (Franco et al., 2013).

Environmental properties are intrinsically dynamic (Kunkel and Radke, 2011; Matamoros and Rodriguez, 2017) and the use of constant values would not properly describe realistic conditions. Hence, the use of external measurement inputs was allowed for the following properties:

- Sunlight intensity
- Water pH
- Air temperature
- Water temperature
- Wind speed

Irrigation is, also, seasonal dependent (Portmann et al., 2008). For this reason, the flow splitters, used to simulate the water withdrawal, were, thus, enhanced to describe a dynamic pattern throughout the year.

SI.1.2 Coupled soil-plant model (CSPM)

Figure S1 shows the modelled flows mainly responsible for CECs traslocation inside the plants.

Figure S1. Modelled CECs flows inside the plants. In red the new phloem flux added.

As multiple crops are commonly grown on the same field within the same year (Borrelli et al., 2014), the model was extended to describe the succession of two different crops within the same year. Furthermore, an extra input was added to the model to account for the water and CECs mass added to the first soil layer through irrigation and the time-step used in the solution scheme was refined to prevent numerical issues.

Soil pH is not constant with depth, especially in agricultural field in which the surface layers are extensively plowed and fertilized (Zhang et al., 2017b). Therefore, a pH value was given to each of the five soil layers present in the model. Moreover, soil temperatures vary throughout the year due to the fluctuations of air temperature and solar radiation. As depth increases, temperatures are more uniform throughout the year, compared to the overlying layers (Xing and Spitler, 2017). Hence, a daily temperature was added for each of the layers 2-5, while the surface layer temperature was assumed equal to the air above.

SI.2 Chemicals

Table S1. PNECs of selected CECs.

		PNEC value (ng/L)	Reference
clarithromycin	CLA	200	Zhao et al. (2017b)
sulfamethoxazole	SMX	890	Huang et al. (2018)
diclofenac	DCF	9800	Zhao et al. (2017b)
ibuprofen	IBU	18	Huang et al. (2018)
paracetamol	PAR	367	Riva et al. (2019)
carbamazepine	CBZ	9000	Zhao et al. (2017b)
furosemide	FUR	45100	Riva et al. (2019)
17- β estradiol	E2	2	Caldwell et al. (2012)
estrone	E1	6	Caldwell et al. (2012)
17- α ethinylestradiol	EE2	0.1	Caldwell et al. (2012)
perfluorooctanoic acid	PFOA	1.07E06	Gredelj et al. (2018)
perfluorooctane sulfonate	PFOS	1.6E04	Gredelj et al. (2018)
triclosan	TCS	2.6	Gredelj et al. (2018)

SI.2.1 CECs physicochemical properties and degradation rates

Table S2 presents the physicochemical values used during the simulations. QSAR estimates of CLogP and ECOSAR of PFOS and PFOA were discarded as not reliable (Arp et al., 2006).

Table S2. Physicochemical properties of the selected CECs. All properties from Advanced Chemistry Development Inc. (2015), unless noted. (b) indicates a basic pK_a

CEC	Mol mass (g/mol)	pK _a	Vapor P (mmHg)	Log(Sw) (M)	Log(Kow)	Log(Koc)	Log(K _{HSA})
CLA	747.95	8.5 (b)	0	-3.29	3.2		2.77
SMX	253.28	5.7	1.9E-09	-2.62	0.89	2.34 ²⁸ , 1.79 ²⁸ , 1.3 ²⁸ , 1.4 ²⁸ , 2.06 ²⁸ , 3.47 ²⁸	2.86

Table S2. Cont.

DCF	296.15	4.4	1.59E-07	-5.1	4.4		4.91
IBU	206.28	4.3	1.39E-04	-3.99	3.4		4.74
PAR	151.06	10.2	1.43E-06	-0.99	0.46	1.55 ¹	2.43
CBZ	236.27	-	5.78E-07	-3.33	2.3		3.91
FUR	330.74	3.5	0	-3.66	2.03		5.27
E2	272.38	10	9.8E-09	-4.64	3.36		4.41
E1	270.37	9.9	1.54E-08	-3.95	3.13		1.06
EE2	296.40	10	3.7E-09	-4.7	3.54		3.55
PFOA	414.07	2.4, 0.5, 3.8 ⁵ , 0.5 ⁶ , 1.31 ⁷ , 1.01 ⁴ , <1 ¹⁵ , 2.5 ¹² , 2.8 ³ , 0.9 ⁸ , 0 ⁹ , 0 ¹³ , -0.21 ¹⁰ , 2.14 ¹¹	3.14E-02 ¹³ , 3.11E-03 ¹³ , 3.14E-02 ¹³ , 2.4E-03 ¹⁴ , 2.25E-04 ¹⁴ , 9.02E-02 ¹⁹ , 1.5E-01 ¹⁹	-4.54, -1.98 ² , -2.01 ¹¹ , -2.09 ² , -1.64 ² , -2.73 ⁸	4.6, 7.75, 4.3 ¹⁴ , 5.3 ⁸ , 4.59 ¹⁴ , 1.92 ¹³ , 6.26 ¹⁵	2.06 ¹⁷ , 1.9 ¹⁷ , 2.17 ¹⁷ , 1.1 ¹⁵ , 3.2 ¹⁵ , 1.47 ²⁷	5.02, 4.0 ²⁰ , 3.4 ²⁰ , 3.42 ²¹ , 4.34 ²² , 3.57 ²³ , 4.38 ²⁴ , 5.16 ²⁴ , 4.5 ²⁴
PFOS	500.13	-5.7, -3.7, -3.41 ⁸ , 0.14 ¹⁰ , -3.3 ¹²	2.6E-02 ¹⁴ , 2.54E-01 ¹⁴ , 2.34E-03 ¹⁹ , 1.13E-01 ¹⁵	-5.05, -3.84 ⁸ , - 3.79 ¹⁵	2.59, 7.03, 5.25 ¹⁴ , 5.26 ¹⁴ , 2.45 ¹⁶ , 4.67 ¹⁵	1.6 ¹⁵ , 4.8 ¹⁵ , 2.57 ¹⁷ , 2.57 ¹⁷ , 3.1 ¹⁷ , 1.82 ¹⁸	5.02, 7.1 ²⁰ , 3.4 ²⁰ , 3.51 ²⁵ , 3.6 ²⁵ , 3.66 ²³ , 5.12 ²³ , 3.7 ²⁶
TCS	289.54	8.8	4E-06	-4.46	4.76		4.81

References: 1: ECHA (2006), 2: Jensen et al. (2008), 3: Brace (1962), 4: Igarashi and Yotsuyanagi (1992), 5: Burns et al. (2009), 6: Vierke et al. (2013), 7: López-Fontán et al. (2005), 8: Wang et al. (2011), 9: Goss (2008), 10: Ahrens et al. (2012), 11: Nielsen (2012), 12: EFSA Scientific Committee (2008), 13: Ding and Peijnenburg (2013), 14: Arp et al. (2006), 15: Rayne and Forest (2009b), 16: Jing et al. (2009), 17: Higgins and Luthy (2006), 18: Stevens and Coryell (2007), 19: Bhattacharai and Gramatica (2011), 20: Beesoon and Martin (2015), 21: Han et al. (2003), 22: Hebert and MacManus-Spencer (2010), 23: Chen and Guo (2009), 24: Bischel et al. (2010), 25: Wu et al. (2009), 26: Li et al. (2009), 27: Rahman et al. (2014), 28: Barron et al. (2009).

Table S3. CLA degradation rates.

Near-surface photodegradation		Biodegradation					
$K_{pho,nearsurf}$ (d ⁻¹)	Reference	Sediments		Water		Soil	
		$K_{bio, sed}$ (d ⁻¹)	Reference	$K_{bio, wat}$ (d ⁻¹)	Reference	$K_{bio, soil}$ (d ⁻¹)	Reference
0.23	Calza et al. (2013)	2.31E-03 – 2.31E-04	JRC (2003)	Negligible	Alexy et al. (2004)	< 6.93E-4 – 7.80E-3	Kodešová et al. (2016)
0.13	Gozlan and Koren (2016)			< 6.6E-03	Calza et al. (2013)		
				< 9.9E-02	Hanamoto et al. (2013)		
				3.56E-3 – 1.39E-3	Falås et al. (2013)		

Table S4. SMX degradation rates. (an) for anaerobic or anoxic conditions.

Near-surface photodegradation		Biodegradation					
$K_{pho,nearsurf}$ (d ⁻¹)	Reference	Sediments		Water		Soil	
		$K_{bio, sed}$ (d ⁻¹)	Reference	$K_{bio, wat}$ (d ⁻¹)	Reference	$K_{bio, soil}$ (d ⁻¹)	Reference.
4.62 – 5.33	Baena-Nogueras et al. (2017)	6.86E-2 – 0.14	Lai and Hou (2008)	8.66E-3	Poirier-Larabie et al. (2016)	6.08E-02 – 7.7E-02, 3.79E-2 – 4.53E-2 (an)	Lin and Gan (2011)
1.28E-2	Poirier-Larabie et al. (2016)	0.63	Radke et al. (2009)	Negligible	Baena-Nogueras et al. (2017)	4.5E-03 – 0.15	Kodešová et al. (2016)
0.71 – 4.33	Andreozzi et al. (2003)	4.15E-02 (an)	Zhang et al. (2013)	1.24E-2 – 1.38E-2	Adamek et al. (2016)	0.2 – 0.73	Li et al. (2015b)
11.55, 2.77 – 5.78	Lam and Mabury (2005)	2.04E-02	Li et al. (2015c)	2.54E-3	Radke et al. (2009)	0.35	Liu et al. (2010)
9.90 – 17.33	Ryan et al. (2011)	3.65E-2	Lam et al. (2004)	Negligible	Al-Ahmad et al. (1999)		
23.1 – 2.89	Niu et al. (2013)	3.09E-2 – 5.42E-02	Xu et al. (2011)	6.08E-2	Zhang et al. (2013)		
7.22E-02 – 0.75	Boreen et al. (2004)	4.53E-03	Su et al. (2016)				
2.39 – 6.93	Bahn Müller et al. (2014)						
2.17 – 2.77	Oliveira et al. (2019)						
0.39 – 1.73	Bonvin et al. (2013)						
4.08	Willach et al. (2018)						

Table S5. DCF degradation rates. (an) for anaerobic or anoxic conditions.

Near-surface photodegradation		Biodegradation					
$K_{pho,nearsurf}$ (d ⁻¹)	Reference	Sediments		Water		Soil	
		$K_{bio, sed}$ (d ⁻¹)	Reference	$K_{bio, wat}$ (d ⁻¹)	Reference	$K_{bio, soil}$ (d ⁻¹)	Reference
17.33	Buser et al. (1998)	3.73E-03 – 0.13	Kunkel and Radke (2008)	Negligible	Baena-Nogueras et al. (2017)	2.34E-2 – 0.14	Lin and Gan (2011)
86.64	Poiger et al. (2001)	3.45E-02, 1.54E-2 – 1.84E-02 (an)	Koumaki et al. (2017)	4.08E-3	Buser et al. (1998)	3.39E-2 – 0.23	Xu et al. (2009)
1.73	Poirier-Larabie et al. (2016)	2.39E-02 – 7.7E-02	Radke and Maier (2014)	9.9E-03	Poirier-Larabie et al. (2016)	1.73	Grossberger et al. (2014)
99.02 – 115.52	Baena-Nogueras et al. (2017)	8E-3 – 2.6E-2	Gröning et al. (2007)				
34.66	Packer et al. (2003)						
0.35 – 3.47	Andreozzi et al. (2003)						
2.31 – 23.1	Koumaki et al. (2015)						
231.05	Zhang et al. (2017a)						
266.59 – 277.26	Zhang et al. (2011)						
6.93	Vochezer (2010)						
11.55	Radke et al. (2010)						

Table S6. IBU degradation rates. (an) for anaerobic or anoxic conditions.

Near-surface photodegradation		Biodegradation					
$K_{pho,nearsurf}$ (d ⁻¹)	Reference	Sediments		Water		Soil	
		$K_{bio, sed}$ (d ⁻¹)	Reference	$K_{bio, wat}$ (d ⁻¹)	Reference	$K_{bio, soil}$ (d ⁻¹)	Reference
1.68E-03 – 2.77E-02	Yamamoto et al. (2009)	3.65E-02 – 9.9E-02, 1.27E-03 – 3.35E-03 (an)	Conkle et al. (2012)	1.2E-02	Baena-Nogueras et al. (2017)	4.56E-02 – 6.66E-02	Lin and Gan (2011)
7.97E-02	Fono et al. (2006)	0.43, 9.52E-03 – 2.11E-02 (an)	Koumaki et al. (2017)	3.47E-02	Buser et al. (1999)	0.11 – 0.76	Xu et al. (2009)
5.13E-02	Baena-Nogueras et al. (2017)	0.14 – 0.28	Kunkel and Radke (2008)	3.47E-02 – 3.69E-02	Yamamoto et al. (2009)	2.28E-02 – 4.06E-04	Carr et al. (2011)
0.58 – 0.72	Packer et al. (2003)	8.56E-02 – 2.26, 1.47e-02	Radke and Maier (2014)			0.86	Grossberger et al. (2014)
6.30	Peuravuori and Pihlaja (2009)	> 0.12					
1.81E-02	Koumaki et al. (2015)	0.36 – 0.39	Li et al. (2015c)				
1.12	Lin and Reinhard (2005)						
6.93	Jakimska et al. (2014)						

Table S7. PAR degradation rates.

Near-surface photodegradation			Biodegradation					
			Sediments		Water		Soil	
K _{pho,nearsurf} (d ⁻¹)	Reference	K _{bio,sed} (d ⁻¹)	Reference	K _{bio,wat} (d ⁻¹)	Reference	K _{bio,soil} (d ⁻¹)	Reference	
0.58 – 17.32	Baena-Nogueras et al. (2017)	0.77	Lam et al. (2004)	1.2E-02 – 0.33	Yamamoto et al. (2009)	>0.69	Li et al. (2014)	
0.33	Carlos et al. (2012)	0.31	Lin et al. (2010)	2.47E-02	Baena-Nogueras et al. (2017)	3.65 – 3.85	Li et al. (2015b)	
0.3 – 0.46	Yamamoto et al. (2009)	0.22	Löffler et al. (2005)					
Negligible	Kawabata et al. (2013)							
2.69E-02	Peuravuori (2012)							
4.95E-02								
– 0.69	De Laurentiis et al. (2014)							
0.95								
2.47	Li et al. (2017b)							

Table S8. CBZ degradation rates. (an) for anaerobic or anoxic conditions.

Near-surface photodegradation		Biodegradation					
		Sediments		Water		Soil	
K _{pho,nearsurf} (d ⁻¹)	Reference	K _{bio,sed} (d ⁻¹)	Reference	K _{bio,wat} (d ⁻¹)	Reference	K _{bio,soil} (d ⁻¹)	Reference
		2.63E-03 – 4.2E-03,					
0.17	Calza et al. (2013)	1.58E-03 – 2.47E-03 (an)	Conkle et al. (2012)	2.97E-03 – 5.55E-03	Yamamoto et al. (2009)	4E-03 – 1.5E-02	Li et al. (2013)
6.93E-03 – 3.47E-02	Andreozzi et al. (2013)		Radke and Maier (2014)		Baena-Nogueras et al. (2017)	< 6.93E-04 – 5.37E-03	Kodešová et al. (2016)
7.92E-03 – 0.2	Yamamoto et al. (2009)	Negl – 1.2E-02	Löffler et al. (2005)	Negligible	Durán-Álvarez et al. (2015)	< 3.47E-03 – 4.28E-03	Grossberger et al. (2014)
Negligible – 1.39	Baena-Nogueras et al. (2017)		Li et al. (2015c)				
0.34	Carlos et al. (2012)	Negl		< 6.6E-03	Calza et al. (2013)		
0.22 – 0.77		8.45E-03	Lam et al. (2004)				
0.16	Dong et al. (2015)						
1.71E-02	Durán-Álvarez et al. (2015)						
0.87	Peuravuori and Pihlaja (2009)						
0.14 – 2.77	Doll and Frimmel (2003)						
1.26E-02	Lam and Mabury (2005)						
	De Laurentiis et al. (2012)						

Table S9. FUR degradation rates.

Near-surface photodegradation		Biodegradation					
		Sediments		Water		Soil	
K _{pho,nearsurf} (d ⁻¹)	Reference	K _{bio, sed} (d ⁻¹)	Reference	K _{bio, wat} (d ⁻¹)	Reference	K _{bio, soil} (d ⁻¹)	Reference
86.64	Hanamoto et al. (2013)	4.33E-02	Li et al. (2015c)	< 0.1	Hanamoto et al. (2013)	4.91E-03	Polesel et al. (2015b)
		4.62E-02	Radke and Maier (2014)			1.65E-02	
69.31	Yagi et al. (1991)	3.47E-02	–				
		4.95E-02		8.11E-04*	Khan and Ongerth (2004)		
	Jakimska et al. (2014)	1.33E-02	–				
34.66		4.62E-02	Fenner et al. (2016)				

*: calculated correcting degradation rate with canal TSS

Table S10. E2 degradation rates. (an) for anaerobic or anoxic conditions.

Near-surface photodegradation		Biodegradation					
		Sediments		Water		Soil	
K _{pho,nearsurf} (d ⁻¹)	Reference	K _{bio,sed} (d ⁻¹)	Reference	K _{bio,wat} (d ⁻¹)	Reference	K _{bio,soil} (d ⁻¹)	Reference
		5.1E-02 – 5.33E-02,					
2.77 – 17.32	Liu et al. (2017)	1.75E-02 – 2.6E-02 (an)	Robinson et al. (2017)	~ 1E-03	Bradley et al. (2009)	6.3	Colucci and Topp (2002)
2.17 – 2.77	Leech et al. (2009)	2.57	Mashtare et al. (2013)	7.97E-02 – 3.47	Jurgens et al. (2002)	0.17 – 0.46	Carr et al. (2011)
	Chowdhury et al. (2011)		Czajka and Londry (2006)	0.12 – 0.23	Liu et al. (2011)	0.46 – 3.15	Colucci et al. (2001)
2.77 – 2.77	Whidbey et al. (2012)	3.3E-02					
	Lin and Reinhard (2005)	6.3, 1.05 – 1.87 (an)	Jurgens et al. (2002)				
8.66	Jurgens et al. (2002)	>1E-02	Bradley et al. (2009)				
0.13		1.87 – 7.7	Zhang et al. (2016)				
0.83 – 1.24	Li et al. (2017a)						
2.89 – 5.78	Silva et al. (2016b)						

Table S11. EE2 degradation rates.

Near-surface photodegradation		Biodegradation					
		Sediments		Water		Soil	
$K_{pho,nearsurf}$ (d ⁻¹)	Reference	$K_{bio, sed}$ (d ⁻¹)	Reference	$K_{bio, wat}$ (d ⁻¹)	Reference	$K_{bio, soil}$ (d ⁻¹)	Reference
0.16	Matamoros et al. (2009)	6.42E-03	Wu et al. (2015a)	Negligible	Wu et al. (2015a)	0.83	Colucci and Topp (2002)
3.85	Ren et al. (2016)	0.16	Mashtare et al. (2013)	4.08E-02	Jurgens et al. (2002)	3.34E-03 – 3.07E-02	Carr et al. (2011)
0.72	Zuo et al. (2013)	Negligible	Czajka and Londry (2006)	6.42E-03	Zuo et al. (2013)	2.15E-02 – 0.37	Colucci and Topp (2001)
3.46	Whidbey et al. (2012)			< 4.95E-02	Matsuoka et al. (2005)		
6.93	Lin and Reinhard (2005)			Negligible	Liu et al. (2011)		
0.13	Jurgens et al. (2002)						
0.73 -3.46	Ren et al. (2017a)						
2.4E-02 – 0.5	Atkinson et al. (2011)						
0.89 – 7.7	Silva et al. (2016b)						
0.77	Grzybowski and Szydlowski (2014)						
0.63 – 0.69	Ren et al. (2017b)						
1.39	Wu et al. (2015a)						

Table S12. E1 degradation rates. (an) for anaerobic or anoxic conditions.

Near-surface photodegradation		Biodegradation					
		Sediments		Water		Soil	
$K_{pho,nearsurf}$ (d ⁻¹)	Reference	$K_{bio, sed}$ (d ⁻¹)	Reference	$K_{bio, wat}$ (d ⁻¹)	Reference	$K_{bio, soil}$ (d ⁻¹)	Reference
2.31	Atkinson et al. (2011)	6.8E-02 – 0.24, 3.14E-02 – 4.41E-02 (an)	Robinson et al. (2017)	6.36E-02 – 6.93	Jurgens et al. (2002)	1.22E-02 – 0.22	Carr et al. (2011)
4.33 – 7.7	Silva et al. (2016a)	1.93E-02	Mashtare et al. (2013)	0.17 – 3.47	de Mes et al. (2005)	0.41 – 1.14	Colucci et al. (2001)
2.48 – 4.33	Caupos et al. (2011)	1.65, 4.88E-02 – 6.03E-02 (an)	Jurgens et al. (2002)				
13.86 – 34.66	Chowdhury et al. (2011)	>1E-02	Bradley et al. (2009)				
2.48 – 4.33	Lin and Reinhard (2005)						

*: calculated correcting degradation rate with canal TSS

Table S13. PFOA degradation rates.

Near-surface photodegradation		Biodegradation					
		Sediments		Water		Soil	
$K_{pho,nearsurf}$ (d ⁻¹)	Reference	$K_{bio, sed}$ (d ⁻¹)	Reference	$K_{bio, wat}$ (d ⁻¹)	Reference	$K_{bio, soil}$ (d ⁻¹)	Reference
4.22E-04	Vaalgaama et al. (2011)	Negligible	Liou et al. (2010)	Negligible	Liou et al. (2010)	Negligible	Liou et al. (2010)
<1.98E-03	Giesy et al. (2010)						

Table S14. PFOS degradation rates.

Near-surface photodegradation		Biodegradation					
		Sediments		Water		Soil	
$K_{pho,nearsurf}$ (d ⁻¹)	Reference	$K_{bio, sed}$ (d ⁻¹)	Reference	$K_{bio, wat}$ (d ⁻¹)	Reference	$K_{bio, soil}$ (d ⁻¹)	Reference
1642	Brooke et al. (2004)	Negligible	Avendano et al. (2015)	Negligible	Avendano et al. (2015)	Negligible	Avendano et al. (2015)

Table S15. TCS degradation rates.

Near-surface photodegradation		Biodegradation					
		Sediments		Water		Soil	
$K_{pho,nearsurf}$ (d ⁻¹)	Reference	$K_{bio, sed}$ (d ⁻¹)	Reference	$K_{bio, wat}$ (d ⁻¹)	Reference	$K_{bio, soil}$ (d ⁻¹)	Reference
23.1	Baena-Nogueras et al. (2017)	7.2E-03	Wu et al. (2015a)	Negligible	Wu et al. (2015a)	3.85E-02	Wu et al. (2015a)
115.52		6.36E-02,					
23.1	Koumaki et al. (2015)	Negligible (an)	Koumaki et al. (2018)	1.92E-02	Baena-Nogueras et al. (2017)	4.42E-02 – 5.48E-02	Xu et al. (2009)
3.15	Durán-Álvarez et al. (2015)	1.22E-02 – 1.29E-02	NICNAS (2009)	0.69E-02 – 2.1E-02	Durán-Álvarez et al. (2015)	9.78E-03 – 1.74E-03	Carr et al. (2011)
3.3	Latch et al. (2005)					1.2E-02 – 2.17E-02	Wu et al. (2009)
69.31	Lindström et al. (2002)						
23.1	Singer et al. (2002)						
34.66							
69.31	Tixier et al. (2002)						
4.62	Mezcua et al. (2004)						
3.01	Martínez-Zapata et al. (2013)						
231.04	Sanchez-Prado et al. (2006)						
0.2	Wu et al. (2015a)						

SI.3 Model parameters and model inputs

The environmental conditions used in RWQM1s1_MP are shown in Table S16. All values were taken from ARPA (Agenzia Regionale Protezione Ambiente) measurements databases for the year 2016. Regarding the CSPM, the model layers were adapted to the stratigraphy of the various soils with the parameters shown in Table S17. Photodegradation in rice paddies was assumed to be negligible due the sediment in suspension and plants shading. Eventual surface runoff of water was not modeled.

Table S16. Measurements and parameters used to characterize environmental conditions for the RWQM1s1_MP model

Parameter	Source	Notes
Wind speed at 10 m above ground	ARPA	Hourly measurements
Air temperature	ARPA	Hourly measurements
Sunlight intensity	ARPA	Hourly measurements
Water temperature	ARPA	Monthly measurements at end of canal
pH	ARPA	Canal measurements average ± 1 daily fluctuation (Simonsen and Harremoës, 1978)
TWW entering the canal	Mazzini et al. (2013)	Daily flow variations and rain events neglected
TWW CECs concentrations	Castiglioni et al. (2018a, 2018b)	Converted dissolved to total concentrations thanks to equations in Chapra (1997)
Algae, heterotrophs and nitrifying bacteria concentrations		Initially left as default and then calibrated
Algae and bacterial growth parameters	Vanrollenghem et al. (2001)	
Phosphorous sorption and desorption	Chapra (1997)	

Table S17. Environmental parameters and sources used for the CSPM. LOSAN is a publicly available soil database provided by ERSAF (Ente Regionale per i Servizi all'Agricoltura e alle Foreste), CRA-CMA stands for Unità di Ricerca per la Climatologia e la Meteorologia applicate all'Agricoltura

Parameter	Source	Notes
Soil organic carbon	LOSAN	
Soil layer pH	LOSAN	
FC and PWP	LOSAN	Estimated from granulometries using correlations by Rawls et al. (1982)
Soil layers bulk density	LOSAN	Estimated from horizon-specific pedo-transfer functions from SOILQUALIMON report (Brenna et al., 2010)
Air temperature	ARPA	
Rainfall	ARPA	
Relative humidity	ARPA	
Potential evaporation	CRA-CMA	Obtained thanks to correction of potential evapotranspiration using $K_{c,ini}$ (1.05 rice, 0.3 other crops) (Legind et al., 2012; Polesel et al., 2015; Allen et al., 1998)
Soil layer temperature	CRA-CMA	Extrapolated from measurements at depths of 10 and 50 cm thanks to the sinusoidal functions in Xing and Spitler (2017).

SI.3.1 Soil properties

Table S18. Modeled soil properties.

Horizon bottom depth (cm)	pH	Organic carbon (%)	Field capacity (L/L)	Wilting point (L/L)	Bulk density (g/cm ³)
Soil 1: silage maize and ryegrass					
13	7.3	4.9	0.36	0.14	1.09
38	6.9	0.64	0.23	0.10	1.52
60	7.8	0.64	0.31	0.13	1.32
75	7.6	0.5	0.38	0.21	1.80
100	7.5	0.37	0.32	0.14	1.59
Soil 2: silage maize and ryegrass, winter wheat					
20	6.6	1.27	0.24	0.09	1.45
40	6.6	1.27	0.24	0.09	1.45
60	6.7	0.16	0.18	0.09	1.48
80	6.7	0.16	0.18	0.09	1.48
100	6.7	0.16	0.18	0.09	1.48
Soil 3: rice and ryegrass, winter wheat					
35	6.1	2	0.26	0.11	1.33
45	6.9	0.72	0.21	0.08	1.49
67.5	7.2	0.63	0.16	0.06	1.41
90	7.2	0.63	0.16	0.06	1.41
100	7.4	0.44	0.23	0.12	1.49

Pseudo-steady state concentrations of the layers in each modeled soil were obtained with median concentration in TWW as irrigation concentration and median soil degradation rate and used as initial concentrations. For the crop rotation consisting of rice and ryegrass and winter wheat, the soil concentrations at the end of the rice and ryegrass years were used as represented the worst case soil concentration.

SI.3.2 Plant parameters

Table S19. Parameters used for crop modeling. Parameters not noted were left as default.

	Value	Reference	Notes
Maize			
Irrigation	400 mm/season	Borrelli et al. (2014)	
Germination date	20-05	Borrelli et al. (2014)	
Harvest date	30-09	Borrelli et al. (2014)	
Transpiration coefficient	60 L/kg fw	Legind et al. (2012)	
Water content roots	0.84 g/g	Wang et al. (1991)	
Final root mass	2.58 kg fw/m ²	Legind et al. (2012)	
Growth rate	0.081 1/d	Legind et al. (2012)	
Final stem mass	3.61 kg fw/m ²	Legind et al. (2012)	
Final leaf mass	0.96 kg fw/m ²	Legind et al. (2012)	
Final fruit mass	1.78 kg fw/m ²	Legind et al. (2012)	
Water content fruit	0.76	USDA (2018)	
Ryegrass			
Irrigation	210 mm/season	FAO (2007)	Calculated subtracting rainfall from water requirements
Germination date	9-12 with silage maize – 30/10 with rice	Borrelli et al. (2014), Nelli and Sodi (2007)	
Harvest date	14-05 with silage maize – 31/03 with rice	Borrelli et al. (2014), Nelli and Sodi (2007)	
Final root mass	2.35 kg fw/m ²		Assumed equal to winter wheat since same family (Poaceae)
Final stem mass	1.8 kg fw/m ²	Tabaglio et al. (2007), Abraha and Savage (2008)	Calculated assuming water content = 0.8
Final leaf mass	2.2 kg fw/m ²	Tabaglio et al. (2007), Abraha and Savage (2008)	Calculated assuming water content = 0.8
Initial fruit mass	1E-05 kg fw/m ²	Martiniello (1999)	To halt fruit development

Table S19. Cont.

Winter wheat			
Irrigation	100 mm/season	Polesel et al. (2015)	
Germination date	1-03	Legind et al. (2012)	
Harvest date	15-07	Legind et al. (2012)	
Final root mass	2.35 kg fw/m ²	Legind et al. (2012)	
Final stem mass	0.78 kg fw/m ²	Legind et al. (2012)	
Final leaf mass	1.81 kg fw/m ²	Legind et al. (2012)	
Final fruit mass	1.45 kg fw/m ²	Legind et al. (2012)	
Rice			
Irrigation	21.6 mm/day	Nelli and Sodi (2007)	
Germination date	6-04	Nelli and Sodi (2007)	
Harvest date	30-09	Nelli and Sodi (2007)	
Transpiration coefficient	52	Van der Vorm (1980)	
Roots/Stem/Leaf growth rate	0.068	Confalonieri and Bocchi (2005)	
Fruit growth rate	0.13		Doubled compared to plant growth rate similarly to what observed for winter wheat and to match model ripening with agronomic data
Final root mass	3.30	Confalonieri and Bocchi (2005)	
Final stem mass	1.76	Confalonieri and Bocchi (2005)	
Final leaf mass	1.92	Confalonieri and Bocchi (2005)	
Final fruit mass	0.94	Confalonieri and Bocchi (2005)	
Water content root	0.95	Zhang and Webster (2002); Henry et al. (2012)	Calculated based on fresh weight measurements and roots to shoots ratio
Water content fruit	0.23	Rice Knowledge Bank (2017)	

The ratio of phloem to xylem flux for PAR, SMX, IBU and FUR was set to 0.05, a median value between day and night. For the other CECs such ratio was set to 0 as considered non phloem-systemic based on the distinction by Kleier and Hsu (1996) relying on CECs Kow and pK_a. The phloem systemicity of DCF and PFOA was treated as uncertain as potentially affected by plant-specific parameters.

SI.4 Uncertainty propagation

The preliminary uncertainty analysis (UA) of the IUWS_MP model for traditional pollutants was carried out varying the parameters in Table S20. Parameters uncertainty was based on Reichert and Vanrollenghem (2001) considering two classes of confidence: poorly known parameters were varied by 50% (Class 2), while better known parameters were varied by 20% (Class 1). CBZ, DCF and TCS, CECs with different physicochemical characteristics, were chosen amongst the thirteen selected to be tested. Random parameters combinations were obtained through 1000 Latin hypercube samplings for each CEC.

Table S20. Uncertainty classes of parameters for the uncertainty analysis of the traditional pollutants submodel.

Parameter	Class	Parameter	Class
Heterotrophic bacteria input concentration	2	First stage nitrifiers (AOBs) input concentration	2
Phosphate adsorption rate	2	Algae input concentration	2
Saturation coefficient for aerobic growth of heterotrophs on dissolved organic substrate	2	Mean particle diameter	2
Slope	1	Saturation coefficient for growth of AOBs on ammonia	2
Maximum aerobic specific growth rate for heterotrophic biomass	1	Manning n value	1
Top canal width	1	Maximum aerobic specific growth rate for AOBs	1
Depth of canal filled to bank	1		

Successively, the uncertainty regarding the organic micropollutants IUWS_MP submodel and CSPM parameters on CECs concentrations were assessed through a Monte Carlo UA varying the parameters in Table S21. When no anoxic sediment biodegradation rates were found in literature, the aerobic values decreased by an order of magnitude were used. In case only one or two values were found, uniform distributions were used considering either a 10% variation or the range between the two values. The uncertainty of IUWS_MP was estimated thanks to 1000 runs for each water water withdraw scenario. The choice of the number of CSPM runs was based on the comparison of the percentiles from simulations consisting of an increasing amount of runs to the ones resulting from a 5000 runs simulation. As the percentiles from a 2000 runs simulation with Latin hypercube sampling differed negligibly from the 5000 runs simulation, the mentioned settings were chosen for the remaining simulations.

Table S21. Varied parameters during the IUWS_MP and CSPM Monte Carlo uncertainty analyses

Parameter	Distribution	Source	Notes
TWW CEC concentrations	Triangular	Castiglioni et al. (2018a, 2018b)	Inf: minimum value/LOD Mode: median value/0.5*LOQ Sup: maximum value/LOQ
Water aerobic biodegr. rate	Empirical, triangular	Tables S4 – S16	Choice based on the quantity of experimental values found. Values outside 1.5*IQR (interquartile range) considered as outliers. Triangular distributions based on extreme values and medians
Sediment aerobic biodegr. rate			
Sediment anoxic biodegr. rate			
Soil dissipation rate			
Photodegradation rate			
pK _a	Empirical, triangular	Table S3	Choice based on the quantity of experimental values found
K _{ow}			
Solubility			
Vapor pressure			
K _{HSA}			
K _{aw}			
(only perfluorinated chemicals)			
Irrigation concentration	Triangular	Surface water model	Inf: 2.5 th percentile canal conc. Mode: modal canal conc. Sup: 97.5 th percentile canal conc.
pH in cytosol	Uniform	Felle (2001), Felle et al.	10% variation is comparable to observed fluctuations
pH in vacuole		(2005), Johannes et al.	
pH in phloem		(2001), Tournaire-Roux et al. (2003)	
pH in xylem			
Transpiration coeff.	Uniform		Variation equal to 10% to avoid numerical errors
Root mass			
Root growth rate			
Stem growth rate			
Leaves growth rate			
Fruits growth rate	Lognormal	EFSA (2007)	Median: 1.099 95 th percentile: 1.144
Arrhenius temperature correction coeff.			

Other parameters were varied during the analysis of specific CECs to either prevent numerical errors (SMX, PAR, E2), include uncertainty (PFOA, DCF) and improve model results (CLA, PFOS):

- PFOA and DCF phloem flux was set in 50% of the runs as 0;
- CLA's K_{oc} was set to 10176 L/kg to match the value from the equations in Franco and Trapp (2008);
- SMX and PFOS K_{OC} was varied as the other physicochemical properties;

- The soil structure during PAR simulations was modified to 3 layers of 33 cm whose soil properties were calculated as weighted average of the original layers. The dissipation rate was, also, set to the minimum value and the Koc from ECHA (2006) was used;
- The dissipation rate of E2 was limited to 1 d⁻¹.

Successively, the monthly medians of the total CECs concentrations taken at four locations along the canal were fitted with a first-order dissipation rate thanks to the use of Matlab R2016a and the estimated traveling times provided by IUWS_MP model. Yearly best- and worst-case attenuation rates were calculated for all the modeled CECs and compared to literature as a further validation.

Such attenuation rates and influent CECs concentrations (Castiglioni et al., 2018a) were used to estimate the expected concentrations along the canal in case of the absence of the upstream WWTP. Best- and worst-case scenarios were calculated respectively using low concentrations and high rates, and high concentrations and low rates. The median scenarios were obtained thanks to the median concentrations and the mean between the two attenuation rates.

The fruit CECs concentrations in case of the absence of the WWTP were estimated thanks to the expected median concentrations at the end of the canal which were used to reach a pseudo-steady state in the coupled soil-plant model. Successively, the median CECs concentration of the first layer at pseudo-steady state was multiplied by the median or the 97.5th percentile BCFs, calculated from the previous Monte Carlo uncertainty analyses, to estimate of the median and 97.5th percentile fruit concentration.

SI.5 Sensitivity analyses (SA)

Figure S2. Deviation caused by increments of the parameters in the traditional pollutants submodel. S_MP, S_MP_DOC and X_MP indicate respectively the following fractions of a CEC: dissolved, sorbed on colloids and sorbed on TSS.

	C fruit			C leaves			C stem			C roots			C soil, L1			C soil, L2			C soil, L3			C soil, L4			C soil, L5			Leaching		
	CBZ	DCF	TCS	CBZ	DCF	TCS	CBZ	DCF	TCS	CBZ	DCF	TCS	CBZ	DCF	TCS	CBZ	DCF	TCS	CBZ	DCF	TCS	CBZ	DCF	TCS	CBZ	DCF	TCS	CBZ	DCF	TCS
Temp correction coeff			1			1			1			1			1			1			1			1			1			1
Initial water content L2-5																														
Soil field capacity																														
Soil pH																														
pH inside cytosol																														
pH inside phloem																														
pH inside xylem																														
Final root mass																														
Fruit growth rate																														
Leaf growth rate																														
Root growth rate																														
Soil organic carbon																														
Transpiration coeff																														
Water % stem and leaves																														
Ratio Phloem to xylem flux																														
Water % roots																														
Layers thickness																														
Soil bulk density																														

Figure S3. Coupled soil-plant model sensitive parameters in presence or absence of degradation.

SI.6 Fate analysis

SI.6.1 Traditional pollutants

Figure S4. Traditional pollutants calibrated concentrations and measured values in canal. NH = ammonium, NO2 = nitrite, NO3 = nitrate, Sol P = soluble phosphorus, Part P = particulate phosphorus.

SI.6.2 Uncertainty of CECs concentrations in canal

Figure S5. PFOS midday predicted concentration ranges. Days starting from January 1st until December 31st.

Figure S6. TCS midday predicted concentration ranges. Days starting from January 1st until December 31st.

Figure S7. CBZ midday predicted concentration ranges. Days starting from January 1st until December 31st.

Figure S8. E1 midday predicted concentration ranges. Days starting from January 1st until December 31st.

Figure S9. EE2 midday predicted concentration ranges. Days starting from January 1st until December 31st.

Figure S10. IBU midday predicted concentration ranges. Days starting from January 1st until December 31st.

Figure S11. PAR midday predicted concentration ranges. Days starting from January 1st until December 31st.

Figure S12. PFOA midday predicted concentration ranges. Days starting from January 1st until December 31st.

Figure S13. SMX midday predicted concentration ranges. Days starting from January 1st until December 31st.

Figure S14. E2 midday predicted concentration ranges. Days starting from January 1st until December 31st.

SI.6.3 Attenuation along the canal

Figure S15. Predicted median attenuation rates compared to attenuation rates from literature. References: Acuña et al. (2015), Bester (2005), Fono et al. (2006), Guillet et al. (2019), Kunkel and Radke (2011), Lin et al. (2006), Morrall et al. (2004), Matamoros and Rodríguez (2017), Radke et al. (2010), Sabaliunas et al. (2003), Tixier et al. (2003), Williams et al. (2003), Writer et al. (2012, 2013).

SI.6.4 Bioconcentration factors

Figure S16. Fruit BCFs (gdw/gdw) for silage maize, winter wheat, ryegrass and rice with respect to maximum soil concentrations and literature (lit.) values. All simulated soils are shown. BioS indicates experimental studies conducted with the use of biosolids. Red borders indicate the use of an LOD value in the calculation of the BCF. References (including fruit BCFs in Figure 3): Blaine et al. (2013, 2014), Christou et al. (2017), Goldstein et al. (2014), Lechner and Knapp (2011), Mordechay et al. (2018), Navarro et al. (2017), Pannu et al. (2012), Prosser et al. (2014), Sabourin et al. (2012), Shenker et al. (2011), Stahl et al. (2009), Wen et al. (2014), Wu et al. (2010, 2012), Yager et al. (2014)

Figure S17. Leaf BCFs (gdw/gdw) for silage maize, winter wheat, ryegrass and rice with respect to maximum soil concentrations and literature (lit.) values. All simulated soils are shown. BioS indicates experimental studies conducted with the use of biosolids. Red borders indicate the use of an LOD value in the calculation of the BCF. References (including leaf BCFs in Figure 3): Aryal and Reinhold (2011), Bizkarguenaga et al. (2016), Blaine et al. (2013, 2014), Carter et al. (2014), Chitescu et al. (2012), Fu et al. (2016), Goldstein et al. (2014), Holling et al. (2012), Hurtado et al. (2017), Karnjanapiboonwong et al. (2011), Lechner and Knapp (2011), Macherius et al. (2012), Malchi et al. (2014), Mordechay et al. (2018), Navarro et al. (2017), Pannu et al. (2012), Prosser et al. (2014), Shenker et al. (2011), Stahl et al. (2009), Wen et al. (2014, 2016), Winker et al. (2010), Wu et al. (2010, 2012), Yoo et al. (2011), Zhao et al. (2014, 2017).

Figure S18. Root BCFs (g_{dw}/g_{dw}) for silage maize, winter wheat, ryegrass and rice with respect to maximum (a) or median (b) soil concentrations and literature (lit.) values. All simulated soils are shown. BioS indicates experimental studies conducted with the use of biosolids. Red borders indicate the use of an LOD value in the calculation of the BCF. References: Aryal and Reinhold (2011), Bizkarguenaga et al. (2016), Blaine et al. (2014), Carter et al. (2014), Fu et al. (2016), Holling et al. (2012), Hurtado et al. (2017), Karnjanapiboonwong et al. (2011), Lechner and Knapp (2011), Macherius et al. (2012), Malchi et al. (2014), Pannu et al. (2012), Prosser et al. (2014), Sabourin et al. (2012), Shenker et al. (2011), Wen et al. (2014, 2016), Winker et al. (2010), Wu et al. (2010, 2012), Yager et al. (2014), Zhao et al. (2014, 2017).

Figure S19. Stem BCFs (g_{dw}/g_{dw}) for silage maize, winter wheat, ryegrass and rice with respect to maximum (a) or median (b) soil concentrations and literature (lit.) values. All simulated soils are shown. BioS indicates experimental studies conducted with the use of biosolids. References: Aryal and Reinhold (2011), Blaine et al. (2013, 2014), Carter et al. (2014), Chitescu et al. (2012), Lechner and Knapp (2011), Navarro et al. (2017), Prosser et al. (2014), Shenker et al. (2011), Stahl et al. (2009), Wen et al. (2014, 2016), Winker et al. (2010), Wu et al. (2010), Yager et al. (2014), Yoo et al. (2011), Zhao et al. (2014, 2017).

Figure S20. Total leached mass during each crop growing season.

SI.7 Risk assessment

SI.7.1 Environmental risk – Results

Table S22. Maximum, median and minimum RQs at the end of the canal and monthly frequency to exceed a given RQ.

CEC	Max/Median/Min RQ	Monthly frequency of exceedance (%)											
		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
RQ above 0.1													
SMX	0.42/0.16/0.018	74.1	73.9	73.5	73.5	73.4	72.7	72.4	72.6	73.7	73.9	74.2	74.1
IBU	0.15/0.07/0.021	16.9	16.7	16.5	16.4	16.3	15.9	15.8	15.9	16.2	16.4	16.7	16.8
E1	0.25/0.14/0.046	91.7	91.5	91.3	90.8	90.5	89.2	88.4	88.8	90.6	91.2	91.5	91.7
RQ above 1													
CLA	2.63/1.34/0.48	80	79.8	79.7	79.4	79	78.3	77.9	78	78.9	79.3	80	80.4
EE2	45.88/26.74/12.86	100	100	100	100	100	100	100	100	100	100	100	100
E2	2.57/1.48/0.65	95.8	95.6	95.4	95.1	94.9	94.1	93.6	93.8	95	95.3	95.7	95.8
TCS	2.56/1.43/0.44	77.9	77.7	77.6	77.3	77.1	76.5	76.5	76.6	77.3	77.6	77.7	77.9

SI.7.2 Human health risk – Methods

Following the precautionary principle, human health risk was calculated with the assumption of no degradation between harvest and consumption and the complete bioaccessibility and bioavailability of the ingested CECs.

To calculate the ingestion required to exceed the threshold of toxicological concern (*TTC*), *TTC* was set equal to 30, 9.1, 1.5 and 0.0025 $\mu\text{g kg}_{\text{bw}}^{-1} \text{d}^{-1}$, depending on the Cramer classes (I, II and III or genotoxic alerts) which the CEC belongs to (Kroes et al., 2004; Malchi et al., 2014).

Regarding the calculation of the hazard quotient (HQ), the ADI values were either found in Prosser and Sibley (2015) or in the reports produced by EFSA (2008), Environment Protection and Heritage Council of Australia (2008) and EMEA (2000) and, eventually, corrected as reported in Malchi et al. (2015).

SI.7.3 Human health risk – Results

Table S23. Consumption (kg/d) required to exceed *TTC* for crops irrigated with water withdrawn at the end of the canal. P97.5 stands for values calculated with the 97.5th percentile CEC concentration.

CEC	Infants				Adults			
	Rice		Wheat		Rice		Wheat	
	Median	P97.5	Median	P97.5	Median	P97.5	Median	P97.5
SMX	0.07	0.01	1.92	0.23	0.38	0.05	11.2	1.32
DCF	> 100	73.51	> 100	> 100	> 100	> 100	> 100	> 100
IBU	> 100	> 100	> 100	> 100	> 100	> 100	> 100	> 100
PAR	3.06	2.08	72.22	44.84	17.83	12.15	> 100	> 100
CBZ	8.59	5.77	19.03	15.06	50.12	33.65	> 100	87.84
FUR	> 100	> 100	> 100	> 100	> 100	> 100	> 100	> 100
PFOA	> 100	> 100	> 100	> 100	> 100	> 100	> 100	> 100
PFOS	> 100	> 100	> 100	> 100	> 100	> 100	> 100	> 100
TCS	> 100	> 100	> 100	> 100	> 100	> 100	> 100	> 100

Table S24. HQs and His for crops irrigated with water withdrawn at the end of the canal. P97.5 stands for values calculated with the 97.5th percentile CEC concentration.

CEC	Rice		Wheat		Total	
	Median	P97.5	Median	P97.5	Median	P97.5
Infants						
SMX	2.94E-03	2.03E-02	1.86E-04	1.58E-03	3.12E-03	2.19E-02
DCF	2.66E-06	6.35E-04	6.14E-08	1.26E-05	2.72E-06	6.48E-04
IBU	1.12E-09	5.27E-07	1.21E-10	5.94E-08	1.24E-09	5.86E-07
PAR	7.12E-07	1.05E-06	5.64E-08	9.08E-08	7.69E-07	1.14E-06
CBZ	2.62E-03	3.91E-03	2.21E-03	2.80E-03	4.84E-03	6.71E-03
FUR	5.37E-08	8.49E-05	1.25E-07	9.82E-05	1.79E-07	1.83E-04
EE2	2.77E-02	2.57E-01	1.95E-03	3.85E-02	2.97E-02	2.95E-01
E2	1.07E-05	2.53E-05	8.72E-07	2.02E-06	1.15E-05	2.73E-05
E1	5.23E-04	4.82E-03	4.06E-05	7.80E-04	5.63E-04	5.60E-03
PFOS	6.98E-06	2.75E-04	5.44E-06	1.24E-04	1.24E-05	4.00E-04
PFOA	2.23E-08	1.22E-04	1.17E-06	2.66E-04	1.19E-06	3.88E-04
TCS	6.19E-09	1.79E-08	2.57E-09	1.04E-08	8.76E-09	2.83E-08
Hazard Index					3.82E-02	3.31E-01
Adults						
SMX	4.45E-04	3.08E-03	8.88E-05	7.52E-04	5.34E-04	3.83E-03
DCF	4.03E-07	9.63E-05	2.93E-08	6.03E-06	4.33E-07	1.02E-04
IBU	1.70E-10	7.99E-08	2.93E-08	6.03E-06	2.27E-10	1.80E-07
PAR	1.08E-07	1.59E-07	2.69E-08	4.33E-08	1.35E-07	2.20E-07
CBZ	3.98E-04	5.93E-04	1.06E-03	1.33E-03	1.45E-03	1.93E-03
FUR	8.14E-09	1.29E-05	5.96E-08	4.69E-05	6.78E-08	5.97E-05
EE2	4.20E-03	3.89E-02	9.32E-04	1.84E-02	5.13E-03	5.73E-02
E2	1.62E-06	3.84E-06	4.16E-07	9.63E-07	2.03E-06	4.80E-06
E1	7.93E-05	7.30E-04	1.94E-05	3.72E-04	9.86E-05	1.10E-03
PFOS	1.06E-06	1.31E-04	2.60E-06	5.94E-05	3.65E-06	1.90E-04
PFOA	3.38E-09	1.85E-05	5.58E-07	1.27E-04	5.62E-07	1.46E-04
TCS	9.38E-10	2.71E-09	1.23E-09	4.96E-09	2.17E-09	7.67E-09
Hazard Index					7.22E-03	6.47E-02

SI.8 References

- Abraha, M. G., Savage, M. J. (2008). The soil water balance of rainfed and irrigated oats, Italian rye grass and rye using the CropSyst model. *Irrigation Science*, 26(3), 203–212.
- Acuña, V., von Schiller, D., García-Galán, M. J., Rodríguez-Mozaz, S., Corominas, L., Petrovic, M., ... Sabater, S. (2015). Occurrence and in-stream attenuation of wastewater-derived pharmaceuticals in Iberian rivers. *Science of the Total Environment*, 503–504, 133–141.
- Adamek, E., Baran, W., Sobczak, A. (2016). Assessment of the biodegradability of selected sulfa drugs in two polluted rivers in Poland: Effects of seasonal variations, accidental contamination, turbidity and salinity. *Journal of Hazardous Materials*, 313, 147–158.
- Ahrens, L., Harner, T., Shoeib, M., Lane, D. A., Murphy, J. G. (2012). Improved characterization of gas-particle partitioning for per- and polyfluoroalkyl substances in the atmosphere using annular diffusion denuder samplers. *Environmental Science and Technology*, 46(13), 7199–7206.
- Al-Ahmad, A., Daschner, F. D., Kümmerer, K. (1999). Biodegradability of cefotiam, ciprofloxacin, meropenem, penicillin G, and sulfamethoxazole and inhibition of waste water bacteria. *Archives of Environmental Contamination and Toxicology*, 37(2), 158–163.
- Alexy, R., Kümpe, T., Kümmerer, K. (2004). Assessment of degradation of 18 antibiotics in the Closed Bottle Test. *Chemosphere*, 57(6), 505–512.
- Allen, R., Pereira, L. S., Raes, D., Smith, M. (1998). *Crop Evapotranspiration - Guidelines for computing crop water requirements* - FAO Irrigation and Drainage paper. Rome, Italy.
- Andreozzi, R., Marotta, R., Paxéus, N. (2003). Pharmaceuticals in STP effluents and their solar photodegradation in aquatic environment. *Chemosphere*, 50(10), 1319–1330.
- Arp, H. P. H., Niederer, C., Goss, K. (2006). Predicting the Partitioning Behavior of Various Highly Fluorinated. *Environmental Science and Technology*, 40(23), 7298–7304.
- Aryal, N., Reinhold, D. M. (2011). Phytoaccumulation of antimicrobials from biosolids : Impacts on environmental fate and relevance to human exposure. *Water Research*, 45(17), 5545–5552.
- Atkinson, S. K., Marlatt, V. L., Kimpe, L. E., Lean, D. R. S., Trudeau, V. L., Blais, J. M. (2011). Environmental factors affecting ultraviolet photodegradation rates and estrogenicity of estrone and ethinylestradiol in natural waters. *Archives of Environmental Contamination and Toxicology*, 60(1), 1–7.

- Baena-Nogueras, R. M., González-Mazo, E., Lara-Martín, P. A. (2017). Degradation kinetics of pharmaceuticals and personal care products in surface waters: photolysis vs biodegradation. *Science of the Total Environment*, 590–591, 643–654.
- Bahn Müller, S., von Gunten, U., Canonica, S. (2014). Sunlight-induced transformation of sulfadiazine and sulfamethoxazole in surface waters and wastewater effluents. *Water Research*, 57, 183–192.
- Barron, L., Havel, J., Purcell, M., Szpak, M., Kelleher, B., Paull, B. (2009). Predicting sorption of pharmaceuticals and personal care products onto soil and digested sludge using artificial neural networks. *Analyst*, 134(4), 663–670.
- Beesoon, S., Martin, J. W. (2015). Isomer-Specific Binding Affinity of Perfluorooctanesulfonate (PFOS) and Perfluorooctanoate (PFOA) to Serum Proteins. *Environmental Science & Technology*, 49(9), 5722–5731.
- Ben Mordechay, E., Tarchitzky, J., Chen, Y., Shenker, M., Chefetz, B. (2018). Composted biosolids and treated wastewater as sources of pharmaceuticals and personal care products for plant uptake: A case study with carbamazepine. *Environmental Pollution*, 232, 164–172.
- Bester, K. (2005). Fate of triclosan and triclosan-methyl in sewage treatment plants and surface waters. *Archives of Environmental Contamination and Toxicology*, 49(1), 9–17.
- Bhhatrai, B., Gramatica, P. (2011). Prediction of Aqueous Solubility, Vapor Pressure and Critical Micelle Concentration for Aquatic Partitioning of Perfluorinated Chemicals. *Environmental Science & Technology*, 45(19), 8120–8128.
- Bischel, H. N., MacManus-Spencer, L. A., Luthy, R. G. (2010). Noncovalent Interactions of Long-Chain Perfluoroalkyl Acids with Serum Albumin. *Environmental Science & Technology*, 44(13), 5263–5269.
- Bizkarguenaga, E., Zabaleta, I., Mijangos, L., Iparraguirre, A., Fernández, L. A., Prieto, A., Zuloaga, O. (2016). Uptake of perfluorooctanoic acid, perfluorooctane sulfonate and perfluorooctane sulfonamide by carrot and lettuce from compost amended soil. *Science of the Total Environment*, 571, 444–451.
- Blaine, A. C., Rich, C. D., Hundal, L. S., Lau, C., Mills, M. A., Harris, K. M., Higgins, C. P. (2013). Uptake of perfluoroalkyl acids into edible crops via land applied biosolids: Field and greenhouse studies. *Environmental Science and Technology*, 47(24), 14062–14069.

- Blaine, A. C., Rich, C. D., Sedlacko, E. M., Hundal, L. S., Kumar, K., Lau, C., ... Higgins, C. P. (2014). Perfluoroalkyl acid distribution in various plant compartments of edible crops grown in biosolids-amended soils. *Environmental Science and Technology*, 48(14), 7858–7865.
- Bonvin, F., Omlin, J., Rutler, R., Schweizer, W. B., Alaimo, P. J., Strathmann, T. J., ... Kohn, T. (2013). Direct photolysis of human metabolites of the antibiotic sulfamethoxazole: Evidence for abiotic back-transformation. *Environmental Science and Technology*, 47(13), 6746–6755.
- Boreen, A. L., Arnold, W. A., McNeill, K. (2004). Photochemical fate of sulfa drugs in then aquatic environment: Sulfa drugs containing five-membered heterocyclic groups. *Environmental Science and Technology*, 38(14), 3933–3940.
- Borrelli, L., Castelli, F., Ceotto, E., Cabassi, G., Tomasoni, C. (2014). Maize grain and silage yield and yield stability in a long-term cropping system experiment in northern Italy. *European Journal of Agronomy*, 55, 12–19.
- Brace, N. O. (1962). Long Chain Alkanoic and Alkenoic Acids with Perfluoroalkyl Terminal Segments. *Journal of Organic Chemistry*, 27(12), 4491–4498.
- Bradley, P. M., Barber, L. B., Chapelle, F. H., Gray, J. L., Kolpin, D. W., McMahon, P. B. (2009). Biodegradation of 17 β -estradiol, estrone and testosterone in stream sediments. *Environmental Science & Technology*, 43(6), 1902–1910.
- Brenna, S., Rocca, A., Sciacaluga, Ma., Valagussa, M. (2010). SOILQUALIMON - Sistema di monitoraggio della qualità dei suoli in Lombardia. In *Quaderni della Ricerca* (Vol. 110).
- Brooke, D., Footitt, A., Nwaogu, T. A. (2004). Environmental Risk Evaluation Report: Perfluorooctanesulphonate (PFOS). In Environment Agency.
- Burns, D. C., Ellis, D. A., Webster, E., McMurdo, C. J. (2009). Response to Comment on “Experimental p K_a Determination for Perfluorooctanoic Acid (PFOA) and the Potential Impact of p K_a Concentration Dependence on Laboratory-Measured Partitioning Phenomena and Environmental Modeling.” *Environmental Science & Technology*, 43(13), 5152–5154.
- Buser, H. R., Poiger, T., Muller, M. D. (1999). Occurrence and environmental behavior of the chiral pharmaceutical drug ibuprofen in surface waters and in wastewater. *Environmental Science and Technology*, 33(15), 2529–2535.

- Buser, H. R., Poiger, T., Müller, M. D. (1998). Occurrence and fate of the pharmaceutical drug diclofenac in surface waters: Rapid photodegradation in a lake. *Environmental Science and Technology*, 32(22), 3449–3456.
- Caldwell, D. J., Mastrocco, F., Anderson, P. D., Länge, R., Sumpter, J. P. (2012). Predicted-no-effect concentrations for the steroid estrogens estrone, 17 β -estradiol, estriol, and 17 α -ethinylestradiol. *Environmental Toxicology and Chemistry*, 31(6), 1396–1406.
- Calza, P., Medana, C., Padovano, E., Giancotti, V., Minero, C. (2013). Fate of selected pharmaceuticals in river waters. *Environmental Science and Pollution Research*, 20(4), 2262–2270.
- Carlos, L., Mártire, D. O., Gonzalez, M. C., Gomis, J., Bernabeu, A., Amat, A. M., Arques, A. (2012). Photochemical fate of a mixture of emerging pollutants in the presence of humic substances. *Water Research*, 46(15), 4732–4740.
- Carr, D. L., Morse, A. N., Zak, J. C., Anderson, T. A. (2011). Biological degradation of common pharmaceuticals and personal care products in soils with high water content. *Water, Air, and Soil Pollution*, 217(1–4), 127–134.
- Carter, L. J., Harris, E., Williams, M., Ryan, J. J., Kookana, R. S., Boxall, A. B. A. (2014). Fate and Uptake of Pharmaceuticals in Soil–Plant Systems. *Journal of Agricultural and Food Chemistry*, 62(4), 816–825.
- Castiglioni, S., Davoli, E., Riva, F., Palmiotto, M., Camporini, P., Manenti, A., Zuccato, E. (2018a). Data on occurrence and fate of emerging contaminants in a urbanised area. *Data in Brief*, 17, 533–543.
- Castiglioni, S., Davoli, E., Riva, F., Palmiotto, M., Camporini, P., Manenti, A., Zuccato, E. (2018b). Mass balance of emerging contaminants in the water cycle of a highly urbanized and industrialized area of Italy. *Water Research*, 131, 287–298.
- Caupos, E., Mazellier, P., Croue, J. P. (2011). Photodegradation of estrone enhanced by dissolved organic matter under simulated sunlight. *Water Research*, 45(11), 3341–3350.
- Challis, J. K., Hanson, M. L., Friesen, K. J., Wong, C. S. (2014). A critical assessment of the photodegradation of pharmaceuticals in aquatic environments: defining our current understanding and identifying knowledge gaps. *Environmental Science: Processes & Impacts*, 16(4), 672–696.
- Chapra, S. C. (1997). *Surface Water-Quality Modeling*. In McGraw-Hill Series in Water Resources and Environmental Engineering.

- Chen, Y. M., Guo, L. H. (2009). Fluorescence study on site-specific binding of perfluoroalkyl acids to human serum albumin. *Archives of Toxicology*, 83(3), 255–261.
- Chitescu, C. L., Nicolau, A. I., Stolker, A. A. M. (2013). Uptake of oxytetracycline, sulfamethoxazole and ketoconazole from fertilised soils by plants. *Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment*, 30(6), 1138–1146.
- Chowdhury, R. R., Charpentier, P. A., Ray, M. B. (2011). Photodegradation of 17 β -estradiol in aquatic solution under solar irradiation: Kinetics and influencing water parameters. *Journal of Photochemistry and Photobiology A: Chemistry*, 219(1), 67–75.
- Chowdhury, R. R., Charpentier, P., Ray, M. B. (2010). Photodegradation of estrone in solar irradiation. *Industrial and Engineering Chemistry Research*, 49(15), 6923–6930.
- Christou, A., Karaolia, P., Hapeshi, E., Michael, C., Fatta-Kassinos, D. (2017). Long-term wastewater irrigation of vegetables in real agricultural systems: Concentration of pharmaceuticals in soil, uptake and bioaccumulation in tomato fruits and human health risk assessment. *Water Research*, 109(1–2), 24–34.
- Colucci, M. S., Bork, H., Topp, E. (2001). Persistence of estrogenic hormones in agricultural soils: I. 17 β -estradiol and estrone. *J Environ Qual*, 30(6), 2070–2076.
- Colucci, M. S., Topp, E. (2002). Dissipation of part per trillion concentrations of estrogenic hormones from agricultural soils. *Canadian Journal of Soil Science*, 82(3), 335–340.
- Colucci, M. S., Topp, E. (2001). Persistence of Estrogenic Hormones in Agricultural Soils: II. 17 α -Ethinylestradiol. *Journal of Environment Quality*, 30(6), 2077–2080.
- Confalonieri, R., Bocchi, S. (2005). Evaluation of CropSyst for simulating the yield of flooded rice in northern Italy. *European Journal of Agronomy*, 23(4), 315–326.
- Conkle, J. L., Gan, J., Anderson, M. A. (2012). Degradation and sorption of commonly detected PPCPs in wetland sediments under aerobic and anaerobic conditions. *Journal of Soils and Sediments*, 12(7), 1164–1173.
- Czajka, C. P., Londry, K. L. (2006). Anaerobic biotransformation of estrogens. *Science of the Total Environment*, 367(2–3), 932–941.

- De Laurentiis, E., Chiron, S., Kouras-Hadef, S., Richard, C., Minella, M., Maurino, V., ... Vione, D. (2012). Photochemical fate of carbamazepine in surface freshwaters: Laboratory measures and modeling. *Environmental Science and Technology*, 46(15), 8164–8173.
- De Laurentiis, E., Prasse, C., Ternes, T. A., Minella, M., Maurino, V., Minero, C., ... Vione, D. (2014). Assessing the photochemical transformation pathways of acetaminophen relevant to surface waters: Transformation kinetics, intermediates, and modelling. *Water Research*, 53, 235–248.
- de Mes, T., Zeeman, G., Lettinga, G. (2005). Occurrence and fate of estrone, 17 β -estradiol and 17 α -ethynylestradiol in STPs for domestic wastewater. *Reviews in Environmental Science and Bio/Technology*, (4), 275–311.
- Ding, G., Peijnenburg, W. J. G. M. (2013). Physicochemical Properties and Aquatic Toxicity of Poly- and Perfluorinated Compounds. *Critical Reviews in Environmental Science and Technology*, 43(6), 598–678.
- Doll, T. E., Frimmel, F. H. (2003). Fate of pharmaceuticals - Photodegradation by simulated solar UV-light. *Chemosphere*, 52(10), 1757–1769.
- Dong, M. M., Trenholm, R., Rosario-Ortiz, F. L. (2015). Photochemical degradation of atenolol, carbamazepine, meprobamate, phenytoin and primidone in wastewater effluents. *Journal of Hazardous Materials*, 282, 216–223.
- Durán-Álvarez, J. C., Prado, B., González, D., Sánchez, Y., Jiménez-Cisneros, B. (2015). Environmental fate of naproxen, carbamazepine and triclosan in wastewater, surface water and wastewater irrigated soil - Results of laboratory scale experiments. *Science of the Total Environment*, 538, 350–362.
- ECHA. (2006). Paracetamol - Environmental fate & pathways. Retrieved December 18, 2018, from <https://echa.europa.eu/registration-dossier/-/registered-dossier/12532/5/5/2>
- EFSA. (2008). Perfluorooctane sulfonate (PFOS), perfluorooctanoic acid (PFOA) and their salts. *The EFSA Journal*, 653, 1–131.
- EFSA. (2007). Opinion on a request from EFSA related to the default Q10 value used to describe the temperature effect on transformation rates of pesticides in soil. *The EFSA Journal*, 622, 1–32.
- EMA Committee for Veterinary Medicinal Products (2000). Furosemide – Summary Report. Technical Report September 1999.

- Environment Protection and Heritage Council of Australia (2008). Australian Guidelines for Water Recycling: Managing Health and Environmental Risks (Phase 2) - Augmentation of Drinking Water Supplies. Technical Report 22.
- European Commission Joint Research Centre. (2003). Technical guidance document on risk assessment in support of Commission Directive 93/67/ EEC on risk assessment for new notified substances and Commission Regulation (EC) No. 1488/94 on risk assessment for existing substances. Part II. EUR 20418 EN/2. European Chemicals Bureau, Part II, 7–179.
- Falås, P., Longrée, P., La Cour Jansen, J., Siegrist, H., Hollender, J., Joss, A. (2013). Micropollutant removal by attached and suspended growth in a hybrid biofilm-activated sludge process. *Water Research*, 47(13), 4498–4506.
- FAO (2007). ECOCROP - Data Sheet - *Lolium multiflorum*. URL: <http://ecocrop.fao.org/ecocrop/srv/en/dataSheet?id=7389>. Last accessed: 05-10-2018.
- Felle, H. H., Herrmann, A., Hückelhoven, R., Kogel, K. H. (2005). Root-to-shoot signalling: Apoplastic alkalization, a general stress response and defence factor in barley (*Hordeum vulgare*). *Protoplasma*, 227(1), 17–24.
- Felle, H. H. (2001). pH: Signal and messenger in plant cells. *Plant Biology*, 3(6), 577–591.
- Fenner, K., Honti, M., Stamm, C., Varga, L., Bischoff, F. (2016). Suitability of laboratory simulation tests for the identification of persistence in surface waters.
- Fono, L. J., Kolodziej, E. P., Sedlak, D. L. (2006). Attenuation of wastewater-derived contaminants in an effluent-dominated river. *Environmental Science and Technology*, 40(23), 7257–7262.
- Franco, A., Ferranti, A., Davidsen, C., Trapp, S. (2010). An unexpected challenge: Ionizable compounds in the REACH chemical space. *International Journal of Life Cycle Assessment*, 15(4), 321–325.
- Franco, A., Struijs, J., Gouin, T., Price, O. R. (2013). Evolution of the sewage treatment plant model SimpleTreat: Applicability domain and data requirements. *Integrated Environmental Assessment and Management*, 9(4), 560–568.
- Franco, A., Trapp, S. (2008). Estimation of the soil-water partitioning coefficient normalized to organic carbon for ionizable organic chemicals. *Environmental Toxicology and Chemistry*, 27(10), 1995–2004.

- Franco, A., Trapp, S. (2010). A multimedia activity model for ionizable compounds: Validation study with 2,4-dichlorophenoxyacetic acid, aniline, and trimethoprim. *Environmental Toxicology and Chemistry*, 29(4), 789–799.
- Fu, Q., Wu, X., Ye, Q., Ernst, F., Gan, J. (2016). Biosolids inhibit bioavailability and plant uptake of triclosan and triclocarban. *Water Research*, 102, 117–124.
- Giesy, J. P., Naile, J. E., Khim, J. S., Jones, P. D., Newsted, J. L. (2010). Aquatic Toxicology of Perfluorinated Chemicals. In D. M. Whitacre (Ed.), *Reviews of Environmental Contamination and Toxicology* (Vol. 202).
- Goldstein, M., Shenker, M., Chefetz, B. (2014). Insights into the Uptake Processes of Wastewater-Borne Pharmaceuticals by Vegetables. *Environmental Science & Technology*, 48, 5593–5600.
- Goss, K. U. (2008). The pKa values of PFOA and other highly fluorinated carboxylic acids. *Environmental Science and Technology*, 42(2), 456–458.
- Gozlan, I., Koren, I. (2016). Identification, Mechanisms and Kinetics of Macrolide Degradation Product Formation under Controlled Environmental Conditions. *Journal of Environmental Analytical Chemistry*, 03(01), 1–9.
- Gredelj, A., Barausse, A., Grechi, L., Palmeri, L. (2018). Deriving predicted no-effect concentrations (PNECs) for emerging contaminants in the river Po, Italy, using three approaches: Assessment factor, species sensitivity distribution and AQUATOX ecosystem modelling. *Environment International*, 119(June), 66–78.
- Gröning, J., Held, C., Garten, C., Claußnitzer, U., Kaschabek, S. R., Schlömann, M. (2007). Transformation of diclofenac by the indigenous microflora of river sediments and identification of a major intermediate. *Chemosphere*, 69(4), 509–516.
- Grossberger, A., Hadar, Y., Borch, T., Chefetz, B. (2014). Biodegradability of pharmaceutical compounds in agricultural soils irrigated with treated wastewater. *Environmental Pollution*, 185, 168–177.
- Grzybowski, W., Szydlowski, J. (2014). The impact of chromophoric dissolved organic matter on the photodegradation of 17 α -ethinylestradiol (EE2) in natural waters. *Chemosphere*, 111, 13–17.

- Guillet, G., Knapp, J. L. A., Merel, S., Cirpka, O. A., Grathwohl, P., Zwiener, C., Schwientek, M. (2019). Fate of wastewater contaminants in rivers: Using conservative-tracer based transfer functions to assess reactive transport. *Science of The Total Environment*, 656, 1250–1260.
- Han, X., Snow, T. A., Kemper, R. A., Jepson, G. W. (2003). Binding of perfluorooctanoic acid to rat and human plasma proteins. *Chemical Research in Toxicology*, 16(6), 775–781.
- Hanamoto, S., Nakada, N., Yamashita, N., Tanaka, H. (2013). Modeling the photochemical attenuation of down-the-drain chemicals during river transport by stochastic methods and field measurements of pharmaceuticals and personal care products. *Environmental Science and Technology*, 47(23), 13571–13577.
- Hebert, P. C., MacManus-Spencer, L. A. (2010). Development of a Fluorescence Model for the Binding of Medium- to Long-Chain Perfluoroalkyl Acids to Human Serum Albumin Through a Mechanistic Evaluation of Spectroscopic Evidence. *Analytical Chemistry*, 82(15), 6463–6471.
- Henry, A., Cal, A. J., Batoto, T. C., Torres, R. O., Serraj, R. (2012). Root attributes affecting water uptake of rice (*Oryza sativa*) under drought. *Journal of Experimental Botany*, 63(13), 4751–4763.
- Henze, M., Gujer, W., Mino, T., Loosdrecht, M. van. (2002). Activated Sludge Models ASM1, ASM2, ASM2d and ASM3. In IWA Task Group on Mathematical Modelling for Design and Operation of Biological Wastewater Treatment. IWA Publishing.
- Higgins, C. P., Luthy, R. G. (2006). Sorption of perfluorinated surfactants on sediments. *Environmental Science and Technology*, 40(23), 7251–7256.
- Holling, C. S., Bailey, J. L., Heuvel, V., Kinney, C. A. (2012). Uptake of human pharmaceuticals and personal care products by cabbage (*Brassica campestris*) from fortified and biosolids-amended soils. *Journal of Environmental Monitoring*, 14, 3029–3036.
- Huang, Q., Bu, Q., Zhong, W., Shi, K., Cao, Z., Yu, G. (2018). Derivation of aquatic predicted no-effect concentration (PNEC) for ibuprofen and sulfamethoxazole based on various toxicity endpoints and the associated risks. *Chemosphere*, 193, 223–229.
- Hurtado, C., Cañameras, N., Domínguez, C., Price, G. W., Comas, J., Bayona, J. M. (2017). Effect of soil biochar concentration on the mitigation of emerging organic contaminant uptake in lettuce. *Journal of Hazardous Materials*, 323(April 2016), 386–393.

- Igarashi, S., Yotsuyanagi, T. (1992). Homogeneous liquid-liquid extraction by pH dependent phase separation with a fluorocarbon ionic surfactant and its application to the preconcentration of porphyrin compounds. *Mikrochimica Acta*, 106(1–2), 37–44.
- Jakimska, A., Śliwka-Kaszyńska, M., Reszczyńska, J., Namieśnik, J., Kot-Wasik, A. (2014). Elucidation of transformation pathway of ketoprofen, ibuprofen, and furosemide in surface water and their occurrence in the aqueous environment using UHPLC-QTOF-MS *Euroanalysis XVII. Analytical and Bioanalytical Chemistry*, 406(15), 3667–3680.
- Jensen, A. A., Poulsen, P. B., Bossi, R. (2008). Survey and environmental/health assessment of fluorinated substances in impregnated consumer products and impregnating agents. In *Survey of Chemical Substances in Consumer Products*.
- Jing, P., Rodgers, P. J., Amemiya, S. (2009). High lipophilicity of perfluoroalkyl carboxylate and sulfonate: Implications for their membrane permeability. *Journal of the American Chemical Society*, 131(6), 2290–2296.
- Johannes, E., Collings, D. a, Rink, J. C., Allen, N. S. (2001). Cytoplasmic pH dynamics in maize pulvinal cells induced by gravity vector changes. *Plant Physiology*, 127(1), 119–130.
- Jurgens, M. D., Holthaus, K. I. E., Johnson, A. C., Smith, J. J. L., Hetheridge, M., Williams, R. J. (2002). The potential for estradiol and ethinyloestradiol degradation in English Rivers. *Environmental Toxicology and Chemistry*, 21(3), 480–488.
- Karnjanapiboonwong, A., Chase, D. A., Cañas, J. E., Jackson, W. A., Maul, J. D., Morse, A. N., Anderson, T. A. (2011). Uptake of 17 α -ethinyloestradiol and triclosan in pinto bean, *Phaseolus vulgaris*. *Ecotoxicology and Environmental Safety*, 74(5), 1336–1342.
- Kawabata, K., Sugihara, K., Sanoh, S., Kitamura, S., Ohta, S. (2013). Photodegradation of pharmaceuticals in the aquatic environment by sunlight and UV-A, -B and -C irradiation. *The Journal of Toxicological Sciences*, 38(2), 215–223.
- Khan, S. J., Ongerth, J. E. (2004). Modelling of pharmaceutical residues in Australian sewage by quantities of use and fugacity calculations. *Chemosphere*, 54(3), 355–367.
- Kleier, D. A., Hsu, F. C. (1996). Phloem Mobility of Xenobiotics. VII. The Design of Phloem Systemic Pesticides. *Weed Science*, 44(3), 749–756.

- Kodešová, R., Kočárek, M., Klement, A., Golovko, O., Koba, O., Fér, M., ... Grabic, R. (2016). An analysis of the dissipation of pharmaceuticals under thirteen different soil conditions. *Science of the Total Environment*, 544, 369–381.
- Koumaki, E., Mamais, D., Noutsopoulos, C. (2018). Assessment of the environmental fate of endocrine disrupting chemicals in rivers. *Science of the Total Environment*, 628–629, 947–958.
- Koumaki, E., Mamais, D., Noutsopoulos, C. (2017). Environmental fate of non-steroidal anti-inflammatory drugs in river water/sediment systems. *Journal of Hazardous Materials*, 323, 233–241.
- Koumaki, E., Mamais, D., Noutsopoulos, C., Nika, M. C., Bletsou, A. A., Thomaidis, N. S., ... Stratogianni, G. (2015). Degradation of emerging contaminants from water under natural sunlight: The effect of season, pH, humic acids and nitrate and identification of photodegradation by-products. *Chemosphere*, 138, 675–681.
- Kroes, R., Renwick, A., Cheeseman, M., Kleiner, J., Mangelsdorf, I., Piersma, A., Schilter, B., Schlatter, J., van Schothorst, F., Vos, J., and Würtzen, G. (2004). Structure-based thresholds of toxicological concern (TTC): guidance for application to substances present at low levels in the diet. *Food and Chemical Toxicology*, 42(1):65–83.
- Kunkel, U., Radke, M. (2008). Biodegradation of Acidic Pharmaceuticals in Bed Sediments : Insight from a Laboratory Experiment. *Environmental Science and Technology*, 42(19), 7273–7279.
- Kunkel, U., Radke, M. (2011). Reactive tracer test to evaluate the fate of pharmaceuticals in rivers. *Environmental Science and Technology*, 45(15), 6296–6302.
- Lai, H. T., Hou, J. H. (2008). Light and microbial effects on the transformation of four sulfonamides in eel pond water and sediment. *Aquaculture*, 283(1–4), 50–55.
- Lam, M. W., Mabury, S. A. (2005). Photodegradation of the pharmaceuticals atorvastatin, carbamazepine, levofloxacin, and sulfamethoxazole in natural waters. *Aquatic Sciences*, 67(2), 177–188.
- Lam, M. W., Young, C. J., Brain, R. A., Johnson, D. J., Hanson, M. A., Wilson, C. J., ... Mabury, S. A. (2004). Aquatic persistence of eight pharmaceuticals in a microcosm study. *Environmental Toxicology and Chemistry*, 23(6), 1431–1440.

- Latch, D. E., Packer, J. L., Stender, B. L., VanOverbeke, J., Arnold, W. A., McNeill, K. (2005). Aqueous photochemistry of triclosan: Formation of 2,4-dichlorophenol, 2,8-dichlorodibenzo-p-dioxin, and oligomerization products. *Environmental Toxicology and Chemistry*, 24(3), 517–525.
- Lechner, M., Knapp, H. (2011). Carryover of Perfluorooctanoic Acid (PFOA) and Perfluorooctane Sulfonate (PFOS) from Soil to Plant and Distribution to the Different Plant Compartments Studied in Cultures of Carrots (*Daucus carota* ssp. *Sativus*), Potatoes (*Solanum tuberosum*), and Cucu. *Journal of Agricultural and Food Chemistry*, 59(20), 11011–11018.
- Leech, D. M., Snyder, M. T., Wetzel, R. G. (2009). Natural organic matter and sunlight accelerate the degradation of 17 β -estradiol in water. *Science of the Total Environment*, 407(6), 2087–2092.
- Legind, C. N., Rein, A., Serre, J., Brochier, V., Haudin, C. S., Cambier, P., ... Trapp, S. (2012). Simultaneous Simulations of Uptake in Plants and Leaching to Groundwater of Cadmium and Lead for Arable Land Amended with Compost or Farmyard Manure. *PLoS ONE*, 7(10), 1–13.
- Li, F. H., Yao, K., Lv, W. Y., Liu, G. G., Chen, P., Huang, H. P., Kang, Y. P. (2015). Photodegradation of ibuprofen under UV-VIS irradiation: Mechanism and toxicity of photolysis products. *Bulletin of Environmental Contamination and Toxicology*, 94(4), 479–483.
- Li, J., Zhang, Y., Huang, Q., Shi, H., Yang, Y., Gao, S., ... Yang, X. (2017). Degradation of organic pollutants mediated by extracellular peroxidase in simulated sunlit humic waters: A case study with 17B-estradiol. *Journal of Hazardous Materials*, 331, 123–131.
- Li, J., Dodgen, L., Ye, Q., Gan, J. (2013). Degradation kinetics and metabolites of carbamazepine in soil. *Environmental Science and Technology*, 47(8), 3678–3684.
- Li, J., Ye, Q., Gan, J. (2014). Degradation and transformation products of acetaminophen in soil. *Water Research*, 49, 44–52.
- Li, J., Ye, Q., Gan, J. (2015). Influence of organic amendment on fate of acetaminophen and sulfamethoxazole in soil. *Environmental Pollution*, 206, 543–550.
- Li, L., Xu, Z. S., Song, G. W. (2009). Study on the Langmuir aggregation of fluorinated surfactants on protein. *Journal of Fluorine Chemistry*, 130(2), 225–230.
- Li, Y., Pan, Y., Lian, L., Yan, S., Song, W., Yang, X. (2017). Photosensitized degradation of acetaminophen in natural organic matter solutions: The role of triplet states and oxygen. *Water Research*, 109, 266–273.

- Li, Z., Sobek, A., Radke, M. (2015). Flume experiments to investigate the environmental fate of pharmaceuticals and their transformation products in streams. *Environmental Science and Technology*, 49(10), 6009–6017.
- Lin, A. Y. C., Plumlee, M. H., Reinhard, M. (2006). Natural attenuation of pharmaceuticals and alkylphenol polyethoxylate metabolites during river transport: Photochemical and biological transformation. *Environmental Toxicology and Chemistry*, 25(6), 1458–1464.
- Lin, A. Y. C., Reinhard, M. (2005). Photodegradation of common environmental pharmaceuticals and estrogens in river water. *Environmental Toxicology and Chemistry*, 24(6), 1303–1309.
- Lin, A. Y. C., Lin, C. A., Tung, H. H., Chary, N. S. (2010). Potential for biodegradation and sorption of acetaminophen, caffeine, propranolol and acebutolol in lab-scale aqueous environments. *Journal of Hazardous Materials*, 183(1–3), 242–250.
- Lin, K., Gan, J. (2011). Sorption and degradation of wastewater-associated non-steroidal anti-inflammatory drugs and antibiotics in soils. *Chemosphere*, 83(3), 240–246.
- Lindström, A., Buerge, I. J., Poiger, T., Bergqvist, P. A., Müller, M. D., Buser, H. R. (2002). Occurrence and environmental behavior of the bactericide triclosan and its methyl derivative in surface waters and in wastewater. *Environmental Science and Technology*, 36(11), 2322–2329.
- Liou, J. S. C., Szostek, B., DeRito, C. M., Madsen, E. L. (2010). Investigating the biodegradability of perfluorooctanoic acid. *Chemosphere*, 80(2), 176–183.
- Liu, F., Ying, G.-G., Yang, J.-F., Zhou, L.-J., Tao, R., Wang, L., ... Peng, P.-A. (2010). Dissipation of sulfamethoxazole, trimethoprim and tylosin in a soil under aerobic and anoxic conditions. *Environmental Chemistry*, 7(4), 370–376.
- Liu, Y., Sun, W., Ni, J. (2011). Biodegradation of bisphenol A, 17 β -estradiol, and 17 α -ethynylestradiol in river water. *International Journal of Environment and Pollution*, 45, 225–236.
- Liu, Y., Sun, H., Zhang, L., Feng, L. (2017). Photodegradation behaviors of 17 β -estradiol in different water matrixes. *Process Safety and Environmental Protection*, 112, 335–341.
- Löffler, D., Römbke, J., Meller, M., Ternes, T. A. (2005). Environmental fate of pharmaceuticals in water/sediment systems. *Environmental Science and Technology*, 39(14), 5209–5218.

- López-Fontán, J. L., Sarmiento, F., Schulz, P. C. (2005). The aggregation of sodium perfluorooctanoate in water. *Colloid and Polymer Science*, 283(8), 862–871.
- Macherius, A., Eggen, T., Lorenz, W. G., Reemtsma, T., Winkler, U., Moeder, M. (2012). Uptake of Galaxolide, Tonalide, and Triclosan by Carrot, Barley, and Meadow Fescue Plants. *Journal of Agricultural and Food Chemistry*, 60(32), 7785–7791.
- Malchi, T., Maor, Y., and Chefetz, B. (2015). Comments on “Human health risk assessment of pharmaceuticals and personal care products in plant tissue due to biosolids and manure amendments, and wastewater irrigation”. *Environment International*, 82:110–112.
- Malchi, T., Maor, Y., Tadmor, G., Shenker, M., Chefetz, B. (2014). Irrigation of Root Vegetables with Treated Wastewater: Evaluating Uptake of Pharmaceuticals and the Associated Human Health Risks. *Environmental Science & Technology*, 48(16), 9325–9333.
- Martínez-Zapata, M., Aristizábal, C., Peñuela, G. (2013). Photodegradation of the endocrine-disrupting chemicals 4n-nonylphenol and triclosan by simulated solar UV irradiation in aqueous solutions with Fe(III) and in the absence/presence of humic acids. *Journal of Photochemistry and Photobiology A: Chemistry*, 251, 41–49.
- Martiniello, P. (1999). Effects of irrigation and harvest management on dry-matter yield and seed yield of annual clovers grown in pure stand and in mixtures with graminaceous species in a Mediterranean environment. *Grass and Forage Science*, 54, 52–61.
- Mashtare, M. L., Lee, L. S., Nies, L. F., Turco, R. F. (2013). Transformation of 17 α -Estradiol, 17 β -estradiol, and estrone in sediments under nitrate- and sulfate-reducing conditions. *Environmental Science and Technology*, 47(13), 7178–7185.
- Matamoros, V., Duhec, A., Albaigés, J., Bayona, J. M. (2009). Photodegradation of carbamazepine, ibuprofen, ketoprofen and 17 α -ethinylestradiol in fresh and seawater. *Water, Air, and Soil Pollution*, 196(1–4), 161–168.
- Matamoros, V., Rodríguez, Y. (2017). Influence of seasonality and vegetation on the attenuation of emerging contaminants in wastewater effluent-dominated streams. A preliminary study. *Chemosphere*, 186, 269–277.

- Matsuoka, S., Kikuchi, M., Kimura, S., Kurokawa, Y., Kawai, S. (2005). Determination of Estrogenic Substances in the Water of Muko River Using in Vitro Assays, and the Degradation of Natural Estrogens by Aquatic Bacteria. *Journal of Health Science*, 51(2), 178–184.
- Mazzini, R., Pedrazzi, L., Lazarova, V. (2013). Production of high quality recycled water for agricultural irrigation in Milan. In V. Lazarova, T. Asano, A. Bahri, J. Anderson (Eds.), *Milestones in Water Reuse* (pp. 179–190).
- McArdell, C. S., Molnar, E., Suter, M. J. F., Giger, W. (2003). Occurrence and Fate of Macrolide Antibiotics in Wastewater Treatment Plants and in the Glatt Valley Watershed, Switzerland. *Environmental Science and Technology*, 37(24), 5479–5486.
- Mejia Avendaño, S., Zhong, G., Liu, J. (2015). Comment on “Biodegradation of perfluorooctanesulfonate (PFOS) as an emerging contaminant.” *Chemosphere*, 138, 1037–1038.
- Mezcua, M., Gómez, M. J., Ferrer, I., Aguera, A., Hernando, M. D., Fernández-Alba, A. R. (2004). Evidence of 2,7/2,8-dibenzodichloro-p-dioxin as a photodegradation product of triclosan in water and wastewater samples. *Analytica Chimica Acta*, 524(1–2 SPEC. ISS.), 241–247.
- Morrall, D., McAvoy, D., Schatowitz, B., Inauen, J., Jacob, M., Hauk, A., Eckhoff, W. (2004). A field study of triclosan loss rates in river water (Cibolo Creek, TX). *Chemosphere*, 54(5), 653–660.
- Navarro, I., de la Torre, A., Sanz, P., Porcel, M. Á., Pro, J., Carbonell, G., Martínez, M. D. L. Á. (2017). Uptake of perfluoroalkyl substances and halogenated flame retardants by crop plants grown in biosolids-amended soils. *Environmental Research*, 152(October 2016), 199–206.
- Nelli, E. and Sodi, F. (2007). *Riso - Oryza sativa L. - Atlante delle coltivazioni erbacee - Piante industriali*. URL: <http://www.agraria.org/coltivazionierbacee/riso.htm>. Last accessed: 5-10-2018.
- NICNAS. (2009). *Triclosan, Priority Existing Chemical Assessment Report*.
- Nielsen, C. J. (2012). *PFOA Isomers, Salts, and Precursors*.
- Niu, J., Zhang, L., Li, Y., Zhao, J., Lv, S., Xiao, K. (2013). Effects of environmental factors on sulfamethoxazole photodegradation under simulated sunlight irradiation: Kinetics and mechanism. *Journal of Environmental Sciences*, 25(6), 1098–1106.

- Oliveira, C., Lima, D. L. D., Silva, C. P., Calisto, V., Otero, M., Esteves, V. I. (2019). Photodegradation of sulfamethoxazole in environmental samples: The role of pH, organic matter and salinity. *Science of the Total Environment*, 648, 1403–1410.
- Packer, J. L., Werner, J. J., Latch, D. E., McNeill, K., Arnold, W. A. (2003). Photochemical fate of pharmaceuticals in the environment: Naproxen, diclofenac, clofibric acid, and ibuprofen. *Aquatic Sciences*, 65(4), 342–351.
- Pannu, M. W., Toor, G. S., O'Connor, G. A., Wilson, P. C. (2012). Toxicity and bioaccumulation of biosolids-borne triclosan in food crops. *Environmental Toxicology and Chemistry*, 31(9), 2130–2137.
- Peuravuori, J., Pihlaja, K. (2009). Phototransformations of selected pharmaceuticals under low-energy UVA-vis and powerful UVB-UVA irradiations in aqueous solutions-the role of natural dissolved organic chromophoric material. *Analytical and Bioanalytical Chemistry*, 394(6), 1621–1636.
- Peuravuori, J. (2012). Aquatic photochemistry of paracetamol in the presence of dissolved organic chromophoric material and nitrate. *Environmental Science and Pollution Research*, 19(6), 2259–2270.
- Poiger, T., Buser, H. R., Müller, M. D. (2001). Photodegradation of the pharmaceutical drug diclofenac in a lake: Pathway, field measurements, and mathematical modeling. *Environmental Toxicology and Chemistry*, 20(2), 256–263.
- Poirier-Larabie, S., Segura, P. A., Gagnon, C. (2016). Degradation of the pharmaceuticals diclofenac and sulfamethoxazole and their transformation products under controlled environmental conditions. *Science of the Total Environment*, 557–558, 257–267.
- Polesel, F., Plósz, B. G., Trapp, S. (2015). From consumption to harvest: Environmental fate prediction of excreted ionizable trace organic chemicals. *Water Research*, 84, 85–98.
- Portmann, F., Siebert, S., Bauer, C., Döll, P. (2008). Global dataset of monthly growing areas of 26 irrigated crops. *Frankfurt Hydrology Paper*, 06, 400.
- Prosser, R. S., Lissemore, L., Topp, E., Sibley, P. K. (2014). Bioaccumulation of triclosan and triclocarban in plants grown in soils amended with municipal dewatered biosolids. *Environmental Toxicology and Chemistry*, 33(5), 975–984.

- Prosser, R. S. and Sibley, P. K. (2015). Human health risk assessment of pharmaceuticals and personal care products in plant tissue due to biosolids and manure amendments, and wastewater irrigation. *Environment International*, 75:223–233.
- Radke, M., Lauwigi, C., Heinkele, G., Mürdter, T. E., Letzel, M. (2009). Fate of the antibiotic sulfamethoxazole and its two major human metabolites in a water sediment test. *Environmental Science and Technology*, 43(9), 3135–3141.
- Radke, M., Maier, M. P. (2014). Lessons learned from water/sediment-testing of pharmaceuticals. *Water Research*, 55, 63–73.
- Radke, M., Ulrich, H., Wurm, C., Kunkel, U. (2010). Dynamics and attenuation of acidic pharmaceuticals along a river stretch. *Environmental Science and Technology*, 44(8), 2968–2974.
- Rahman, M. F., Peldszus, S., Anderson, W. B. (2014). Behaviour and fate of perfluoroalkyl and polyfluoroalkyl substances (PFASs) in drinking water treatment: A review. *Water Research*, 50, 318–340.
- Rawls, W. J., Brakensiek, D. L., Saxton, K. E. (1982). Estimation of soil water properties. *American Society of Agricultural and Biological Engineers*, 25(5), 1316–1320.
- Rayne, S., Forest, K. (2009). Congener-specific organic carbon-normalized soil and sediment-water partitioning coefficients for the C 1 through C 8 perfluoroalkyl carboxylic and sulfonic acids. *Journal of Environmental Science and Health, Part A*, 44(13), 1374–1387.
- Rayne, S., Forest, K. (2009). Perfluoroalkyl sulfonic and carboxylic acids : A critical review of physicochemical properties , levels and patterns in waters and wastewaters , and treatment methods Perfluoroalkyl sulfonic and carboxylic acids : A critical review of physicochemical prop. 4529.
- Reichert, P., Vanrolleghem, P. (2001). Identifiability and uncertainty analysis of the River Water Quality Model No. 1 (RWQM1). *Water Science and Technology*, 43(7), 329–338.
- Ren, D., Bi, T., Gao, S., Li, X., Huang, B., Pan, X. (2016). Photodegradation of 17 α -ethynylestradiol in nitrate aqueous solutions. *Environmental Engineering Research*, 21(2), 188–195.
- Ren, D., Huang, B., Xiong, D., He, H., Meng, X., Pan, X. (2017). Photodegradation of 17 α -ethynylestradiol in dissolved humic substances solution: Kinetics, mechanism and estrogenicity variation. *Journal of Environmental Sciences*, 54, 196–205.

- Ren, D., Huang, B., Yang, B., Pan, X., Dionysiou, D. D. (2017). Mitigating 17 α -ethynylestradiol water contamination through binding and photosensitization by dissolved humic substances. *Journal of Hazardous Materials*, 327, 197–205.
- Rice Knowledge Bank. (2017). Measuring moisture content. Retrieved October 15, 2018, from <http://www.knowledgebank.irri.org/step-by-step-production/postharvest/milling/milling-and-quality/measuring-moisture-content-in-milling>
- Robinson, J. A., Ma, Q., Staveley, J. P., Smolenski, W. J., Ericson, J. (2017). Degradation and transformation of 17 α -estradiol in water–sediment systems under controlled aerobic and anaerobic conditions. *Environmental Toxicology and Chemistry*, 36(3), 621–629.
- Ryan, C. C., Tan, D. T., Arnold, W. A. (2011). Direct and indirect photolysis of sulfamethoxazole and trimethoprim in wastewater treatment plant effluent. *Water Research*, 45(3), 1280–1286.
- Sabaliunas, D., Webb, S. F., Hauk, A., Jacob, M., Eckhoff, W. S. (2003). Environmental fate of Triclosan in the River Aire Basin, UK. *Water Research*, 37(13), 3145–3154.
- Sabourin, L., Duenk, P., Bonte-Gelok, S., Payne, M., Lapen, D. R., Topp, E. (2012). Uptake of pharmaceuticals, hormones and parabens into vegetables grown in soil fertilized with municipal biosolids. *Science of the Total Environment*, 431, 233–236.
- Sanchez-Prado, L., Llompart, M., Lores, M., García-Jares, C., Bayona, J. M., Cela, R. (2006). Monitoring the photochemical degradation of triclosan in wastewater by UV light and sunlight using solid-phase microextraction. *Chemosphere*, 65(8), 1338–1347.
- Shenker, M., Harush, D., Ben-ari, J., Chefetz, B. (2011). Chemosphere Uptake of carbamazepine by cucumber plants – A case study related to irrigation with reclaimed wastewater. *Chemosphere*, 82(6), 905–910.
- Silva, C. P., Lima, D. L. D., Otero, M., Esteves, V. I. (2016). Photosensitized Degradation of 17 β -estradiol and 17 α -ethynylestradiol: Role of Humic Substances Fractions. *Journal of Environment Quality*, 45(2), 693–700.
- Silva, C. P., Lima, D. L. D., Groth, M. B., Otero, M., Esteves, V. I. (2016). Effect of natural aquatic humic substances on the photodegradation of estrone. *Chemosphere*, 145, 249–255.

- Simonsen, J. F., Harremoës, P. (1978). Oxygen and pH fluctuations in rivers. *Water Research*, 12(7), 477–489.
- Singer, H., Müller, S., Tixier, C., Pillonel, L. (2002). Triclosan: Occurrence and fate of a widely used biocide in the aquatic environment: Field measurements in wastewater treatment plants, surface waters, and lake sediments. *Environmental Science and Technology*, 36(23), 4998–5004.
- Stahl, T., Heyn, J., Thiele, H., Hüther, J., Failing, K., Georgii, S., Brunn, H. (2009). Carryover of perfluorooctanoic acid (PFOA) and perfluorooctane sulfonate (PFOS) from soil to plants. *Archives of Environmental Contamination and Toxicology*, 57(2), 289–298.
- Stevens, J. B., Coryell, A. (2007). Surface Water Quality Criterion for Perfluorooctane Sulfonic Acid.
- Su, T., Deng, H., Benskin, J. P., Radke, M. (2016). Biodegradation of sulfamethoxazole photo-transformation products in a water/sediment test. *Chemosphere*, 148, 518–525.
- Tabaglio, V., Ligabue, M., Reggiani, R., Piazza, C., Tassi, D., Ruozzi, F. (2007). Quale loglio italiano scegliere per la campagna 2007-08. *L'informatore Agrario*, 35, 40–43.
- Tixier, C., Singer, H. P., Canonica, S., Müller, S. R. (2002). Phototransformation of triclosan in surface waters: A relevant elimination process for this widely used biocide - Laboratory studies, field measurements, and modeling. *Environmental Science and Technology*, 36(16), 3482–3489.
- Tixier, C., Singer, H. P., Oellers, S., Müller, S. R. (2003). Occurrence and Fate of Carbamazepine, Clofibric Acid, Diclofenac, Ibuprofen, Ketoprofen, and Naproxen in Surface Waters. *Environmental Science & Technology*, 37(6), 1061–1068.
- Tournaire-Roux, C., Sutka, M., Javot, H., Gout, E., Gerbeau, P., Luu, D. T., ... Maurel, C. (2003). Cytosolic pH regulates root water transport during anoxic stress through gating of aquaporins. *Nature*, 425(6956), 393–397.
- USDA (2018). National Nutrient Database for Standard Reference Legacy Release - Basic Report: 11167, Corn, sweet, yellow, raw. URL: <https://ndb.nal.usda.gov/ndb/foods/show/11167>. Last accessed: 18-10-2018
- Vaalgamaa, S., Vähätalo, A. V., Perkola, N., Huhtala, S. (2011). Photochemical reactivity of perfluorooctanoic acid (PFOA) in conditions representing surface water. *Science of the Total Environment*, 409(16), 3043–3048.

- Van der Vorm, P. D. J. (1980). Uptake of Si by five plant species, as influenced by variations in Si-supply. *Plant and Soil*, 56(1), 153–156.
- Vanrolleghem, P., Borchardt, D., Henze, M., Rauch, W., Reichert, P., Shanahan, P., Somlyódy, L. (2001). River Water Quality Model no. 1 (RWQM1): III. Biochemical submodel selection. *Water Science and Technology*, 43(5), 31–40.
- Vierke, L., Berger, U., Cousins, I. T. (2013). Estimation of the acid dissociation constant of perfluoroalkyl carboxylic acids through an experimental investigation of their water-to-air transport. *Environmental Science and Technology*, 47(19), 11032–11039.
- Vochezer, K. (2010). Modelling of Carbamazepine and Diclofenac in a River Network by Modelling of Carbamazepine and Diclofenac in a River Network. Swedish University of Agricultural Sciences, Uppsala.
- Wang, X. -L, Canny, M. J., McCully, M. E. (1991). The water status of the roots of soil-grown maize in relation to the maturity of their xylem. *Physiologia Plantarum*, 82(2), 157–162.
- Wang, Z., MacLeod, M., Cousins, I. T., Scheringer, M., Hungerbühler, K. (2011). Using COSMOtherm to predict physicochemical properties of poly- and perfluorinated alkyl substances (PFASs). *Environmental Chemistry*, 8(4), 389–398.
- Wen, B., Li, L., Zhang, H., Ma, Y., Shan, X., Zhang, S. (2014). Field study on the uptake and translocation of per fluoroalkyl acids (PFAAs) by wheat (*Triticum aestivum* L.) grown in biosolids-amended soils. *Environmental Pollution*, 184, 547–554.
- Wen, B., Wu, Y., Zhang, H., Liu, Y., Hu, X., Huang, H. (2016). The roles of protein and lipid in the accumulation and distribution of per fluoroctane sulfonate (PFOS) and per fluoroctanoate (PFOA) in plants grown in biosolids-amended soils *. *Environmental Pollution*, 216, 682–688.
- Whidbey, C. M., Daumit, K. E., Nguyen, T. H., Ashworth, D. D., Davis, J. C. C., Latch, D. E. (2012). Photochemical induced changes of in vitro estrogenic activity of steroid hormones. *Water Research*, 46(16), 5287–5296.
- Willach, S., Lutze, H. V., Eckey, K., Löppenberg, K., Lüling, M., Wolbert, J. B., ... Schmidt, T. C. (2018). Direct Photolysis of Sulfamethoxazole Using Various Irradiation Sources and Wavelength Ranges -

- Insights from Degradation Product Analysis and Compound-Specific Stable Isotope Analysis. *Environmental Science and Technology*, 52(3), 1225–1233.
- Williams, R. J., Johnson, A. C., Smith, J. J. L., Kanda, R. (2003). Steroid estrogens profiles along river stretches arising from sewage treatment works discharges. *Environmental Science and Technology*, 37(9), 1744–1750.
- Winker, M., Clemens, J., Reich, M., Gulyas, H., Otterpohl, R. (2010). Ryegrass uptake of carbamazepine and ibuprofen applied by urine fertilization. *Science of The Total Environment*, 408(8), 1902–1908.
- Writer, J. H., Antweiler, R. C., Ferrer, I., Ryan, J. N., Thurman, E. M. (2013). In-stream attenuation of neuro-active pharmaceuticals and their metabolites. *Environmental Science and Technology*, 47(17), 9781–9790.
- Writer, J. H., Ryan, J. N., Keefe, S. H., Barber, L. B. (2012). Fate of 4-nonylphenol and 17 β -estradiol in the Redwood River of Minnesota. *Environmental Science and Technology*, 46(2), 860–868.
- Wu, C., Huang, X., Lin, J., Liu, J. (2015). Occurrence and Fate of Selected Endocrine-Disrupting Chemicals in Water and Sediment from an Urban Lake. *Archives of Environmental Contamination and Toxicology*, 68(2), 225–236.
- Wu, C., Spongberg, A. L., Witter, J. D., Sridhar, B. B. M. (2012). Transfer of wastewater associated pharmaceuticals and personal care products to crop plants from biosolids treated soil. *Ecotoxicology and Environmental Safety*, 85, 104–109.
- Wu, C., Spongberg, A. L., Witter, J. D. (2009). Adsorption and Degradation of Triclosan and Triclocarban in Soils and Biosolids-Amended Soils. *Journal of Agricultural and Food Chemistry*, 57, 4900–4905.
- Wu, C., Spongberg, A. L., Witter, J. D., Fang, M., Czajkowski, K. P. (2010). Uptake of Pharmaceutical and Personal Care Products by Soybean Plants from Soils Applied with Biosolids and Irrigated with Contaminated Water. *Environmental Science & Technology*, 44(16), 6157–6161.
- Wu, L., Gao, H.-W., Gao, N., Chen, F.-F., Chen, L. (2009). Interaction of perfluorooctanoic acid with human serum albumin. *BMC Structural Biology*, 9(1), 31.
- Xing, L., Spitler, J. D. (2017). Prediction of undisturbed ground temperature using analytical and numerical modeling. Part I: Model development and experimental validation. *Science and Technology for the Built Environment*, 23(5), 787–808.

- Xu, B., Mao, D., Luo, Y., Xu, L. (2011). Sulfamethoxazole biodegradation and biotransformation in the water-sediment system of a natural river. *Bioresource Technology*, 102(14), 7069–7076.
- Xu, J., Wu, L., Chang, A. C. (2009). Degradation and adsorption of selected pharmaceuticals and personal care products (PPCPs) in agricultural soils. *Chemosphere*, 77(10), 1299–1305.
- Yager, T. J. B., Furlong, E. T., Kolpin, D. W., Kinney, C. A., Zaugg, S. D., Burkhardt, M. R. (2014). Dissipation of Contaminants of Emerging Concern in Biosolids Applied to Nonirrigated Farmland in Eastern Colorado. *JAWRA Journal of the American Water Resources Association*, 50(2), 343–357.
- Yagi, N., Kenmotsu, H., Sekikawa, H., Takada, M. (1991). Studies on the Photolysis and Hydrolysis of Furosemide in Aqueous Solution. *Chemical & Pharmaceutical Bulletin*, 39, 454–457.
- Yamamoto, H., Nakamura, Y., Moriguchi, S., Nakamura, Y., Honda, Y., Tamura, I., ... Sekizawa, J. (2009). Persistence and partitioning of eight selected pharmaceuticals in the aquatic environment: Laboratory photolysis, biodegradation, and sorption experiments. *Water Research*, 43(2), 351–362.
- Yoo, H., Washington, J. W., Jenkins, T. M., Ellington, J. J. (2011). Quantitative determination of perfluorochemicals and fluorotelomer alcohols in plants from biosolid-amended fields using LC/MS/MS and GC/MS. *Environmental Science and Technology*, 45(19), 7985–7990.
- Zhang, N., Li, J. M., Liu, G. G., Chen, X. L., Jiang, K. (2017a). Photodegradation of diclofenac in aqueous solution by simulated sunlight irradiation: Kinetics, thermodynamics and pathways. *Water Science and Technology*, 75(9), 2163–2170.
- Zhang, Y., Biswas, A., Adamchuk, V. I. (2017b). Implementation of a sigmoid depth function to describe change of soil pH with depth. *Geoderma*, 289, 1–10.
- Zhang, N., Liu, G., Liu, H., Wang, Y., He, Z., Wang, G. (2011). Diclofenac photodegradation under simulated sunlight: Effect of different forms of nitrogen and Kinetics. *Journal of Hazardous Materials*, 192(1), 411–418.
- Zhang, W., Webster, E. P. (2002). Shoot and root growth of rice (*Oryza sativa*) in response to V-10029. *Weed Technology*, 16(4), 768–772.
- Zhang, Y., Xu, J., Zhong, Z., Guo, C., Li, L., He, Y., ... Chen, Y. (2013). Degradation of sulfonamides antibiotics in lake water and sediment. *Environmental Science and Pollution Research*, 20(4), 2372–2380.

- Zhang, Y., Sangster, J. L., Gauza, L., Bartelt-Hunt, S. L. (2016). Impact of sediment particle size on biotransformation of 17 β -estradiol and 17 β -trenbolone. *Science of the Total Environment*, 572, 207–215.
- Zhao, S., Fan, Z., Sun, L., Zhou, T., Xing, Y., Liu, L. (2017a). Interaction effects on uptake and toxicity of perfluoroalkyl substances and cadmium in wheat (*Triticum aestivum* L.) and rapeseed (*Brassica campestris* L.) from co-contaminated soil. *Ecotoxicology and Environmental Safety*, 137(July 2016), 194–201.
- Zhao, S., Fang, S., Zhu, L., Liu, L., Liu, Z., Zhang, Y. (2014). Mutual impacts of wheat (*Triticum aestivum* L.) and earthworms (*Eisenia fetida*) on the bioavailability of per fluoroalkyl substances (PFASs) in soil. *Environmental Pollution*, 184, 495–501.
- Zhao, W., Wang, B., Wang, Y., Deng, S., Huang, J., Yu, G. (2017b). Deriving acute and chronic predicted no effect concentrations of pharmaceuticals and personal care products based on species sensitivity distributions. *Ecotoxicology and Environmental Safety*, 144(February), 537–542.
- Zuo, Y., Zhang, K., Zhou, S. (2013). Determination of estrogenic steroids and microbial and photochemical degradation of 17 α -ethinylestradiol (EE2) in lake surface water, a case study. *Environmental Sciences: Processes and Impacts*, 15(8), 1529–1535.