Bacterial community analysis for investigating bacterial transfer from tonsils to the pig carcass

Jakobsen, Anne Mette; Bahl, Martin Iain; Buschhardt, Tasja; Hansen, Tina Beck; Al-Soud, Waleed Abu; Brejnrod, Asker D.; Sørensen, Søren J.; Nesbakken, Truls; Aabo, Søren

Published in:
International Journal of Food Microbiology

Link to article, DOI:
[10.1016/j.ijfoodmicro.2019.02.003](https://doi.org/10.1016/j.ijfoodmicro.2019.02.003)

Publication date:
2019

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Jakobsen, A. M., Bahl, M. I., Buschhardt, T., Hansen, T. B., Al-Soud, W. A., Brejnrod, A. D., Sørensen, S. J., Nesbakken, T., & Aabo, S. (2019). Bacterial community analysis for investigating bacterial transfer from tonsils to the pig carcass. *International Journal of Food Microbiology*, 295, 8-18.
<https://doi.org/10.1016/j.ijfoodmicro.2019.02.003>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 Bacterial community analysis for investigating bacterial transfer from tonsils to the pig carcass

2

3 Anne Mette Jakobsen^a, Martin Iain Bahl^a, Tasja Buschhardt^a, Tina Beck Hansen^a, Waleed Abu Al-Soud^c,
4 Asker D. Brejnrod^c, Søren J. Sørensen^c, Truls Nesbakken^b and Søren Aabo^{a*}

5 ^a Technical University of Denmark – National Food Institute, Kemitorvet, Building 202, DK-2800 Kgs. Lyngby,
6 Denmark

7 ^b Norwegian University of Life Sciences, Faculty of Veterinary Medicine, Campus Oslo, Dept. of Food Safety
8 and Infection Biology, P. O. Box 8146 Dep., 0033 Oslo, Norway

9 ^c University of Copenhagen – Department of Biology, Section of Microbiology, Universitetsparken 15, DK-
10 2100 København Ø

11 *Corresponding author

12

13

14

15

16 Keywords: Pig slaughter, indicator, microbiota, 16S rRNA gene sequencing, cross-contamination

17 **Abstract**

18 Tonsils in the oral cavity are an important source of contamination during pig slaughter, but have not
19 received as much attention as faecal contamination. In the present study, ten pigs were sampled from
20 tonsils, faeces and three different areas on each carcass. The samples were analysed by both culturing of
21 *Escherichia coli* and *Yersinia enterocolitica* and by 16S rRNA gene sequencing to characterize the bacterial
22 communities. Comparing culture data from deep tonsil tissue and tonsil surface showed similar numbers of
23 *E. coli* but significantly higher numbers of *Y. enterocolitica* in the deep tissue samples. Microbiota analysis
24 showed similar bacterial communities in the two sample types at phylum level, while comparison at genus
25 level showed significant differences between the relative abundance of several genera in the two sample
26 types. The finding of a significantly higher relative abundance of *Yersinia* in tonsil tissue compared to tonsil
27 surface supported the culture analysis. The microbiota analysis also investigated characteristics of the
28 bacterial community that could discriminate bacterial transfer from tonsils and faeces to the carcass during
29 slaughter. The microbiota analyses demonstrated that Fusobacteria and Proteobacteria are the most
30 abundant phyla in tonsils, while Firmicutes showed the highest relative abundance in faeces. The dominating
31 phylum on carcasses was Proteobacteria. Besides Proteobacteria, the swabbing area on the forepart of the
32 carcass, showed a higher relative abundance of Firmicutes and Fusobacteria compared to swabbing areas
33 on the rear part and mid-section of the carcass. Principal coordinate analysis showed clear clustering of
34 samples based on sample source (tonsils, faeces and carcass). Carcass swab samples from the forepart
35 tended to cluster closer to the tonsil samples compared to carcass swab samples from the rear part and mid-
36 section. Identification of the genera *Fusobacterium*, *Moraxella*, *Actinobacillus* and non-*E. coli* genera of the
37 family Enterobacteriaceae in carcass swabs could indicate tonsil contamination, while faecal contamination
38 would more likely include higher prevalence of bacteria belonging to the class of Clostridia. The present
39 study supports that it is possible to identify bacterial groups that are indicative for either tonsil or faecal
40 carcass contamination. The level and composition of Enterobacteriaceae on the carcasses did, however,
41 indicate that other sources of meat contamination than tonsils and faeces may be important, such as the
42 process environment.

43

44 1 Introduction

45 During pig slaughter, cross-contamination and transfer of bacteria can easily occur from e.g. the
46 gastrointestinal tract or the oral cavity to the rest of the carcass (Borch et al., 1996). The risk of
47 contamination from the gastrointestinal tract is commonly considered important, due to the very high
48 bacterial load in this compartment. However, it is also important to consider the oral cavity, containing the
49 palatine tonsils, as a source of bacterial contamination of the carcass. Until now, methods to distinguish
50 between the various sources of bacterial contamination of carcasses during slaughter have been restricted
51 to culture-based techniques. These have inherent low resolution due to limitations in culturing and typing of
52 many bacterial groups leading to a lack in information on the relative contribution of the different sources.

53 The tonsils are known to be colonized by a diverse bacterial community, which includes both pathogenic and
54 non-pathogenic species and serve the role of presenting the immune system to foreign substances including
55 bacteria (Fredriksson-Ahomaa et al., 2009; Horter et al., 2003; Lowe et al., 2011). Bacterial colonisation of
56 the tonsils may further facilitate entry into the animal (Horter et al., 2003). During slaughter and dressing of
57 carcasses, it is common practise to recommend that the tonsils are not cut through (Anonymous, 2014).
58 Consequently, the tonsils should follow the plucks (the red offal) and not remain on the head/carcass after
59 evisceration, since the splitting procedure might lead to contamination of the head and potentially transfer of
60 bacteria to other areas of the carcass. Specific pathogenic bacteria such as *Salmonella enterica*, *Listeria*
61 *monocytogenes*, *Campylobacter* spp. and *Yersinia enterocolitica* have all previously been isolated from pig
62 tonsils by use of culture-dependent methods (Fredriksson-Ahomaa et al., 2009; Horter et al., 2003;
63 Nesbakken et al., 2003). Recently, the use of culture-independent DNA sequencing methods has provided
64 much deeper knowledge concerning the natural microbial diversity of tonsils. Accordingly, Lowe et al. (2012)
65 defined the 'core microbiome' of the palatine tonsils in pigs to include the bacterial families Pasteurellaceae,
66 Moraxellaceae, Fusobacteriaceae, Veillonellaceae, Peptostreptococcaceae and Streptococcaceae.

67 In addition to pathogens as *Salmonella* spp. and *Y. enterocolitica*, often described in relation to pig slaughter,
68 the tonsils have also been shown to contain a significant proportion of bacteria belonging to the phylum
69 Fusobacteria, which include members known to have the potential of causing illness in both humans and
70 animals (Aliyu et al., 2004; Fredriksson-Ahomaa et al., 2009; Langworth, 1977; Lowe et al., 2012).

71 The main objective of the present study was to investigate whether bacterial transfer from tonsils and faeces
72 to predefined areas of the pig carcass during pig slaughter could be elucidated by comparing systematic
73 differences between bacterial communities in faeces and tonsils to the composition of the microbiota on the
74 carcass. We further explored the possibility to identify bacterial communities, which can indicate the origin of
75 the contamination. To enable this, the bacterial communities in samples from tonsils, faeces and three
76 different areas on the carcass from the same pig were analysed by 16S rRNA gene sequencing. In parallel,
77 specific bacterial species, namely *Y. enterocolitica* and *E. coli*, were also enumerated by culturing. Carcass
78 contamination from the tonsils could differ, depending on whether tonsils are damaged or not during
79 slaughter. Therefore, differences between the bacterial community on the surface and in the deep tissue of
80 the tonsils were also investigated.

81 **2 Material and methods**

82 **2.1 Collection of samples**

83 The investigation was carried out during October and November 2013 and consisted of two sampling rounds.
84 First, for comparison of the microbiota within the tonsil tissue and on the tonsil surface, respectively, tonsils
85 were collected from pigs in two Norwegian slaughterhouses with a slaughter capacity of approximately 120 –
86 150 pigs per hour. Each slaughterhouse was visited once. Between three and 26 pigs were sampled from
87 each of five different herds. Tongue and tonsils were taken from the pluck using sterile equipment just after
88 evisceration and placed into sterile plastic bags. All samples were kept under cold conditions (approximately
89 5 °C) in the slaughterhouse and during transport to the laboratory. A total of 35 tonsils were included for
90 analysis following visual inspection of integrity in the laboratory.

91 Second, for the comparison of the microbial community of faeces, tonsils and three different carcass areas,
92 samples were collected from ten additional pigs, two pigs from each of five herds, in a single Norwegian
93 slaughterhouse. Tonsils were retrieved as described above. Faecal samples were collected by cutting off a
94 10 – 15 cm section of the rectum and placing it in a sterile plastic bag. For carcass swabs, sterile gauze
95 swabs (4 ply) were placed in sterile stomacher bags and pre-moistened with 7 ml peptone water (1 g Bacto
96 Peptone (Difco, 0118-17, Fisher Scientific, Denmark) and 8.5 g NaCl per litre). Carcass swabs were taken
97 just before cooling from three areas on one half of each carcass (Figure 1). Area A: approx. 400 cm² located

98 on the inner side of the rear section of the carcass and around the rectum, Area B: approx. 1,000 cm²
99 stretching from the forepart to the rear section on the inner side of the carcass along the backbone on the cut
100 surface and area C: approx. 400 cm² on the inner side of the forepart around the jaw and throat. The areas
101 were selected due to the putative higher possibility of contamination from the oral cavity on area C,
102 contamination from the saw cutting through the head on area B and contamination from the rectum on area
103 A. A single swab sample was additionally taken from the saw used to cut the carcass. All samples were
104 placed in a cooling box with cooling elements and kept cold until further analyses in the laboratory.

105 **2.2 Bacteriological analysis**

106 **2.2.1 Sample preparation**

107 In the laboratory, both palatine tonsils were removed from the root of the tongue using sterile instruments
108 and transferred to a stomacher bag with filter (BagFilter® No. 8, WWR, Denmark). Peptone water was added
109 to the stomacher bag (1:1 w/v) and following 20 s of stomaching, the tonsils were removed from the
110 stomacher bag and the liquid collected for analysis. To expose bacteria located in the deep tonsil tissue, the
111 same tonsils were subsequently sliced thoroughly with a sterile scalpel. The slices were transferred to a new
112 stomacher bag with filter. Peptone water was added (1:1 w/v) and the samples stomached for 20 s and liquid
113 collected for analysis. For the samples from the ten pigs, tonsil surface was prepared for analysis as
114 described above. Faeces were collected from the rectum sections and diluted with peptone water (1:10 w/v)
115 and transferred to stomacher bags with filter, and stomached for 20 s. For carcass and saw swabs, a volume
116 of 20 ml peptone water was added to the stomacher bags and samples were homogenised for 20 s.

117 **2.2.2 Culture dependent analysis for *Yersinia enterocolitica***

118 All samples were analysed quantitatively for *Y. enterocolitica* by direct plating on cefsulodin-irgasan-
119 novobiocin (CIN) agar plates (CM653 + SR109 Oxoid, Fisher Scientific, Denmark) and *Y. enterocolitica*
120 chromogenic medium (YeCM) (CHROMagar, SmithMed, Norway). In parallel, ten-fold dilutions from all
121 samples were enriched in irgasan-ticarcillin-potassium chlorate (ITC) broth (Sigma–Aldrich, Denmark) and
122 analysed semi-quantitatively by plating on CIN and YeCM agar. The qualitative and the quantitative analysis
123 followed essentially the standard procedures of NMKL 117 3rd edition but were modified by using YeCM as
124 an additional indicative medium.

125

126 **2.2.3 Identification and characterisation of *Yersinia enterocolitica***

127 Presumptive colonies were sub-cultured on bromthymol-blue sucrose agar on which *Y. enterocolitica*
128 produce yellow colonies due to the fermentation of sucrose. Further biotyping was based on the revised
129 scheme of Wauters et al. (1987) using tests for esculin hydrolysis and indole production from tryptophan
130 together with a test for urea formation on urea agar (Difco; 0283-02-6 Bacto Urea agar Base, 0140-01 Bacto
131 Agar, Fisher Scientific, Denmark). Finally, isolates of Biotype 4 were serotyped by use of slide agglutination
132 with commercial O:3 antiserum (Bio-Rad; Statens Serum Institute, Denmark).

133 **2.2.4 Culture analysis for *Escherichia coli***

134 Samples obtained from tonsils, faeces and carcass swabs were analysed quantitatively for *E. coli* by use of
135 the SimPlate method (AOAC International, 2005). From appropriate 10-fold dilutions, 1 ml was placed in the
136 centre of the plating device (SimPlate Devices, Part No: 65009-20, Biocontrol, Germany) and 9 ml of blue
137 mixed nutrient agar (Multi Test, Part No: 66008-500, Biocontrol, Germany) was added in the same spot. The
138 sample was mixed gently in the SimPlate device according to the instruction of the supplier and the plates
139 were incubated at 37 °C for 24 – 28 h. Wells with a change in colour from blue to red and fluorescent when
140 exposed to UV-light at 366 nm, were counted as positives. Quantification of *E. coli* per ml was calculated
141 based on the most probable number (MPN) principle.

142 **2.3 Bacterial community analysis by 16S rRNA gene sequencing**

143 From each sample, 10 ml homogenate was transferred to a 15 ml centrifuge tube and centrifuged at 10 °C
144 for 10 min at 4500 rpm (Beckman Coulter, Allegra X-22R centrifuge, Minnesota, US). The supernatant was
145 discarded and the pellet was carefully re-suspended in 1 ml LB medium containing 15 % glycerol. One ml of
146 the cell suspension was transferred to a 1.5 ml Eppendorf tube and stored at -80 °C until DNA extraction.
147 DNA was extracted from the thawed cell suspensions using the PowerLyzer® PowerSoil® DNA Isolation Kit
148 (Mo Bio Laboratories, Carlsbad, CA, US) following the manufacturer instructions, modified by using a Mixer
149 Mill MM 300 (Retsch GmbH, Haan, Germany) for 10 min at 30 cycles per second for cell lysis. DNA extracts
150 were kept frozen at -20 °C until further analyses (approx. after two months).

151 The 16S rRNA gene amplification procedure followed a two-step strategy, in which the first amplified the V3-
152 V4 variable regions from the 16S rRNA gene and the second attached Illumina sequencing sample specific
153 adaptors. In the first PCR reaction, the universal primers PBU (5'-CCT ACG GGA GGC AGC AG-3')

154 (Tulstrup et al., 2015) and PBR-long (5'-GGA CTA CCA GGG TAT CTA ATC-3') (developed in this study)
155 were used to amplify approximately 466 base pairs (V3-V4 region) of the 16S rRNA gene. The reaction
156 mixture consisting of 4 µl 10X AccuPrime PCR Buffer II, 0.8 U AccuPrime Taq DNA Polymerase (Invitrogen,
157 Life technologies, CA, US), 0.4 µM primer PBU, 0.4 µM primer PBR-long, 0.8 µg bovine serum albumin
158 (BSA) and 10 µl template DNA, in a total volume of 40 µl per sample. The DNA extracts from the faecal
159 samples were diluted ten-fold before use as templates in the PCR reaction. The PCR program consisted of
160 an initial denaturation at 94 °C for 2 min, followed by 30 cycles amplification at 94 °C for 20 sec, annealing at
161 58 °C for 20 sec and extension at 68 °C for 1 min, followed by a final extension at 68 °C for 5 min and
162 cooling to 4 °C. The size and quality of the PCR products were assessed by gel electrophoreses, and
163 purified with QIAquick® PCR Purification kit (Qiagen, Valencia, CA, US) according to the manufacturer
164 instructions. The DNA concentration of the purified PCR products was measured with a Qubit® fluorometer
165 (Thermo Fisher Scientific) and adjusted to 5 ng/µl DNA for use as template for the second PCR amplification.

166 The second PCR amplification was carried out in 96-well microtiter plates using sequencing primers
167 originally published by Yu et al. (2005) and modified by Sundberg et al. (2013) as follows: 341F (5'-CCT
168 AYG GGR BGC ASC AG-3') and 806R (5'-GGA CTA CNN GGG TAT CTA AT-3'), both of which with 5'
169 Illumina sequencing adaptors and index-tags as previously described (Mortensen et al., 2016; Thorsen et al.,
170 2016). The reaction mix contained 2.0 µl of the diluted amplicons (5 ng/µl) 2 µl 10x AccuPrime PCR Buffer II,
171 0.6U AccuPrime Taq DNA Polymerase (Invitrogen, Life technologies, CA, US) and 0.5 µM fusion 341F and
172 0.5 µM fusion 806R in a total volume of 20 µl. The PCR was run as above, with exception of a reduced
173 annealing temperature of 56 °C and 15 amplification cycles.

174 **2.3.1 DNA sequencing**

175 The amplification products of the second PCR were purified with Agencourt AMPure XP Beads (Beckman
176 Coulter Genomics, MA, US) following the manufacturer instructions using 0.7 X volume beads and quantified
177 as described above. Equimolar amounts of all the amplification products were pooled together in a single
178 tube and concentrated using the DNA Clean & Concentrator TM-5 Kit (Zymo Research, Irvine, CA, US)
179 following the manufacturer instructions. The concentration of the pooled library was determined using the
180 Quant-iT™ High-Sensitivity DNA Assay Kit and Qubit® fluorometer (Life Technologies) following the
181 manufacturer instructions. Amplicon sequencing was performed on the Illumina MiSeq Desktop Sequencer

182 (Illumina Inc., CA, US). For each run, a 1 – 5 % PhiX DNA was added. All reagents used were from the
183 MiSeq Reagent Kits v2 (Illumina Inc., CA, US). Automated cluster generation and 250 paired-end
184 sequencing with dual-index reads were performed. Up to 192 samples were sequenced per run. Technical
185 replicates of the sequencing were not carried out in this study. In parallel studies, the variation between
186 technical replicates has been addressed and found to be minimal (data not shown).

187

188 **2.3.2 Bioinformatics on bacterial community data**

189 Raw fastq-files demultiplexed using the MiSeq Controller Software (Illumina Inc., CA, US) were processed
190 with BioDSL (<https://github.com/maasha/BioDSL>). Primers and diversity spacers were identified and
191 truncated with the function trim_primer(). Paired ends were mated using the function assemble_pairs(). Any
192 sequences shorter than 100 bp were discarded and remaining sequence reads were dereplicated using
193 dereplicate_seq. The sequences were clustered at 97 % identity with cluster_otus(), a wrapper around
194 USEARCH ver. 7.0.1090 (Edgar, 2010). Chimeras were checked and removed with uchime_ref() against the
195 GOLD database (Haas et al., 2011). The resulting sequences were classified using classify_seq against the
196 Ribosomal Database Project (RDP) trainset9 database (032012) available from the Mothur website.
197 Align_seq_mothur (Schloss et al., 2009), a wrapper around Mothurs align.seq() function for aligning
198 Operational Taxonomic Units (OTUs) against an RDP template. The Mothur version used was v.1.33.3.
199 Downstream processing of sequence data in the generated OTU table was performed in QIIME (Caporaso et
200 al., 2010) using the core_diversity_analysis.py script. For diversity analysis, a phylogenetic tree was
201 generated in QIIME using the make_phylogeny.py script based on a PyNAST alignment of all identified
202 OTUs including an archaea for rooting. The OTU-table was first filtered to include only those OTUs
203 annotated to Bacteria and to exclude OTUs with a number of sequences below 0.005 % of the total number
204 of sequences (Bokulich et al., 2012). Samples were rarefied to 10,000 reads per sample for analysis and
205 those samples with fewer reads were excluded. The relative abundance at phylum level was calculated
206 using the collapse_samples.py function. Phyla present in less than 1 % in any sample type was compiled in
207 the phyla 'other'. Alpha diversity, describing the bacterial diversity within a sample, was calculated for each
208 sample by the Shannon index and species diversity (based on OTU richness). Assessment of beta diversity,
209 describing the overall diversity in the community composition between sample types, was performed by
210 Principal Coordinate Analysis (PCoA) based on unweighted UniFrac distances. Variation within and between

the different sample types were calculated based on the unweighted UniFrac distances by use of the `make_distance_boxplots.py` function. Heatmaps were used to visualize relative bacterial abundances at the genus level for individual samples across sample types. The heatmaps were based on the relative abundance of the genera present in at least 35 % of all samples, with filtering performed with the `filter_otus_from_otu_table.py` script. Additionally, bacteria which could not be classified at the genus level were excluded from the heatmaps. Similarity between community composition at the operational taxonomic unit (OTU) level in tonsils, faeces and the three areas on the carcass was assessed using the `shared_phylotypes.py` script.

2.5 Statistics

For the culture data, statistics were conducted on $\log_{10}$ transformed data in the Microsoft Excel (2010) Data analysis add-in. A paired t-test was used for the comparison of the number of *E. coli* and *Y. enterocolitica* present in samples from tonsil surface and deep tonsil tissue and for the comparison of recoveries from the growth media CIN and YeCM. A single factor analysis of variance (ANOVA) was conducted for comparison of *E. coli* numbers detected in tonsils, faeces and on the carcass of ten pigs. A significance level threshold of $p < 0.05$ was applied in general.

Most statistical tests related to the sequencing data were conducted in GraphPad Prism (version 7, GraphPad Software Inc., La Jolla, CA, US). *P*-values calculated for multiple comparisons were adjusted by the False Discovery Rate (FDR) (Benjamini and Hochberg, 1995). A paired t-test was used for the overall comparison of the bacterial community composition on phylum level of samples from tonsil surface and deep tonsil tissue, while an unpaired t-test was used to compare the alpha diversity measures (Shannon index and species diversity) calculated for the two sample types. One-way ANOVA with multiple comparisons (Tukey's multiple comparisons test) was used to compare the alpha diversity measures (Shannon index and species diversity) calculated for the paired samples from tonsils, faeces and carcass swabs. The test was also used to assess the variation within and between different samples types and for the comparison of the mean relative abundance of specific genera in the paired samples from tonsils, faeces and carcass swabs. A one-way ANOVA of repeated measures and with multiple comparisons was used for comparison of the shared OTUs found in tonsils, faeces and on the three carcass swab areas.

For the heatmap comparing the community on tonsil surface to that of tonsil tissue (Figure 5), pairwise comparisons of the $\log_{10}$ transformed relative abundances were conducted by use of permutation tests with 10,000 iterations and calculation of the FDR-adjusted p -values. These calculations were conducted in the software R (ver. 2.15.1, R Core Team 2014, Vienna, Austria) and RStudio (ver. 0.96.331) using in-house scripts.

3. Results

3.1 Culture analyses

The bacterial community structure of tonsil surface and deep tonsil tissue was investigated in tonsils from 35 pig carcasses by both culture- and 16S rRNA gene sequencing-based analysis. The occurrence of *E. coli* and *Y. enterocolitica* on surface and in deep tissue of the tonsils is listed in Table 1. The number of *Y. enterocolitica* was found to be significantly higher in the deep tonsil tissue ($p < 0.05$) compared to surface. All *Y. enterocolitica* suspected colonies were identified as biotype 4, serotype O:3. As no significant difference ($p = 0.67$, paired t-test) was seen between the number of *Y. enterocolitica* found on CIN and YeCM, only results from CIN are presented.

From the three different areas on the carcass, mean levels of *E. coli* and *Y. enterocolitica* were between 0.1 and 4.0 CFU/cm², while the mean numbers in tonsils and faecal samples were 4.9 and 5.4 $\log_{10}$ CFU/g, respectively (Table 2). Significantly lower levels of *E. coli* were found on area B than on both area A and C. *Y. enterocolitica* was detected only in one faecal sample whereas seven tonsil samples had levels above the detection level (Table 2). *Y. enterocolitica* was not detected in any swab samples from area A, B and C by direct plating or following selective enrichment.

3.2 Comparison of bacterial communities on tonsil surface and in deep tonsil tissue

Analyses of the bacterial community structure by 16S rRNA gene sequencing were performed with a mean sequencing depth of $27,671 \pm 12,234$ reads per sample. Samples were subsampled to 10,000 reads for further analysis. None of the samples from tonsil surface and deep tonsil tissue were, thereby, discarded from the analysis.

At the phylum level, no significant differences in relative abundance between the six phyla found on tonsil surface and in deep tonsil tissue were seen (Figure 2a). The bacterial community in tonsils were estimated to

265 consist of approximately 30 – 40 % Proteobacteria, 30 % Firmicutes, 20 % Fusobacteria and 10 – 20 %
 266 Bacteroidetes. The phyla Spirochaetes and Actinobacteria had mean relative abundances below 3 %. No
 267 significant difference in alpha diversity was seen between the surface of tonsils and the deep tonsil tissue
 268 (Shannon index; $p = 0.59$ and observed species; $p = 0.56$) (Figure 3a-b). Analysis of differences in bacterial
 269 community between sample types (beta diversity) by principal coordinate analysis (PCoA) based on
 270 unweighted UniFrac distances, indicated some separation between the two sample types (Figure 4a), which
 271 was supported by statistical analysis of the differences in UniFrac distances within and between sample
 272 types ($p < 0.05$) (Figure 4b). The distribution of relative abundance of 63 different genera in tonsil surface
 273 samples and samples from deep tonsil tissue were visualised in a heatmap (Figure 5). Subsequent statistical
 274 analysis revealed 18 genera to be significantly different between the two sample types, including a significant
 275 higher relative abundance of *Yersinia* spp., and *Pasteurella* spp. in deep tonsil tissue compared to tonsil
 276 surfaces and significantly higher relative abundance of *Actinobacillus* spp. on tonsil surface. (FDR adjusted- p
 277 < 0.05).

278 **3.3 Comparison of bacterial communities in tonsils and faeces to communities in carcass swab** 279 **samples from 10 pigs**

280 When using a cut-off value of 10,000 reads, two faecal samples and one carcass swab sample from area A
 281 and B, respectively, were discarded from down-stream analysis. The remaining samples had an average
 282 sequencing depth of $28,440 \pm 11,710$ reads per sample.

283 The average distribution of phyla in the tonsil samples (Figure 2b) appeared similar to what was found in the
 284 35 samples from tonsil surface and deep tonsil tissue (Figure 2a). The faecal samples showed a mean
 285 relative abundance of Firmicutes and Bacteroidetes of approximately 80 % and 20 %, respectively,
 286 representing almost all sequences. The carcass swab samples showed increasing mean relative
 287 abundances of Proteobacteria from area C (50 %), to area B (71 %) to area A (86 %) while the relative
 288 abundance of both Firmicutes and Bacteroidetes were lowest in samples from area A (10 % and 2 %, respectively)
 289 and higher in samples from area B (15 % and 11 %, respectively) and highest in area C (27 % and 13 %, respectively). The mean relative abundance of Fusobacteria was also highest in samples from
 290 area C (6 %).

292 The alpha diversity (Shannon-index) showed an overall significant difference ($p < 0.05$) between the sample
293 types with the diversity in sample area A found to be significantly lower than areas B and C (Figure 3c). No
294 significant difference ($p = 0.08$) was seen between the richness of observed species in the different sample
295 types (Figure 3d).

296 Principal coordinates analysis (PCoA) of the bacterial communities at the OTU level in tonsils, faeces and in
297 the three different carcass swab samples showed clustering of the different communities (Figure 4c). In the
298 PCoA plot, especially, the faecal and tonsil samples clustered separately while the carcass swabs clustered
299 together. Visually, samples from area C clustered slightly closer to the tonsil samples compared to samples
300 from the two other sample areas of the carcass. The single sample from the saw was included in the PCoA.
301 Faecal samples clustered more closely together than other sample types, which is supported by low intra-
302 sample average distance (Figure 4d). No significant differences were found between mean distances within
303 the tonsil samples and within samples from area A and B, respectively, while a significant difference was
304 found between the tonsil samples and the samples from area C ($p < 0.05$). A significant difference was also
305 found between the mean distances within samples from area B and area C ($p = 0.02$). Although, the overall
306 mean distances varied within the samples, average distances within all sample types were found to be
307 significantly lower than the overall mean distances between all sample types ($p < 0.05$).

308 Looking at the distribution of OTUs shared between sample sites, we found that on average 43 % of the
309 OTUs present in tonsils were also present in samples from area C around the head (Figure 6a), 45 % were
310 found in samples from area A around rectum and around 30 % were found in samples from area B along the
311 backbone on the cut surface. Significantly less tonsil OTUs were shared with the OTU's from faecal samples
312 compared to the sharing with the carcass swab samples ($p < 0.05$). From the faecal samples on average 48
313 % of the OTUs were found in samples from area A around the rectum, around 40 % in samples from area B
314 and 36 % in samples from area C around the head (Figure 6b).

315 The relative distribution of the bacterial communities in the different sample types were visualised at the
316 genus level in a heatmap (Figure 7). When compared to the other sample types, the tonsil samples showed
317 higher relative abundance of genera such as *Fusobacterium*, *Pasteurella*, *Moraxella* and *Actinobacillus*. In
318 the faecal samples we noted higher relative abundances for several of the genera belonging to the class
319 Clostridia (Firmicutes) including *Blautia* and *Roseburia*. Among the three carcass areas, area C showed the

highest relative abundance of *Yersinia*, *Fusobacterium*, and *Actinobacillus* but also most genera within the class Clostridia. Area A showed higher relative abundance of for *Escherichia/Shigella* compared to the two other areas. Overall, the sample from the saw showed low relative abundance of the phyla Actinobacteria, Bacteroidetes and Firmicutes and high relative abundances of genera within Proteobacteria. The sample from the saw e.g. showed the highest relative abundance of *Salmonella* and *Escherichia/Shigella* among all sample types. Significantly higher mean relative abundance of *Yersinia* was found in samples from area C compared to the other sample types ($p < 0.05$) (Figure 8a), while the mean relative abundance of *Escherichia/Shigella* in samples from area A was significantly higher than what was found in samples from tonsils, faeces and area B ($p < 0.023$) (Figure 8b). The tonsils were observed to have a significantly higher mean relative abundance of *Fusobacterium* ($p \leq 0.05$) compared to all the other sample types (Figure 8c).

4. Discussion

Tonsil surface and deep tonsil tissue showed overall comparable communities at the phylum level (Figure 2a). In contrast, when analysing at the more detailed genus level, 18 of the 63 compared genera were found to be significantly different in relative abundance, between the two sample types (Figure 5), which was supported by analysis of unweighted UniFrac distances (Figure 4a). The significant difference in relative abundance of *Yersinia* was consistent with the significantly higher number of *Y. enterocolitica* cultured in the deep tonsil tissue samples compared to tonsil surface. This is coherent with the ability of *Y. enterocolitica* to colonise the tonsils (Thibodeau et al., 1999). Lowe et al. (2012) also compared the bacterial communities from pig tonsil tissue and brush samples from the surface of tonsils. They primarily found differences between the communities in the species within the Enterobacteriaceae family and the obligate anaerobes, which they found more abundant in tissue samples compared to surface brush samples. This is essentially in accordance with our findings. When comparing surface associated bacteria with the deep tissue associated bacteria we did not apply surface disinfection of the tonsil surface before the analysis of the deep tissue community. A disinfectant applied on the surface may possibly have gained access to the bacterial community in the deeper tonsil tissue, thus biasing the community analysis. Also by our procedure, we cannot exclude that some bacteria, firmly attached to the surface, has been released during the second stomaching of the tonsil tissue, thus contributing to bias in the community analysis. Van Damme et al. 2018 compared the number of *E. coli* in tonsil surface to that of tonsil tissue and found similar levels of *E. coli* in tonsil tissue as in this study. In contrast to the present study, however, they recovered considerable lower

349 numbers of *E. coli* by swabbing the surface than we did by stomaching. This could imply that we released
350 more tissue associated *E. coli* into the surface homogenate in our samples. Despite this uncertainty, the
351 microbiota analysis was able to disclose significant differences between tonsil surface and deep tissue
352 associated bacteria at genus level. Collectively, we consider the employed procedure acceptable in relation
353 to the conclusions of the study.

354 Faeces may also be a source of *Y. enterocolitica* contamination, but with much lower prevalence than seen
355 for tonsils (Thibodeau et al., 1999). By culturing, we found *Y. enterocolitica* from only 10 % of the faecal
356 samples while it was found in 60 – 70 % of the tonsil samples. This is in accordance with findings of Van
357 Damme et al. (2013). A difference in carriage in faeces and in tonsils may be driven by a decrease in the
358 proportion of pigs carrying human pathogenic *Y. enterocolitica* in faeces from about 135 days of age, while
359 many of their tonsils do remain positive for *Y. enterocolitica* at the time of slaughter (Nesbakken et al., 2006).
360 Accordingly, the tonsils may constitute a more important source of carcass contamination with human
361 pathogenic *Y. enterocolitica* compared to faeces.

362 *Escherichia coli* can be present in large numbers both in tonsils (Fredriksson-Ahomaa et al., 2009) and in
363 faeces (Nauta et al., 2013) and culturing of these organisms was included in the study in order to be able to
364 compare the bacterial load of *E. coli* in the two sources (Table 2). However, both culturing and sequence-
365 based analysis revealed no difference in load of *Escherichia* per gram between the two sources. The very
366 high level of *E. coli* in tonsils may be due to ingestion of faeces.

367 If bacterial species are distributed differently in tonsils and faeces, their relative abundances may be used as
368 an indicative marker for the source of carcass contamination. Even though *Y. enterocolitica* meet this criteria
369 by being more abundant in tonsil, this study does not support it as a culturable marker for tonsil origin of
370 carcass contamination as it was not detected on any of the swabbing areas by culturing.

371 In order to ease interpretation of results, the swabbing areas A, B and C were located on the inside of the
372 carcass, as the inside is expected to be sterile until the carcass is cut open, and thus all bacteria present on
373 raw muscle tissue must have originated from cross-contamination.

374 Very low levels of *E. coli* were recovered by culturing from the swabbing areas with significantly higher
375 numbers found on area A and C around the rectum and the head, respectively compared to area B on the

376 mid part of the carcass. Thus, *E. coli* does not candidate as culturable marker for defining the relative
377 contribution of faeces and tonsils as source of contamination.

378 By use of 16S rRNA gene sequencing it was possible to analyse the source of contamination in more depth.
379 At phylum level we found the bacterial community of pig tonsils primarily to be dominated by Proteobacteria
380 (30 – 50 %) in line with the “core microbiome” previously described by Lowe et al. (2012) although they
381 found Fusobacteria less abundant. By culture-based analysis, Van Damme et al, (2018) supported this
382 observation indirectly by finding Enterobacteriaceae, being part of Proteobacteria, more dominating in tonsils
383 than in faeces. Faecal samples were vastly dominated by Firmicutes and Bacteroidetes (Figure 2b), which is
384 consistent with previous studies by Kim et al. (2015) and Pajarillo et al. (2014). At phylum level the bacterial
385 community of carcass area C mimicked the community of tonsil quite closely (Figure 2b). Like area C, the
386 areas A and B also contained a large proportion of Proteobacteria, but they did only contain a very small
387 population of Fusobacteria. Even though the relative abundances of Bacteroidetes and Firmicutes in area A
388 and B are low, the almost absence of Fusobacteria indicate faeces to be the dominating source of area A
389 rather than the tonsils. At the same time, the high abundance of Proteobacteria indicate that a third source,
390 not covered by this study, contribute to the carcass contamination. This could be the process environment.

391 As carcasses are hanged with heads down, the community at lower parts could potentially be more diverse
392 because the faecal community may blend in with the tonsil community at the forepart. A significant lower
393 sequence diversity (lower alpha diversity (Shannon index)) in samples obtained from area A compared to
394 those from area B and C (Figure 3c) supported this. Somewhat contradicting, however, no difference in the
395 number of observed species between areas A, B and C were found (Figure 3d).

396 Investigating for systematic patterns in OTU sequence data by PCoA analysis, the carcass swab samples
397 clustered together, with a tendency that samples from areas A and C clustered closer to the tonsil samples
398 than area B (Figure 4c). This, together with the fact that around 45 % of the OTUs present in the bacterial
399 community of tonsils were also found in samples from area A and C (Figure 6a) could indicate that tonsils
400 may play a role in contamination of both the forepart (area C) and of the ham area (area A). When
401 comparing sequences from the ham area (area A) to sequences from faeces and tonsils almost 50 % of the
402 OTUs from area A were present in faecal samples while only approximately 25 % were found in tonsil
403 samples (Figure 6b). The latter indicate that area A mainly is contaminated with faeces but with evidence of

404 a contribution from tonsils. The finding of 45 % of tonsil OTU's in the ham region (area A) is likely a
405 combination of contamination from tonsils and a high overlap between faecal and tonsil communities in
406 general.

407 From the heatmap (Figure 7) it appears that tonsils compared to the other sample types, were dominated
408 with genera within the family Pasteurellaceae (*Actinobacillus*, *Haemophilus* and *Pasteurella*) and the genera
409 *Moraxella*, *Alkanindiges* as well as *Fusobacterium*. As these genera were part of the 'core microbiome' in
410 porcine tonsils defined by Lowe et al. (2012) their presence could represent a general property of tonsils and
411 potentially be indicative of contamination originating from tonsils. Contamination of the area C seems to be
412 dominated by the flora from the tonsils, which is supported by a high relative abundance of *Fusobacterium*
413 and *Actinobacillus* in tonsils and in area C, *Actinobacillus* is also detected from area A and B in low levels
414 indicating tonsils as source of contamination (Figure 7).

415 The swabbing areas showed higher relative abundances of Enterobacteriaceae genera (other than *E. coli*)
416 compared to samples from tonsils and faeces, which indicates a source of contamination not fully explained
417 by faeces and tonsils as sources. If psychrotrophic Enterobacteriaceae establish and grow in the process
418 environment, as may also be the case for the genus *Aeromonas* (Figure 7), a subsequent transfer to the
419 carcass, could potentially explain the increased occurrence of Enterobacteriaceae in swab samples. The
420 distribution of shared OTUs (Figure 6) also suggests a different source than faeces and tonsils to contribute
421 to carcass contamination. The swab sample from the saw showed high relative abundances of several
422 genera within the Enterobacteriaceae family and could be an example of process equipment where
423 Enterobacteriaceae dominates e.g. due to growth.

424 The bacterial community composition of tonsils and faeces found in this study were similar to reports by Kim
425 et al. (2015); Lowe et al. (2011, 2012), and Pajarillo et al. (2014). However, it is important to mention that
426 differences in sample preparation, DNA extraction, and data handling can lead to differences in classification
427 and may challenge the ability to compare sequencing results between studies. When specifically trying to
428 compare culture-based cell counts for *E. coli* and *Y. enterocolitica* to the relative abundance of the
429 corresponding genera obtained by sequencing, the detection limit for the sequencing method is also of great
430 importance. Despite *E. coli* being detected in higher absolute numbers in faeces compared to carcass
431 samples, the sequencing did not show a higher relative abundance of *Escherichia/Shigella* in faecal samples

432 compared to the other samples (Figure 8b). This is most likely due to a higher total bacterial load in the
433 faecal samples and, thus, a higher detection limit for *E. coli* and lower deduced relative abundance when
434 sequencing.

435 The present study supports the hypothesis that it is possible to identify bacterial groups that are
436 characteristic for either tonsil or faecal carcass contamination. The level and composition of
437 Enterobacteriaceae on the carcasses indicates that other sources of contamination than tonsils and faeces
438 may be important. Future studies could include more samples from the working environment and process
439 equipment to provide a more in depth understanding of the specific sources. Our conclusions are based on
440 data from ten animals and future investigations will be needed to determine if the same systematic patterns
441 can be observed at population level. Inclusion of more slaughterhouses will also be relevant. If our findings
442 hold true on population level, it would enable a more general deduction of the sources behind bacteria found
443 on carcasses and may thus provide a more thorough basis for hygiene intervention at slaughter.

444 **Acknowledgements**

445 The present study was financed by the Danish Council for Strategic Research (DCSR, process 12-131417)
446 as a part of collaboration with the Strategic Research Collaboration in Food Science in the State of São
447 Paulo, Brazil and Denmark.

448 The Norwegian slaughterhouses are acknowledged for their kindness and help in relation to sampling.

Table 1. Prevalence and mean number for *Escherichia coli* and *Yersinia enterocolitica* on the surface and in the deep tissue of tonsils (n=35) (Results for *Yersinia* only shown for CIN-agar).

	Tonsil surface		Tonsil tissue	
	Prevalence	Mean concentration	Prevalence	Mean concentration
	[%]	[log CFU/g]± sd	[%]	[log CFU/g] ± sd
<i>E. coli</i>	100	5.5 ± 0.6	100	5.6 ± 0.4
<i>Y. enterocolitica</i> (CIN)	70.7	3.5 ± 1.2	68.3	4.2 ± 0.9

449

450

Table 2. Numbers of *Escherichia coli* (ECC) from faeces, tonsils and from swabbing areas A, B and C (see Figure 1), detected by use of the SimPlate method (AOAC International, 2005), and numbers of *Yersinia enterocolitica* (YeC) detected by culturing on CIN from the same sampling sites.

Carcass no.	Faeces		Area A		Area B		Area C		Tonsils	
	[log CFU/g]				[log CFU/cm ²]				[log CFU/g]	
	ECC	YeC	ECC	YeC	ECC	YeC	ECC	YeC	ECC	YeC
1	4.8	4.91	0.27	<-0,2	- 0.8	<-0,5	1.7	<-0,2	5.9	3.0
2	4.8	<1,0	<- 1.2*	<-0,2	- 1.5	<-0,5	1.2	<-0,2	4.5	3.0
3	5.2	<1,0	- 0.39	<-0,2	- 1.3	<-0,5	0.3	<-0,2	4.9	1.8
4	5.0	<1,0	- 0.87	<-0,2	- 1.5	<-0,5	- 0.1	<-0,2	5.1	<1.3
5	6.7	<1,0	<- 1.2	<-0,2	- 1.3	<-0,5	<- 1.2	<-0,2	4.2	1.6
6	< 4.0**	<1,0	<- 1.2	<-0,2	<- 1.5	<-0,5	<- 1.2	<-0,2	5.1	<1.3
7	5.7	<1,0	- 0.57	<-0,2	- 0.6	<-0,5	0.8	<-0,2	4.9	<1.3
8	6.9	<1,0	- 0.03	<-0,2	- 1.3	<-0,5	- 0.4	<-0,2	4.7	<1.3
9	5.9	<1,0	- 0.27	<-0,2	<- 1.5	<-0,5	- 0.4	<-0,2	4.1	3.6
10	5.2	<1,0	<- 1.2	<-0,2	<- 1.5	<-0,5	- 0.6	<-0,2	5.9	4.7
Mean	5.4 ^a		- 0.8 ^b		- 1.4 ^c		-0.1 ^b		4.9 ^a	2.2
sd***	0.9		0.6		0.4		1.0		0.6	1.3

*<: below detection limit.
** Plating started from 10⁻⁴ dilution.
*** For figures below detection limit log10 (0,5X detection limit) is used for calculation of standard deviation (sd).
For mean values different letters indicate a significant difference (5 % significance level)

453 **References**

- 454 Aliyu, S.H., Marriott, R.K., Curran, M.D., Parmar, S., Bentley, N., Brown, N.M., Brazier, J.S., Ludlam, H.,
455 2004. Real-time PCR investigation into the importance of *Fusobacterium necrophorum* as a cause of
456 acute pharyngitis in general practice. J. Med. Microbiol. 53, 1029–1035. doi:10.1099/jmm.0.45648-0
- 457 Anonymous, 2014. Commission Regulation (EU) No 218/2014 of 7 March 2014, amending Annexes to
458 Regulations (EC) No 853/2004 and (EC) No 854/2004 of the European Parliament and of the Council
459 and Commission Regulation (EC) No 2074/2005. Off. J. Eur. Communities.
- 460 AOAC International, 2005. Official Method 2005.03 Detection and Confirmed Quantitation of Coliforms and
461 *E. coli* in Foods SimPlate Coliform and *E. coli* Color Indicator. Available at:
462 http://www.aoac.org/omarev1/2005_03.pdf.
- 463 Benjamini, Y., Hochberg, Y., 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach
464 to Multiple Testing. J. R. Stat. Soc. Ser. B 57, 289–300.
- 465 Bokulich, N.A., Subramanian, S., Faith, J.J., Gevers, D., Gordon, J.I., Knight, R., Mills, D.A., Caporaso, J.G.,
466 2012. Quality-filtering vastly improves diversity estimates from Illumina amplicon sequencing. Nat.
467 Methods 10, 57–59. doi:10.1038/nmeth.2276
- 468 Borch, E., Nesbakken, T., Christensen, H., 1996. Hazard identification in swine slaughter with respect to
469 foodborne bacteria. Int. J. Food Microbiol. 30, 9–25. doi:10.1016/0168-1605(96)00988-9
- 470 Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Peña,
471 A.G., Goodrich, J.K., Gordon, J.I., Huttley, G. a, Kelley, S.T., Knights, D., Koenig, J.E., Ley, R.E.,
472 Lozupone, C. a, Mcdonald, D., Muegge, B.D., Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J.,
473 Walters, W. a, Widmann, J., Yatsunenko, T., Zaneveld, J., Knight, R., 2010. QIIME allows analysis of
474 high- throughput community sequencing data. Nat. Methods 7, 335–336. doi:10.1038/nmeth0510-335
- 475 Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics 26, 2460–
476 2461. doi:10.1093/bioinformatics/btq461

477 Fredriksson-Ahomaa, M., Gerhardt, M., Stolle, A., 2009. High bacterial contamination of pig tonsils at
478 slaughter. *Meat Sci.* 83, 334–6. doi:10.1016/j.meatsci.2009.06.004

479 Horter, D.C., Yoon, K.-J., Zimmerman, J.J., 2003. A review of porcine tonsils in immunity and disease. *Anim.*
480 *Health Res. Rev.* 4, 143–155. doi:10.1079/AHR200358

481 Haas, B.J., Gevers, D., Earl, A.M., Feldgarden, M., Ward, D. V, Giannoukos, G., Ciulla, D., Tabbaa, D.,
482 Highlander, S.K., Sodergren, E., Methé, B., DeSantis, T.Z., Petrosino, J.F., Knight, R., Birren, B.W.,
483 2011. Chimeric 16S rRNA sequence formation and detection in Sanger and 454-pyrosequenced PCR
484 amplicons. *Genome Res.* 21, 494–504. doi:10.1101/gr.112730.110

485 Kim, J., Nguyen, S.G., Guevarra, R.B., Lee, I., Unno, T., 2015. Analysis of swine fecal microbiota at various
486 growth stages. *Arch. Microbiol.* 197, 753–9. doi:10.1007/s00203-015-1108-1

487 Langworth, B.F., 1977. *Fusobacterium necrophorum*: its characteristics and role as an animal pathogen.
488 *Bacteriol. Rev.* 41, 373–390.

489 Lowe, B.A., Marsh, T.L., Isaacs-Cosgrove, N., Kirkwood, R.N., Kiupel, M., Mulks, M.H., 2011. Microbial
490 communities in the tonsils of healthy pigs. *Vet. Microbiol.* 147, 346–57.
491 doi:10.1016/j.vetmic.2010.06.025

492 Lowe, B.A., Marsh, T.L., Isaacs-Cosgrove, N., Kirkwood, R.N., Kiupel, M., Mulks, M.H., 2012. Defining the
493 “core microbiome” of the microbial communities in the tonsils of healthy pigs. *BMC Microbiol.* 12, 20.
494 doi:10.1186/1471-2180-12-20

495 Mortensen, M. S., Brejnrod, A. D., Roggenbuck, M., Abu Al-Soud, W., Balle, C., Krogfelt, K. A., Stokholm, J.,
496 Thorsen, J., Waage, J., Rasmussen, M. A., Bisgaard, H., Sørensen, S. J. (2016). The developing
497 hypopharyngeal microbiota in early life. *Microbiome*, 4(1), 70. doi:10.1186/s40168-016-0215-9

498 Nauta, M., Barfod, K., Hald, T., Sørensen, A.H., Emborg, H.-D., Aabo, S., 2013. Prediction of *Salmonella*
499 carcass contamination by a comparative quantitative analysis of *E. coli* and *Salmonella* during pig
500 slaughter. *Int. J. Food Microbiol.* 166, 231–7. doi:10.1016/j.ijfoodmicro.2013.07.014

501 Nesbakken, T., Eckner, K., Høidal, H.K., Røtterud, O.J., 2003. Occurrence of *Yersinia enterocolitica* and
502 *Campylobacter* spp. in slaughter pigs and consequences for meat inspection, slaughtering, and
503 dressing procedures. *Int. J. Food Microbiol.* 80, 231–240.

504 Nesbakken, T., Iversen, T., Eckner, K., Lium, B., 2006. Testing of pathogenic *Yersinia enterocolitica* in pig
505 herds based on the natural dynamic of infection. *Int. J. Food Microbiol.* 111, 99–104.
506 doi:10.1016/j.ijfoodmicro.2006.04.019

507 Pajarillo, E.A.B., Chae, J.P., Balolong, M.P., Kim, H.B., Seo, K.-S., Kang, D.-K., 2014. Pyrosequencing-
508 based analysis of fecal microbial communities in three purebred pig lines. *J. Microbiol.* 52, 646–51.
509 doi:10.1007/s12275-014-4270-2

510 Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski, R.A., Oakley,
511 B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B., Thallinger, G.G., Van Horn, D.J., Weber, C.F.,
512 2009. Introducing mothur: open-source, platform-independent, community-supported software for
513 describing and comparing microbial communities. *Appl. Environ. Microbiol.* 75, 7537–41.
514 doi:10.1128/AEM.01541-09

515 Sundberg, C., Al-Soud, W.A., Larsson, M., Alm, E., Yekta, S.S., Svensson, B.H., Sørensen, S.J., Karlsson,
516 A., 2013. 454 pyrosequencing analyses of bacterial and archaeal richness in 21 full-scale biogas
517 digesters. *FEMS Microbiol. Ecol.* 85, 612–626. doi:10.1111/1574-6941.12148

518 Thibodeau, V., Frost, E.H., Chenier, S., Quessy, S., 1999. Presence of *Yersinia enterocolitica* in tissues of
519 orally-inoculated pigs and the tonsils and feces of pigs at slaughter. *Can. J. Vet. Res.* 63.

520 Thorsen, J., Brejnrod, A., Mortensen, M., Rasmussen, M. A., Stokholm, J., Al-Soud, W. A., Sørensen, S.,
521 Bisgaard, H., Waage, J. (2016). Large-scale benchmarking reveals false discoveries and count
522 transformation sensitivity in 16S rRNA gene amplicon data analysis methods used in microbiome
523 studies. *Microbiome*, 4(1), 62. doi:10.1186/s40168-016-0208-8

524 Tulstrup, M.V.-L., Christensen, E.G., Carvalho, V., Linninge, C., Ahrné, S., Højberg, O., Licht, T.R., Bahl,
525 M.I., 2015. Antibiotic Treatment Affects Intestinal Permeability and Gut Microbial Composition in

526 Wistar Rats Dependent on Antibiotic Class. PLoS One 10, e0144854.
527 doi:10.1371/journal.pone.0144854

528 Van Damme, I., Berkvens, D., Baré, J., De Zutter, L., 2013. Influence of isolation methods on the occurrence
529 of plasmid-carrying *Yersinia enterocolitica* serotype O:3 in slaughter pig tonsils, faeces and carcass
530 surface swabs. Int. J. Food Microbiol. 164, 32–35. doi:10.1016/j.ijfoodmicro.2013.03.019

531 Van Damme, I., mattheus, W., Bertrand, S., Zutter, L.D. (2018), Quantification of hygiene indicators and
532 *Salmonella* in the tonsils, oral cavity and rectal content samples of pigs during slaughter. Food
533 Microbiol. 71, 120-128

534 Wauters, G., Kandolo, K., Janssens, M., 1987. Revised biogrouping scheme of *Yersinia enterocolitica*.
535 Contrib. Microbiol. Immunol. 9, 14–21.

536 Yu, Y., Lee, C., Kim, J., Hwang, S., 2005. Group-specific primer and probe sets to detect methanogenic
537 communities using quantitative real-time polymerase chain reaction. Biotechnol. Bioeng. 89, 670–679.
538 doi:10.1002/bit.2034

539

540 **Figure captions**

541

542 Figure 1. Illustration of the three swabbing areas A, B and C. Source: www.21food.com.

543

544 Figure 2. Bacterial community composition in samples at phylum level. Mean relative abundance of the
545 phyla; a) found in samples from tonsil surface and tonsil tissue, b) in samples from faeces, area A, area B,
546 area C (see location in Figure 1) and tonsil surface.

547

548 Figure 3. Alpha diversity of bacterial communities calculated as Shannon index (a,c) and number of
549 observed OTUs (b,d) in samples of tonsil surface and deep tonsil tissue (a-b) and in samples from tonsils,
550 faeces, and carcass areas A, B and C (c-d). Columns show means with error bars indicating standard error
551 of means (SEM). Asterisks indicate significant differences, ****, $p < 0.0001$; ns, not significant.

552

553 Figure 4. Principal Coordinates Analysis (PCoA) plots (unweighted UniFrac distances) (a,c) and mean
554 UniFrac distances within and between different sample types (b, d) for samples of tonsil surface ($n = 35$) and
555 deep tonsil tissue ($n = 32$) (a-b) and in samples from tonsils ($n = 10$), faeces ($n = 8$), and carcass areas A (n
556 $= 9$), B ($n = 9$), C ($n = 10$) and the saw ($n = 1$), (c-d). In bar plots (b, d) error bars indicate standards error of
557 the means (SEM) and asterisks indicate significant differences, ****, $p < 0.0001$.

558

559 Figure 5. Heatmap of relative abundance of bacterial genera for individual samples obtained from tonsil
560 surface ($n = 35$) and tonsil tissue ($n = 32$). Colours indicate $\log_{10}$ relative abundance. A colour indication of
561 e.g. -2 means an abundance of 1% of the total community. Only bacterial genera with 35 % prevalence
562 across all samples are included. Asterisks indicate significant differences, with red indicating higher relative
563 abundance in tonsil surface and green indicating higher abundance in tonsil tissue (FDR corrected
564 permutation based t-test); *, $p < 0.05$; **, $p < 0.01$. The left-hand colour bar shows the classification of
565 bacterial genera (yellow: Actinobacteria, red: Bacteroidetes, cyan: Bacilli (Firmicutes), dark blue: Clostridia
566 (Firmicutes), light blue: Erysipelotrichia (Firmicutes), purple: Fusobacteria, salmon: β -Proteobacteria, light-
567 green: ϵ -Proteobacteria, dark-green: γ -Proteobacteria, orange: SR1, Spirochaetes and Tenericutes).

568

569 Figure 6. Mean fraction of those OTUs identified in tonsils (a) and faeces (b) also found in other
570 compartments including area A (around the head), area B (along the backbone on the cut surface) and area
571 C (around the head). Columns show means with error bars indicating standard error of means (SEM).
572 Asterisks indicate significant differences (FDR-adjusted p -values); ns, not significant; *, $p < 0.05$; **, $p < 0.01$.

573

574 Figure 7. Heatmap of relative abundance of bacterial genera in individual from tonsils (n = 10), faeces (n = 8)
575 and carcass swabs from area A (n = 9), B (n = 9) and C (n = 10). Colours indicate log₁₀ relative abundance..
576 Only bacterial genera with 35 % prevalence across all samples are included. The left-hand colour bar shows
577 the classification of bacterial genera (yellow: Actinobacteria, red: Bacteroidetes, cyan: Bacilli (Firmicutes),
578 dark blue: Clostridia (Firmicutes), light blue: Erysipelotrichia (Firmicutes), purple: Fusobacteria, salmon: β-
579 Proteobacteria, light-green: ε-Proteobacteria, dark-green: γ-Proteobacteria, orange: SR1, Spirochaetes and
580 Tenericutes).

581

582 Figure 8. Relative abundance of *Yersinia* (a), *Escherichia/Shigella* (b) and *Fusobacterium* (c) in samples
583 from tonsils, faeces and area A, B and C. Columns show mean values with error bars indicating standard
584 error of means.

585

586

587 Figure 1

588

589

Fig. 2

Fig. 2

590

591

592 Figure 3

593

594

595

596

597

598

599

600

601

602

603

604 Figure 4

605

606

607

Figure 5

Figure 5

608

609

610

611 Figure 6

612

613

Figure 7

Figure 7

614

615

616

Highlights

- Major phyla in pig tonsils were Proteobacteria and Fusobacteria
- Clear differences in microbiota at genus level between tonsil surface and deep tonsil tissue
- The genera *Moraxella*, *Fusobacterium*, and *Actinobacillus* are potential markers of carcass contamination from tonsils