



## The use of egg and soy phosphatidylcholines with attached caffeic acid enhances oxidative stability of high fat emulsion

Yesiltas, Betül; García Moreno, Pedro Jesús; Sørensen, Ann-Dorit Moltke; Anankanbil, Sampson; Banerjee, Chiranjib; Guo, Zheng; Ogilby, Peter R.; Jacobsen, Charlotte

*Publication date:*  
2019

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Yesiltas, B., García Moreno, P. J., Sørensen, A-D. M., Anankanbil, S., Banerjee, C., Guo, Z., Ogilby, P. R., & Jacobsen, C. (2019). *The use of egg and soy phosphatidylcholines with attached caffeic acid enhances oxidative stability of high fat emulsion*. Abstract from 30th Nordic Lipidforum Symposium 2019, Horsens, Denmark.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## **The use of egg and soy phosphatidylcholines with attached caffeic acid enhances oxidative stability of high fat emulsion**

Betül Yesiltas, Pedro J. García-Moreno, Ann-Dorit M. Sørensen, Sampson Anankanbil, Chiranjib Banerjee, Guo Zheng, Peter R. Ogilby, Charlotte Jacobsen

This study was carried out to investigate the effects of combined use of sodium caseinate (CAS), commercial phosphatidylcholine (PC) and modified PCs on physical and oxidative stability of 70% fish oil-in-water emulsions. Caffeic acid was covalently attached to both modified PCs (egg PC and soy PC; PCs originated from eggs and soy, respectively) in order to increase the antioxidant activity of PCs and investigate the advantage of bringing the antioxidant activity to the close proximity of oil-water interface. Results showed that the oxidative stability was improved when part of the PC was substituted with modified egg PC or soy PC. Emulsions containing a low concentration of modified PCs (10% wt of total PC) resulted in a prooxidative effect on the formation of hydroperoxides (peroxide value) compared to emulsions with free caffeic acid added. On the other hand, a decrease in the formation of volatile oxidation products was observed for emulsions containing higher levels of modified PCs (10% wt of total PC) compared to the emulsions with free caffeic acid added at its equivalent concentration. Increased concentration of modified PCs provided higher oxidative stability in high fat emulsions independently of the type of the modified PCs. Moreover, it was shown that the oxidation did not propagate from one droplet to another droplet by the use of microscopy. In this case, oxidation was initiated by singlet oxygen produced in a single droplet using a focused laser. Oxidation propagation was followed using the fluorescence of unoxidized and oxidized C11-BODIPY.