



New hatchery methods for efficient spore use and seedling production of *Palmaria palmata* (dulse)

Schmedes, Peter Søndergaard; Nielsen, Mette Møller

Published in:
Journal of Applied Phycology

Link to article, DOI:
[10.1007/s10811-019-01998-0](https://doi.org/10.1007/s10811-019-01998-0)

Publication date:
2020

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Schmedes, P. S., & Nielsen, M. M. (2020). New hatchery methods for efficient spore use and seedling production of *Palmaria palmata* (dulse). *Journal of Applied Phycology*, 32, 2183-2193.
<https://doi.org/10.1007/s10811-019-01998-0>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 New hatchery methods for efficient spore use and seedling production of *Palmaria palmata* (dulse)

2

3 Peter Søndergaard Schmedes^{1,a}, Mette Møller Nielsen¹

4 ¹Technical University of Denmark, DTU Aqua, Øroddevej 80, 7900 Nykøbing Mors, Denmark

5 <https://orcid.org/0000-0001-5772-4737>

6 Key words: Dispersal, fertilization, hatchery, tetraspores, reattachment, spore seeding

7 ^aCorresponding author: peson@aqu.dtu.dk

8

9 Abstract

10 *Palmaria palmata* (dulse) is a high valued rhodophyte; nevertheless, its hatchery methods are
11 underdeveloped. New hatchery methods are required to improve the spore use efficiency and seeding
12 quality, which are important benchmarks for a viable cultivation. This study investigated a method using
13 vertical seeding tanks (exp. 1), hemispherical agitation and flow-through conditions to improve spore
14 dispersal on net substrates. Tanks were inoculated with different amounts of sori tissue, which sporulated in
15 three consecutive seeding periods. The results demonstrated significant effect of seeding period where 5-15
16 g FW sori could be used to seed three nets (~126 m rope) over the course of nine days providing a density
17 up to 10 seedlings cm⁻¹ after 32 days. The effluent spores were collected in detaining tanks and germinated
18 into a propagule mix of female and male gametophytes. The propagule mix was efficient as a secondary
19 seeding inoculum, as propagules were able to reattach to substrates up to 39 days after their release as
20 spores (exp.2). Adding male gametes to the propagule mix and spore seeded ropes was tested as a relevant
21 hatchery step to activate female gametophytes and significantly resulted in more than a doubling of
22 seedlings (exp.3). This study present new methods and strategies to improve spore use efficiency and to
23 obtain an equal spore dispersal on net substrates for hatchery production of *P. palmata*.

24

25 1. Introduction

26 The rhodophyte *Palmaria palmata* (L.) F. Weber and D. Mohr has traditionally been used for human
27 consumption with records dating back to the 9th century (Mouritsen et al. 2013). The emerging evidence of its
28 umami flavor (Mouritsen et al. 2013), bioactive and health properties of protein hydrolysates (Harnedy et al.
29 2014; Admassu et al. 2018), water extracts (Lee et al. 2017) and biorefined compounds (Schiener et al.
30 2017) has renewed interest in the use of *P. palmata*, resulting in increased focus on its cultivation. *P.*
31 *palmata* can be cultivated in tanks using vegetative growth (Morgan and Simpson 1981; Pang and Lüning
32 2004; Matos et al. 2006; Corey et al. 2014), but the prospects of cultivating the species from tetraspores has
33 lately received increasing attention in Europe (Edwards and Dring 2011; Werner and Dring 2011; Sanderson
34 et al. 2012). However, despite studies founding the essential understanding of the life cycle (van der Meer
35 and Chen 1979; van der Meer and Todd 1980) and cultivating *P. palmata* from spores (Browne 2001;
36 Sanderson 2006; Werner and Dring 2011; Grandorf Bak 2019; Schmedes et al. 2019), the methodology for
37 large-scale hatchery production in Europe is still in its infancy.

38 Essentially, the use of spores for cultivation of *P. palmata* is based on three steps; 1) the collection or
39 induction of sori, 2) the release and dispersal of spores and 3) the spore attachment and growth; and it is the
40 efficiency and success of these three steps that determines the usefulness of a given hatchery protocol. As
41 the steps are interlinked, the initial handling of sori, which represent only 8-10% of the total frond area during
42 the peak season of fertility (Werner and Dring 2011), is crucially important. Currently, the most commonly
43 used hatchery protocol to produce *P. palmata* seeded substrates is a flat tank using a 1:1 areal coverage of
44 substrate with sori and a three days spore release duration. However, this method requires a large amount
45 of sori, and furthermore, results in poor dispersal and high mortality (60-90%) of the spores (Werner and
46 Dring 2011). Also, by the using this method, it is often observed that a considerable amount of the spores
47 settle on tank surfaces instead of the intended cultivation substrate (personal observation).

48 Taking into account these results and observations, it is clear that there is a bottleneck in spore use
49 efficiency within *P. palmata* hatcheries. This particular matter was addressed in a recent study suggesting a
50 GMA-seeding method (Germination, Maceration, and Agitation method) as an alternative seeding method for
51 *P. palmata* (Schmedes et al. 2019). This study found that germinated propagules (i.e. a mix of spores and
52 seedlings) of *P. palmata* were able to establish discoid reattachment to a substrate after forced de-
53 attachment and maceration, resulting in high settlement efficiency as well as a high dispersal of the
54 propagules. However, the extensive use of this seeding method is still unknown. Besides the use of this
55 seeding strategy, the use of male gametes of *P. palmata* to fertilize female gametophytes has been
56 suggested as a potential hatchery step to optimize the hatchery production (Mine and Tatewaki 1994; Le
57 Gall et al. 2004; Schmedes et al. 2019). When using sori for spore-seeding, a mixture of male and female
58 spores develop after attachment, however, only the males will develop into a harvestable thallus unless a
59 fertilization step, enabling zygote formation in the female gametophyte, is included. Theoretically, this would
60 double the seedling density, as the crustose-like female gametophyte will develop into a sporophytic thallus
61 after the zygote. Yet, this is still uninvestigated under relevant hatchery conditions.

62 Because of the continued challenges in producing *P. palmata* sporophytes and gametophytes, this present
63 study investigated hatchery methods and strategies based on Schmedes et al. (2019) to further improve the

64 utilization of spores for cultivation of *P. palmata* on substrates. Five strategies to improve the hatchery
65 protocol were tested: 1) A vertical seeding tank in flow-through condition agitated with air bubbles to promote
66 dispersal of spores. 2) The effect of different amount of sori during the seeding phase. 3) The effect of using
67 the same sori material in three consecutive seeding periods on the spore germination and seedling density.
68 4) The use of detained spores as a propagule seeding inoculum, based on the GMA-method. 5) The effect of
69 fertilization on the number of attached seedlings under relevant hatchery conditions.

70


71 2. Materials and methods

72 Three experiments were set up to test the five improvements in hatchery techniques described above.

73 Experiment 1 investigated a new flow-through seeding system for improved seeding dispersal, additionally
74 using different sori amounts in consecutive seeding periods as a means to improve sori use. This experiment
75 addressed the first three steps of hatchery improvements. Experiment 2 covered step four, optimizing the
76 use of detained spores, and finally, experiment 3 exploited the opportunity to include fertilization as a means
77 to improve spore use, addressing step five of hatchery improvement techniques.

78 2.1. Experiment 1: spore seeding and sori use.

79 Nine conical tanks (polyethylene, 30 L) were set up as seeding tanks in a parallel flow-through system with a
80 water flow of 0.5 L min⁻¹ (fig. 1A) in a 10 °C cold room. Air bubbles from the bottom (1.2 L min⁻¹) bubbled
81 through the sori package near the surface (fig. 1C), generated a hemispherical circulation of the water
82 allowing the spores to disperse and aerated the tank volume. Water for the entire system was circulated from
83 a reservoir tank with a circulation pump and exchanged (10% of total volume) on a daily basis. No nutrients
84 were added. From each seeding tank, an outlet, placed diagonal to the inlet, led the effluent water directly
85 into a 2 L spore-detaining tank (SDT) (fig. 1B). Here, effluent spores aggregated on the bottom, while water
86 was led out in the top and back to the reservoir tank through a nylon filter (15 µm) for UV treatment (60 W).
87 Spores detained in the SDTs were used in experiment 2.




88

Figure 1.

89 Twenty-seven pieces of net (polypropylene, 0.25*1.40 m, ϕ =5-7mm) were prepared as cultivation substrate
90 by soaking them in lukewarm tap water a month before, then kept in clean seawater for seven days before
91 use. At the experimental start, the nets were submerged in the seeding tanks as a two-layered net spiral (fig.
92 1B).

93 Fertile *P. palmata* tetrasporophytes and male gametophytes were collected 4 January 2019 in the intertidal
94 zone near Fornæs light house, Denmark (56.443534N, 10.958985E) and kept in running seawater (4 °C) in
95 dim natural light ($\sim 15 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$, PAR). On 10 January, 218 fertile fronds were rinsed in 0.2 μm
96 filtered and sterilized seawater (fig. 2A) before being desiccated for 20 hours at 5 °C in darkness. Presence
97 of ripe sporangia was verified by inspection (fig. 2B). Triplicates of three groups of sori amount (5, 10 or 15 g
98 FW; fresh weight) were prepared as sori packages by placing the tissue between two layers of plastic net
99 (15*15 cm, fig. 2C).



100 Figure 2.

101 The experiment started 11th of January 2019 by adding a net and a sori package (3 replicates of 3 sori
102 amounts) to each of the nine tanks. The seeding ran over a course of nine days, keeping the same sori
103 package in the tanks for all nine days but exchanging the nets with new ones every third day to test the
104 potential of consecutive seeding periods (Day 0-3, Day 3-6, Day 6-9) using the same sori. During seeding,
105 the surface irradiance was 15-22 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$, 10:14 h L: D. At the end of each consecutive seeding
106 period, the nine nets were labelled and transferred to nursery tanks containing 300 L enriched seawater (0.2
107 μm filtered); 10% strength of Varicon Aqua Cell-Hi F2P, a F/2 nutrient media with vitamins based on Guillard
108 and Ryther (1962). Air stones provided water agitation in the tanks after three days. Irradiance was 35 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ (10:14 h L: D), but raised to 50 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ from week 2.

110

111 *Data acquisition*

112 The net-seeding efficiency using sori in the flow-through system was calculated by equation 1:


113
$$\text{Net seeding efficiency} = \frac{\# \text{ spores on net}}{\# \text{ Total spores}(\text{net} + \text{detained})} * 100\%$$

114 This was carried out by counting the number of attached spores and seedlings on a 4 cm net subsample for
115 each of the 9 nets (n=3) three days after each seeding period. Additionally, the nets were counted at day 19
116 and 32 after each of the consecutive seeding periods to compare spore and seedling density within and
117 between groups over time. The subsamples were taken from the same position regarding the in, - and outlet
118 of the tanks on each net spiral. To verify an even spore dispersal on the entire net, additional subsamples
119 were taken from the 10 g sori batch in the first seeding period (Day 0-3). From each net (n=3), three
120 subsamples were taken from the bottom and top part of the net (N=9) and tested for unequal variance using
121 Levene's test. Similarly, by counting spores on subsamples (N=21) from all around one net spiral in the
122 same vertical level the spore dispersal was assessed. By verification, we extrapolated the spore and
123 seedling densities for each counting day and calculated the germination success by equation 2:

124
$$\text{Germination (\%)} = \frac{\# \text{ Seedlings}}{\# \text{Total propagules (spores + seedlings)}} * 100\%$$

125 The total number of detained spores in each SDT after a seeding event (fig. 3A) was estimated based on
126 subsample counting, by the following process: 1) A homogenous spore mix (fig. 3B) was prepared by
127 reducing the SDT volume, dislodging the spore aggregates and macerating the solution for 30 seconds with
128 a kitchen blender. 2) A 10 mL subsample was transferred to a gridded petri dishes (fig. 3C) and added 20
129 mL enriched (10% F/2) seawater. 3) The spores were kept at 5 °C and $5 \pm 2 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ (10:14 h
130 L:D) for three days before five random fields (each 0.0035 mm^2) were photographed and counted. The total
131 number of spores was estimated by factor multiplying the mean count of subsamples ($\text{STD}_{\text{VOLUME}} * 10 \text{ mL}^{-1}$
132 subsample and petri dish area * 0.0035 mm^2).

133



134 Figure 3.


135

136 2.2. Experiment 2: Mixed propagules as seeding inoculum

137 In Experiment 2, the propagules germinated from detained spores from experiment 1 were tested as seeding
138 inoculum, according to the GMA-method (Schmedes et al. 2019). After subsampling for petri dish cultures in
139 experiment 1, the rest of the macerated spore solution from the last seeding period (Day 6-9) was poured
140 into a plexiglas containers (5 L), settled on the bottom (fig. 4A) and cultivated at $20 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$
141 (adjusted to $5 \pm 2 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$, after ten days) and 5 °C. After one day, enriched seawater and
142 germanium-oxide (10% F/2 + 1 mg L⁻¹ GeO) was added and after six days, an air stone was put for aeration.
143 The detained spores germinated into a mixture of propagules (i.e. spores and gametophytes) at these
144 conditions before use.

145 Twenty-seven days after the initiation of Day 6-9 seeding period, some of the propagules on the bottom were
146 dislodged (Fig. 4B), suspended in 400 mL enriched (10% F/2) seawater and macerated to break propagule

147 aggregates (fig. 4C) according to Schmedes et al. (2019). The propagules were photographed before and
148 after the maceration treatment (fig. 4D, E). A total of 20 mL of the propagule mix was added to each of 10
149 beakers (n=10) containing 400 mL enriched (10% F/2) seawater and a piece of rope (10 cm) standing
150 upright in the beaker to test the ability of the propagules to reattach. The rope was made by untangling
151 several net meshes and cut and cut into pieces. Immediately, 1 mL subsamples were withdrawn and the
152 concentration of macerated propagules was estimated to 259 ± 8 spores mL^{-1} and 422 ± 13 seedlings mL^{-1}
153 (mean \pm SE, n=4). The beakers were constantly agitated from the bottom via air bubbles (2.5 L min^{-1}) to
154 promote the dispersal of the propagules.
155



156 Figure 4.
157


158 After 3 days inoculation (“agitation period day0-2”), the maximum density of attached spores and seedlings
159 was estimated by counting the three densest subparts (0.5 cm) for each rope, after which the ropes were
160 transferred to 20 L tanks at the similar conditions until they were counted again at day 10. After transferring
161 the ropes pieces out of the beakers, the propagule mix that maintained in the beakers under same conditions
162 until day 10, where concentration (mean of technical replicate \pm SE) of spores (237 ± 7) and seedlings
163 (394 ± 9) seedlings was similar to the starting concentration (two-sampled t-tests: $t(4)$, $p=0.086$, $p=0.131$,
164 respectively). Then, new pieces of rope were added for 3 days inoculation to see how long after the
165 dislodgement and maceration the propagules would be able to reattach (“agitation period day10-12”). Also,
166 at day 10, new beakers (n=10) were set up under same conditions using freshly dislodged and macerated
167 propagules to see if the age of the propagules (27 vs. 37 days) at dislodgement affected the ability to
168 reattach. The concentration of propagules (250 ± 20 spores and 396 ± 22 seedlings) was similar (two-sampled
169 t-tests: $t(4)$: $p = 0.584$, $p = 0.341$) with the other start concentrations.

170 2.3. Experiment 3: Including fertilization in hatchery

171 The effect of fertilizing female gametophytes was tested by adding male gametes to propagules cultured in
172 petri dishes and spore-seeded ropes cultured in beakers. A solution of propagules (349 ± 26 propagules mL^{-1} ,
173 $n = 9$, technical replicates) was prepared from the same detained spore batch as used in experiment 2 by
174 dislodging and macerating after 11 days of germination. A total of 15 mL propagule solution was then poured
175 into each of 18 petri dishes and filled with additional 10 mL enriched (10% F/2) seawater and kept in 15 ± 3
176 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ PAR (12:12 h L:D) irradiance at 5 °C.

177 At day 1, a solution of male gametes was prepared and 15 mL of this solution was added to nine petri
178 dishes, containing the propagule solution. Nine other dishes received 15 mL seawater (10% F/2) as a
179 control. The male gamete solution was prepared by desiccating (3 h, dark at 5 °C) thirty fertile male
180 gametophytic fronds (22.4 g FW, see fig. 5A and 5B) before being rehydrated (1 h, $5 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$
181 PAR) in an agitated volume of 400 mL seawater + 0.4 mL GeO (1g/L). The male gametes ($\varnothing \sim 5 \mu\text{m}$) were
182 visible using high magnification (fig. 5C and 5D). The concentration of gametes was estimated to be ~ 1000
183 gametes mL^{-1} , using a Neubauer cell counting chamber. After adding, the cultures were mixed on a stirring
184 table for 30 min at 160 rpm. At day 4 and 6, the petri dish volumes were exchanged with new additions of
185 male gamete solution (5 mL, $\sim 10^6$ gametes mL^{-1}).

186 For each petri dish, spores and seedlings were counted in five random fields (each 0.0035 mm^2) at day 0, 3,
187 10 and 19 and the number of seedlings were adjusted to the spore count of previous counting day.



188 Figure 5.

189 In addition, we assessed the effect of fertilization on the seeded rope pieces (Exp. 2). This was carried out in
190 beakers (1.8 L) kept at 5 °C, in which 4 cm pieces of seeded rope were tumbled by air bubble agitation
191 ($n=11$). The pieces were excised from “Day 3-6 and 10 g sori” nets (Exp. 1) and nursed for 16 days at 10 °C,
192 before being transferred to the beakers. This was followed by adding 40 mL solution containing male
193 gametes to the beakers, while a control group was added the same enriched seawater (10% F/2) with no
194 gametes ($n=11$). The numbers of spores and seedlings was counted at day 1, 3, 6, and 12 and the seedling
195 number was presented as relative to spore number.

196 2.4. Statistics

197 For all data sets, Shapiro Wilks test was used to check normality and Levene test was used to check
198 variance homogeneity. The analysis was carried out using SAS, JMP 13, using a significance level of 0.05.
199 When sufficient, data were log-transformed to ensure homogeneity of variance and normal distributed
200 residuals and ANOVAs were conducted to compare the main effects. In case of none normal distribution or
201 homogeneity of variance, the Wilcoxon two-sample test was used. All data are given as mean \pm standard
202 error (SE), unless stated otherwise.

203 Experiment 1. In a factorial design, nets were manipulated to be in one of nine groups forming the
204 combination of consecutive seeding periods (3 levels; Day 0-3, Day 3-6, Day 6-9) and sori amount (3 levels;
205 5, 10, 15 g FW), and 2-way ANOVA (2w-an), including the interaction term, using log-transformed data was
206 conducted to compare effects on spore and seedling numbers attached to the nets, the number of detained
207 spores in STDs, the spore-seeding efficiency as well as the germination success (%) between all groups,
208 followed by Tukey's HSD post hoc test. The Wilcoxon two-sample test was used to assess for any significant
209 change in mean densities of spores and seedlings across the counting days within each factorial group.

210
211 Experiment 2. Datasets were log-transformed to ensure normal distribution, but spore counts did not obey
212 homogeneity of variance (Levene, $p < 0.0208$). Hence, the Wilcoxon test was used to assess for significant
213 difference in mean settlement density at each count day.

214 Experiment 3. For the petri dish cultures, the effect of adding male gametes on the number of seedlings
215 adjusted to the previous spore count was compared using a two-tailed Student's t test ($n=11$). For the rope
216 cultures, the univariate repeated measures ANOVA was used to test effect of the between factor (two levels;
217 male gametes vs. control) and the within factors (time; four levels) on the adjusted number of seedlings.
218 Datasets were normal distributed and displayed equal variance after log-transformation. The Mauchly's
219 sphericity test: $\chi^2(5) = 8.3513$, $p = 0.1379$ allowed to report the p-value of the F test.

220
221 3. Results

222 3.1. Experiment 1: spore seeding and sori use

223 Spores settled on all nets during the consecutive seeding periods and for all amount of sori used (fig. 6A-C).
224 Spores were present on all counting days, while seedlings (fig. 6D-F) were only present from counting day
225 19 and onwards. On counting day 3, the number of spores attached to the nets was significantly affected by
226 seeding periods (2w-an: $F_{2,8} = 7.2461$, $p = 0.0049$), the second (fig. 6B) and third (fig. 6.C) seeding period
227 (day 3-6 and day 6-9) showing similar and significant higher spore densities (Tukey's: $p = 0.0036$) than the
228 first seeding period (day 0-3) (fig. 6A). Also, the sori amount used for seeding, significantly affected the
229 number of attached spores (2w-an: $F_{2,8} = 4.0989$, $p = 0.0341$), showing significant difference between high
230 and low sori amounts (Tukey's, $p = 0.0109$). On counting day 19, the number of spores (fig. 6A-C) and
231 seedlings (fig. 6D-F) as well as the spore germination success (fig. 6G-I) was similar for all groups (2w-an:
232 $p > 0.5606$). On counting day 32, the spore settlement was still similar across all groups (2w-an: $p > 0.1179$)

233 (fig. 6A-C), whereas the number of seedlings (fig. 6D-F) was significant affected by seeding period (2w-an:
 234 $F_{2,8} = 6.1718$, $p = 0.0091$), showing similar seedling density in second and third period (Tukey's, $p = 0.8110$),
 235 both significantly higher than the first seeding period (Tukey's, $p = 0.0105$, $p = 0.0383$, respectively). At day
 236 32, the average seedling density was 36 ± 8 seedlings per 4 cm. However, the germination success (fig. 6G-I)
 237 up to 50-80% turned out to be similar between seeding periods (2w-an: $F_{2,8} = 3.0829$, $p = 0.0706$) and sori
 238 groups (2w-an: $F_{2,8} = 0.1873$, $p = 0.83$). Within each seeding period (fig. 6A-C), the spore count decreased
 239 significantly over time (Chi-squared: $\chi^2(9) = 10.14$, $p = 0.0063$), whereas the increase in seedling density
 240 was only significantly on nets produced in second (Mann-Whitney two-sample test: $p=0.0030$) and third ($p =$
 241 0.0295) seeding periods.




Figure 6.

242

243 The additional rope pieces sampled to assess whether settlement was homogeneous dispersed across the
 244 entire net, showed that this was the case. Both spores and seedlings showed equal variance between top
 245 and bottom of the nets (spores: $p = 0.17$, seedlings: $p = 0.40$) and means (Mann-Whitney two-sample test: p
 246 $= 0.40$). Furthermore, the densities showed equal variance horizontally around the net spiral (spores: $p =$
 247 0.23 , seedlings: $p = 0.33$). Hence, the number of spores settled on rope pieces cut from the nets was
 248 extrapolated to whole nets (fig. 7A-C) and summed with the number of detained spores (fig. 7D-F) to
 249 calculate the total number of released spores. This number was then used to estimate a net seeding
 250 efficiency of the system (fig. 7G-I) by equation 1. The net seeding efficiency of the agitated, flow-through
 251 seeding tank system using sori packages was in average $16 \pm 1.6\%$ (fig.7G-I) and not affected by either
 252 seeding period or sori amount (2w-an: $p > 0.28$). This indicates that app. 84 % of the released tetraspores
 253 were washed out and detained (fig. 7D-F).

254 Effluent spores aggregated as red plumages on the bottom of all the SDTs, and were visible within 3 hours
 255 during the first seeding period (Day 0-3). The number of detained spores from the flow-through system (fig.
 256 7D-F) was significantly affected by seeding period (2w-an: $F_{2,8} = 5.66$, $p = 0.0124$), where second period
 257 provided higher detained spore yield compared to the third period (Tukey's: $p = 0.0117$) and similar spore
 258 yield as first period ($p = 0.6593$). The amount of sori used also affected the number of detained spores

259 significantly (2w-an: $F_{2,8} = 5.07$, $p = 0.0178$) where 15 g sori provided higher yield compared to 5 g sori
 260 (Tukey's: $p = 0.0175$). In total, the amount of released spores (detained + attached to nets) from the three
 261 different sori packages during nine days of sporulation was significantly higher for the 15 g sori group
 262 (1,300,563±71,639), compared to the 10 g sori (959,697±69,618) and the 5 g sori group (522,230±61,869)
 263 spores.
 264




Figure 7.

265
 266
 267

3.2. Experiment 2: Mixed propagules as seeding inoculum

268 On counting day 3, the number of reattached spores (fig.8A vs. 8B) and seedlings (fig. 8C vs. 8D) was
 269 significantly higher, when the inoculum was dislodged at day 27-29 compared to day 37-39 (Mann-Whitney
 270 two-sample test (10), $p < 0.0001$). Furthermore, the number of reattached spores and seedlings on ropes
 271 exposed to different agitation periods (Day 0-2 vs. Day 10-12) was significantly different ($p < 0.0001$, $p =$
 272 0.0012 , respectively). After nursing these ropes for additional seven days (counting day 10), the effect of
 273 agitation period on the density of spores and seedlings remained significant ($p = 0.0002$, $p = 0.0009$). In
 274 contrast, the effect of inoculum age at dislodgement was insignificant on the spore density ($p = 0.0697$) and
 275 seedling density ($p = 0.9938$).





Figure 8.

276

277 3.3 Experiment 3: Including fertilization in hatchery

278 After 19 days in petri dish culture, the adjusted seedling number was 50.25 % higher (fig. 9A) with male
 279 gametes solution added, compared to the control (two-tailed Student's t test: $t(10)$, $p < 0.0001$), whereas no
 280 significant differences were found at the other days ($p > 0.98$), except day 3, where the control group showed
 281 a slightly higher number of seedlings ($p = 0.0042$).

282 The effect of adding male gametes to the rope cultures (fig. 9B) was significant from day 3 (MANOVA: $F_{1,19} =$
 283 121.445 , $p < 0.0001$) with higher number of seedlings at all counting days (253.2 % at day 12). Yet, the effect
 284 of time was insignificant on the number of seedlings (MANOVA: $F_{3,17} = 0.5462$, $p = 0.6573$), but caused a
 285 significant interaction term of gamete addition * time: $F_{3,17} = 4.0842$, $p = 0.0235$, because of the higher
 286 seedling count after day 0.



287
288 Figure 9.

289 Discussion

290 The results of this study demonstrate effective methods and strategies in the pursuit of optimizing the
 291 hatchery production of *P. palmata* by using vertical seeding tanks, consecutive use of sori packages and
 292 agitation during the seeding phase. The strategy of using a secondary seeding inoculum of germinated
 293 propagules, based on collecting the effluent spores, obviously increase the spore use efficiency. With
 294 presented seeding system, 9 nets were spore-seeded, with an equivalent linear length of ~ 126 meter of
 295 rope, using as little as 5 g FW sori, which result in an average of 9 seedling cm^{-1} after 32 days. The use of
 296 spores for seeding substrates is prospective for larger scale cultivation (Browne 2001; Edwards 2007;
 297 Werner and Dring 2011). The latter argued that an initial seeding density of 100 spores cm^{-1} is required to
 298 obtain a final seedling density of ~ 6-8 seedlings cm^{-1} as a mortality rate of spores of 60-80% took place. On
 299 top, only male gametophytes developed a thallus, representing 50% of the total amount of spores. The initial
 300 spore density encountered in the present study (Exp. 1) on the nets was lower than 100 spores cm^{-1} ,
 301 however, we estimated a spore germination success (reciprocal to mortality) of 50-80% on the nets similar to
 302 what was achieved in a previous study (Le Gall et al., 2004). This indicates that the seeding and nursery
 303 conditions presented here were good, though it might be overestimated, as we were not able to count the
 304 dying spores. In contrast, we observed a germination success of only 8-14 % in petri dishes cultured in
 305 stagnant seawater under the same conditions (unpublished work), which is in the range of previous report
 306 (Edwards 2007; Edwards and Dring 2011). By using higher amounts of sori in the seeding phase of net, the
 307 nets showed higher spore density after 3 days, but decreased to similar levels for all sori amount used, after
 308 32 days of nursery, in line with previous findings (Edwards 2007; Werner and Dring 2011). In present study,
 309 the average seedling density after 32 days of nursery was not significantly affected by the sori amounts
 310 tested. Besides testing efficient ways to handle sori tissue for optimal spore yield, it is important that future
 311 hatchery trials consider the sori-to-substrate density, as high spore density seem to even out during nursery,
 312 thus the sori could have been used more efficient. A current hatchery protocol for *P. palmata*, found that a
 313 sori-to-substrate ratio of 150 g FW sori 84 m^{-1} substrate was required to secure sufficient seedling density,

314 which converts to ~130 kg FW fronds to seed one long-line of 100 m (Werner and Dring 2011). In
315 comparison, several kilometers of substrate can be seeded with motile zoospores of *Saccharina latissima* by
316 using 150 g sori, due to the multifold number of biflagellate zoospores released within 1 hour and their
317 capability of high dispersal (personal observation). We suggest that a lower amount of sori is sufficient for
318 seeding, while considering the sori-to-substrate ratio.

319 Three times the amount of seeded substrate was produced by using three consecutive
320 seeding periods, compared to the conventional 3-days seeding phase. The highest seedling density after 32
321 days for nets seeding was found in the second (~8 seedlings cm⁻¹ and third seeding period (~10 seedlings cm⁻¹).
322 In the first seeding period (Day 0-3), our tank setup provided a spore-seeding efficiency of 16%, meaning
323 that ~84 % of the released tetraspores were detained in the down-stream detaining tanks. The amount of
324 detained spores was highest after the second seeding period (Day 3-6). Even 38 days after finalizing the last
325 seeding period we observed red tetraspore aggregates in some of the spore-detaining tanks. This
326 observation supports previous findings, where sori was observed to release for 21 days (Schmedes et al.
327 2019) to 40 days (Wood 2018).

328 Several bottlenecks have been identified by using the current hatchery protocol for *P. palmata*
329 for large-scale cultivation – a protocol where sori is placed above the cultivation substrates (1:1 areal
330 coverage) in horizontal tanks, which impose a high sori requirement. Facing other challenges when hatching
331 rhodophytes, such as seasonal variation in spore availability (Kain 1986; Le Gall et al. 2004), a relative low
332 spore release yield (Edwards 2007), poor spore dispersal before settlement (Edwards and Dring 2011) and
333 low survival of spores (Sanderson 2006; Werner and Dring 2011), all impose a low spore use efficiency and
334 little control of seedling quality. Overall, an even spore dispersal and good seedling density was found in this
335 study using a flow-through system and relative high aeration, agitating the water. Nevertheless, further
336 investigations of the previous mentioned challenges are highly relevant to optimize before commercial
337 hatcheries can be established. Hence, the recently developed GMA-method (Schmedes et al. 2019) to
338 improve the spore efficiency for hatching *P. palmata* was investigated. This method was applied by
339 dislodging and maceration of detained spores, which then germinated into propagules, which again were
340 macerated to break aggregates of spores and tiny seedlings into single and small groups of spores and
341 seedlings, before used as a secondary seeding inoculum. The level of water agitation of 2.5 L min⁻¹
342 dispersed the propagules, yet, did not compromise the establishment of a discoid reattachment on the ropes
343 (Exp. 2). Results demonstrated that the reattachment was negatively affected by the biological age at
344 dislodgement and amount of days in agitation, which adds knowledge to the extent of which macerated *P.*
345 *palmata* propagules can be used as seeding inoculum. This is in agreement with findings in other red
346 seaweed species, such as *Chondracanthus chamissoi* (Gigartinales), which forms a substrate reattachment
347 by the production of secondary attachment discs (Sáez et al. 2008). The thallus fragments showed a
348 decreasing reattachment probability over time (Fonck et al. 2008). Also, the red seaweed *Gelidium chilense*
349 (Montagne) formed bundles of rhizoids in agitated water (Santelices and Varela 1994), while absent in
350 *Gelidium coulteri* cultured in stagnant water conditions (Macler and West 1987). In comparison, a direct
351 seeding method for cultivating the brown macroalgae species *S. latissima*, as a way to optimize hatchery
352 duration and costs, is currently used by the Hortimare company, as a seeding technique for commercial

353 cultivation in Europe. Here, tiny germinated and activated sporophytes are applied to substrates and
354 establish firm attachment with their developing haptera organs, with a potential benefit of using a glue
355 (Kerrison et al. 2018). Whether the propagule seeding method of *P. palmata* would benefit by using glue as a
356 means to maximize seedling density, is of high interest due to the high commercial value of the biomass.

357 The inclusion of a fertilization step proved to be highly relevant to increase the number of
358 seedlings in the hatchery production, by releasing male gametes and adding this male gamete solution to the
359 female gametophytes that are developing their trichogynes (Mine and Tatewaki 1994; Le Gall et al. 2004).
360 This resulted in at least a doubling of the number of seedlings on spore-seeded rope and suggests that male
361 gametes can be added 10-22 days after seeding substrates to increase the overall spore-use-efficiency of *P.*
362 *palmata*.

363

364 4. Conclusions

365 A new method is reported here to handle fertile sori of *P. palmata* for efficient release and dispersal of
366 tetraspores using sori packages in vertical flow-through tanks. This achieves a sufficient seedling density ~9
367 seedlings cm⁻¹ rope. Consecutive use of sori greatly improves the spore use efficiency and was found to
368 have a positive effect on seeding quality, even for different sori amounts used in the sporulation phase. The
369 spore inoculation success (net-seeding efficiency ~16%) was relative low, however, the ~84 % effluent
370 spores was used in propagule inoculation by using the GMA-method. Hence, to increase the use of spores
371 for efficient inoculation, the importance of addressing following parameters to further improve the spore-use
372 efficiency for seeding *P. palmata* is needed. The following are of importance, i.e., tank typology, water
373 motion for optimal spore dispersal and settlement, intermittent batch conditions to increase net seeding
374 efficiency, substrate-to-volume density, sori position, and flow-through rate. It was demonstrated that effluent
375 spores from the SDTs were suited as a secondary seeding inoculum, using the Germinate-Macerate-Agitate
376 seeding method (Schmedes et al. 2019).

377

378 References

- 379 Admassu H, Gasmalla MAA, Yang R, Zhao W (2018) Bioactive peptides derived from seaweed protein and
380 their health benefits : Antihypertensive, antioxidant, and antidiabetic properties. J Food Sci 83(1).
- 381 Browne KL (2001) Mariculture of the edible red alga, *Palmaria palmata*. PhD thesis, Queen's University of
382 Belfast.
- 383 Corey P, Kim JK, Duston J, Garbary DJ (2014) Growth and nutrient uptake by *Palmaria palmata* integrated
384 with Atlantic halibut in a land-based aquaculture system. Algae 29(1):35–45.
- 385 Edwards MD (2007) The cultivation of the edible red alga, *Palmaria palmata* for aquaculture, PhD thesis,
386 The Queen's University of Belfast.
- 387 Edwards MD, Dring MJ (2011) Open-Sea cultivation trial of the red alga, *Palmaria palmata* from seeded
388 tetraspores in Strangford Lough, Northern Ireland. Aquaculture 317(1–4):203–209.
- 389 Fonck E, Martínez R, Vásquez J, Bulboa C (2008) Factors that affect the re-Attachment of *Chondracanthus*
390 *chamissoi* (Rhodophyta, Gigartinales) thalli. J Appl Phycol 20(3):311–14.
- 391 Grandorf Bak U (2019) Seaweed cultivation in the Faroe Islands : An investigation of the biochemical
392 composition of selected macroalgal species, optimised seeding technics, and open-ocean cultivation
393 methods from a commercial perspective. PhD thesis, Technical University of Denmark.
- 394 Guillard RRL and Ryther JH (1962) Studies on Marine Planktonic Diatoms I. *Cyclotella nana* Hustedt and
395 *Detonula confervacea* (Cleve) Gran. Can J Microbiol 8:229-239.
- 396 <https://doi.org/10.1139/m62-029>
- 397 Harnedy PA, Soler-Vila A, Edwards MD, FitzGerald RJ (2014). The effect of time and origin of harvest on the
398 in vitro biological activity of *Palmaria palmata* protein hydrolysates. Food Res Int 62:746–52.
- 399 Kain JM (1986) Plant size and reproductive phenology of six species of Rhodophyta in subtidal Isle of Man.
400 Br. Phycol J 21(2):129–38.
- 401 Kerrison PD, Stanley MS, Hughes AD (2018) Textile substrate seeding of *Saccharina latissima* sporophytes
402 using a binder: An effective method for the aquaculture of kelp. Algal Res 33:352–57.
- 403 Lee D, Mizuho N, Yutaka S, Hiroki S (2017) Anti-inflammatory effects of dulse (*Palmaria palmata*) resulting
404 from the simultaneous water-extraction of phycobiliproteins and chlorophyll a. Food Res Int 100:514–
405 21.
- 406 Le Gall L, Pien S, Rusig AM (2004) Cultivation of *Palmaria palmata* (Palmariales, Rhodophyta) from isolated
407 spores in semi-controlled conditions. Aquaculture 229(1–4):181–91.
- 408 Macler BA, West JA (1987) Life history and physiology of the red alga, *Gelidium coulteri*, in unialgal culture.
409 Aquaculture 61(3–4):281–93.

- 410 Matos J, Costa S, Rodrigues A, Pereira R, Pinto IS (2006) Experimental integrated aquaculture of fish and
411 red seaweeds in Northern Portugal. *Aquaculture* 252(1):31–42..
- 412 Mine I, Tatewaki M (1994) Attachment and fusion of gametes during fertilization of *Palmaria* sp.
413 (Rhodophyta). *J Phycol* 30:55–66.
- 414 Morgan KC, Simpson FJ (1981) The cultivation of *Palmaria palmata*: Effect of light intensity and temperature
415 on growth and chemical composition. *Bot Mar* 24(10):547–52.
- 416 Mouritsen OG, Dawczynski C, Duelund L, Jahreis G, Vetter W, Schröder M (2013) On the human
417 consumption of the red seaweed dulse (*Palmaria palmata* (L.) Weber & Mohr). *J Appl Phycol*
418 25(6):1777–91.
- 419 Pang S, Lüning K (2004) Tank cultivation of the red alga *Palmaria palmata*: Effects of intermittent light on
420 growth rate, yield and growth kinetics. *J Appl Phycol* 16(2):93–99.
- 421 Sáez F, Macchiavello J, Fonck E, Bulboa C (2008) The role of the secondary attachment disc in the
422 vegetative propagation of *Chondracanthus chamissoi* (Gigartinales; Rhodophyta). *Aquat Bot* 89 (1):63–
423 65.
- 424 Sanderson JC, Dring MJ, Davidson K, Kelly MS (2012) Culture, yield and bioremediation potential of
425 *Palmaria palmata* (Linnaeus) Weber & Mohr and *Saccharina latissima* (Linnaeus) C.E. Lane, C. Mayes,
426 Druehl & G.W. Saunders adjacent to fish farm cages in Northwest Scotland. *Aquaculture* 354–
427 355:128–135.
- 428 Sanderson JC (2006) Reducing the environmental impact of sea-cage fish farming through cultivation of
429 seaweed. PhD thesis, UHI Millennium Institute & Scottish Association of Marine Science (SAMS)
- 430 Santelices B, Varela D (1994) Abiotic control of reattachment in *Gelidium chilense* (Montagne) Santelices &
431 Montalva (Gelidiales; Rhodophyta). *J Exp Mar Biol Ecol* 177:145–55.
- 432 Schiener P, Zhao S, Theodoridou K, Carey M, Mooney-McAuley K, Greenwell C (2017) The nutritional
433 aspects of biorefined *Saccharina latissima*, *Ascophyllum nodosum* and *Palmaria palmata*. *Biomass*
434 *Conversion and Biorefinery* 7(2):221–35.
- 435 Schmedes PS, Nielsen MM, Petersen JK (2019) Improved *Palmaria palmata* hatchery methods for
436 tetraspore release, even settlement and high seedling survival using strong water agitation and
437 macerated propagules. *Algal Res* 40. doi: 10.1016/j.algal.2019.101494
- 438 van der Meer JP, Chen LCM (1979) Evidence for sexual reproduction in the red algae *Palmaria palmata* and
439 *Halosaccion ramentaceum*. *Can J Bot* 57(21):2452–59.
- 440 van der Meer JP, Todd ER (1980) The life history of *Palmaria palmata* in culture. A new type for the
441 Rhodophyta. *Can J Bot* 58:1250–56.
- 442 Werner A, Dring MJ (2011) Aquaculture explained: Cultivating *Palmaria palmata*. Irish Sea Fisheries Board,

443 report no. 27 - project PBA/SW/07/001. Dublin.

444 Wood E (2018) Cultivation of *Palmaria palmata*: spore release, vegetative propagation and decontamination
445 techniques. Scottish Association for Marine Science (SAMS) and University of the Highlands and Islands.

446

447 Acknowledgements

448 We sincerely thank Pascal David Alain Barreau and Kasper Lenda Andersen for their technical help installing
449 the experimental setup, as well as for system maintenance and data collection. The study was funded by the
450 Joint Doctoral Degree agreement between the National Institute of Aquatic Resources (DTU Aqua) at
451 Technical University of Denmark and the Norwegian University of Science and Technology (NTNU), Norway,
452 as well as the Tang.nu project (under Grant Agreement No. 13744, Velux Foundation) and the MacroSea
453 project, Grant no. 254883, funded by the Research Council of Norway.

454 Compliance with Ethical Standards

455 Informed consent of the document with no conflict of interest. We did not conduct research on human or
456 animals.

457

458 Figure 1. A: Parallel flow-through setup of conical seeding tanks (30 L) used for experiment 1. B: The sori
459 packages were fixed centrally above the net spirals and 2 cm below water surface. Effluent spores were
460 detained in spore-detaining tanks (SDTs). C: Aeration (1.2 L min^{-1}) from the bottom provided hemispherical
461 water circulation and dispersal of released spores.

462 Figure 2. A: Fertile tetrasporophytic fronds were cleaned and the presence of dark-red sporangia were
463 validated (B). C: Sori packages (5, 10, or 15 g FW) were prepared by placing the sori between two layers of
464 green plastic mesh ($n=3$).

465 Figure 3. A: Dark-red aggregates of spores on the bottom of spore-detaining tanks (STDs) accumulated
466 during sporulation. B: A well-mixed spore solution was obtained by dislodging the aggregates and
467 macerating them. C: The number spores were counted in five random fields (each of 0.0035 mm^2) and used
468 as a basis to estimate the total amount of detained spores.

469 Figure 4. A: Detained and macerated spores of *P. palmata* at the bottom of a 5 L Plexiglas tank. B: Plexiglas
470 tank after dislodgement of propagules on the left side of the tank. C: Propagule mix after dislodgement and
471 maceration. The solution was inspected before (D) and after (E) the maceration treatment. F: Single
472 individual seedling displaying hair-like proliferations from the basal disc area. Scale bar represent 500 μm .

473 Figure 5. A: Fertile male gametophytes inspected by stereomicroscopy. B: cross-section microscopy was
474 used to verify reproductive appearance of the males. C, D: Spherical male gamete released in a solution.

475 Figure 6. The number of spores (A-C) and seedlings (D-F) attached to the nets and the spore germination
476 percentage (G-I). 4 cm net pieces were counted on day 3, 19 and 32 after each of the three consecutive
477 periods (Day 0-3, Day 3-6, Day 6-9) with the use of different sori amount (5, 10, 15 g). Data is presented as
478 $\text{mean} \pm \text{SE}$, $n=3$.

479 Figure 7. A-I: The extrapolated number of spores attached to the nets and (D-F) the total number of detained
480 spores collected in the spore-detaining tanks (SDTs) as a function of three consecutive seeding periods and
481 the use of three different amount of sori (5, 10, 15 g FW). G-I: Net seeding efficiency, calculated from
482 equation 1. Data is presented as $\text{mean} \pm \text{SE}$, $n=3$.

483 Figure 8. Attached spores (A-B) and seedlings (C-D) on rope pieces seeded with macerated propagule
484 solution at day 3 and 10 and as a function of age at dislodgement (day 27-29 vs. day 37-39) and agitation
485 period (day 0-2 vs. day 10-12). Data is presented as outlier box plots (1st, 3rd quartile whisker) based on
486 three subpart counts on each of ten pieces of rope ($n=10$, $N=30$).

487 Figure 9. A: The normalized seedling count in petri dish cultures ($n=10$) of macerated *P. palmata* propagules
488 as an effect of adding male gametes. B: The normalized seedling count on spore-seeded rope cultures
489 ($n=11$) as an effect of adding male gametes. Data is presented as outlier box plots (1st, 3rd quartile whisker).