



Seasonal variation of drop size distribution: influence on rainfall kinetic energy

Tilg, Anna-Maria; Vejen, Flemming; Nielsen, Morten; Hasager, Charlotte Bay

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Tilg, A-M., Vejen, F., Nielsen, M., & Hasager, C. B. (2019). *Seasonal variation of drop size distribution: influence on rainfall kinetic energy*. Abstract from EGU General Assembly 2019, Vienna, Austria.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Seasonal variation of drop size distribution: influence on rainfall kinetic energy

Anna-Maria Tilg (1), Flemming Vejen (2), Morten Nielsen (1), and Charlotte Hasager (1)

(1) Technical University of Denmark, DTU Wind Energy, Roskilde, Denmark (anmt@dtu.dk), (2) Danish Meteorological Institute, Copenhagen, Denmark

The rainfall kinetic energy (RKE) influences erosion processes to a high degree. The most prominent example is soil erosion (Wischmeier and Smith 1958) but also the erosion of the leading edges of turbine blades of wind turbines is related to rainfall kinetic energy (Bech et al. 2018). Based on drop size distribution (DSD) measurements of disdrometers, the RKE can be calculated directly. It is known that DSD and therefore RKE vary during an event but there are fewer studies on the seasonal variation of these parameters. To estimate seasons with higher or lower erosion potential, it is important to know more about the seasonal behaviour of DSD and RKE.

For our analysis we used five years of quality controlled Thies disdrometer data. The disdrometer was installed in connection with the Danish hydrological observatory project – HOBE (Jensen and Illangasekare 2011) in the western part of Denmark.

The results show that the mean concentration of drops and RKE are higher in summer and autumn and lower in winter and spring. Furthermore, the data indicates that drops with a diameter larger than 1 mm are more frequent in the seasons June to August (JJA) and September to November (SON). Whereas the concentration of drops decreases with increasing diameter for all seasons, RKE reaches a maximum at a mean drop diameter between 1.125 and 1.625 mm depending on the season. In the season December to February (DJF) raindrops with a mean diameter of 1.125 mm provide most kinetic energy. In contrast, raindrops with a mean diameter of 1.625 mm are responsible for most of RKE in the season June to August (JJA).

With these results, we conclude that the potential of erosion is higher in summer and autumn because of higher concentration of raindrops, higher percentage of larger raindrops and therefore higher RKE.

References

- Bech, J. I., Hasager, C. B., & Bak, C. (2018). Extending the life of wind turbine blade leading edges by reducing the tip speed during extreme precipitation events. *Wind Energy Science*, (in press), 729–748. doi:10.5194/wes-2017-62
- Jensen, K. H., & Illangasekare, T. H. (2011). HOBE: A Hydrological Observatory. *Vadose Zone Journal*, 10(1), 1. doi:10.2136/vzj2011.0006
- Wischmeier, W. H., & Smith, D. D. (1958). Rainfall energy and its relation to soil loss. *Transactions of the American Geophysical Union*, (39), 285–291. doi:10.1029/TR039i002p00285