The inflammatory response to bone infection A review based on animal models and human patients

Lüthje, Freja Lea; Jensen, Louise Kruse; Elvang Jensen, Henrik; Skovgaard, Kerstin

Published in:
APMIS

Link to article, DOI:
[10.1111/apm.13027](https://doi.org/10.1111/apm.13027)

Publication date:
2020

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Lüthje, F. L., Jensen, L. K., Elvang Jensen, H., & Skovgaard, K. (2020). The inflammatory response to bone infection A review based on animal models and human patients. *APMIS*, 128(4), 275-286.
<https://doi.org/10.1111/apm.13027>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The inflammatory response to bone infection

A review based on animal models and human patients

Freja Lea Lüthje^{a,b}(frejal@sund.ku.dk), Louise Kruse Jensen^a(louise-k@sund.ku.dk), Henrik Elvang Jensen^a(elvang@sund.ku.dk), Kerstin Skovgaard^b(kesk@dtu.dk)

^a Department of Veterinary and Animal Science, University of Copenhagen, 1870 Frederiksberg C, Denmark

^b Department of Biotechnology and Biomedicine, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark

Running head: The inflammatory response to bone infection

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/APM.13027](https://doi.org/10.1111/APM.13027)

This article is protected by copyright. All rights reserved

Summary

Authors: Freja Lea Lüthje (FLL), Louise Kruse Jensen (LKJ), Henrik Elvang Jensen (HEJ), Kerstin Skovgaard (KS)

Title: The inflammatory response to bone infection

Subtitle: A review based on animal models and human patients

Bone infections are difficult to diagnose and treat, especially when a prosthetic joint replacement or implant is involved. Bone loss is a major complication of osteomyelitis, but the mechanism behind has mainly been investigated in cell cultures and has not been confirmed in human settings. Inflammation is important in initiating an appropriate immune response to invading pathogens. However, many of the signaling molecules used by the immune system can also modulate bone remodeling and contribute to bone resorption during osteomyelitis. Our current knowledge of the inflammatory response relies heavily on animal models as research based on human samples is scarce. *Staphylococcus aureus* is one of the most common causes of bone infections and is the pathogen of choice in animal models. The regulation of inflammatory genes during prosthetic joint infections and implant associated osteomyelitis has only been studied in rodent models. It is important to consider the validity of an animal model when results are extrapolated to humans, and both bone composition and the immune system of pigs has been shown to be more similar to humans, than **to** rodents. Here *in vivo* studies on the inflammatory response to prosthetic joint infections and implant associated osteomyelitis is reviewed.

Keywords: (max 5) osteomyelitis, inflammation, prosthetic joint infection, cytokines, immune response

Corresponding author: Freja Lea Lüthje (Address: Ridebanevej 3, 1870 Frederiksberg C, Denmark; Mail: frejal@sund.ku.dk; Phone: +45 2229 5904)

Introduction

Each year joint replacements increase the quality of life for millions of people. In Scandinavia, more than 150,000 knee arthroplasties were carried out from 1999 to 2007 with a continuous increase every year (1). In the U.S., the number of hip and knee arthroplasties is expected to rise from around 600,000 procedures in 2003 to more than 4 million in 2030 (2). The increasing number of joint replacements has resulted in an increase in prosthetic joint infections (PJI) (3). Along with joint prosthetics, other types of bone implants are also frequently used for fixation of bone fractures and infections of such implants are called fracture related infections or implant associated osteomyelitis (IAO). Bone infections like PJI and IAO are severe long-term conditions and recurrence of infection is common (4,5). When a chronic implant-related bone infection is established, it is unlikely to be resolved by antibiotic treatment alone and, therefore, requires surgical intervention (6). Furthermore, careful debridement of necrotic bone is necessary to avoid recurrent infections, and two-stage exchange is recommended in PJI for the same reason (7). This entails surgical removal of the infected implant and necrotic bone, followed by 4-6 weeks of antibiotics, before a new implant can be inserted. Thus, the current treatment is a prolonged and painful affair. In addition, it can also be difficult to diagnose PJI as it can be hard to distinguish from aseptic implant failure (8). Bone infections are very complex diseases where both local bone cells and the infecting pathogen can modify the immune response. Therefore, it is highly relevant to study the local osseous immune response in animal models and osteomyelitis patients in order to develop better treatments and diagnostic tools. This review aims to describe how bone remodeling can be affected by the immune response raised towards bone infection and to summarize the regulation of inflammatory genes found *in vivo*. Animal studies have been limited to those involving PJI and IAO models, while all studies of human bone infections have been included.

Normal bone homeostasis

Bone is a dynamic tissue under constant remodeling to adapt to changes in mechanical loading and to repair microcracks. The remodeling consists of bone resorption carried out by osteoclasts and bone formation by osteoblasts. Osteoclasts are the primary cells to degrade mineralized bone and are thus essential in handling damaged bone. They belong to the hematopoietic lineage like macrophages. Fully differentiated osteoclasts are giant polykaryons. Osteoblasts belong to the mesenchymal lineage and their primary function is to form bone *i.e.* osteoid (9). Terminally differentiated osteoblasts surrounded by mineralized osteoid become osteocytes, which are the most numerous cell type in bone tissue and are believed to play a pivotal role in sensing bone damage and initiating remodeling (10).

Osteoblasts also have an important role in regulating osteoclast activity. Osteoclast precursors require binding of receptor activator NF kappa B ligand (RANKL) to the receptor activator NF kappa B (RANK) to initiate the formation of multinuclear cells by cell fusions (11). RANKL is produced by osteoblasts, which can also synthesize the RANKL decoy receptor osteoprotegerin (OPG) that inhibits RANK-RANKL interactions (Figure 1). Thus by expressing RANKL or OPG osteoblasts can increase or decrease osteoclast activity, respectively, and the ratio between these molecules is therefore used as an indicator of bone erosion (12).

Osteoclast activity is also regulated by many other molecules (for an extensive review see Loi et al., 2016), and several inflammatory genes can affect osteoclastogenesis. To complicate matters further, RANK/RANKL are also involved in regulating the immune system (14).

Apart from regulating bone metabolism osteoclasts and osteoblasts also interact with the immune system. In fact, the close interplay between the immune system and the skeletal system has led to a new interdisciplinary research field called osteoimmunology (15). Osteoclasts have been shown to produce interleukin (IL)-10, IL-6, transforming growth factor beta and tumor necrosis factor (TNF) in response to lipopolysaccharide and to function as antigen presenting cells *in vitro* (16). **In addition osteoblast cultures have been reported to produce IL-6, IL-12, CXCL8 and CCL2 in response to *Staphylococcus aureus***, and thus take part in immune activation (17–19). Osteoblasts have also been shown to activate T lymphocytes by expressing MHC class II molecules, which are normally only expressed by professional antigen-presenting cells (20). In addition, the antimicrobial peptides human beta-defensin-1, -2 and -3 can be produced by osteoblasts and are expressed in human bone (21–23). However, Chang et al. argues that it is actually resident tissue macrophages, called osteomacs, that respond to bacterial stimuli in osteoblast cultures (24).

Osteolysis

Bone loss is a major complication of osteomyelitis. Experiments in immune deficient mice with periodontitis indicate that it is mainly the immune system and not the bacteria that is responsible for osteolysis in bacterial bone infections (25,26). Thus, the bone loss can be considered collateral damage caused by excessive inflammation. *In vitro* studies suggest that bone infection leads to both increased osteoclast activity and decreased osteoblast activity. *Staphylococcus aureus* and *S. aureus* biofilm conditioned media reduces osteoblast viability and inhibits bone formation. At the same time, the secretion of RANKL is upregulated, leading to increased bone resorption (27,28) (Figure 1). *Staphylococcus aureus* infected osteoblasts also upregulate cell surface receptors that can induce apoptosis (29) and

downregulate the expression of OPG, which can lead to increased osteoclast activity. *S. aureus* also affect osteoclasts directly by promoting cellular fusion (30). Therefore, osteolysis in osteomyelitis has typically been ascribed to an increase in RANKL-induced osteoclastogenesis (28). However, the increase in RANKL expression and osteoclast activity has not been confirmed in human settings.

The only *in vivo* study of local RANKL expression during PJI revealed no difference in RANKL-expression between patients with PJI and aseptic loosening of implants. A moderate upregulation of RANKL at the osteolytic site compared to a distant site was also reported (31). In addition, there was no difference in the serum level of RANKL in patients with PJI and aseptic loosening (32).

The assumption that osteoclasts are solely responsible for the bone loss during osteomyelitis probably also builds on the false notion that osteoclast are the only cell type to degrade bone (33). However, monocytes, fibroblasts and neutrophils have all been shown able to degrade bone matrix (34–37).

Staphylococcus aureus

Staphylococcus aureus, one of the most common etiologies of bone infections (38–40), is the preferred pathogen in animal models of osteomyelitis (Table 1). The persistence of *S. aureus* infections is primarily attributed to the formation of biofilms. Biofilms consist of microorganisms organized in a matrix of extracellular polymeric substances. The matrix contains polysaccharides, proteins, lipids and nucleic acids excreted by bacteria (41) but can also include compounds from the host (42). **Both diagnosis and treatment of biofilm infections can be difficult and requires special clinical attention(43).** Biofilm formation is a survival strategy that protects the bacteria from UV radiation, pH stress, chemical exposure, dehydration and antibiotic treatment (41). The tolerance to antibiotics has been reported to increase 100- to 1,000-fold in biofilms compared to the same planktonic bacterial strain (44). Biofilms also protect bacteria from phagocytosis by macrophages **and neutrophils** (45,46).

It is also well established, that *S. aureus* can be taken up by osteoblasts *in vitro* and there are a few reports of *in vivo* internalization in human osteocytes (47,48). It has been speculated that internalization is a virulence factor protecting the bacterium from the immune system and antibiotics (49,50). However, the prevalence of *S. aureus* uptake by osteoblasts and osteocytes *in vivo* is unknown and, therefore, it is difficult to assess the importance in the recalcitrance of bone infections.

In addition, *S. aureus* have several strategies to evade both the innate and adaptive immune response (51,52): e.g. it can prevent complement activation (53), induce apoptosis in B cells by staphylococcal protein A (54) and in macrophages by degrading neutrophil extracellular traps (55).

Inflammation

Inflammation is a fundamental defense response to injury and infection, and is essential for pathogen defense and tissue healing. However, it is important that the inflammatory response is tightly regulated as immune dysregulation can cause tissue damage. As the host in many cases is unable to clear the bone infection, the infection turns chronic and subsequently the inflammatory processes leads to osteolysis (56). **Clinically ,there is no clearly defined line between acute and chronic osteomyelitis (57). Acute osteomyelitis describes recent bone infections causing systemic inflammation (58). In contrast, chronic osteomyelitis has a limited systemic response and is associated with avascular necrotic bone (7).**

The host immune response to PJI and IAO is affected by multiple contributing factors. These include the bone damage from implant insertion, the implant *per se* and the pathogens. The implant insertion procedure causes osteonecrosis due to high temperatures (59) while the implant as an foreign body also elicits a molecular immune response (60). Following implantation plasma proteins immediately coats the implant, the compliment and the coagulation systems are activated, immune cells are recruited and cytokines released. All these action are part of the foreign body reaction (61).

Even though immune cells, especially neutrophils and macrophages (61,62), migrate to the implant, the presence of a foreign body lowers the number of pathogens necessary to cause an infection (63,64). This can be explained by reduced phagocyte function at the implant site. As the neutrophils are unable to phagocytose the implant they release pro-inflammatory cytokines and neutrophil extracellular traps, which can cause neutrophil apoptosis (62). Macrophages, upon adhesion to the implant, releases reactive oxygen intermediates, degradative enzymes and acids in an attempt to degrade the foreign body. The macrophages are left unsuccessful and exhausted unable to produce more biocidal agents (61). In addition, the surface chemistry of the implant, especially hydrophilic and anionic surfaces, can induce apoptosis in macrophages (65). **Moreover, oxygen availability is important for the neutrophilic microbicidal function (46) and the oxygen supply is limited during bone infections (66). The vascular channels in bone are compressed by inflammatory processes, such as pus formation, leading to restricted blood perfusion (67). At the same time, both neutrophils and the infecting microorganism can consume large amounts of oxygen resulting in hypoxia around biofilm infections (68).**

Several pro-inflammatory and anti-inflammatory cytokines have been examined during bone infections in animal models and human patients (Tables 1 and 2). The results of these studies are reviewed in the following two sections. Apart from modulating the inflammatory response many of the cytokines have been shown to affect bone metabolism *in vitro* (69). The known *in vitro* effect on bone cells of the cytokines upregulated *in vivo* is summarized in Figure 2.

Pro-inflammatory cytokines in osteomyelitis

TNF is a pro-inflammatory cytokine that together with IL-1 and IL-6 is essential for initiating an inflammatory response to trauma and infection (70). TNF is upregulated in the bone of patients with PJI compared to patients with aseptic loosening of implants (31) and in the bone of several murine IAO models (71–73). *In vitro* TNF stimulates osteoclastogenesis (74), inhibits osteoblast differentiation (75,76) and induces IL-6 production in osteoblasts (77).

IL-1 β is highly elevated in bones during IAO in mice (72,73) and in humans with PJI (78). IL-1 β -deficient mice with PJI showed increased bacterial burdens and fewer neutrophils in the infected joint compared to wild type (WT) mice (79). Apart from attracting neutrophils and maintaining inflammation, IL-1 β also affects bone remodeling. Like TNF, IL-1 β inhibits osteoblast differentiation (76), while it stimulates the production of IL-6 and RANKL, leading to increased osteoclast activity (77,80). In addition, IL-1 α , which binds to the same receptors as IL-1 β , can directly stimulate osteoclast development in cultures without osteoblasts (81), and IL-1 α is upregulated in the serum of osteomyelitis patients (82).

IL-6 has a multitude of both innate and adaptive immune functions and have both pro-inflammatory and anti-inflammatory properties (83). In an *ex vivo* study of fetal mouse calvaria IL-6 was shown to increase bone resorption and osteoclast numbers (77). However, IL-6 have also been shown to inhibit osteoclastogenesis by skewing monocytic cells away from osteoclast differentiation and towards the macrophage lineage (84). Two studies of IL-6 deficient mice with periapical infection showed opposing results, one found a greater bone loss in the IL-6 deficient mice than in WT mice (83), while the other failed to find any bone loss in the IL-6 deficient infected mice (85). Thus, the effect of IL-6 *in vivo* during bone infections is still unclear and awaits further studies. The expression of IL-6 is upregulated in several mouse models of IAO (71,72,86). Both human and murine osteoblasts increase the production of IL-6 in response to *S. aureus* (17,18,87). In patients with posttraumatic osteomyelitis the expression of IL-6 was found to be higher both in serum and in the infected bones of acute cases compared to chronic cases (58). Asensi et al. (2004) also found a higher level of IL-6 in serum from osteomyelitis patients (etiology not specified) compared to healthy controls.

IL-17A is a proinflammatory cytokine involved in the immune response against bacterial infections (88) and it is highly upregulated in murine IAO models (71,73). IL-17A induces the differentiation of osteoblasts and the production of RANKL both *in vitro* and *in vivo*, thus promoting both bone formation and degradation (89–91).

The chemokine CXCL8 (a.k.a. IL-8) is upregulated locally in PJI patients compared to patients with aseptic loosening of implants (31,92). In addition, human osteoblast cell lines exposed to *S. aureus* upregulate the expression of CXCL8 (17,93). CXCL8 is an important chemoattractant for neutrophils during infection and enhances osteoclastogenesis (94). Despite the central role of CXCL8 in the early immune defense of pig and human, this chemokine has not been identified in mice and is an example of differences in human and murine innate immune systems (95).

The chemokine CXCL2 is also upregulated in patients with PJI compared to patients with aseptic loosening (31). This upregulation has also been shown in a murine model of IAO (78). In addition, CCL5, colony stimulating factor 3 and IL-12 are all upregulated in murine IAO models (71,78), and CCL5, CXCL9, CXCL10 and CXCL11 are upregulated in infected bone samples of patients with PJI compared to samples from uninfected bone (47).

Anti-inflammatory cytokines in osteomyelitis

IL-4 inhibits osteoclast differentiation *in vitro* (96) and reduced the number of osteoclasts in an murine model (97). This anti-inflammatory cytokine was upregulated at day 14 in a murine model of IAO (72) and in the bone during aseptic bone healing (73).

IL-10, like IL-4, suppresses the development of osteoclasts *in vitro* (98). IL-10 was upregulated in a murine IAO model from day 5, and it was concluded that IL-10 enhanced the persistence of biofilm infection with *S. aureus* in C57BL/6 mice after 14 days (99). In contrast a study comparing the immune response in C57BL/6 and BALB/c mice with IAO concluded that BALB/c mice benefited from a higher IL-4 and IL-10 expression after 49 days (100). Thus, the length of the experiment greatly affects the overall conclusion. In the comparison of Th2 biased BALB/c mice and Th1-biased C57BL/6 mice, Prabhakara *et al.* found no difference between the mouse strains at day 14, where the study by Heim *et al.* ended. However, at day 21 more than half of the BALB/c mice (with an upregulation of IL-4 and IL-10 compared to the C57BL/6 mice) had cleared the infection spontaneously. In addition, biofilm was only formed on the implants of C57BL/6 mice. As Heim *et al.* measured the bacterial burden via plate counts dormant cells which are common in biofilms was not considered. Since the study by Heim *et al.* only measured bacteria in a culturable state the results are mainly applicable for acute infections with planktonic bacteria, while the

study by Prabhakara *et al.* indicates that upregulation of IL-4 and IL-10 can prevent development of chronic biofilm infections in mice. Another possible explanation for the opposing conclusions is that they used different bacterial strains (Table 1) that could contain different virulence **genes**.

Animal models

As seen from Table 1, all *in vivo* studies of cytokine expression in IAO and PJI have been carried out in mice or rats, so apart from studies using samples from human patients (Table 2), all knowledge about cytokine production and pathways involved in bone infections relies on rodent models. Therefore, it is important to consider how the rodent bone physiology and immune system differ to humans, when assessing the validity of rodent models. Aerssens et al. (1998) compared the composition and density of human, dog, pig, cow, sheep, chicken, and rat cortical femoral bone, and found rat bone to resemble human bone the least. The immune system of mice also differs from the human counterpart in several aspects, e.g. the human genes IL-26, CXCL8 and CXCR1 has no homolog in mice and neutrophils are the dominating leukocyte in peripheral blood in humans while lymphocytes dominate in mice (102,103). In addition, the transcriptional response to inflammatory conditions in mice does not always mimics the human response well. In fact, no correlation was found between the genes regulated under inflammatory conditions in humans and the orthologue genes in relevant murine models (104). Furthermore, inbred strains are preferred in rodent studies and there can be major differences in the development of disease between different mouse strains (100). This means that extrapolation of results from rodents to humans should be done with care. Unfortunately, human bone samples of especially healthy control tissue are very difficult to obtain (Table 2). Therefore, more comparable animal models should be considered.

The use of porcine models of infectious diseases are becoming increasingly popular (105), and pigs have several advantages compared to mice as models for osteomyelitis. Their size makes it possible to operate with the same tools as in humans and adequate quantities of tissue can be sampled, so pooling of samples is unnecessary. In addition, the bone anatomy and physiology of pigs are similar to humans (106) and the porcine immune system also resembles humans more than that of rodents (95,107). **The limitations of porcine models includes fewer available molecular tools, considerably less scientific literature regarding the porcine genome and immune system compared to rodents, and a relatively high cost – often resulting in small sample sizes (108).**

In a recent study of a porcine IAO model, **we saw an upregulation in gene expression of many of the same inflammatory markers** - including TNF, IL-1 β , IL-6 and IL-17A (Lüthje et al., unpublished data) - as

are seen in rodent models and patients with osteomyelitis (Table 1 and 2). Furthermore, a significant upregulation of IL-26, which has no homolog in rodents but is present in humans (103), was found.

Regardless of the animal model, new results of the pro- and anti-inflammatory orchestration of bone infection must be confirmed in human samples whenever possible. Currently, the descriptions of the immune response during human osteomyelitis are at best incomplete, making it difficult to predict good biomarkers and treatment targets. Furthermore, most animal model studies are done in relatively young animals while it is mainly elderly people that require prosthetic joint replacements. It is known that both bone metabolism, inflammation and the immune system are affected by age (109,110). In a rat IAO model the cytokine expression was very different in young and old rats (111), consequently, age is a factor that further complicates the translation of results from the laboratory to the clinic.

Conclusion

The immune response towards an infection is a complex process, and this is especially true in bone tissue where the immune activation and signaling also affect bone remodeling. Thus, a delicate balance is necessary to regulate inflammation in bone infections. An inflammatory response is necessary to control the invading pathogens and dampening the inflammation can increase the bacterial burden (99).

However, a too strong response can damage the host tissue, creating devitalized bone for *S. aureus* to develop biofilm on (56). Clearing of the infection is also complicated by the fact that classical immune cells are poorly equipped to degrade dead and infected bone tissue. Therefore, regulation of bone degradation is very important at the site of infection and many pro-inflammatory cytokines also promote osteolysis (69). However, the effect of cytokines on bone metabolism is mostly based on in vitro studies, which rarely provide the complete picture. Only a few studies of the human immune response to osteomyelitis have been conducted, and rarely with samples from bone (Table 2). This is because it is difficult to take bone samples, especially healthy control tissue, without harming the patients. Therefore, it is important that the chosen animal models mimic the human immune response as much as possible. Furthermore, researchers should be aware of known differences in the immune response between the chosen animal species and humans.

Acknowledgement

This paper was financed by the European Union's Horizon 2020 research and innovation program under NOMORFILM project grant agreement No 634588 and by the Danish Medical Research Council under the grant no. 4005-00035B.

Literature

1. Robertsson O, Bizjajeva S, Fenstad AM, Furnes O, Lidgren L, Mehnert F, et al. Knee arthroplasty in Denmark, Norway and Sweden. *Acta Orthop*. 2010 Feb 24;81(1):82–9.
2. Kurtz S. Projections of Primary and Revision Hip and Knee Arthroplasty in the United States from 2005 to 2030. *J Bone Jt Surg*. 2007 Apr 1;89(4):780.
3. Kurtz SM, Lau E, Schmier J, Ong KL, Zhao K, Parvizi J. Infection Burden for Hip and Knee Arthroplasty in the United States. *J Arthroplasty*. 2008 Oct;23(7):984–91.
4. Lee J, Kang CI, Lee JH, Joung M, Moon S, Wi YM, et al. Risk factors for treatment failure in patients with prosthetic joint infections. *J Hosp Infect*. 2010;75(4):273–6.
5. Marculescu CE, Berbari EF, Hanssen AD, Steckelberg JM, Harmsen SW, Mandrekar JN, et al. Outcome of Prosthetic Joint Infections Treated with Debridement and Retention of Components. *Clin Infect Dis*. 2006;42:471–8.
6. Del Pozo J, Patel R. Infection Associated with Prosthetic Joints. *N Engl J Med*. 2009;361(8):787–94.
7. Lew DP, Waldvogel FA. Osteomyelitis. *Lancet*. 2004;364:369–79.
8. Tande AJ, Patel R. Prosthetic joint infection. *Clin Microbiol Rev*. 2014;27(2):302–45.
9. Rauner M, Stein N, Hofbauer LC. Basics of Bone Biology. Pietschmann P, editor. *Principles of Osteoimmunology: Molecular Mechanisms and Clinical Applications*. Vienna: Springer Vienna; 2012. 1–26 p.
10. Bellido T. Osteocyte-driven bone remodeling. *Calcif Tissue Int*. 2014;94(1):25–34.
11. Hofbauer LC, Heufelder AE. Role of receptor activator of nuclear factor- κ B ligand and osteoprotegerin in bone cell biology. *J Mol Med*. 2001 Jun 3;79(5–6):243–53.

12. Matsuo K, Irie N. Osteoclast–osteoblast communication. *Arch Biochem Biophys*. 2008;473(2):201–9.
13. Loi F, Córdova LA, Pajarinen J, Lin T, Yao Z, Goodman SB. Inflammation, fracture and bone repair. *Bone*. 2016;86:119–30.
14. Guerrini MM, Takayanagi H. The immune system, bone and RANKL. *Arch Biochem Biophys*. 2014 Nov 1;561:118–23.
15. Arron JR, Choi Y. Bone versus immune system. *Nature*. 2000 Nov 30 408(6812):535–6.
16. Li H, Hong S, Qian J, Zheng Y, Yang J, Yi Q. Cross talk between the bone and immune systems: osteoclasts function as antigen-presenting cells and activate CD4+ and CD8+ T cells. *Blood*. 2010 Jul 15;116(2):210–7.
17. Dapunt U, Giese T, Stegmaier S, Moghaddam A, Hänsch GM. The osteoblast as an inflammatory cell: production of cytokines in response to bacteria and components of bacterial biofilms. *BMC Musculoskelet Disord*. 2016;17(1):243.
18. Bost KL, Ramp WK, Nicholson NC, Bento JL, Marriott I, Hudson MC. Staphylococcus aureus infection of mouse or human osteoblasts induces high levels of interleukin-6 and interleukin-12 production. *J Infect Dis*. 1999 Dec 1;180(6):1912–20.
19. Bost KL, Bento JL, Petty CC, Schrum LW, Hudson MC, Marriott I. Monocyte chemoattractant protein-1 expression by osteoblasts following infection with Staphylococcus aureus or Salmonella. *J Interferon Cytokine Res*. 2001 May;21(5):297–304.
20. Schrum LW, Bost KL, Hudson MC, Marriott I. Bacterial infection induces expression of functional MHC class II molecules in murine and human osteoblasts. *Bone*. 2003 Nov;33(5):812–21.
21. Warnke PH, Springer IN, Russo PAJ, Wiltfang J, Essig H, Kosmahl M, et al. Innate immunity in human bone. *Bone*. 2006;38(3):400–8.
22. Varoga D, Tohidnezhad M, Paulsen F, Wruck CJ, Brandenburg L, Mentlein R, et al. The role of human beta-defensin-2 in bone. *J Anat*. 2008 Dec;213(6):749–57.
23. Varoga D, Wruck CJ, Tohidnezhad M, Brandenburg L, Paulsen F, Mentlein R, et al. Osteoblasts participate in the innate immunity of the bone by producing human beta defensin-3. *Histochem*

Cell Biol. 2009;131(2):207–18.

24. Chang MK, Raggatt L, Alexander KA, Kuliwaba JS, Fazzalari NL, Schroder K, et al. Osteal Tissue Macrophages Are Intercalated throughout Human and Mouse Bone Lining Tissues and Regulate Osteoblast Function In Vitro and In Vivo. *J Immunol*. 2008 Jul 7;181(2):1232–44.
25. Teng YTA, Nguyen H, Gao X, Kong YY, Gorczynski RM, Singh B, et al. Functional human T-cell immunity and osteoprotegerin ligand control alveolar bone destruction in periodontal infection. *J Clin Invest*. 2000;106(6):59–67.
26. Baker PJ, Howe L, Garneau J, Roopenian DC. T cell knockout mice have diminished alveolar bone loss after oral infection with *Porphyromonas gingivalis*. *FEMS Immunol Med Microbiol*. 2002;34(1):45–50.
27. Widaa A, Claro T, Foster TJ, O'Brien FJ, Kerrigan SW. *Staphylococcus aureus* Protein A Plays a Critical Role in Mediating Bone Destruction and Bone Loss in Osteomyelitis. *PLoS One*. 2012 Jul 11;7(7):e40586.
28. Sanchez CJ, Ward CL, Romano DR, Hurtgen BJ, Hardy SK, Woodbury RL, et al. *Staphylococcus aureus* biofilms decrease osteoblast viability, inhibits osteogenic differentiation, and increases bone resorption in vitro. *BMC Musculoskelet Disord*. 2013 Dec 14;14(1):187.
29. Young AB, Cooley ID, Chauhan VS, Marriott I. Causative agents of osteomyelitis induce death domain-containing TNF-related apoptosis-inducing ligand receptor expression on osteoblasts. *Bone*. 2011;48(4):857–63.
30. Trouillet-Assant S, Gallet M, Nauroy P, Rasigade J-P, Flammier S, Parroche P, et al. Dual impact of live *Staphylococcus aureus* on the osteoclast lineage, leading to increased bone resorption. *J Infect Dis*. 2015 Feb 15;211(4):571–81.
31. Dapunt U, Giese T, Lasitschka F, Lehner B, Ewerbeck V, Hänsch GM. Osteoclast Generation and Cytokine Profile at Prosthetic Interfaces: A Study on Tissue of Patients with Aseptic Loosening or Implant-Associated Infections. *Eur J Inflamm*. 2014 Jan 1;12(1):147–59.
32. Friedrich MJ, Wimmer MD, Schmolders J, Strauss AC, Ploeger MM, Kohlhof H, et al. RANK-ligand and osteoprotegerin as biomarkers in the differentiation between periprosthetic joint infection and aseptic prosthesis loosening. *World J Orthop*. 2017;8(4):342–9.

33. Clarke B. Normal Bone Anatomy and Physiology. Clin J Am Soc Nephrol. 2008 Nov 1;3(Supplement 3):S131–9.
34. Pap T, Claus A, Ohtsu S, Hummel KM, Schwartz P, Drynda S, et al. Osteoclast-independent bone resorption by fibroblast-like cells. Arthritis Res Ther. 2003 Mar 26;5(3):R163–73.
35. Mundy GR, Altman AJ, Gondek MD, Bandelin JG. Direct Resorption of Bone by Human Monocytes. Science (80-). 1977;196(4294):1109–11.
36. Nanes MS, Pacifici R. Inflammatory Cytokines. In: Bronner F, Farach-Carson MC, Rubin J, editors. Bone Resorption. vol 2. London: Springer; 2005. p. 67–90.
37. Ohtsu S, Yagi H, Nakamura M, Ishii T, Kayaba S, Soga H, et al. Enhanced neutrophilic granulopoiesis in rheumatoid arthritis. Involvement of neutrophils in disease progression. J Rheumatol. 2000;27(6):1341–51.
38. Arciola C, An Y, Campoccia D, Donati M, Montanaro L. Etiology of implant orthopedic infections: A survey on 1027 clinical isolates. Int J Artif Organs. 2005 Nov;28(11):1091–100.
39. Wright JA, Nair SP. Interaction of staphylococci with bone. Int J Med Microbiol. 2010;300(2):193–204.
40. Gundtoft PH, Pedersen AB, Schønheyder HC, Møller JK, Overgaard S. One-year incidence of prosthetic joint infection in total hip arthroplasty: a cohort study with linkage of the Danish Hip Arthroplasty Register and Danish Microbiology Databases. Osteoarthr Cartil. 2017;25(5):685–93.
41. Gupta P, Sarkar S, Das B, Bhattacharjee S, Tribedi P. Biofilm, pathogenesis and prevention-a journey to break the wall: a review. Arch Microbiol. 2016 Jan;198(1):1–15.
42. Nishitani K, Sutipornpalangkul W, de Mesy Bentley KL, Varrone JJ, Bello-Irizarry SN, Ito H, et al. Quantifying the natural history of biofilm formation in vivo during the establishment of chronic implant-associated Staphylococcus aureus osteomyelitis in mice to identify critical pathogen and host factors. J Orthop Res. 2015 Sep33(9):1311–9.
43. Høiby N, Bjarnsholt T, Moser C, Bassi GL, Coenye T, Donelli G, et al. ESCMID guideline for the diagnosis and treatment of biofilm infections 2014. Clin Microbiol Infect. 2015 May 1;21:S1–25.
44. Ceri H, Olson ME, Stremick C, Read RR, Morck D, Buret a. The Calgary Biofilm Device : New

Technology for Rapid Determination of Antibiotic Susceptibilities of Bacterial Biofilms The Calgary Biofilm Device : New Technology for Rapid Determination of Antibiotic Susceptibilities of Bacterial Biofilms. *J Clin Microbiol.* 1999;37(6):1771.

45. Hanke ML, Kielian T. Deciphering mechanisms of staphylococcal biofilm evasion of host immunity. *Front Cell Infect Microbiol.* 2012;2(May):1–12.
46. Hajdamowicz NH, Hull RC, Foster SJ, Condliffe AM. The Impact of Hypoxia on the Host-Pathogen Interaction between Neutrophils and *Staphylococcus aureus*. *Int J Mol Sci.* 2019 Nov 7;20(22):5561.
47. Yang D, Wijenayaka AR, Solomon LB, Pederson SM, Findlay DM, Kidd SP, et al. Novel Insights into *Staphylococcus aureus* Deep Bone Infections: the Involvement of Osteocytes. *MBio.* 2018 Apr 24 ;9(2):e00415-18.
48. Bosse MJ, Gruber HE, Ramp WK. Internalization of bacteria by osteoblasts in a patient with recurrent, long-term osteomyelitis: A case report. *J Bone Jt Surg - Ser A.* 2005;87(6):1343–7.
49. Ellington JK, Harris M, Hudson MC, Vishin S, Webb LX, Sherertz R. Intracellular *Staphylococcus aureus* and antibiotic resistance: Implications for treatment of staphylococcal osteomyelitis. *J Orthop Res.* 2006;24(1):87–93.
50. Josse J, Velard F, Gangloff SC. *Staphylococcus aureus* vs. Osteoblast: Relationship and Consequences in Osteomyelitis. *Front Cell Infect Microbiol.* 2015;5(November):1–17.
51. Ricciardi BF, Muthukrishnan G, Masters E, Ninomiya M, Lee CC, Schwarz EM. *Staphylococcus aureus* Evasion of Host Immunity in the Setting of Prosthetic Joint Infection: Biofilm and Beyond. *Curr Rev Musculoskelet Med.* 2018 Sep 9;11(3):389–400.
52. Thammavongsa V, Kim HK, Missiakas D, Schneewind O. Staphylococcal manipulation of host immune responses. *Nat Rev Microbiol.* 2015;13(9):529–43.
53. Zipfel PF, Skerka C. *Staphylococcus aureus*: The multi headed hydra resists and controls human complement response in multiple ways. *Int J Med Microbiol.* 2014 Mar 1;304(2):188–94.
54. Goodyear CS, Silverman GJ. Death by a B Cell Superantigen. *J Exp Med.* 2003 May 5;197(9):1125–39.

55. Thammavongsa V, Missiakas DM, Schneewind O. Staphylococcus aureus degrades neutrophil extracellular traps to promote immune cell death. *Science* (80-). 2013 Nov 15;342(6160):863–6.
56. Wagner C, Obst U, Hänsch GM. Implant-associated posttraumatic osteomyelitis: Collateral damage by local host defense? *Int J Artif Organs*. 2005;28(11):1172–80.
57. Zimmerli W, Sendi P. Orthopaedic biofilm infections. *APMIS*. 2017;125:353–64.
58. Klosterhalfen B, Peters KM, Tons C, Hauptmann S, Klein CL, Kirkpatrick CJ. Local and systemic inflammatory mediator release in patients with acute and chronic posttraumatic osteomyelitis. *J Trauma*. 1996;40(3):372–8.
59. Augustin G, Zigman T, Davila S, Udilljak T, Staroveski T, Brezak D, et al. Cortical bone drilling and thermal osteonecrosis. *Clin Biomech*. 2012;27(4):313–25.
60. Lüthje FL, Skovgaard K, Jensen HE, Kruse Jensen L. Pigs are useful for the molecular study of bone inflammation and regeneration in humans. *Lab Anim*. 2018 Dec 13;52(6):630–40.
61. Anderson JM, Rodriguez A, Chang DT. Foreign body reaction to biomaterials. *Semin Immunol*. 2008;20(2):86–100.
62. Jhunjhunwala S, Aresta-DaSilva S, Tang K, Alvarez D, Webber MJ, Tang BC, et al. Neutrophil responses to sterile implant materials. *PLoS One*. 2015;10(9):1–16.
63. Elek SD, Conen PE. The virulence of staphylococcus pyogenes for man. A Study of the Problems of Wound Infection. *Br J Exp Pathol*. 1957;38(6):573–86.
64. Cordero J, Munuera L, Folgueira MD. Influence of bacterial strains on bone infection. *J Orthop Res*. 1996 Jul;14(4):663–7.
65. Brodbeck WG, Shive MS, Colton E, Nakayama Y, Matsuda T, Anderson JM. Influence of biomaterial surface chemistry on the apoptosis of adherent cells. *J Biomed Mater Res*. 2001 Jun 15;55(4):661–8.
66. Mader JT, Brown GL, Guckian JC, Wells CH, Reinartz JA. A mechanism for the amelioration by hyperbaric oxygen of experimental staphylococcal osteomyelitis in rabbits. *J Infect Dis*. 1980;142(6):915–22.
67. Emslie KR, Fenner LM, Nade SML. Acute haematogenous osteomyelitis: II. The effect of a

- metaphyseal abscess on the surrounding blood supply. *J Pathol*. 1984 Feb 1;142(2):129–34.
68. Moser C, Pedersen HT, Lerche CJ, Kolpen M, Line L, Thomsen K, et al. Biofilms and host response – helpful or harmful. *Apmis*. 2017;125(4):320–38.
69. Lorenzo J. The Effects of Immune Cell Products (Cytokines and Hematopoietic Cell Growth Factors) on Bone Cells. In: Lorenzo J, Horowitz MC, Choi Y, Takayanagi H, Schett G, editors. *Osteoimmunology: Interactions of the Immune and Skeletal System*. 2nd ed. Academic Press; 2016. p. 143–67.
70. Baumann H, Gauldie J. The acute phase response. *Immunol Today*. 1994;15(2):74–80.
71. Prabhakara R, Harro JM, Leid JG, Harris M, Shirtliff ME. Murine immune response to a chronic *Staphylococcus aureus* biofilm infection. *Infect Immun*. 2011 Apr 1;79(4):1789–96.
72. Yoshii T, Magara S, Miyai D, Nishimura H, Kuroki E, Furudoi S, et al. Local levels of interleukin-1 β , -4, -6, and tumor necrosis factor α in an experimental model of murine osteomyelitis due to *Staphylococcus aureus*. *Cytokine*. 2002 Jul;19(2):59–65.
73. Rochford ETJ, Sabaté Brescó M, Zeiter S, Kluge K, Poulsson A, Ziegler M, et al. Monitoring immune responses in a mouse model of fracture fixation with and without *Staphylococcus aureus* osteomyelitis. *Bone*. 2016 Feb;83:82–92.
74. Azuma Y, Kaji K, Katogi R, Takeshita S, Kudo A. Tumor necrosis factor-alpha induces differentiation of and bone resorption by osteoclasts. *J Biol Chem*. 2000 Feb 18;275(7):4858–64.
75. Gilbert L, He X, Farmer P, Boden S, Kozlowski M, Rubin J, et al. Inhibition of Osteoblast Differentiation by Tumor Necrosis Factor- α . *Endocrinology*. 2000;141(11):3956–64.
76. Lacey DC, Simmons PJ, Graves SE, Hamilton JA. Proinflammatory cytokines inhibit osteogenic differentiation from stem cells: implications for bone repair during inflammation. *Osteoarthritis Cartilage*. 2009;17(6):735–42.
77. Ishimi Y, Miyaura C, Jin CH, Akatsu T, Abe E, Nakamura Y, et al. IL-6 is produced by osteoblasts and induces bone resorption. *J Immunol*. 1990 Nov 15;145(10):3297–303.
78. Heim CE, Vidlak D, Scherr TD, Hartman CW, Garvin KL, Kielian T. IL-12 promotes myeloid-derived suppressor cell recruitment and bacterial persistence during *Staphylococcus aureus* orthopedic

Accepted Article
implant infection. *J Immunol.* 2015 Apr 15;194(8):3861–72.

79. Bernthal NM, Pribaz JR, Stavrakis AI, Billi F, Cho JS, Ramos RI, et al. Protective role of IL-1 β against post-arthroplasty *Staphylococcus aureus* infection. *J Orthop Res.* 2011 Oct;29(10):1621–6.
80. Hofbauer LC, Lacey DL, Dunstan CR, Spelsberg TC, Riggs BL, Khosla S. Interleukin-1B and tumor necrosis factor- α , but not interleukin-6, stimulate osteoprotegerin ligand gene expression in human osteoblastic cells. *Bone.* 1999;25(3):255–9.
81. Jimi E, Nakamura I, Duong LT, Ikebe T, Takahashi N, Rodan GA, et al. Interleukin 1 Induces Multinucleation and Bone-Resorbing Activity of Osteoclasts in the Absence of Osteoblasts/Stromal Cells. *Exp Cell Res.* 1999 Feb 25;247(1):84–93.
82. Asensi V, Valle E, Meana A, Fierer J, Celada A, Alvarez V, et al. In vivo interleukin-6 protects neutrophils from apoptosis in osteomyelitis. *Infect Immun.* 2004 Jul 1;72(7):3823–8.
83. Balto K, Sasaki H, Stashenko P. Interleukin-6 deficiency increases inflammatory bone destruction. *Infect Immun.* 2001;69(2):744–50.
84. Duplomb L, Baud'huin M, Charrier C, Berreur M, Trichet V, Blanchard F, et al. Interleukin-6 Inhibits Receptor Activator of Nuclear Factor κ B Ligand-Induced Osteoclastogenesis by Diverting Cells into the Macrophage Lineage: Key Role of Serine 727 Phosphorylation of Signal Transducer and Activator of Transcription 3. *Endocrinology.* 2008 Jul 1;149(7):3688–97.
85. Baker PJ, Dixon M, Evans RT, Dufour L, Johnson E, Roopenian DC. CD4(+) T cells and the proinflammatory cytokines gamma interferon and interleukin-6 contribute to alveolar bone loss in mice. *Infect Immun.* 1999;67(6):2804–9.
86. Windolf CD, Meng W, Lögters TT, MacKenzie CR, Windolf J, Flohé S. Implant-associated localized osteitis in murine femur fracture by biofilm forming *Staphylococcus aureus*: a novel experimental model. *J Orthop Res.* 2013 Dec;31(12):2013–20.
87. Ning R, Zhang X, Guo X, Li Q. *Staphylococcus aureus* regulates secretion of interleukin-6 and monocyte chemoattractant protein-1 through activation of nuclear factor kappaB signaling pathway in human osteoblasts. *Brazilian J Infect Dis.* 2011 May;15(3):189–94.
88. Chamoun MN, Blumenthal A, Sullivan MJ, Schembri MA, Ulett GC. Bacterial pathogenesis and interleukin-17: interconnecting mechanisms of immune regulation, host genetics, and microbial

virulence that influence severity of infection. *Crit Rev Microbiol*. 2018;44(4):465–86.

89. Huang H, Kim HJ, Chang E-J, Lee ZH, Hwang SJ, Kim H-M, et al. IL-17 stimulates the proliferation and differentiation of human mesenchymal stem cells: implications for bone remodeling. *Cell Death Differ*. 2009;16:1332–43.
90. Ono T, Okamoto K, Nakashima T, Nitta T, Hori S, Iwakura Y, et al. IL-17-producing $\gamma\delta$ T cells enhance bone regeneration. *Nat Commun*. 2016;7:1–9.
91. Kotake S, Udagawa N, Takahashi N, Matsuzaki K, Itoh K, Ishiyama S, et al. IL-17 in synovial fluids from patients with rheumatoid arthritis is a potent stimulator of osteoclastogenesis. *J Clin Invest*. 1999;103(9):1345–52.
92. Dapunt U, Giese T, Maurer S, Stegmaier S, Prior B, Hansch GM, et al. Neutrophil-derived MRP-14 is up-regulated in infectious osteomyelitis and stimulates osteoclast generation. *J Leukoc Biol*. 2015;98(4):575–82.
93. Wright KM, Friedland JS. Differential regulation of chemokine secretion in tuberculous and staphylococcal osteomyelitis. *J Bone Miner Res*. 2002 Sep;17(9):1680–90.
94. Kopesky P, Tiedemann K, Alkekha D, Zechner C, Millard B, Schoeberl B, et al. Autocrine signaling is a key regulatory element during osteoclastogenesis. *Biol Open*. 2014 Aug 15;3(8):767–76.
95. Dawson H. A Comparative Assessment of the Pig, Mouse and Human Genomes. In: *The Minipig in Biomedical Research*. CRC Press; 2011. p. 323–42.
96. Riancho JA, Zarrabeitia MT, Gonzalez-Macias J. Interleukin-4 modulates osteoclast differentiation and inhibits the formation of resorption pits in mouse osteoclast cultures. *Biochem Biophys Res Commun*. 1993;196(2):678–85.
97. Pesanti EL, Lorenzo JA. Osteoclasts and effects of interleukin 4 in development of chronic osteomyelitis. *Clin Orthop Relat Res*. 1998;(355):290–9.
98. Owens JM, Gallagher AC, Chambers TJ. IL-10 modulates formation of osteoclasts in murine hemopoietic cultures. *J Immunol*. 1996;157(2):936–40.
99. Heim CE, Vidlak D, Kielian T. Interleukin-10 production by myeloid-derived suppressor cells contributes to bacterial persistence during *Staphylococcus aureus* orthopedic biofilm infection. *J*

Leukoc Biol. 2015 Dec 1;98(6):1003–13.

100. Prabhakara R, Harro JM, Leid JG, Keegan AD, Prior ML, Shirtliff ME. Suppression of the inflammatory immune response prevents the development of chronic biofilm infection due to methicillin-resistant *Staphylococcus aureus*. *Infect Immun*. 2011 Dec 1;79(12):5010–8.
101. Aerssens J, Boonen S, Lowet G, Dequeker J. Interspecies Differences in Bone Composition, Density, and Quality: Potential Implications for in Vivo Bone Research. *Endocrinology*. 1998;139(2):663–70.
102. Mestas J, Hughes CCW. Of mice and not men: differences between mouse and human immunology. *J Immunol*. 2004 Mar 1;172(5):2731–8.
103. Donnelly RP, Sheikh F, Dickensheets H, Savan R, Young HA, Walter MR. Interleukin-26: an IL-10-related cytokine produced by Th17 cells. *Cytokine Growth Factor Rev*. 2010 Oct;21(5):393–401.
104. Seok J, Warren HS, Cuenca AG, Mindrinos MN, Baker H V, Xu W, et al. Genomic responses in mouse models poorly mimic human inflammatory diseases. *Proc Natl Acad Sci U S A*. 2013 Feb 26;110(9):3507–12.
105. Jensen LK, Henriksen NL, Jensen HE. Guidelines for porcine models of human bacterial infections. *Lab Anim*. 2019;53(2):125–36.
106. Peric M, Dumic-Cule I, Grcevic D, Matijasic M, Verbanac D, Paul R, et al. The rational use of animal models in the evaluation of novel bone regenerative therapies. *Bone*. 2015;70:73–86.
107. Meurens F, Summerfield A, Nauwynck H, Saif L, Gerdts V. The pig: a model for human infectious diseases. *Trends Microbiol*. 2012;20(1):50–7.
108. Mair KH, Sedlak C, Käser T, Pasternak A, Levast B, Gerner W, et al. The porcine innate immune system: An update. *Dev Comp Immunol*. 2014;45(2):321–43.
109. Becerikli M, Jaurich H, Schira J, Schulte M, Döbele C, Wallner C, et al. Age-dependent alterations in osteoblast and osteoclast activity in human cancellous bone. *J Cell Mol Med*. 2017 Nov 1;21(11):2773–81.
110. Ventura MT, Casciaro M, Gangemi S, Buquicchio R. Immunosenescence in aging: between immune cells depletion and cytokines up-regulation. *Clin Mol Allergy*. 2017 Dec 14;15(21):1–8.
111. García-Alvarez F, Navarro-Zorraquino M, Castro A, Grasa JM, Pastor C, Monzón M, et al. Effect of

age on cytokine response in an experimental model of osteomyelitis. *Biogerontology*. 2009 Oct10(5):649–58.

112. Marriott I, Gray DL, Tranguich SL, Fowler VG, Stryjewski M, Scott Levin L, et al. Osteoblasts express the inflammatory cytokine interleukin-6 in a murine model of *Staphylococcus aureus* osteomyelitis and infected human bone tissue. *Am J Pathol* . American Society for Investigative Pathology; 2004;164(4):1399–406.
113. Marriott I, Gray DL, Rati DM, Fowler VG, Stryjewski ME, Levin LS, et al. Osteoblasts produce monocyte chemoattractant protein-1 in a murine model of *Staphylococcus aureus* osteomyelitis and infected human bone tissue. *Bone*. 2005;37(4):504–12.
114. Heim CE, Vidlak D, Scherr TD, Kozel JA, Holzapfel M, Muirhead DE, et al. Myeloid-derived suppressor cells contribute to *Staphylococcus aureus* orthopedic biofilm infection. *J Immunol*. American Association of Immunologists; 2014 ;192(8):3778–92.
115. García-Alvarez F, Monzón M, Grasa JM, Laclériga A, Amorena B, García-Alvarez I, et al. Interleukin-1, interleukin-6, and interleukin-10 responses after antibiotic treatment in experimental chronic *Staphylococcus aureus* osteomyelitis. *J Orthop Sci*. 2006;11(4):370–4.
116. Littlewood-Evans A, Hattenberger M, Luscher C, Pataki A, Zak O, O'Reilly T. Local expression of tumor necrosis factor alpha in an experimental model of acute osteomyelitis in rats. *Infect Immun*. 1997;65(8):3438–43.
117. Gaviria-Agudelo C, Carter K, Tareen N, Pascual V, Copley LA. Gene expression analysis of children with acute hematogenous osteomyelitis caused by methicillin-resistant *Staphylococcus aureus*: Correlation with clinical severity of illness. *PLoS One*. 2014;9(7):1–11.
118. Ferreira GF, Moraes C, Silveira AMS da, Correa-Oliveira R, Teixeira-Carvalho A, Martins-Filho OA, et al. Distinct cytokine profiles of circulating mononuclear cells stimulated with *Staphylococcus aureus* enterotoxin A in vitro during early and late episodes of chronic osteomyelitis. *Mem Inst Oswaldo Cruz*. Fundação Oswaldo Cruz; 2012 ;107(3):348–55.
119. Fullilove S, Jellis J, Hughes SPF, Remick DG, Friedland JS. Local and systemic concentrations of tumour necrosis factor- α , interleukin-6 and interleukin-8 in bacterial osteomyelitis. *Trans R Soc Trop Med Hyg*. 2000;94(2):221–4.

120. Wagner C, Kondella K, Bernschneider T, Heppert V, Wentzensen A, Hänsch GM. Post-Traumatic Osteomyelitis: Analysis of Inflammatory Cells Recruited into the Site of Infection. *Shock*. 2003;20(6):503–10.
121. Wagner C, Heck D, Lautenschläger K, Iking-Konert C, Heppert V, Wentzensen A, et al. T Lymphocytes in Implant-Associated Posttraumatic Osteomyelitis: Identification of Cytotoxic T Effector Cells at the Sites of Infection. *Shock*. 2006;25(3):241–6.
122. Dapunt U, Maurer S, Giese T, Gaida MM, Hänsch GM. The macrophage inflammatory proteins MIP1 (CCL3) and MIP2 (CXCL2) in implant-associated osteomyelitis: Linking inflammation to bone degradation. *Mediators Inflamm*. 2014;2014.
123. Jin T, Lu Y, He QX, Wang H, Li BF, Zhu LY, et al. The Role of MicroRNA, miR-24, and Its Target CHI3L1 in Osteomyelitis Caused by *Staphylococcus aureus*. *J Cell Biochem*. 2015;116(12):2804–13.
124. Evans CAW, Jellis J, Hughes SPF, Remick DG, Friedland JS. Tumor Necrosis Factor- α , Interleukin-6, and Interleukin-8 Secretion and the Acute-Phase Response in Patients with Bacterial and Tuberculous Osteomyelitis. *J Infect Dis*. 1998; 177(6):1582–7.
125. Bansal VP, Mittal PK, Ashokraj G. Humoral immune responses in osteomyelitis. *Int Orthop*. 1992;16(3).

Figure legends

Figure 1 Differentiation of mature osteoclast requires stimulation of RANK by RANKL. Osteoblasts produce both RANKL and the decoy receptor OPG, which inhibits the stimulation of RANK. Thus, osteoblasts have an important role in regulating the rate of osteoclastogenesis.

Figure 2 Based on in vivo and clinical studies the mentioned pro- and anti-inflammatory cytokines have been shown upregulated during bone infection. The effects of these cytokines on bone cells (osteoblasts and osteoclasts) are based on in vitro studies and arrows indicate a stimulatory effect while T-bars indicate inhibition.

Table 1 The cytokine response in animal models of *S. aureus* induced implant associated osteomyelitis or prosthetic joint infections

Model	<i>S. aureus</i> strain	Implant	Observation	Ref
BALB/c mice	UAMS-1	none	IL-6↑ in bone and the expression co-localize with osteoblasts.	(112)
BALB/c mice	UAMS-1	none	CCL2 (called MCP-1)↑ in bone and the expression co-localize with osteoblasts.	(113)
BALB/c mice	ATCC 29213	titanium locking plate	IL-6↑ in lavage fluid from soft tissue but IL-6→ in blood.	(86)
C57BL/6 and BALB/c mice	MRSA-M2	insect pin	Th2/Treg response protects against chronic <i>S. aureus</i> infections. IL-4↑ and IL-10↑ in infected BALB/c mice compared to C57BL/6 mice.	(100)
C57BL/6 mice	MRSA-M2, UAMS-1	insect pin	IL-2↑, IL-12 p70↑, TNF↑, IL-6↑ and IL-17↑ in bone.	(71)
C57bl/6 mice	JAR06.01.31	titanium locking plate	IL-17↑ in bone. TNF↑, IL-1β↑ and IL-4↓ in bone and soft tissue.	(73)
C57BL/6 mice	USA300 LAC:: <i>lux</i>	nickel-titanium wire	Myeloid-derived suppressor cells (MDSCs) suppress the immune response to <i>S. aureus</i> biofilm infections	(114)
C57BL/6 mice	USA300 LAC:: <i>lux</i>	nickel-titanium wire	IL-10 production by MDSCs contributes to the persistence of <i>S. aureus</i> biofilm infection.	(99)
C57BL/6NCr mice	USA300 LAC:: <i>lux</i>	nickel-titanium wire	IL-12p40↑, IL-1β↑, TNF↑, CSF3 (called G-CSF)↑, CXCL2↑ and CCL5↑ in tissue around knee joint. Increased infiltration of MDSCs and reduced infiltration of monocytes, macrophages and T-cells.	(78)
CD1 mice	ATCC 49230	silk suture	Osteoclasts do not absorb damaged bone in infected animals but are active at the margins of the infected site, resulting in sequestra formation.	(97)
ICR mice	E-31461	silk suture	IL-1β↑, IL-4↑, IL-6↑ and TNF↑ in bone	(72)
Wistar rats	9213	stainless steel needle	Antibiotic treatment affects the cytokine expression (IL-1α, IL-6 and IL-10).	(115)
Wistar rats	9213	stainless steel needle	IL-2↑ and IL-10↑ in blood of young rats (three months) and IL-6 ↑ in blood of old rats(22 months). IL-10↑ in tibia of young rats and IL-1β↑ in old rats.	(111)
Mandarin rats	1098	dental gypsum	TNF↑ in bone.	(116)

→: no significant change, ↑: increased level, ↓: decreased level

Table 2 The human immune response to osteomyelitis

Patient group	Control	Samples	Observation	Ref
Children with acute hematogenous OS caused by MRSA	healthy age- and gender-match children	blood	Genes related to neutrophil activity, coagulation and inflammation↑ while genes linked to activity of T-cells, NK-cells and B-cells↓.	(117)
acute or chronic OS	blood bank donors	blood	IL-6↑, IL-1α↑, TNF→ in serum.	(82)
chronic staphylococcal OS	healthy volunteers	blood	IL-6↓ in peripheral blood mononuclear cell supernatant and NO↓ in serum.	(118)
chronic OS	orthopedic patients with non-infectious or inflammatory disorders	blood and pus from infected bone	IL-6↑, CXCL8 (called IL-8)↑ and TNF↑ in pus compared to plasma, but no systemic differences to controls	(119)
patients with chronically infected mandibles	patients undergoing plastic surgery	bone	Human beta-defensin-1, -2 and 3↑ in bone.	(21)
acute and chronic posttraumatic OS	patients with fresh fractures without infection	blood and tissue from infected bone	CXCL8↑, IL-6↑, TNF↑, IL-1β↑ and LTB4↑ in acute OS samples compared to chronic bone samples. All measured inflammatory mediators↑ in the plasma of patients with acute OS compared to chronic and control group.	(58)
IAO	healthy donors (only blood)	lavage from infected implant and blood	Lavage contained 1-2 x 10 ⁷ leukocytes, predominantly (75-90 %) activated polymorphonuclear neutrophils	(120)
IAO	healthy donors (only blood)	lavage from infected implant and blood	CD8+ T cells ↑ in blood.	(121)
PJI	patients with aseptic loosening	blood and tissue samples from synovia, tibia, intramedullary space and unaffected muscle	S100A9 (called MRP-14)↑ and CXCL8↑ in synovial and tibial tissue. S100A9↑ in blood.	(92)
PJI	patients with aseptic loosening	soft tissue from osteolytic, adjacent and distal site, and a muscle sample.	CXCL8↑, IL-1β↑ and CXCL2↑ at the osteolytic site.	(31)
PJI	patients with aseptic loosening	Blood, tissue from site of osteolysis and unaffected muscle	CXCL2↑ and CCL3→ at osteolytic site.	(122)
PJI	aseptic loosening	bone	CSF3↑, IL-1β↑, IL-6↑, CXCL8↑ and CD40L↑ in bone.	(78)
PJI in acetabular bone	sample from uninfected iliac	bone	CCL5↑, CXCL9↑, CXCL10↑ and CXCL11↑ in bone and viable <i>S. aureus</i> are present inside osteocytes.	(47)

	wing of the same patients			
OS caused by <i>S. aureus</i>	Healthy volunteers	blood	miR-24↓ in blood.	(123)
bacterial OS	Healthy relatives or patients with non-inflammatory disorders	blood	IL-6→, TNF→, total leukocyte count↑, erythrocyte sedimentation rate↑, C-reactive protein↑ and plasma albumin↓ in blood.	(124)
bacterial OS	healthy age-match individuals	serum	IgA→, IgM→ and IgG→ in serum.	(125)
unspecified OS	healthy tissue from hip replacements	bone	Human β-defensin-2 ↑ in bone.	(22)
unspecified OS	healthy tissue from hip replacements	bone	Human β-defensin-3 → in bone.	(23)

→: no significant change, ↑: increased level, ↓: decreased level, IAO: implant associated osteomyelitis, MRSA: Methicillin resistant *Staphylococcus aureus*, OS: osteomyelitis, PJI: prosthetic joint infection.

apm_13027_f1.tif

apm_13027_f2.tif