Atmospheric Iodine (^{127}I and ^{129}I) Record in Spruce Tree Rings in the Northeast Qinghai-Tibet Plateau

Zhao, Xue; Hou, Xiaolin; Zhou, Weijian

Published in:
Environmental Science and Technology

Link to article, DOI:
[10.1021/acs.est.9b01160](https://doi.org/10.1021/acs.est.9b01160)

Publication date:
2019

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Zhao, X., Hou, X., & Zhou, W. (2019). Atmospheric Iodine (^{127}I and ^{129}I) Record in Spruce Tree Rings in the Northeast Qinghai-Tibet Plateau. *Environmental Science and Technology*, 53(15), 8706-8714.
<https://doi.org/10.1021/acs.est.9b01160>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Atmospheric iodine (^{127}I and ^{129}I) record in spruce tree rings in the northeast Qinghai-Tibet Plateau

Xue Zhao^{a,c,e}, *Xiaolin Hou*^{a,b,c,d*}, *Weijian Zhou*^{a,b,d}

a. State Key Laboratory of Loess and Quaternary Geology, Shaanxi Key Laboratory of Accelerator Mass Spectrometry Technology and Application, Xi'an AMS Center, Institute of Earth Environment, CAS, Xi'an 710061, China

b. CAS Center for Excellence in Quaternary Science and Global Change, Xi'an, 710061, China

c. Technical University of Denmark, Center for Nuclear Technologies, Risø Campus, Roskilde 4000, Denmark

d. Open Studio for Oceanic-Continental Climate and Environment Changes, Pilot National Laboratory for Marine Science and Technology (Qingdao), Qingdao 266100, China

e. University of Chinese Academy of Sciences, Beijing 100049, China

* Corresponding author: Technical university of Denmark, Center for Nuclear Technologies, Frederiksborgvej 399, Roskilde 4000, Denmark. Tel.: +45-21325129; fax: +45-46775347; E-mail address: houxl@ieecas.cn (X. Hou).

ABSTRACT: Atmospheric iodine isotopes have significant impacts on climate change and human health. However, the sources, transport pathway, and transfer process of atmospheric iodine are still not well understood. Tree rings of spruce collected from the east edge of Qinghai-Tibet Plateau were analyzed for iodine isotopes (^{127}I and ^{129}I). The results show that the levels and temporal variation of atmospheric iodine were well recorded in the spruce tree rings, and iodine concentrations in tree rings increased three times from 1960 to 2015, reflecting the increased releases of iodine to the atmosphere in the past decades due to the human activities. The anthropogenic ^{129}I in the tree rings presents the record of the human nuclear activities in the past 55 years. The sources and the transport pathways of radioactive substances could be extracted from the ^{129}I recorded in the tree rings in Qinghai-Tibet region. They are fallout of the global nuclear weapons tests in 1961-1962, releases of the Chinese atmospheric nuclear weapons tests in 1964-1980 transported through the tropospheric northwest wind, the releases of the Chernobyl accident dispersed through Westerlies and the continuous air releases before 1997 and the re-emission of marine discharges from the European nuclear fuel reprocessing plants transported through Westerlies.

Key words: Atmospheric iodine, Iodine-129, source term, transport pathway, Spruce, Tree rings

INTRODUCTION

Iodine is an essential trace element of mammals. Iodine deficiency can result in irreversibly neurological damage and developmental retardation in newborns. The major source of iodine in terrestrial environment is from atmospheric deposition.¹ The atmospheric iodine has a significant impact on climate through tropospheric ozone depletion and aerosol formation, and was

observed several times to be changed in recent decades.^{1,2,3} Although some investigations have been carried out by measuring iodine in soil, sediment and ice cores,^{3,4} the sources, the transport pathway and transfer process of atmospheric iodine in the ecosystem are still not well understood due to its multi-sources and the difficulties on discrimination of the different sources.

¹²⁹I, a long-lived radioisotope of iodine ($T_{1/2}=15.7$ Ma), has been released to the environment by human nuclear activities, including nuclear fuel reprocessing plants (NFRPs), nuclear weapons tests (NWTs) and nuclear accidents (NA). The anthropogenic ¹²⁹I (NFRPs: 7400 kg; NWTs: 150 kg; NAs: 7.2 kg) has highly overwhelmed the natural inventory in surface environment (250 kg, with a pre-nuclear ¹²⁹I/¹²⁷I atomic ratio of 1.5×10^{-12}).^{5,6} Due to unique source of anthropogenic ¹²⁹I, volatile property and biophilic characteristics of iodine, the anthropogenic ¹²⁹I has been widely applied as an environmental tracer for investigation of regional radioactive sources and the related atmospheric transport pathways by determination of ¹²⁹I preserved in time-serial samples (sediment, ice core and coral samples).⁶⁻¹² However, due to the insufficient resolutions of sediment samples caused by low deposition rates in many locations, the sampling difficulties and specified locations of ice core and coral samples, they are not sufficient for the investigation in large areas, especially in mid-low latitude terrestrial environment.

Tree ring is an excellent time-serial material to monitor past climate changes as well as anthropogenic activities due to its wide distribution, easy access, annual resolution, accurate chronology and environmental sensitivity,¹³ which might preserve the environmental radioactivity information both in pre-nuclear and nuclear age and in different locations. Dendrochemistry based on stable isotopic composition and elements concentrations in annual ring as retrospective proxy data has been successfully used to investigate the past environmental

change, such as past temperature and precipitation ($\delta^{18}\text{O}$, $\delta^{13}\text{C}$, etc.),¹³ and anthropogenic release of toxic elements to atmosphere (Pb, Cd, Hg, etc.).¹⁴ A few tentative works on regional radioactive deposition (including sources and transport) have also been reported by determination the concentration of radionuclides, such as ^{40}K , ^{137}Cs and ^{134}Cs .¹⁵ Two investigations on ^{129}I in tree ring samples collected near the West Valley NFRP in USA and the Kalsruhe NFRP in Germany were reported.^{16,17} The variation trend of ^{129}I levels in the ring-porous tree (elm, oak and locast), in which vessels are localized in the outmost growth ring, followed the regional deposition pattern of ^{129}I released from local NFRPs.¹⁶ While, the tree rings of semidiffuse to diffuse-porous wood (cherry and maple), where vessels are disseminated throughout the trunk, showed a less matched ^{129}I pattern to the local NFRP releases.¹⁶ This indicates a critical function of vessels in the iodine cross-ring translocation. However, high values of $^{129}\text{I}/^{127}\text{I}$ atomic ratio up to 3.6×10^{-9} were also measured in pre-nuclear rings from ring-porous tree (elm), indicating a cross-ring translocation of iodine occurred too.¹⁶ It is probably due to the negative hydrostatic pressure in vessels compared to that in adjacent parenchymal cells and tracheid cells.¹⁸ Therefore, coniferous tree might be more suitable for less cross-ring translocation of iodine due to no vessels in this species of tree.

The Qinghai-Tibet plateau (QTP) has an average altitude of 4000 m and cold ecosystem with widely distributed permafrost,¹⁹ it is therefore quite sensitive to environment changes. Meanwhile, this region is also important for the climate and ecosystem of a large area in the world because it acts as the largest fresh water reservoir in Asia including over $2 \times 10^{11} \text{ m}^3$ of glacier resource, $3 \times 10^{10} \text{ m}^3$ of annual runoff and 1800 lakes. In 1964 -1980, 22 atmospheric nuclear tests with the total yield of 22.72 Mt of TNT were conducted at Lop Nor NWTs site in Northwest China.²⁰ Due to the prevailing Westerlies at Lop Nor, the released ^{129}I from the

atmospheric nuclear weapons tests at this site might be transported to the QTP which is located in the 1100 km downwind area of Lop Nor.

This work aims to investigate the sources, transport pathway and transfer of atmospheric iodine in the QTP by determining ^{127}I and ^{129}I in tree rings of spruce. The feasibility using tree rings of spruce to record the temporal variation of iodine isotopes in the atmosphere will be investigated in order to obtain a historic record of ^{129}I and ^{127}I levels in the atmosphere; this is useful for reconstruction of the levels of radioactive fallout and providing fundamental database for regional environmental change research.

MATERIALS AND METHODS

Materials and dendrochronology. Two tree disks of Qinghai spruce PC1 and PC2 were collected from Qunjia (36° 16' 44.04" N, 101° 40' 5.76" E) and Maixiu National Park (35° 16' 26.46" N, 101° 55' 1.26" E) in Qinghai province, China, respectively. The sampling sites are located in the northeast edge of the QTP with average altitude over 3000 m and typical plateau continental climate. The Westerly is the dominant wind in this region.¹⁹

Qinghai spruce (*Picea crassifolia* Kom.), a coniferous species, is the dominant forest species in this area. Due to banned logging in the virgin forests in China in the past two decades, only the trees that naturally fell down can be sampled. In June 2017, one disk (PC1) was sampled from a spruce tree, which fell down in Dec 2015 in the foot of a slope of 10-20° with low canopy density. Another disk (PC2) was sampled from a spruce, which fell down in Jan 2017 in the middle of a slope of 20° with moderate canopy density. The sampling site of PC2 is located

km southeast of PC1. A disk of about 5 cm thickness was taken from the trunk of each tree at a height of 30 cm above the ground and transported to the laboratory for analysis.

The tree disks were naturally air dried in a cool and ventilated room in Xi'an, China. The surface of the wood disks was polished with sandy papers to easily identify the annual ring by different colors between early and late wood (Fig. 1). No visible multiple rings with abnormal thin width or missing rings with discontinuous boundary were observed in the transection of both disks. Therefore, they were dated by counting rings from outside according to date of fall. The age span is from 1961 to 2015 in PC1, and 1961 to 2016 in PC2 (Fig. 1). An uncertainty of ± 1 year for each tree disk was estimated due to the possibility of visual inspection error. The center part of the disk could not be used due to limited massive ring with super thin width. The clearest part with arrayed rings was selected to dissect the annual ring from the outmost layer (Fig. 1). Each dissected sample was dried at 60°C until constant weight, and then cut into small pieces.

Fig. 1 Wood disks of PC1 tree (up) and PC2 tree (bottom), showing the annual rings

Analytical Methods. Iodine in the dried tree ring samples was separated using an oxidative combustion method modified from Hou and Wang.²¹ 5 g of dried sample was combusted in a tube furnace (Radtec LTD, Southampton, UK) with a stepwise temperature increasing protocol as shown in Fig S1. Due to the ignition point of about 270 °C in the major part of this type of wood, a slow increase of temperature during 230-350 °C was used to avoid production of a large amount of combustion gases (CO₂, CO, NO_x) in a short time. A second carbonization was implemented by increasing the temperature from 350 to 550 °C in a ramp speed of 2.5 °C /min to combust the remained residues (including rosin with ignition point of 480-500 °C). Afterwards, the temperature was increased to 800 °C and remained for 3 hours to completely decompose the sample. Iodine released during combustion was collected in a bubbler with 30 ml trap solution of 0.4 mol/l NaOH-0.05 mol/l NaHSO₃. The detailed method was presented in the supporting information. 1.0 ml of iodine trap solution was diluted with deionized water for measurement of ¹²⁷I using ICP-MS (Agilent 8800 ICP-MS). Cs⁺ was used as internal standard, 0.15 mol/L NH₃·H₂O was used as rinse solution. The detection limit of this method for ¹²⁷I was estimated to be 0.012 ng/ml. Due to large dilution (more than 30 times) of iodine trap solution, the iodine concentrations in some samples from PC2 are close to the detection limit (0.006 µg/g), causing a large measurement uncertainty (> 50%). Therefore, data of these samples are not included for further discussion. After addition of stable iodine carrier (prepared from iodine crystal purchased from Woodward company with ¹²⁹I/¹²⁷I atomic ratio <2×10⁻¹⁴) into the remaining trap solution, iodine (iodide) was precipitated as AgI for determination of ¹²⁹I by a 3 MV Tandem AMS system (HVEE) at Xi'an AMS Center, China. The details of AMS system and measurement method for ¹²⁹I have also been reported elsewhere,²² and briefly described in Supporting Information. ¹²⁹I/¹²⁷I ratios in the procedure blanks prepared using the same procedure as the samples were

measured to be below 2×10^{-13} , which are at least one order of magnitudes lower than that of samples.

RESULTS AND DISCUSSION

¹²⁷I and ¹²⁹I level in the tree rings. The measured ¹²⁷I concentrations are 0.01-0.10 µg/g with the most data fall into the range of 0.02-0.05 µg/g in tree rings of PC1, and 0.008-0.025 µg/g with the most fall into the range of 0.01-0.02 µg/g in tree rings of PC2 (Fig. 2 a). The higher iodine concentration from PC1 might be attributed to its sampling location, at forest edge with sufficient water, sunlight and fresh air, confirmed by its larger ring width than that of PC2. The iodine concentrations in the two spruce trees are much lower than the reported levels in deciduous trees, such as 0.1-0.6 µg/g in Black locust collected from Karlsruhe, Germany in 1954-1983, 0.7-1.2 µg/g in Elm from New York State, USA in 1950-1986 and 0.04-0.09 µg/g in Red Oak from New York State, USA in 1961-1987 (Fig. 2 a).^{16,17} This probably caused by low uptake capacity of iodine from environment in this region with low temperature and associated low growth rate.

Significant variations of ¹²⁹I concentrations were observed in two tree rings, i.e. $(0.7-3.5) \times 10^6$ atoms/g in PC1 and $(0.5-3) \times 10^6$ atoms/g in PC2 (Fig. S3, Table S1). The over fivefold difference in 55-56 years might be attributed to the varied sources and concentrations of ¹²⁹I in the environment. In addition, the variation might also result from the varied accumulation/uptake rate of iodine into tree rings in different periods. To overcome this problem, the ¹²⁹I/¹²⁷I atomic ratios were applied to represent the level of ¹²⁹I in the tree rings.

Fig. 2 Comparison of measured ^{127}I concentrations (a) and $^{129}\text{I}/^{127}\text{I}$ ratios (b) in the tree rings in this work with the reported values in other locations^{16,17, 23, 24}

The $^{129}\text{I}/^{127}\text{I}$ atomic ratios of $(6-15)\times 10^{-9}$ were obtained in the tree ring of PC1 and $(4-20)\times 10^{-9}$ in PC2 (Fig. 2 b), which are more than two orders of magnitude higher than the pre-nuclear level of $^{129}\text{I}/^{127}\text{I}$ ratios (1.5×10^{-12} for marine system and 2.0×10^{-11} in terrestrial environment),^{5,25} indicating anthropogenic sources of ^{129}I . However, this level is at least two orders of magnitude lower than the values reported outside the nuclear fuel reprocessing sites in Germany (10^{-7} - 10^{-5}) and USA (10^{-7} - 10^{-6}), but similar to those observed in the areas without direct influenced by nuclear activities (10^{-9} - 10^{-8}) (Fig. 2 b).^{16,17} The average ratio of $^{129}\text{I}/^{127}\text{I}$ in the two rings in 2009-2015 is around 6×10^{-9} , in the same magnitude with the values measured in herbs (4.5×10^{-9}) collected in 2015²³ and surface soils (3×10^{-9}) collected in 2013²⁴ in north China, representing the level of ^{129}I in this region.

Iodine uptake and integration in the tree rings from atmosphere. The measured $^{129}\text{I}/^{127}\text{I}$ ratios (average of 6.6×10^{-9}) during 2009-2015 in the rings of PC1 tree are more than two times higher than the reported values in surface soil (average: 3×10^{-9} , collected in 2009-2013) and river water (1.1×10^{-9} , collected in 2016-2017) in nearby locations (Fig. 2).^{24,26} During 1963-1977, the ratios (average of 9.5×10^{-9}) in the rings of both trees are similar to the reported $^{129}\text{I}/^{127}\text{I}$ ratios (average of 9×10^{-9}) in the atmospheric fallout in Tokyo (35 °N) before the operating of Tokai NFRPs (Fig. 4a).²⁷ No ^{129}I data in atmospheric sample from the study region was reported. Since the ^{129}I level in atmosphere in the study region (36 °N) should be similar with that in the same latitude region as Tokyo, the result might indicate that iodine uptake from atmosphere is the dominant way entering into the tree rings compared to that from soil system.

The Qinghai spruce uptakes inorganic elements and water from soil through root, and translocates it to other parts through the pits on the tracheids cells wall relying on transpiration stream (Fig. S2). This slow process could induce a time lag of months to years.²⁸ Meanwhile, spruce can also uptake the nutrient from atmosphere through foliar absorption, and phloem transport, then to inner xylem through radially aligned ray parenchyma cells (the only radial transport tissue of tree trunk), which is sustainable by photosynthesis energy without much time lag.²⁹

Iodine in soil mainly originates from the atmospheric deposition, decomposition of dead plants and animals (absorbed iodine from atmosphere or soil), and weathering of parent material of soil.³⁰ The anthropogenic ^{129}I presents mainly in the upper 50 cm of soil profiles in the background region (far away from any nuclear facilities),³⁰ and the natural $^{129}\text{I}/^{127}\text{I}$ ratios in the deep soil ($<10^{-11}$) are 2-3 orders of magnitude lower than the present environmental level (e.g. atmosphere, surface soil, etc.).²⁵ Considering the deep root of spruce ($>>50$ cm), lower $^{129}\text{I}/^{127}\text{I}$

ratios observed in the surface soil in this region compared to that in the tree ring, as well as the similar $^{129}\text{I}/^{127}\text{I}$ ratios of the tree rings with that in atmospheric fallout in Japan, iodine in the investigated tree rings should be dominantly taken from atmosphere. The similar pattern of higher $^{129}\text{I}/^{127}\text{I}$ ratios in precipitation (3.8×10^{-9}) than that in surface soil (1.9×10^{-9}) was also reported in Xi'an.³¹ Due to the non-essential element for trees, iodine was also demonstrated to be hardly translocated to the aboveground compartments from the tree root system.²⁹

In atmosphere, iodine presents mainly in gaseous form and partly associated with aerosol.^{32,33} Gaseous iodine can be trapped directly through stomata in the tree leaves. Aerosol deposited on tree leaves might be adhered on the surface through wax on the spruce needles, and the associated iodine can be leached out by water vapor. Then soluble iodine can be taken up to the tree leaves through hydrophilic holes on leaf's surface and/or active absorption by plasmodesmata.³⁴ The absorbed iodine is translocated through phloem, ray parenchyma cells and then fixed in xylem. The Fukushima derived ^{129}I was observed to be well recorded in *Japanese cedar* (coniferous species) through atmosphere absorption.³⁵ Much higher ^{129}I concentration was measured in litter than in surrounding soil collected in nearby area of a NFRP, implying a more direct uptake way of iodine in tree/vegetation from atmosphere by foliage.¹⁷

The translocation pathway of the atmospheric uptake iodine into the xylem relies on the only radial tissue (ray parenchyma cells) and the pits on longitudinal tracheid cell walls (Fig. S2). The ray parenchyma cells, is alive only in sapwood (usually the out most 10-12 rings) but dead in heartwood.³⁶ A considerable amount of radial moving of water solution in alive ray parenchyma cells was detected based on fluorescein dyes.¹⁸ However the migration depth was limited only in the outmost ring of mature xylem from the anatomical observations by scanning electron microscope.¹⁸ A similar low crossing rings movement of water soluble fluorine (similar

chemistry with iodine) was also reported in Norway spruce.³⁷ The pits in longitudinal tracheid cells are usually less developed, both the size and quantity, from late wood than that from early wood, which will primarily limit iodine moving across one single growth ring. The water transport by tracheid cells was only observed in early wood, but not in late wood based on single growing spruce ring.³⁷ Combined with the fluctuated variation of the observed $^{129}\text{I}/^{127}\text{I}$ ratios over rings in both trees (Figure. 3a and b) instead of monotonous gradient distributions, it indicates that the crossing rings translocation of iodine hardly happens in the selected Qinghai spruce.

Fig. 3 Temporal variation of the measured $^{129}\text{I}/^{127}\text{I}$ ratios in the tree rings from two sites in the QTP (a), in comparison with the reported variation of $^{129}\text{I}/^{127}\text{I}$ in atmospheric fallout from Tokyo (b)²⁷, yield of fission NWTs (c), yield of thermonuclear weapons tests at Lop Nor

(d)²⁰, air releases of ¹²⁹I from major NFRPs (e) and marine discharges of ¹²⁹I from NFRPs (f).³³

Historic records of atmospheric iodine level. ¹²⁷I concentrations in the tree rings of PC1 (Fig. 4a) were 0.010-0.033 µg/g with an average of 0.020 µg/g in 1961-1983, and increase to 0.023-0.051 µg/g with an average of 0.039 µg/g in 1984-2001, then to 0.049-0.100 with an average of 0.069 µg/g in 2001-2016. A threefold increase was observed, but no such variation in PC2 (Fig. 4a). This increased iodine concentrations might reflect the increased iodine concentrations in atmosphere based on the inferred atmospheric absorption pathway. A similar three folds elevation of atmospheric iodine has been observed in Alpine ice core,⁴ and Greenland ice core³ from 1950-2000s. Significant correlation of ¹²⁷I concentrations in the tree rings with that in Greenland ice core³ (directly deposition from atmosphere) (R=0.77, P<0.01) are observed (Fig. 4b). However, no such kind of correlation was found in PC2. The lack of partial data in PC2 or/and the sampling site located in the middle of forest with higher canopy density than that in PC1 and lower environmental sensitivity might attribute to this phenomenon.³⁸ Although the uptake capacity of iodine through the foliar absorbing way can be affected by water condition, sunlight, temperature, etc., a less variation of temperature (<1°C of the annual average temperature), fluctuating variation of annual precipitation (from the nearest Xining weather station), and almost steady sunlight condition were reported in the past 50-60 years in the studied time scale,³⁹ excluded effect of these parameters on the uptake of iodine into the tree region in this location.

Fig. 4 Variation of iodine (^{127}I) concentration in PC1 and PC2 tree (a), in comparison with the temporal variation of iodine concentrations in ice core in Greenland³

Iodine in atmosphere might originate from oceanic emissions of gaseous iodine (through biological, chemical and photochemical process), terrestrial emissions of gaseous iodine from salt marches, rice fields, peatland and forest system, and combustion of fossil fuel and biomass, of which the oceanic emission dominates.^{1,3,4} The increased atmospheric iodine level might be attributed to the increased releases from these processes. The oceanic emission of molecular iodine (I_2) was observed to increase with the tropospheric ozone (O_3) content,¹ which increased by a factor of two from the 1950s to early 1990s due to the elevated global combustion of fossil fuel.^{40,41} The oceanic emission iodine was reported to be enhanced by the sub-ice phytoplankton production that associated with the recent thinning of sea ice.³ Meanwhile, the increased

consumption of fossil fuels (from 1000 TWh/y in 1980s to 3200 TWh/y in 2000s) in the past decades could also directly contribute to the increased iodine concentration in the atmosphere, which was estimated to be 5000 tons/y in 2005.^{42, 43}

Records of human nuclear activities in the tree rings. The distribution of $^{129}\text{I}/^{127}\text{I}$ atomic ratios (Fig. 3a and b) in the two tree rings show similar variation trend, e.g. the $^{129}\text{I}/^{127}\text{I}$ peaks in 1962, 1964-1968, 1976, 1996 and 2003 occurred in both tree rings, as well as the peaks in 1972/1973, (1986-1988)/(1986-1987), 1992/1991, 1998 and 2000/1999 occurred in PC1/PC2 respectively. The one-year discrepancy between the records in two tree rings could be caused by the uncertainty of dating (± 1) from counting rings. In addition to that, due to the lack of data in 1980 in PC2 rings compared to an obvious peak in 1980 from PC1 rings, and a separate high values period of 2006-2008 in PC1 rings, might be attributed to the different sampling locations and/or uncertainty of dating results (± 1 year) from counting rings.

The similar peaks in the two tree rings should be attributed to the peak values of ^{129}I occurred in atmosphere during these years, resulting from the anthropogenic releases. Fig. 3 shows the variation of $^{129}\text{I}/^{127}\text{I}$ ratios in two tree rings in comparison with that in atmospheric fallout records in Tokyo (Fig. 3a and b), the time distribution of explosion yields of atmospheric NWTs (Fig. 3c and d), and the annual discharges (air and marine) of ^{129}I from the three European NFRPs which accounts for the most of ^{129}I inventory in current environment (Fig. 3e and f).

Atmospheric nuclear weapons tests. Several peaks of $^{129}\text{I}/^{127}\text{I}$ atomic ratios in 1962, 1964-1968, 1972, 1976 and 1980 were observed in the tree rings of PC1, and in 1961-1962, 1964-1968 and 1976 in the rings of PC2. The atmospheric nuclear weapon tests are the only source of ^{129}I during this period since the release amount from NFRPs were negligible.²⁰

The twofold increased $^{129}\text{I}/^{127}\text{I}$ ratios (8.8×10^{-9}) in 1962 compared to the value of $(3-4) \times 10^{-9}$ in 1961 and 1963 was observed in tree rings of PC1, and 9×10^9 in 1961-1962 compared to 5×10^9 in 1963 in PC2 rings, corresponding with the date of the largest numbers of atmospheric NWTs in 1961-1962. The observed value in the tree rings of 1961-1962 (average: 9×10^9) agrees with the value recorded in atmospheric fallout samples in 1963 from Tokyo (15.5×10^9).²⁷ This signal was widely detected by peaks of $^{129}\text{I}/^{127}\text{I}$ ratio, ^{137}Cs and $^{239,240}\text{Pu}$ activities in north China.^{7,44} The 1-2 years lag among these records could be attributed to the hysteresis effect of samples or/and the dating uncertainty (± 1 year).

A broad peak of $^{129}\text{I}/^{127}\text{I}$ ratios corresponding to 1964-1968 was observed in both tree rings, which matches well with several high fission yield tests at Lop Nor (Fig. 3c), such as the first one in 1964 (20 kt) and the second one in 1965 (40 kt). The highest value of 12×10^9 in 1966 in PC1 rings agrees well with the boosted fission tests carried in 1966 with fission yield of 250 kt of TNT, and the similar high $^{129}\text{I}/^{127}\text{I}$ ratio (12×10^9) was also recorded in atmospheric fallout samples in 1966 collected from Tokyo.²⁷ The constant high ^{129}I level in both trees in 1967-1968, as well as the high value in the atmospheric fallout in Tokyo in 1968, should be attributed to a test of fission weapon in 1967 at Lop Nor and the first two Chinese thermonuclear weapons tests in 1967 and 1968, where a detonator with high ^{235}U fuel load might be applied, causing releases of fission products including ^{129}I .

The radioactive substances from atmospheric NWTs were injected into different height of atmosphere depending on the total explosion yield, the height of burst, and the meteorological condition after the detonation.²⁰ The radioactive substances from fission weapons with relative lower yields and some part from thermonuclear weapons were mainly injected into troposphere to disperse regionally.²⁰ The sampling sites in northeast QTP are located about 1100 km

southeast of Lop Nor. The dominant Westerlies in the troposphere in the Lop Nor region can transport these radioactive substances (including ^{129}I) to the sampling sites in several days and produce $^{129}\text{I}/^{127}\text{I}$ ratio peaks in the tree rings. The simulated forward trajectories of air masses by the HYSPLIT model (Fig. S4) confirmed that the air masses in the troposphere at the Lop Nor after the tests on 16th Oct. 1964, 14th May 1965, 28th Dec. 1966, 17th Jul. 1967, and 28th Dec. 1968 reached to the sampling region in a few days.⁴⁵

One $^{129}\text{I}/^{127}\text{I}$ ratio peak, observed in 1972 in PC1 rings, is in accord with the high fission yield of 170 kt of TNT from a fizzled thermonuclear test conducted on 18th Mar. 1972 at Lop Nor. No significant peak of $^{129}\text{I}/^{127}\text{I}$ ratio was observed in 1972 but a high value in 1973 (8.5×10^{-9}) in PC2, which could be due to the ± 1 year's uncertainty of dating. A similar peak was also measured in atmospheric fallout samples in 1972 collected from Tokyo (11.5×10^{-9}) (Fig. 3a).²⁷ The highest value of $^{129}\text{I}/^{127}\text{I}$ ratios of 20×10^{-9} was observed in 1976 in both tree rings that matches very well with the date of the fizzled thermonuclear test with a high fission yield of 200 kt of TNT on 26th Sep. 1976 at Lop Nor, as well as a similar peak in atmospheric fallout in Tokyo.²⁷ A peak value of $^{129}\text{I}/^{127}\text{I}$ ratios (15.3×10^{-9}) was observed in 1980 in PC1 rings in consistent with the time of the last atmospheric test at Lop Nor. The forward trajectory analysis of air mass also confirmed the stratospheric air masses at the Lop Nor after the tests on 18th Mar. 1972, 26th Sep. 1976, and 16th Oct. 1980 arrived in the sampling region in a few days (Fig. S4)

No significantly high ^{129}I signals were observed after 5 high yield thermonuclear tests conducted on 29th Sep. 1969 (3 Mt), 14th Oct. 1970 (3.4 Mt), 27th Jun. 1973 (3 Mt), 17th Jun. 1974 (1 Mt) and 17th Nov. 1976 (4 Mt) (Fig. 3d), reflecting less ^{129}I release due to the less fission fuel loaded as detonator.

Releases from nuclear accidents. A peak of $^{129}\text{I}/^{127}\text{I}$ ratios covers 3 years in 1986-1988 ($(14.6-18.4) \times 10^9$) in PC1, and 2 years in 1986-1987 ($(12.7-13.8) \times 10^9$) in PC2 were observed. Since atmospheric NWTs have already been ceased from 1980, and the almost steady discharges of ^{129}I from the main NFRPs (air: 13-14 kg/y; marine: 70 kg/y), this might be attributed to the Chernobyl accident on 26th April 1986 with 1.3-6 kg ^{129}I released to the environment.³³ The Chernobyl accident derived radioactive fallout had dispersed to north China through the Westerlies, and the level decreased from northwest towards east and south.⁴⁶ The total β activity in atmospheric fallout collected in Urumchi (1400 km in the northwest of sampling sites, 43° N), and the short lived radioactive ^{131}I activity in milk was observed to elevate over 50 times compared to the background value.^{46,47} A 15-times elevation of the total β activity was also measured in atmospheric fallout in Xi'an (600 km in the east of sampling sites, 33° N).⁴⁷ Although no direct monitor data of radioactivity after the Chernobyl accident is available in the sampling site, it should receive the Chernobyl derived ^{129}I because the sampling site is located between Xi'an and Urumchi. The Chernobyl derived $^{129}\text{I}/^{127}\text{I}$ peak was even observed in sediment samples from Jiaozhou Bay in the similar latitude but 1600 km east to the study area.⁷ A broad ^{129}I peaks in both tree ring profiles (PC1: 3 years; PC2: 2 years) might be contributed to the re-emission of the deposited ^{129}I in the high contaminated soil to atmosphere in the following 2-3 years.

No significant increased $^{129}\text{I}/^{127}\text{I}$ value was observed in 2011 (7.1×10^9) compared to 6.9×10^9 in 2012 and 5.2×10^9 in 2010 in PC1 rings, indicting an insignificant contribution from the Fukushima accident with about 1.2 kg ^{129}I releases (including 0.35 kg direct marine discharge).⁷ The atmospheric releases from Fukushima accident mainly dispersed eastwards and deposited in the North Pacific Ocean, and the marine discharges including ^{129}I were mainly transported

eastwards by North Pacific Current.⁴⁸ Although some radionuclides (mainly ^{131}I , ^{137}Cs and ^{134}Cs) in aerosol samples collected in North China after Fukushima were measurable, but the level is much lower compared to those in North America and Europe, and at least two orders of magnitudes lower than that from Chernobyl.^{49,50,51} Therefore, the contribution to the study area is negligible.

Releases from nuclear fuel reprocessing plants. A few peaks of $^{129}\text{I}/^{127}\text{I}$ ratios were observed since 1991 in the rings of both trees, i.e. 1992 (PC1)/1991 (PC2), 1996 (PC1 and PC2), 1998 and 2000 (PC1)/1999 (PC2) and 2003 (PC1 and PC2) (Fig. 3a and b). Since the atmospheric NWTs ceased in 1980, no reported nearby nuclear facilities and nuclear accidents can release ^{129}I to the sampling region. The most possible source of ^{129}I during this period should be the releases from NFRPs, which dominate the ^{129}I inventory in the present environment.³³ Although the reprocessing plants have been in operation in USA, Russia, Japan, China, Germany, France and UK, the European reprocessing plants in Marcoule (1959-1997), La Hague (1966-), and Sellafield (1951-) are the most important ones due to their high capacities and large scales in reprocessing operation. The contribution from two Chinese reprocessing plants in Guangyuan (1975-1991) and Jiuquan (1968-1984) was insignificant due to the small capacity and ^{129}I release compared to the European ones.²⁰ The investigation of ^{129}I distribution in surface soil from a large area of China did not show any unexpected increase level, except the local region of these two NFRPs (<100 km).⁵² The contribution from Tokai NFRPs (1977-2005) in Japan should be negligible because of the small volume of ^{129}I release (1 kg), the far distance, and non-major direction of wind.²⁰

The total atmospheric release of ^{129}I from the three European NFRPs gradually increased to 10 kg/y in 1976 from lower than 5 kg /y in late 1960s, and then kept at a relative constant high

level of about 15 kg /y until 1996. Afterwards, it gradually decreased to a quite low value of 1.4 kg/y in 2007 due to the shutdown of NFRP at Marcoule in 1997 and the updated system in other reprocessing plants for capture of gaseous radioiodine (Fig. 3e).³³ The high value in 1980 (18 kg) compared to that in 1979 (13 kg) might add a small background value on the peak produced by the last Chinese NWT. The marine discharges, mainly from the two European reprocessing plants at La Hague (France) and Sellafield (UK), gradually increased to 65 kg/y in 1980s from less than 40 kg/y before that, and significantly increased after 1990 and keep at a high value of 250 kg/y (Fig. 3f).

The ¹²⁹I peak observed in both tree rings in 1991/1992 matches well with the first increased total marine discharges (to 100 kg/y in 1991 from 70 kg/y in 1990), when air discharges of 13 kg were steady. The peak observed in 1996 matches with the most drastically elevated marine discharges up to 320 kg/y from below 200 kg/y in 1995, under the almost steady air discharges of (9-12) kg/y. The peaks observed in 1998-2000 in the two tree rings agree with the first peak of marine discharges of 350 kg/y in 1998-1999, while the air discharges continuously reduced by half (1997: 9 kg; 1999: 5 kg). The peak in 2003 matched well with the high value of 360 kg/y of marine discharges in 2002 compared to the value of 280 kg/y in 2001 and 2003, under the almost steady air discharges of 4 kg/y (Fig. 3e and f).

Some exceptional events were not recorded in tree rings, such as the elevation of total air discharges from 16 kg/y in 1983 to 19 kg/y in 1984, and the three-time elevation of total marine discharges from 100 kg in 1991/1992 to 325 kg in 1996. The small increment could be responsible for the former one. The decreased air discharges from Marcoule NFRPs from 8 kg/y in 1991/1992 to 2 kg/y in 1996 could explain the later one due to the closer latitude of 43 °N compared to that of La Hague (49° N) and Sellafield (54° N).

The marine discharges of ^{129}I can be re-emitted to the atmosphere in gaseous form confirmed by the similar high $^{129}\text{I}/^{127}\text{I}$ ratios observed in aerosols and rainwater samples collected in the North Europe (Denmark, Germany and Sweden) and the contaminated seawater samples ($^{129}\text{I}/^{127}\text{I} = 10^{-8}\text{-}10^{-6}$).^{32,53} The re-emitted ^{129}I can enter into lower troposphere, and to higher altitude by intensive vertical motions, then further transport eastwards through the Westerlies (around 1500 m-12000 m, mainly in 35° N - 65°N) to East Asia. The European NFRPs derived ^{129}I signal has been observed in sediment,^{6,7} precipitation,⁵⁴ and ground water²⁶ in Westerly area of north China (> 33° N). The ^{129}I was finally taken up by tree leaves and recorded as $^{129}\text{I}/^{127}\text{I}$ ratio peaks in both tree rings within 1 year in consideration of the long-distance transport.

Above all, the sources, transport pathway and transfer process of atmospheric iodine in the northeast part of the QTP were well traced. The terrestrial iodine from northwest upwind area (traced by ^{129}I released from NWTs at Lop Nor and from Chernobyl Accident), and marine iodine from the mid-high latitude of Atlantic Sea (traced by ^{129}I released from NFRPs at Sellafield, La Hague and Marcoule) can be transported to the studied area through the Westerlies, which contribute significantly for the iodine in atmosphere and then ecosystem.

SUPPORTING INFORMATION

The detailed chemical separation procedure, preparation and measurement of ^{127}I and ^{129}I , the tissues, the microscopic structures of selected wood species are presented; The analytical results of ^{129}I and ^{127}I concentration in two tree rings (Table S1), protocol of combustion for iodine separation from wood (Fig. S1), diagram of tree rings and microstructure of nonporous and porous tree (Fig. S2), temporal variations of ^{129}I concentrations in PC1 and PC2 (Fig. S3), and simulated forward trajectories of air mass after the major Chinese atmospheric nuclear weapons tests (Fig. S4) are included.

423

424 ACKNOWLEDGEMENTS

425 This work was supported by the Ministry of Science and Technology of China (No.
426 2015FY110800), the National Natural Science Foundation of China (No. 11875261, 91643206,
427 11605206, 41603125), Chinese Academy of Sciences (132B61KYSB20180003), National
428 Research Program for Key Issues in Air Pollution Control (DQGG0105-02). AMS measurement
429 of ^{129}I was performed by Dr. Qi Liu in the Xi'an AMS Center. X. Zhao thanks Xuhong Jiang,
430 Xiaohu Xiong, Jing Li and Miao Fang from Institute of Earth Environment for their helps in the
431 sampling and sample preparation and Dr. Jixin Qiao from the Technical University of Denmark
432 for her constructive suggestions in the manuscript preparation. The authors gratefully
433 acknowledge the NOAA Air Resources Laboratory (ARL) for the provision of the HYSPLIT
434 transport and dispersion model (<http://www.ready.noaa.gov>) used in this publication.

435

436 References

- 437 (1) Carpenter, L. C.; MacDonald, S. M.; Shaw, M. D.; Kumar, R.; Saunders, R. W.; Parthipan, R.;
438 Wilson, J.; Plane, L. M. Atmospheric iodine levels influenced by sea surface emissions of inorganic
439 iodine. *Nature Geoscience*. **2013**, 6(2), 108-111; DOI 10.1038/ngeo1687.
- 440 (2) Gantt, B.; Sarwar, G.; Xing, J.; Simon, H.; Schwede, D.; Hutzell, W. T.; Mathur, R.; Saiz-Lopez, A.
441 The impact of iodide-mediated ozone deposition and halogen chemistry on surface ozone
442 concentrations across the continental United States. *Environ. Sci. Technol.* **2017**, 51(3), 1458-1466;
443 DOI 10.1021/acs.est.6b03556.
- 444 (3) Cuevas, C.; Maffezzoli, N.; Corella, J.; Spolaor, A.; Vallelonga, P.; Kjer, H. A.; Simonsen, M.;
445 Winstrup, M.; Vinther, B.; Horvat, C.; Fernandez, R. P.; Kinnison, D.; Lamarque, J.; Barbante, C.;

- Saiz-Lopez, A. Rapid increase in atmospheric iodine levels in the North Atlantic since the mid-20th century. *Nature Communications*. **2018**, 9(1), 1452-1458; DOI 10.1038/s41467-018-03756-1.
- (4) Legrand, M.; McConnell, J. R.; Preunkert, S.; Arienzo, M.; Chellman, N.; Gleason, K.; Sherwen, T.; Evans, M.; Carpenter, L. J. Alpine ice evidence of a three-fold increase in atmospheric iodine deposition since 1950 in Europe due to increasing oceanic emissions. *Proceedings of the National Academy of Sciences*. **2018**, 115(48), 12136-12141; DOI 10.1073/pnas.1809867115.
- (5) Moran, J. E.; Fehn, U.; Teng, R. T. D. Variations in $^{129}\text{I}/^{127}\text{I}$ ratios in recent marine sediments: evidence for a fossil organic component. *Chemical Geology*. **1998**, 152(1-2), 193-203; DOI 10.1016/S0009-2541(98)00106-5.
- (6) Zhao, X.; Hou, X.; Du, J.; Fan, Y. Anthropogenic ^{129}I in the sediment cores in the East China Sea: Sources and transport pathways. *Environmental Pollution*. **2019**, 245, 443-452; DOI 10.1016/j.envpol.2018.11.018.
- (7) Fan, Y.; Hou, X.; Zhou, W.; Liu, G. ^{129}I record of nuclear activities in marine sediment core from Jiaozhou Bay in China. *Journal of Environmental Radioactivity*. **2016**, 154, 15-24; DOI 10.1016/j.jenvrad.2016.01.008.
- (8) Bautista, Vii A. T.; Miyake, Y.; Matsuzaki, H.; Lizuka, Y.; Horiuchi, K. High-resolution ^{129}I bomb peak profile in an ice core from SE-Dome site, Greenland. *Journal of Environmental Radioactivity*. **2018**, 184-185, 14-21; DOI 10.1016/j.jenvrad.2017.12.015.
- (9) Biddulph, D. L.; Beck, J. W.; Burr, G. S.; Donahue, D. J. Two 60-year records of ^{129}I from coral skeletons in the South Pacific Ocean. *Radioactivity in the Environment*. **2006**, 8, 592-598; DOI 10.1016/S1569-4860(05)08047-2.
- (10) Bautista, Vii A. T.; Matsuzaki, H.; Siringan, F. P. Historical record of nuclear activities from ^{129}I in corals from the northern hemisphere (Philippines). *Journal of Environmental Radioactivity*. **2016**, 164, 174-181; DOI 10.1016/j.jenvrad.2016.07.022.
- (11) Chang, C. C.; Burr, G. S.; Jull, A. J. T.; Russell, J. L.; Biddulph, D.; White, L.; Prouty, N. G.; Chen, Y.; Shen, C.; Zhou, W.; Lam, D. D. Reconstructing surface ocean circulation with ^{129}I time

- series records from corals. *Journal of Environmental Radioactivity*. **2016**, 165, 144-150; DOI 10.1016/j.jenvrad.2016.09.016.
- (12) Prouty, N. G.; Roark, E. B.; Mohon, L. M.; Chang, C. C. Uptake and distribution of organo-iodine in deep-sea corals. *Journal of Environmental Radioactivity*. **2018**, 187, 122-132; DOI 10.1016/j.jenvrad.2018.01.003.
- (13) Trevor, J. P.; Michael, F. P.; Steven, V. K.; Thomas, W. D. Climatic signals in $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ of tree-rings from White Spruce in the Mackenzie Delta Region, Northern Canada. *Arctic, Antarctic, and Alpine Research*. **2009**, 41(4), 479-505; DOI 10.1657/1938-4246-41.4.497.
- (14) Watmough, S. A. Monitoring historical changes in soil and atmospheric trace metal levels by dendrochemical analysis. *Environmental Pollution*. **1999**, 106(3), 391-403; DOI 10.1016/S0269-7491(99)00102-5.
- (15) Haas, G.; Schupfner, R.; Müller, A. Radionuclide uptake and long term behavior of ^{137}Cs , ^{134}Cs and ^{40}K in tree rings of spruce. *Journal of Radioanalytical & Nuclear Chemistry*. **1995**, 194(2), 277-282; DOI 10.1007/BF02038424.
- (16) Rao, U.; Fehn, U.; Muramatsu, Y.; McNeil, H.; Sharma, P.; Elmore, D. Tracing the history of nuclear releases: determination of ^{129}I in tree rings. *Environmental Science & Technology*. **2002**, 36(6), 1271-1275; DOI 10.1021/es011045i.
- (17) Hauschild, J. and Aumann, D. C. Iodine-129 and natural iodine in tree rings in the vicinity of a small nuclear fuels reprocessing plant. *Die Naturwissenschaften*. **1995**, 72(5), 270-272; DOI 10.1007/BF00448689.
- (18) Sebastian, P.; Justine, R.; Tjoelker, M.; Salih, A. Phloem as capacitor: Radial transfer of water into xylem of tree stems occurs via symplastic transport in ray parenchyma. *Plant Physiology*. **2015**, 167(3), 963-971; DOI 10.1104/pp.114.254581.
- (19) Wang, G.; Li, Y.; Wu, Q.; Wang, Y. Impacts of permafrost changes on alpine ecosystem in Qinghai-Tibet Plateau. *SCIENCE IN CHINA SERIES D-EARTH SCIENCES*. **2006**, 49(11), 1156-1169; DOI 10.1007/s11430-006-1156-0.

- (20) UNSCEAR, 2000: *Sources, Effects and Risk of Ionizing Radiation. UNSCEAR 2000 Report to the General Assembly, Annex C: Exposures from Man-made Sources of Radiation*; United Nations Scientific Committee on the Effect of Atomic Radiation: New York, 2000; www.unscear.org/docs/publications/2000/UNSCEAR_2000_Annex-C-CORR.pdf.
- (21) Hou, X. and Wang, Y. Determination of ultra-low level ^{129}I in vegetation using pyrolysis for iodine separation and accelerator mass spectrometry measurement. *Journal of Analytical Atomic Spectrometry*. **2016**, 31(6), 1298-1310; DOI 10.1039/C6JA00029K.
- (22) Hou, X.; Zhou, W.; Chen, N.; Zhang, L.; Liu, Q.; Fan, Y.; Liang, W.; Fu, Y. Determination of ultralow level $^{129}\text{I}/^{127}\text{I}$ in natural samples by separation of microgram carrier free Iodine and Accelerator Mass Spectrometry detection. *Analytical Chemistry*. **2010**, 82, 7713-7721; DOI 10.1021/ac101558k.
- (23) Wang, Y. Investigation of ^{129}I in vegetation samples from Northwest of China. M.S. Dissertation, Chinese Academy of Sciences (Institute of Earth Environment Science), Beijing, 2016.
- (24) Zhang, D. Iodine and Plutonium isotope distribution and tracer study in Northern China. Ph.D. Dissertation, Chinese Academy of Sciences (Institute of Earth Environment Science), Beijing, 2018.
- (25) Fan, Y.; Zhou, W.; Hou, X. Pre-nuclear level of ^{129}I in Chinese Loess-Paleosol sections: a search for the natural ^{129}I level for dating in terrestrial environments. *Geochimica Et Cosmochimica Acta*. **2018**, 231, 64-72; DOI 10.1016/j.gca.2018.04.014.
- (26) Chen, N. Analysis method of iodine isotope in water samples and its application in environmental tracing. Ph.D. Dissertation, Chinese Academy of Sciences (Institute of Earth Environment Science), Beijing, 2018.
- (27) Toyama, C.; Muramatsu, Y.; Igarashi, Y.; Aoyama, M.; Matsuzaki, H. Atmospheric fallout of ^{129}I in Japan before the Fukushima accident: regional and global contributions (1963-2005). *Environmental Science & Technology*. **2013**, 47, 8383-8390; DOI 10.1021/es401596z.

- 522 (28) Roulier, M.; Bueno, M.; Thiry, Y.; Coppin, F.; Redon, P.; Hecho, I. L.; Pannier, F. Iodine
523 distribution and cycling in a beech (*Fagus sylvatica*) temperate forest. *Science of the Total*
524 *Environment*. **2018**, 645, 431-440; DOI 10.1016/j.scitotenv.201807.039.
- 525 (29) Momoshima, N. and Bondietti, E. A. Cation binding in wood: applications to understanding
526 historical changes in divalent cation availability to red spruce. *Canadian Journal of Forest Research*.
527 **1990**, 20(12), 1840-1849; DOI 10.1139/x90-247.
- 528 (30) Luo, M.; Hou, X.; Zhou, W.; He, C.; Chen, N.; Liu, Q.; Zhang, L. Speciation and migration of
529 ^{129}I in soil profiles. *Journal of Environmental Radioactivity*. **2012**, 118, 30-39; DOI
530 10.1016/j.jenvrad.2012.11.011.
- 531 (31) Zhang, L.; Zhou, W.; Hou, X.; Chen, N.; Liu, Q.; He, C.; Fan, Y.; Luo, M.; Wang, Z.; Fu, Y.
532 Level and source of ^{129}I of environmental samples in Xi'an region, China. *Science of the Total*
533 *Environment*. **2011**, 409(19), 3780-3788; DOI 10.1016/j.scitotenv.2011.06.007.
- 534 (32) Hou, X.; Aldahan, A.; Nielsen, S. P.; Possnert, G. Time series of ^{129}I and ^{127}I speciation in
535 precipitation from Denmark. *Environmental Science & Technology*. **2009**, 43, 6522-6528; DOI
536 10.1021/es9012678.
- 537 (33) Hou, X.; Hansen, V.; Aldahan, A.; Possnert, G.; Lind, O. C.; Lujaniene, G. A review on
538 speciation of iodine-129 in the environmental and biological samples. *Analytica Chimica Acta*. **2009**,
539 632, 181-196; DOI 10.1016/j.aca.2008.11.013.
- 540 (34) Li, Y.; Li, X.; Xiao, Y.; Zhao, B.; Wang, L. Advances in study on mechanism of foliar nutrition
541 and development of foliar fertilizer application. *Scientia Agricultura Sinica*. **2009**, 42(1), 162-172;
542 DOI 10.3864/j.issn.0578-1752.2009.01.020.
- 543 (35) Xu, S.; Cook, G. T.; Cresswell, A. J.; Dunbar, E.; Freeman, S. P.; Hou, X.; Jacobsson, P.; Kinch,
544 H. R.; Naysmith, P.; Sanderson, D. C.; Tripney, B. G. Radiocarbon Releases from the 2011
545 Fukushima Nuclear Accident. *Sci.Rep*. **2016**, 6, 36947-369752; DOI 10.1038/srep36947.

- (36) Wang, X.; Wang, C.; Zhang, Q.; Li, S.; Li, G. Growth characteristics of heartwood and sapwood of the major tree species in Northeast China. *Science Silvae Sinicae*. **2008**, *44*(5), 102-108; DOI 10.11707/j.1001-7488.20080520.
- (37) Eder, M.; Jungnikl, K.; Burgert, I. A close-up view of wood structure and properties across a growth ring of Norway spruce (*Picea abies*[L] Karst.). *Trees*. **2009**, *23*(1), 79-84; DOI 10.1007/s00468-008-0256-1.
- (38) Molder, I. European beech grows better and is less drought sensitive in mixed than in pure stands: tree neighbourhood effects on radial increment. *Trees*. **2014**, *28*(3), 777-792; DOI 10.1007/s00468-014-0991-4.
- (39) *NOAA: State of the Climate Report, Climate Monitoring*; National Oceanic and Atmospheric Administration: Washington, DC; <https://www.ncdc.noaa.gov>.
- (40) Young, P.; Archibald, A.; Bowman, K. Pre-industrial to end ²¹th century projections of tropospheric ozone from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP). *Atmospheric Chemistry and Physics*. **2013**, *13*(4), 2063-2090; DOI 10.5194/acp-13-2063-2013.
- (41) Cooper, O.; Parrish, D.; Ziemke, J. Global distribution and trends of tropospheric ozone: An observation-based review. *Elem Sci Anth*. **2014**, *2*, 29-39; DOI 10.12952/journal.elementa.000029.
- (42) Esen, A. N.; et al.. An overview of energy technologies for a sustainable future. In *Energy Systems and Management*; Bilge A., Toy A., Gunay M., Eds.; Springer, Cham: Istanbul 2014; pp 13.
- (43) Wu, X.; Gao, Y.; Guo, Y.; Wu, J. Pollution characteristics of bromine and iodine as trace element in atmospheric particles in Shanghai Caohejing region. *Jornal of Shanghai Normal University (Nature Science)*. **2009**, *38*(2), 203-209; DOI 10.3969/j.issn.1000-5137.2009.02.016.
- (44) Liao, H.; Bu, W.; Zheng, J.; Wu, F.; Yamada, M. Vertical distributions of radionuclides (²³⁹⁺²⁴⁰Pu, ²⁴⁰Pu/²³⁹Pu, and ¹³⁷Cs) in sediment cores of Lake Bosten in Northwestern China. *Environmental Science & Technology*. **2014**, *48*(7), 3840-3846; DOI 10.1021/es405364m.

- (45) Stein, A. F.; Draxler, R. R.; Rolph, G. D.; Stunder, B. J. B.; Cohen, M. D.; Ngan, F. NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *Bull. Amer. Meteor. Soc.*, **2015**, 96, 2059-2077; DOI <http://dx.doi.org/10.1175/BAMS-D-14-00110.1>.
- (46) Xu, H.; Wei, J.; Yan, Z.; Zou, W. Influence evaluation of Chernobyl accident in Xinjiang by ^{131}I measured in milk, breast milk and pasture. *Chinese journal of public health*. **1989**, 5(7), 18-20.
- (47) Zhang, J.; Zou, W.; Xu, H.; Wei, J. The total β activity of atmospheric fallouts and aerosols in Xinjiang Province after Chernobyl accident. *Environmental Protection of Xinjiang*. **1987**, 2, 50-57.
- (48) Hou, X.; Povinec, P.; Zhang, L.; Shi, K.; Biddulph, D.; Chang, C.; Fan, Y.; Golser, R.; Hou, Y.; Jeskovsky, M.; Jull, A. J.; Liu, Q.; Luo, M.; Steier, P.; Zhou, W. Iodine-129 in seawater offshore Fukushima: distribution, inorganic speciation, sources, and budget. *Environmental Science & Technology*. **2013**, 47(7), 3091-3098; DOI 10.1021/es304460k.
- (49) Tang, M.; Tsuang, B.; Kuo, P. H. Dose estimation for nuclear power plant 4 accident in Taiwan at Fukushima nuclear meltdown emission level. *Journal of Environmental Radioactivity*. **2016**, 155-156, 71-83; DOI 10.1016/j.jenvrad.2016.01.022.
- (50) Wang, L.; Zheng, G.; Zhao, S.; Kuo, S.; Xu, M.; Wang, X. Impact of Fukushima Nuclear Accident to China's mainland environment. *Radiation Protection*. **2012**, 32(6), 325-335.
- (51) Zheng, J.; Togami, K.; Watanabe, Y.; Uchida, S.; Aono, T.; Nobuyoshi, I.; Yoshida, S.; Kubota, Y.; Fuma, S.; Ihara, S. Isotopic evidence of plutonium release into the environment from the Fukushima DNPP accident. *Scientific Reports*. **2012**, 2(3), 304-312; DOI 10.1038/srep00304.
- (52) Fan, Y. Spatial distribution of ^{129}I in Chinese surface soil and preliminary study on the ^{129}I chronology. Ph.D. Dissertation, Chinese Academy of Sciences (Institute of Earth Environment Science), Beijing, 2013.
- (53) Englund, E.; Aldahan, A.; Hou, X.; Possnert, G.; Soderstrom, C. Iodine (^{129}I and ^{127}I) in aerosols from northern Europe. *Nuclear Instruments & Methods in Physics Research*. **2010**, 268, 1139-1141; DOI 10.1016/j.nimb.2009.10.118.

(54) Jiang, X. A study on the level and variation of ^{129}I in precipitation in Xi'an region and its application as environmental tracer. M.S. Dissertation. Chinese Academy of Sciences (Institute of Earth Environment Science), Beijing, 2017.

Caption of the figures

Fig. 1 Wood disks of PC1 tree (up) and PC2 tree (bottom), showing the annual rings

Fig. 2 Comparison of measured ^{127}I concentrations (a) and $^{129}\text{I}/^{127}\text{I}$ ratios (b) in the tree rings in this work with the reported values in other locations^{16,17, 23, 24}

Fig. 3 Temporal variation of the measured $^{129}\text{I}/^{127}\text{I}$ ratios in the tree rings from two sites in the QTP (a), in comparison with the reported variation of $^{129}\text{I}/^{127}\text{I}$ in atmospheric fallout from Tokyo (b)²⁷, yield of fission NWTs (c), yield of thermonuclear weapons tests at Lop Nor (d)²⁰, air releases of ^{129}I from major NFRPs (e) and marine discharges of ^{129}I from NFRPs (f).³³

Fig. 4 Variation of iodine (^{127}I) concentration in PC1 and PC2 tree (a), in comparison with the temporal variation of iodine concentrations in ice core in Greenland³