



Investigating the use of peracetic acid to reduce saprolegniasis in Atlantic salmon RAS culture

Good, Christopher; Davidson, John; May, Travis; Crouse, Curtis; Lepine, Christine; Redman, Natalie; Murray, Megan; Summerfelt, Steven; Straus, David L.; Harper, Susan

Total number of authors:
15

Published in:
5th NordicRAS Workshop on Recirculating Aquaculture Systems Berlin, Germany, 7-8 October 2019: Book of Abstracts

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Good, C., Davidson, J., May, T., Crouse, C., Lepine, C., Redman, N., Murray, M., Summerfelt, S., Straus, D. L., Harper, S., Marancik, D., Welch, T. J., Peterson, B., Pedersen, L-F., & Phuntumart, V. (2019). Investigating the use of peracetic acid to reduce saprolegniasis in Atlantic salmon RAS culture. In J. Dalsgaard (Ed.), *5th NordicRAS Workshop on Recirculating Aquaculture Systems Berlin, Germany, 7-8 October 2019: Book of Abstracts* (pp. 41-41). DTU Aqua.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Investigating the use of peracetic acid to reduce saprolegniasis in Atlantic salmon RAS culture

Christopher Good^{1*}, John Davidson¹, Travis May¹, Curtis Crouse¹, Christine Lepine¹, Natalie Redman¹, Megan Murray¹, Steven Summerfelt², David Straus³, Susan Harper⁴, David Marancik⁵, Timothy Welch⁶, Brian Peterson⁷, Lars-Flemming Pedersen⁸, Vipaporn Phuntumart⁹

¹The Conservation Fund's Freshwater Institute, Shepherdstown, WV, USA; ² Superior Fresh, Hixton, WI, USA; ³USDA-ARS, Stuttgart, AR, USA; ⁴USDA-ARS, Beltsville, MD, USA; ⁵St. George's University School of Veterinary Medicine, Grenada, West Indies; ⁶USDA-ARS, Leetown, WV, USA; ⁷USDA-ARS, Franklin, ME, USA; ⁸Danish Technical University, Hirtshals, Denmark; ⁹Bowling Green State University, Bowling Green, OH, USA

Abstract

Despite numerous biosecurity benefits provided by land-based, closed-containment water recirculation aquaculture system (RAS) operations, opportunistic pathogens, such as *Saprolegnia* spp. oomycetes (causative agents of saprolegniasis) can still represent disease threats if environmental conditions are unfavorable, during vulnerable fish life-stages, or following stressful procedures. Saprolegniasis in general is associated with enormous losses in Atlantic salmon *Salmo salar* aquaculture, especially during the early rearing fry stage, smoltification, and following vaccination. We have investigated strategies to reduce saprolegniasis during each of these high-risk production cycle stages; however, at the time of abstract submission, only one study has been completed, while the remaining two are ongoing and will be presented at the workshop. To assess post-vaccination saprolegniasis, we applied daily peracetic acid (PAA) bath treatments at three dosages while evaluating the effects of these treatments on biofilter performance in replicated (12) experimental-scale RAS stocked with Atlantic salmon parr (200 fish per RAS, 94g mean weight) following intracoelomic injection vaccination. The PAA doses assessed were 0.2, 0.5, and 1.0 mg/L, and these were applied to each culture tank for a period of six weeks post-vaccination. Survival, waterborne *Saprolegnia* spp. colony counts, gill, spleen, and kidney histopathology, and biofiltration function (measured by total ammonia nitrogen (TAN) removal efficiency), were assessed. Visible skin lesions and fin erosion, hemorrhage, and observable *Saprolegnia* spp. infection were also evaluated. While no major post-vaccination saprolegniasis occurred in this study, survival was statistically ($p < 0.05$) lower in control salmon, and visible saprolegniasis was significantly more prevalent in the control group. Despite less saprolegniasis being associated with PAA treatment, all treatment groups were associated with significantly lower fish weight by study's end. Biofilter TAN removal efficiency was not impacted by PAA administration at all dosages. Results suggest that low-dose PAA might be effective in reducing post-vaccination saprolegniasis while not significantly impacting RAS biofiltration. Further research, however, is necessary to confirm these findings, especially in commercial settings experiencing significant issues with saprolegniasis.

Acknowledgements: The post-vaccination research was funded through the US Department of Agriculture, Agriculture Research Service (USDA-ARS); the fry and smolt studies were funded through University of Wisconsin Sea Grant, part of the National Sea Grant College Program overseen by the National Oceanographic and Atmospheric Administration (NOAA).

*cgood@conservationfund.org