



Piscine orthoreovirus-3 (PRV-3), a new pathogen for farmed rainbow trout

Vendramin, Niccolò; Sørensen, Juliane; Cuenca, Argelia; Dahle, Maria; Olsen, Anne Berit; Iburg, Tine; Rimstad, Espen; Olesen, Niels Jørgen

Published in:

5th NordicRAS Workshop on Recirculating Aquaculture Systems Berlin, Germany, 7-8 October 2019: Book of Abstracts

Publication date:

2019

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Vendramin, N., Sørensen, J., Cuenca, A., Dahle, M., Olsen, A. B., Iburg, T., Rimstad, E., & Olesen, N. J. (2019). Piscine orthoreovirus-3 (PRV-3), a new pathogen for farmed rainbow trout. In J. Dalsgaard (Ed.), *5th NordicRAS Workshop on Recirculating Aquaculture Systems Berlin, Germany, 7-8 October 2019: Book of Abstracts* (pp. 39-39). DTU Aqua.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Piscine orthoreovirus-3 (PRV-3), a new pathogen for farmed rainbow trout

Niccolò Vendramin^{1*}, Juliane Sørensen², Argelia Cuenca¹, Maria Dahle³, Anne Berit Olsen⁴, Tine Iburg¹, Espen Rimstad², Niels Jørgen Olesen¹

¹DTU AQUA, Technical University of Denmark, Kgs. Lyngby, Copenhagen; ²Department of Food Safety and Infection Biology, Norwegian University of Life Sciences, Oslo, Norway; ³Norwegian Veterinary Institute, Oslo, Norway; ⁴Norwegian Veterinary Institute, Bergen, Norway

Abstract

Piscine orthoreovirus – PRV have emerged as relevant pathogens for salmonid aquaculture worldwide. Three different subtypes of this viral species have so far been described:

- PRV-1 is the causative agent of heart and skeletal muscle inflammation (HSMI) in Atlantic salmon and is associated with jaundice syndrome in farmed Chinook salmon
- PRV-2 causes erythrocytic inclusion body syndrome (EIBS) in Coho salmon
- PRV-3 causes heart pathology resembling HSMI in rainbow trout

PRV-3 was firstly discovered in 2013 in Norway during disease outbreaks affecting farmed rainbow trout. The Norwegian PRV-3 isolate has been characterized by performing full genome sequencing and demonstrating causative relationship between the infection and the development of heart pathology in Rainbow trout.

An experimental infection study with purified virus demonstrated that PRV-3 infects rainbow trout and induces pathological heart lesions similar to Heart and Skeletal Muscle Inflammation (HSMI)

During 2017 the presence of PRV-3 was also reported in different countries in Europe. Interestingly, these viral isolates appear to be genetically distinct from the Norwegian isolate leading to proposition of two separate clades within PRV-3 viral type (PRV-3a and PRV-3b).

In Denmark the virus has been associated with severe disease outbreaks in recirculating aquaculture systems. Clinical signs are represented by reduced appetite followed by uncoordinated swimming behavior and increased mortality; necropsy findings include severe anemia and ascites. Such outbreaks are complex disease cases where different bacterial (including *Flavobacterium psychrophilum* and *Renibacterium salmoninarum*) and viral pathogens (IPNV) are present at the farms. Notably PRV-3 load increases in the target organs (heart, spleen) before the clinical disease appear, whereas the other pathogens are not detected in a systematic pattern.

In 2018 in cooperation with the Danish aquaculture industry, the presence of the virus has been mapped in the country, and a comparative study including the 2 PRV-3 subtypes conducted.

The preliminary results show suggests that PRV-3b is capable of faster replication than PRV-3a inducing higher innate inflammatory response and more severe heart pathology.

Results will be presented and discussed.

Acknowledgements: This research was funded by Henriksens Fond.

*niven@aqua.dtu.dk